


This article was downloaded by: [Helmholtz-Zentrum Muenchen]

On: 13 April 2015, At: 22:33

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Molecular & Cellular Oncology

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/kmco20>

Glutathione peroxidase 4 (Gpx4) and ferroptosis: what's so special about it?

Marcus Conrad^a & José Pedro Friedmann Angeli^a

^a Institute of Developmental Genetics; Helmholtz Zentrum München; München, Germany

Accepted author version posted online: 30 Jan 2015.



[Click for updates](#)

To cite this article: Marcus Conrad & José Pedro Friedmann Angeli (2015) Glutathione peroxidase 4 (Gpx4) and ferroptosis: what's so special about it?, *Molecular & Cellular Oncology*, 2:3, e995047, DOI: [10.4161/23723556.2014.995047](https://doi.org/10.4161/23723556.2014.995047)

To link to this article: <http://dx.doi.org/10.4161/23723556.2014.995047>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Versions of published Taylor & Francis and Routledge Open articles and Taylor & Francis and Routledge Open Select articles posted to institutional or subject repositories or any other third-party website are without warranty from Taylor & Francis of any kind, either expressed or implied, including, but not limited to, warranties of merchantability, fitness for a particular purpose, or non-infringement. Any opinions and views expressed in this article are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor & Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

It is essential that you check the license status of any given Open and Open Select article to confirm conditions of access and use.

Glutathione peroxidase 4 (Gpx4) and ferroptosis: what's so special about it?

Marcus Conrad* and José Pedro Friedmann Angeli

Institute of Developmental Genetics; Helmholtz Zentrum München; München, Germany

Keywords: cell death, lipid peroxidation, non-apoptotic cell death, regulated necrosis

The system X_C^- /glutathione/glutathione peroxidase 4 (Gpx4) axis pivotally controls ferroptosis, a recently described form of regulated non-apoptotic cell death. Compelling evidence has established that this route of cell death is not only of high relevance for triggering cancer cell death, but also proves to be amenable for therapeutic intervention to halt ischemia/reperfusion-related diseases.

Glutathione peroxidase 4 (Gpx4) is 1 of 8 known glutathione peroxidases in mammals. With the exception of Gpx5, Gpx7, and Gpx8 (and Gpx6 in rodents), glutathione peroxidases (Gpx) are selenoproteins.¹ Selenoproteins contain the 21st amino acid selenocysteine, which differs from cysteine by a single atom, selenium replacing for sulfur. Although the advantage of using selenocysteine instead of cysteine is not really understood, it is generally believed that selenocysteine provides higher reactivity and superior efficiency in redox reactions. All members of the Gpx family share the same basic function of the reduction of peroxides at the expense of glutathione (GSH), or other thiol containing compounds in the case of the monomeric Gpx4, Gpx7, and Gpx8. Despite structural and catalytic similarities, Gpx4 is unique among Gpx isoforms as it is the only enzyme capable of reducing esterified oxidized fatty acids and cholesterol hydroperoxides. The relevance of this unique function is highlighted by *in vivo* studies using knockout (KO) models for this enzyme. The first KO of *Gpx4* was reported in 2003² and found to cause early embryonic lethality. Therefore, investigations into Gpx4 functions in adult animals are only possible using conditional KO systems.

In 2008, we generated the first animal carrying a conditional *Gpx4* allele.³ Using mouse embryonic fibroblasts with an inducible *Gpx4* disruption, we were able to demonstrate that a non-apoptotic form of cell death is elicited upon *Gpx4* deletion and that this cell death is preceded by lipid oxidation.³ Subsequent studies with these animals revealed that Gpx4 is essential for maintaining tissue homeostasis by preventing cell demise and tissue damage in several organs including brain, skin, and endothelium.^{3,4} Nevertheless, some tissues did not appear to be directly affected by loss of Gpx4, such as the heart (unpublished observation), indicating that specific conditions might be required to trigger this type of cell death.

In 2012, Stockwell and colleagues described a novel form of cell death that was elicited by inhibiting the cystine/glutamate antiporter, system X_C^- , a transporter that feeds cells with cysteine to be used for protein and, in particular, GSH biosynthesis⁵ (Fig. 1). Following up on this, the same group identified the most important GSH-dependent enzyme in this pathway as GPX4.⁶ Yet these studies have largely focused on the therapeutic possibility of triggering this type of cell death in cancer cells, leaving the option of pharmacological inhibition of this

pathway in a relevant pathological situation unaddressed. With this in mind, we have generated animals with inducible deletion of *Gpx4* as an ideal tool to reveal the adult tissues most sensitive to ferroptosis-relevant cell death. We found that the animals died within 2 weeks of Gpx4 loss as a result of massive renal tubule cell death and acute kidney failure.⁷ Pharmacological targeting of this pathway *in vivo* was further shown to be possible by the development of a novel class of drug-like small molecules named liproxstatins that are able to extend the survival of *Gpx4* null mice by approximately 35%. Further evidence that ferroptosis could indeed be targeted *in vivo* under pathologically relevant conditions was independently provided by 2 groups, who showed that liproxstatin and a second generation of ferrostatins mitigated tissue damage in preclinical models of liver and kidney ischemia/reperfusion damage, respectively.^{7,8}

Despite these recent advances in this intriguing field, questions concerning the downstream mechanisms in the response to *Gpx4* deletion have remained largely unanswered. Our recent work now shows that one of the earliest events upon loss of Gpx4 *in vivo* is cardiolipin oxidation, spreading from there to other classes of

© Marcus Conrad and José Pedro Friedmann Angeli

*Correspondence to: Marcus Conrad; Email: marcus.conrad@helmholtz-muenchen.de

Submitted: 11/28/2014; Revised: 12/01/2014; Accepted: 12/02/2014

<http://dx.doi.org/10.4161/23723556.2014.995047>

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The moral rights of the named author(s) have been asserted.




Figure 1. Biochemical events in the upstream control of ferroptosis. The upstream events controlling ferroptotic signaling are depicted. Cystine, the oxidized form of cysteine (Cys-S-S-Cys), is taken up by the cystine/glutamate antiporter that consists of the 4F2 (Slc3a2) heavy chain and xCT (Slc7a11) light chain. Upon intracellular reduction, cysteine, along with glutamate and glycine, are used for protein and glutathione (GSH) biosynthesis. GSH is synthesized in 2 consecutive steps by the heterodimer γ -glutamylcystine synthetase (γ GCS consists of Gclc and Gclm) and glutathione synthetase (Gss, GS), each step requiring one molecule of ATP. Glutathione peroxidase 4 (Gpx4) is a central GSH-utilizing enzyme and counteracts lipoxygenase (Alox) activities and phospholipid/cardioplin oxidation events. In the absence of Gpx4, cardioplin oxidation followed by phospholipid peroxidation causes ferroptosis through still unknown mechanisms. Small-molecule inhibitors of the different players are shown on the right. Abbreviations: BSO, L-buthionine sulfoximine; GSSG, oxidized glutathione; γ Glu-Cys, γ -glutamylcystine.

phospholipids such as phosphatidylethanolamine and phosphatidylcholine.⁷ This feature seems to be clearly distinct from apoptosis, in which only cardioplin oxidation via the peroxidase function of cytochrome c (cytochrome c, somatic; Cysc) has been demonstrated.⁹ Although the reasons for this are still unclear, one may speculate that Gpx4 is required to allow a “silent” form of cell death that could

tentatively switch upon Gpx4 loss/malfunction to a highly inflammatory cell death modality culminating in the release of several proinflammatory lipid mediators.⁷ What happens to cells undergoing apoptosis under Gpx4-compromised conditions is still not known and warrants further investigation.

Another point that deserves attention is how a ferroptotic-like cell death can

be induced in an ischemia/reperfusion scenario. It has been reported that proper Gpx4 function is required to inhibit cell death in kidney tubule cells yet this phenotype is observed only under complete Gpx4 loss, which is most likely not the case under a pathologically relevant setting. Notwithstanding, ferroptosis inhibitors present a beneficial effect during ischemia/reperfusion, indicating the engagement of this pathway under these conditions.^{7,8} Therefore, it seems plausible that Gpx4 function and/or expression might be modulated under such conditions, for example by post-translational modifications that would directly influence its catalytic activity such as oxidative degradation by oxygen radicals or proteasome-mediated degradation. Although still speculative, these hypotheses are not without foundation as similar events have already been reported for some other redox-related enzymes. A second consideration is that this modification would most likely occur at a specific site; we tentatively speculate that such events would likely take place at the inner membrane space of mitochondria, as we have shown that eliciting ferroptosis disrupts the outer mitochondrial membrane.⁷ Moreover, this was also identified as the major site of oxygen radical production under ischemic conditions through re-oxidation of succinate via the succinate dehydrogenase complex during reverse electron transport at mitochondrial complex I.¹⁰

We believe that a better understanding of how Gpx4 function is modulated under particular settings such as ischemia/reperfusion and tumorigenesis, and how downstream events orchestrate cellular demise under ferroptotic conditions, will allow us to efficiently prevent and/or trigger this form of cell death in therapeutically responsive scenarios.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

1 Brigelius-Flohe R, Maiorino M. Glutathione peroxidases. *Biochim Biophys Acta* 2013; 1830:3289–303;

PMID:23201771; <http://dx.doi.org/10.1016/j.bbagen.2012.11.020>

2 Yant LJ, Ran Q, Rao L, Van Remmen H, Shibata T, Belter JG, Motta L, Richardson A, Prolla TA. The

selenoprotein GPX4 is essential for mouse development and protects from radiation and oxidative damage insults. *Free Radic Biol Med* 2003; 34:496–502;

- PMID:12566075; [http://dx.doi.org/10.1016/S0891-5849\(02\)01360-6](http://dx.doi.org/10.1016/S0891-5849(02)01360-6)
- 3 Seiler A, Schneider M, Förster H, Roth S, Wirth EK, Culmsee C, Plesnila N, Kremmer E, Rådmark O, Wurst W, et al. Glutathione Peroxidase 4 Senses and Translates Oxidative Stress into 12/15-Lipoxygenase Dependent- and AIF-Mediated Cell Death. *Cell Metab* 2008; 8:237–48; PMID:18762024; <http://dx.doi.org/10.1016/j.cmet.2008.07.005>
 - 4 Wortmann M, Schneider M, Pircher J, Hellfritsch J, Aichler M, Vegi N, Kölle P, Kuhlencordt P, Walch A, Pohl U, et al. Combined deficiency in glutathione peroxidase 4 and vitamin e causes multiorgan thrombus formation and early death in mice. *Circ Res* 2013; 113:408–17; PMID:23770613; <http://dx.doi.org/10.1161/CIRCRESAHA.113.279984>
 - 5 Dixon SJ, Lemberg KM, Lamprecht MR, Skouta R, Zaitsev EM, Gleason CE, Patel DN, Bauer AJ, Cantley AM, Yang WS, et al. Ferroptosis: an iron-dependent form of nonapoptotic cell death. *Cell* 2012; 149:1060–72; PMID:22632970; <http://dx.doi.org/10.1016/j.cell.2012.03.042>
 - 6 Yang WS, SriRamaratnam R, Welsch ME, Shimada K, Skouta R, Viswanathan VS, Cheah JH, Clemons PA, Shamji AF, Clish CB, et al. Regulation of Ferroptotic Cancer Cell Death by GPX4. *Cell* 2014; 156:317–31; PMID:24439385; <http://dx.doi.org/10.1016/j.cell.2013.12.010>
 - 7 Friedmann Angeli JP, Schneider M, Proneth B, Tyurina YY, Tyurin VA, Hammond VJ, Herbach N, Aichler M, Walch A, Eggenhofer E, et al. Inactivation of the ferroptosis regulator Gpx4 triggers acute renal failure in mice. *Nat Cell Biol* 2014 16:1180–91; PMID:25402683; <http://dx.doi.org/10.1038/ncb3064> advance online publication
 - 8 Linkermann A, Skouta R, Himmerkus N, Mulay SR, Dewitz C, De Zen F, Prokai A, Zuchtriegel G, Krombach F, Welz PS, et al. Synchronized renal tubular cell death involves ferroptosis. *Proc Natl Acad Sci U S A* 2014; 111:16836–41; PMID:25385600; <http://dx.doi.org/10.1073/pnas.1415518111>
 - 9 Kagan VE, Tyurin VA, Jiang J, Tyurina YY, Ritov VB, Amoscato AA, Osipov AN, Belikova NA, Kapralov AA, Kini V, et al. Cytochrome c acts as a cardiolipin oxygenase required for release of proapoptotic factors. *Nat Chem Biol* 2005; 1:223–32; PMID:16408039; <http://dx.doi.org/10.1038/nchembio727>
 - 10 Chouchani ET, Pell VR, Gaude E, Aksentijević D, Sundier SY, Robb EL, Logan A, Nadtochiy SM, Ord EN, Smith AC, et al. Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS. *Nature* 2014; 515:431–5; PMID:25383517; <http://dx.doi.org/10.1038/nature13909>