

Testing the Role of Predicted Gene Knockouts in Human Anthropometric Trait Variation

Samuel Lessard^{1,2}, Alisa K. Manning³, Cécile Low-Kam^{1,2}, Paul L. Auer⁴, Ayush Giri⁵, Mariaelisa Graff⁶, Claudia Schurmann⁷, Hanieh Yaghootkar⁸, Jian'an Luan⁹, Tonu Esko^{3,10,11}, Tugce Karaderi¹², NHLBI GO Exome Sequence Project¹³, GOT2D¹³, T2D-GENES¹³, GIANT Consortium¹³, Erwin P. Bottinger⁷, Yingchang Lu⁷, Chris Carlson¹⁴, Mark Caulfield^{15,16}, Marie-Pierre Dubé^{1,2}, Rebecca D. Jackson¹⁷, Charles Kooperberg¹⁴, Barbara McKnight¹⁸, Ian Mongrain¹, Ulrike Peters¹⁴, Alex P. Reiner¹⁴, David Rhoads¹, Nona Sotoodehnia¹⁹, Joel N. Hirschhorn^{3,11,20}, Robert Scott⁹, Patricia B. Munroe^{15,16}, Timothy M. Frayling⁸, Ruth J.F. Loos⁷, Kari E. North⁶, Todd L. Edwards⁵, Jean-Claude Tardif^{1,2}, Cecilia M. Lindgren^{3,12,21}, Guillaume Lettre^{1,2,*}

¹Montreal Heart Institute, Montréal, Québec H1T 1C8, Canada

²Faculté de Médecine, Université de Montréal, Montréal, Québec H3T 1J4, Canada

³Broad Institute of the Massachusetts Institute of Technology and Harvard University, Cambridge, MA 02142, USA

⁴School of Public Health, University of Wisconsin-Milwaukee, Milwaukee, WI 53201-0413, USA

⁵Center for Human Genetics Research, Vanderbilt Epidemiology Center, Department of Medicine, Vanderbilt University, Nashville, TN 37212, USA

⁶University of North Carolina Gillings School of Global Public Health, Chapel Hill, NC 27599, USA

⁷The Charles Bronfman Institute of Personalized Medicine, Icahn School of Medicine at Mount Sinai Hospital, New York, NY 10029, USA

⁸Genetics of Complex Traits, University of Exeter Medical School, University of Exeter, Exeter, EX2 5DW, UK

⁹Medical Research Council Epidemiology Unit, University of Cambridge, Cambridge, CB2 0QQ, UK

¹⁰Estonian Genome Center, University of Tartu, Tartu, Estonia

¹¹Division of Endocrinology, Genetics and Basic and Translational Obesity Research, Children's Hospital Boston, Boston, MA 02115, USA

¹²Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford OX3 7BN, UK

¹³A full list of members and affiliations appears in the Supplemental Text.

¹⁴Fred Hutchinson Cancer Research Center, Division of Public Health Sciences, Seattle, WA 98109-1024, USA

¹⁵Clinical Pharmacology, William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, EC1M 6BQ, UK

¹⁶NIHR Barts Cardiovascular Biomedical Research Unit, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, EC1M 6BQ, UK

¹⁷Division of Endocrinology, Diabetes and Metabolism, Ohio State University, Columbus, Ohio 43210, USA

¹⁸Department of Biostatistics, University of Washington, Seattle, WA 98195, USA

¹⁹Division of Cardiology, Cardiovascular Health Research Unit, University of Washington, Seattle, WA 98195-6422, USA

²⁰Department of Genetics, Harvard Medical School, Boston, MA 02115, USA

²¹The Big Data Institute, University of Oxford, Oxford, UK

*Corresponding author: Guillaume Lettre, Montreal Heart Institute, 5000 Bélanger Street, Montréal, Québec, Canada, H1T 1C8, Phone : 514-376-3330 x2657, Fax : 514-593-2539, Email : guillaume.lettre@umontreal.ca

ABSTRACT

Although the role of complete gene inactivation by two loss-of-function mutations inherited in *trans* is well-established in recessive Mendelian diseases, we have not yet explored how such gene knockouts (KOs) could influence complex human phenotypes. Here, we developed a statistical framework to test the association between gene KOs and quantitative human traits. Our method is flexible, publicly available, and compatible with common genotype format files (*e.g.* PLINK and vcf). We characterized gene KOs in 4,498 participants from the NHLBI Exome Sequence Project (ESP) sequenced at high coverage (>100X), 1,976 French Canadians from the Montreal Heart Institute Biobank sequenced at low coverage (5.7X), and >100,000 participants from the GIANT Consortium genotyped on an exome array. We tested associations between gene KOs and three anthropometric traits: body mass index (BMI), height and BMI-adjusted waist-to-hip ratio (WHR). Despite our large sample size and multiple datasets available, we could not detect robust associations between specific gene KOs and quantitative anthropometric traits. Our results highlight several limitations and challenges for future gene KO studies in humans, in particular when there is no prior knowledge on the phenotypes that might be affected by the tested gene KOs. They also suggest that gene KOs identified with current DNA sequencing methodologies probably do not strongly influence normal variation in BMI, height, and WHR in the general human population.

INTRODUCTION

The identification of complete loss-of-function (LoF) alleles (*i.e.* genetic null or amorphic alleles) is a powerful strategy to characterize gene functions through random (*e.g.* chemical mutagenesis) or targeted (*e.g.* knockout (KO) methodology in the mouse, RNAi) genetic experiments. In contrast to model organisms, humans are not amenable to such genetic manipulations. Yet, there is tremendous biomedical interest in understanding how the complete disruption of both copies of a gene may impact human biology (1). Our complex physiology, interactions with our environment, and gene redundancy within our genome are only few of the reasons highlighting the importance of describing the phenotypic consequences of gene inactivation in humans. From a drug development perspective, the identification of humans with gene KOs also offers naturally occurring genetic experiments to assess the potential pleiotropic effects of candidate target genes (2).

Mendelian diseases, such as sickle cell anemia [MIM 603903] and cystic fibrosis [MIM 219700], offer an entry point into the study of gene functions in humans. Indeed, the study of these conditions continues to yield important insights into human biology in health and disease (3). But only a limited number of genes have been implicated in Mendelian diseases: as of October 13 2015, there were 4,651 genes in the Online Mendelian Inheritance in Man (OMIM) database with phenotype-causing mutations. Furthermore, these mutations are often rare such that it is difficult to assemble sufficiently large cohorts of patients to study their pleiotropic effects. Gene KOs can have strong phenotypic effects on anthropometric traits in the context of Mendelian disorders or syndromes, as evident by mutations causing early-onset morbid obesity (*PCSK1*, *LEPR*) or dwarfism (*GHI* *GHR*, *ATR*) (4-6). These mutations are rare (often private) and unlikely to be involved in anthropometric trait variation in the general population. However,

the possibility that gene KOs of more subtle effect might influence normal variation in anthropometric traits remains to be investigated.

Large-scale whole-exome and -genome sequencing projects are beginning to systematically catalogue coding genetic variation in the human genome, including predicted LoF variants (7-11). On average, there are ~100-200 LoF variants per individual, resulting in ~20 genes that are inactivated through homozygosity or compound heterozygosity (12). These numbers include mostly common variants, which are more likely to be phenotypically neutral given the effect of purifying selection (13). Limiting to variants with a minor allele frequency (MAF) <0.5%, the 1000 Genomes Project estimated that there are 10-20 LoF variants per individual (8). LoF variants are usually defined as variants that truncate protein sequences (nonsense and frameshift insertion-deletion (indel)) or that abrogate splice sites or stop codons (stop-loss) (12). Using this definition of LoF variant, and limiting their analyses to variants with a MAF <2%, Sulem *et al.* found that ~8% of 104,220 Icelanders carry at least one complete gene KO, and that most gene KOs are seen in <5 individuals (14).

Recently, several studies have explored the link between gene KOs and human complex phenotypes, such as chronic diseases (12, 15-17) and autism (18). As mentioned above, it is well-established that rare gene inactivation can cause extreme anthropometric phenotypes in several human recessive disorders. The goal of our study is to extend this observation and determine whether gene KOs of modest phenotypic effect also contribute to anthropometric trait variation in the general human population. We developed a statistical method to test for association between predicted gene KOs and quantitative human phenotypes and characterized the distribution of predicted gene KOs in 2,772 European Americans and 1,726 African Americans from the

National Heart, Lung, and Blood Institute (NHLBI) Exome Sequence Project (ESP). We then applied our method to detect associations between gene KOs and three quantitative anthropometric traits (body mass index (BMI), adult height, and BMI-adjusted waist-to-hip ratio (WHR)) using high coverage whole-exome sequence (WES) data from 4,498 ESP participants, low coverage whole-genome sequence (WGS) data from 1,969 French Canadians, and >100,000 participants from the GIANT Consortium genotyped on an exome array.

RESULTS

Number and distribution of predicted gene KOs in ESP

We identified 18,137 and 21,935 LoF variants in 1,726 African Americans and 2,772 European Americans from ESP, respectively (**Table 1, Table S1**). These LoF variants included protein truncating (nonsense, frameshift indel), stop-loss and splice site variants. On average, we found 65 and 39 rare or low-frequency LoF variants (MAF <5%) per African-American and European-American ESP participant, respectively (**Table 1**). These numbers are higher than some of the previous estimates (12, 16, 18), mostly because we included frameshift indels in our analyses. When excluding frameshift indels, we found on average 26 and 16 LoF variants with MAF <5% per ESP African American and European American, respectively. Descriptive statistics on the number of LoF variants in ESP after excluding frameshift indels are available in **Table S2**. We screened the ESP dataset for individuals who are homozygous or compound heterozygous for LoF variants, and are therefore predicted KOs for a given gene. To detect compound heterozygosity, we used phased genotype information generated with the software Beagle to distinguish between LoF variants inherited in *cis* or *trans* (**Table 1**) (19). The identification of LoF variants depends on the gene annotation used. To address this concern, we re-analyzed the ESP WES data using the GENCODE basic transcripts annotation instead of

RefSeq, and only considered variants that fell within all transcripts for a given gene. We obtained very similar association results between the two annotation software (**Figure S1**). We present below results generated with the RefSeq annotation.

Common LoF variants are responsible for most predicted gene KOs (**Figure 1**, and **Figure S2** for distributions without frameshift indels). For instance, in ESP African Americans, we found on average 25.9 and 2.5 predicted gene KOs per individual when analyzing all or rare/low-frequency LoF variants, respectively (**Table 1**). The corresponding numbers in European Americans are 23.2 and 1.1 for all and rare/low-frequency LoF variants (**Table 1**). While this article was under review, the Exome Aggregation Consortium (ExAC) reported an average of 35 homozygous protein-truncating variants per individual. This number is higher than the average number of homozygous LoF variants that we found in ESP (~21-23/participant, **Table 1**) (20). This difference might simply reflect increased power in ExAC to discover rare mutations owing to its larger sample size (N=60,706 vs N=4,498 for ESP). Because common LoF are more likely to be phenotypically neutral (13), we focused all subsequent analyses on LoF with MAF <5% within ethnic group or sub-study. In the ESP dataset, we found 2,071 and 1,433 genes with both alleles inactivated by such LoF variants in at least one African American or one European American, respectively (**Table 1**). The higher number of predicted gene KOs in African Americans has been previously observed and is consistent with increased genetic diversity in African-ancestry populations (12). Overall, very few individuals shared the same gene KOs, most of them being found in only one individual (**Figure 1**). Homozygosity of LoF variants is responsible for the majority of these KO events as we only found (after taking phase information into account) compound heterozygous individuals for ~8% of the genes with at least

one gene KO (**Table 1**). Stop-loss variants might not be as detrimental as other categories of LoF variants, but they are implicated in less than 0.9% of all gene KOs identified in ESP.

Predicted gene KO associated with anthropometric traits in ESP

We tested our newly developed method (**Figure 2**) on three anthropometric traits (BMI, height, and WHR) that are available in a large number of ESP participants. We stratified our analyses by ethnicity and meta-analyzed association results (**Figure 3**). Assuming that most genes are independent and given the number of genes for which we could find at least one predicted knocked out individual, we used the following Bonferroni-corrected significance threshold to declare significance: $\alpha=2 \times 10^{-5}$. No single genes reached this significance threshold for any of the three tested anthropometric traits after meta-analysis (**Table S3**).

To increase statistical power, we attempted to replicate genes with a nominal $P < 0.05$ in the ESP dataset using the WGS data from the MHI Biobank (N=1,976). We limited our analysis to genes with at least two KO individuals. Although the MHI Biobank dataset results from low-pass WGS, the number of identified LoF variants and gene KOs was similar to the number observed in ESP (**Table S1**), suggesting that the data is sufficiently comprehensive to support these analyses. We found that 30-40% of gene KOs in ESP were also knocked out in the MHI Biobank, highlighting the challenge to replicate such studies in humans. This might particularly be true for gene KOs observed only in ESP African Americans given that the MHI Biobank includes individuals of European ancestry. We combined the ESP and MHI Biobank results but we did not observe any significant associations with quantitative anthropometric traits (**Table S3**). We report results with a meta-analysis $P < 0.005$ in **Table 2**. The most promising gene KO association that we found is between *PKHD1L1* and lower BMI: we found 20 KO individuals for

this gene who have on average a BMI that is 0.8 standard deviation (SD) below the population mean (corresponding to ~ 3.6 kg/m²). *PKHD1L1* may play a role in immunity (21).

While examining the top candidate genes, we noticed that *PKHD1L1* is a large gene (78 exons, coding sequence is ~ 14 kilobases), raising the possibility that our method could favor longer genes. In the ESP dataset, we found, as expected, that the number of LoF variants in a given gene is strongly correlated with the length of the coding sequence or the number of exons (all $P < 1 \times 10^{-67}$). However, the number of individuals who carry a rare gene KO is not correlated with the length of the coding sequence or the number of exons of the gene (all $P > 0.2$), except for a weak correlation observed in ESP African Americans with the length of the coding sequence (Pearson's $r = 0.066$, $P = 0.003$). To exclude the possibility that gene length may influence our results, we tested correlations with association results from the ESP and MHI Biobank combined analyses. With one exception (among 12 correlation tests performed), we found no significant correlations between the length of the coding sequence or the number of exons and association P -values for BMI, height, and WHR (all $P > 0.25$). In ESP African Americans, there was a weak correlation between the length of the coding sequence and the BMI P -values (Pearson's $r = 0.069$, $P = 0.002$), but it was in the opposite direction from our expectations (shorter genes have slightly more significant P -values). Together, these analyses suggest that our method to test association between gene KOs and human quantitative traits is largely insensitive to gene length.

Gene KO identification and association testing using exome array data

Recognizing that the main limitation of our analysis is sample size, we contacted studies that are involved in the GIANT Consortium. Although WES or WGS data is not readily available for most of these studies, they all have genotyped their participants using an exome array that

targets 250,000 – mostly coding – variants. We reasoned that the large sample size offered by the GIANT Consortium could compensate for the limited number of variants present on the exome array. We recruited 22 studies, totaling >102,000 individuals (BMI and height available for all, WHR available for >62,000 individuals). Each study ran the method locally, stratifying all analyses by ethnicity, and we then combined results using meta-analysis methodology (22). The frequency of KO events was similar in ESP and the GIANT studies. However, there were more singletons (genes with a single KO individual) observed in European-ancestry individuals from the GIANT studies because of the very large sample size (**Figure S3**).

We present the BMI, height, and WHR meta-analysis results for the GIANT studies in **Figure 3**. As reported above for the WES sequence datasets, and despite a sample size that is >10-times larger, we could not detect significant associations between gene KOs and quantitative anthropometric traits after accounting for the number of tests performed (**Table 3** and **Table S4**). The most interesting finding pertains to the association between height and inactivation of *GRHPH*: autosomal recessive Mendelian mutations in this gene cause primary hyperoxaluria type 2 [MIM 260000](23). Primary hyperoxaluria type 1 [MIM 259900], a more severe form of the disease caused by mutations in *AGXT*, is characterized by very severe growth failure (24). However, the connection between primary hyperoxaluria type 2 caused by recessive mutations in *GRHPH* and growth in humans has not been as clearly documented, although there is one case report of a child with this disease and short stature (25).

Prioritizing gene KOs using a candidate-gene approach

We next asked whether we would increase power to detect associations between gene KO and anthropometric traits by restricting our analyses to strong candidate genes. We focused on

subsets of genes that are associated with any phenotypes in OMIM, or genes that are intolerant to LoF mutations based on the Residual Variation Intolerance Score (RVIS) or the probability of being LoF Intolerant (pLI) score (20, 26). We observed several genes that deviate from the null when restricting our analyses to these candidate genes, especially for the OMIM genes in the larger GIANT datasets for BMI and WHR (**Figure 4**). We also reasoned that the Mouse Genome Informatics (MGI) database might be a good source of candidate genes for our human KO experiment. We retrieved the human homologues of genes from 30 MGI phenotype categories, and tested them against anthropometric traits (**Figure S5**). Again, we observed inflation of the KO association results when compared to the null distribution, suggesting that some of these genes might influence anthropometric traits when completely inactivated. The most noticeable result was the distribution of test statistics in the GIANT BMI analysis for genes related to taste and olfaction (**Figure S5**). Genes related to this category were significantly enriched for genes with a BMI P -value <0.05 in GIANT (Fisher's exact test $P=0.008$).

DISCUSSION

We developed a simple statistical method to test the association between predicted gene KOs and human quantitative traits. We tested our method on three quantitative anthropometric traits (BMI, height, and WHR) in large DNA sequencing (ESP and MHI Biobank, >6,400 individuals) and genotyping (22 participating GIANT studies, >102,000 individuals) datasets. Despite this large sample size, we did not identify significant genetic associations with predicted gene KOs, although the association between *PKHD1L1* and BMI or *GRHPH* and height are interesting and should be tested for replication. Within the limitations of our study design (sample size, incomplete catalogue of LoF variants), our results suggest that there are no predicted gene KOs with modest-to-large effect size on anthropometric trait variation in the general population.

This conclusion is largely consistent with results from a recent study of homozygous LoF variants in 1,432 individuals (17).

Importantly, our approach and results can guide future gene KO studies in humans. First, our method assumes that all LoF alleles for a given gene will shift the phenotypic mean in the same direction. Although it is a sensitive approach in this first large-scale gene KO experiment for quantitative traits, alternative methods could explore effect on phenotypic variance. Second, in order to maximize our sample size, we combined datasets from different technologies (WES, WGS, exome array). Although we accounted for this technical heterogeneity – gene KO statistics were similar across datasets – this approach could have introduced unanticipated biases. Ideally, high coverage WGS data would be available for gene KO studies. Third, haplotype phasing of DNA sequence data from unrelated individuals (ESP and MHI Biobank), and the lack of phase information for the GIANT ExomeChip studies, has limited our ability to identify compound heterozygous individuals. This could impact our results as nearly 20% of all gene KOs identified in this study were due to compound heterozygosity. We note, however, that excluding compound heterozygotes from the ESP analyses had very limited impact on our results (**Figure S6**). Fourth, we only considered nonsense, splice site, stop-loss and frameshift indels as LoF variants to identify gene KOs. Some of these variants are likely neutral: for instance, genes are more tolerant to non-synonymous variants at the 3' end of a gene, and nearby variants can rescue the effect of LoF alleles (12). Furthermore, we excluded missense variants from our analyses, although functional characterization can lead to the identification of missense alleles with strong phenotypic effect on human complex phenotypes (27, 28).

The main limiting factors of gene KO studies in humans are the sample size and the depth of genetic information available. We have shown that even when the sample size is very large, most gene KOs are identified in single individuals (**Figure S3**). To be successful, we will need to develop tools to prioritize genes or increase the number of gene KOs. One possibility may be to consider only genes expressed in a tissue that is relevant for the phenotype of interest (*e.g.* genes expressed in growth plates for height). Another promising solution may be to consider KOs in biological pathways instead of single genes as the testing unit. For instance, a researcher interested in blood lipid genetics could pool together all individuals that carry a gene KO in any of the enzymes or transporters implicated in lipid metabolism. We illustrated this candidate-gene approach by prioritizing OMIM disease-causing genes, genes intolerant to LoF mutations, and genes with relevant mouse KO phenotypes. In particular for the BMI analysis, the enrichment of genes with mouse homologues that disrupt taste or olfaction when inactivated is of interest (**Figure S5**). Reverse genetic strategies – finding a function to a gene by first disrupting it – have been very successful in model organisms. Despite early challenges, the large-scale identification of LoF variants and characterization of gene KOs promise to also yield interesting insights into human biology.

MATERIALS AND METHODS

Ethics statement

This project was approved by the Ethics Committee of the Montreal Heart Institute (#11-1333, #2013-297, #2013-1438).

NHLBI Exome Sequence Project (ESP)

We conducted our initial analysis on the final whole-exome ESP dataset, which is described elsewhere (9). This dataset was generated from high coverage WES (median depth >100X)(9). All study participants in each of the component studies provided written informed consent for the use of their DNA in studies aimed at identifying genetic risk variants for disease and for broad data sharing. Institutional certification was obtained for each sample to allow deposition of phenotype and genotype data in dbGaP and BAM files in the short-read archive. We excluded individuals based on sex mismatch between clinical database and genotype-inferred sex (N=13), high homozygosity (N=1), high genotyping missing rate (>10%)(N=1), if they appear as population outliers in principal component analyses (N=30), low concordance to genome-wide association study data (N=4), or unresolved participant identifiers (N=4). Moreover, we randomly excluded one member of each pair of duplicates (N=16), and of first- and second-degree relatives (N=108). We also removed individuals with chronic obstructive pulmonary disease or asthma, as these conditions could influence anthropometric traits (N=688). Finally, we removed participants from the CARDIA (N=201) and MESA (N=406) studies, as requested by investigators from these studies. We kept individuals aged between 21 and 80 years old, height between 125 and 240 cm, BMI <75 kg/m², and WHR <1.5. In total, we analyzed anthropometric traits in 1,726 African Americans and 2,772 European Americans (**Table S1**).

Variant quality-control and annotation

We phased variants in the ESP dataset using Beagle 4.0 and the default parameters (19). We define LoF variants as variants that create or remove stop codons (nonsense and stop-loss) that disrupt essential splice sites (two intronic bases at exon-intron boundaries), or that change the reading frame (frameshift indel). We annotated single-base pair variants using in-house custom

scripts and build 37.1 of the human genome reference sequence. We annotated frameshift indels using SeattleSeq (hg19, v.9.03, <http://snp.gs.washington.edu/SeattleSeqAnnotation138/>). We included in our analyses only frameshift indel variants that fall within validated RefSeq genes (release 69). After filtering out variants with a call rate <95% or a Hardy-Weinberg $P < 1 \times 10^{-6}$, we retained in our analyses 18,137 and 21,935 LoF variants in African- and European-ancestry individuals, respectively (**Table S1**). For comparison, we also annotated ESP variants using Ensembl's Variant Effect Predictor (VEP) module and basic transcripts from GENCODE. We obtained very similar results (**Figure S1**).

Replication cohorts with WGS or WES data available

We used low-pass WGS data (mean coverage 5.7X) from 2,002 French-Canadian participants recruited by the Montreal Heart Institute (MHI) Biobank. Genotypes were imputed and phased using Beagle 4.0 using the default parameters (19). Individuals were removed due to low or high inbreeding coefficient ($N=4$). Variants with Hardy-Weinberg $P < 1 \times 10^{-8}$ were excluded. In total, 1,976 MHI Biobank participants with anthropometric traits available were included in the replication analyses (**Table S1**).

GIANT Consortium ExomeChip datasets

We analyzed Illumina ExomeChip genotype data from 22 studies that are members of the Genetic Investigation of ANthropometric Traits (GIANT) Consortium (**Table S1**). In total, 103,838, 102,775, and 62,355 individuals were included in the BMI, height and BMI-adjusted WHR analyses, respectively. Individuals were from European- ($N=90,927$; 19 studies), African- ($N=7,576$; 2 studies), and Hispanic-ancestry ($N=5,335$; 1 study). To increase the number of LoF variants available on the ExomeChip, we broaden our definition of splice-site variants to include

variants located two base pairs on either side of exon-intron boundaries. This is the splice-site definition implemented by dbNSFP (29) and used by GIANT across the Consortium's ExomeChip effort. Using the most severe annotation from ENSEMBL's VEP tool, we found that 17.8% (797/4483) of these splice site-variants disrupt a canonical splice-site, 46.7% (2094/4483) are missense variants, and 31.6% (1419/4483) affect a nucleotide around the splice-site (1-3 bases within exon or 3-8 bases within intron). Phasing information was not available for the GIANT exome array data. Because we focused on rare variants, we assumed that when two rare LoF variants were observed in the same gene in the same individual, they were inherited in *trans* to create a compound heterozygous gene KO.

Statistical analyses

We developed a flexible method to determine if the complete inactivation of genes by LoF variants is associated with human quantitative traits (**Figure 2**). For each gene, our method searches for individuals that are either homozygotes or compound heterozygotes for LoF variants; we refer to these individuals as predicted KOs. For X-linked markers that fall outside of the pseudoautosomal regions, we consider predicted gene KOs in men if they carry a single LoF variant. For compound heterozygosity, we use phase information to distinguish LoF variants that segregate on the same haplotype (in *cis*) or on different haplotypes (in *trans*). When phasing information is not available (*e.g.* GIANT ExomeChip data), we assume that rare LoF variants segregate on different haplotypes. The method then calculates for each gene the phenotypic mean in predicted KO individuals. Finally, it computes statistical significance using phenotype permutations, as follows:

$$P_{left} = \frac{\sum_{i=1}^n \mathbb{1}_{m_i \leq m}}{n}; P_{right} = \frac{\sum_{i=1}^n \mathbb{1}_{m_i \geq m}}{n}$$

$$P_{final} = 2 \times \text{minimum}(P_{left}, P_{right}),$$

where $\mathbb{1}$ is the indicator function, m is the observed mean phenotype in predicted KO individuals, m_i is the i^{th} permuted mean, n is the number of permutations, P_{left} and P_{right} are the left- and right-tail P -values, and P_{final} is the reported two-tailed P -value. Using simulated null phenotypes and the ESP dataset, we showed that the test is well-calibrated (**Figure S4**). This method assumes that gene inactivation results in the same phenotypic effect (increase or decrease trait value) in all predicted KO individuals for a given gene. The current implementation of our method also currently assumes that tested individuals are unrelated and that the phenotypic distributions are symmetrical. It is compatible with standard genotype file formats (*e.g.* PLINK, vcf). The scripts to run our method are publicly available at: <http://www.mhi-humangenetics.org/en/resources>.

Association of rare predicted gene KOs with anthropometric traits

We analyzed BMI, adult height and BMI-adjusted WHR. We stratified all our analyses by ethnic group, and we only considered rare or low-frequency LoF variants with MAF <5%. We used 10,000 permutations to assess statistical significance. For genes with an empirical $P < 2 \times 10^{-4}$ (*i.e.* permuted means were never higher (or lower) than the observed mean among 10,000 permutations), we re-ran the analysis using 100,000 permutations: only two genes fell in that category (*BRPF1* $P_{\text{height}} = 1.8 \times 10^{-4}$; *SPZI* $P_{\text{WHR}} = 2.2 \times 10^{-4}$). For ESP samples, we corrected anthropometric traits for sex, age, ESP phenotype groups, exon capture reagents and the first three principal components, as recommended by the ESP investigators. We then applied inverse normal transformation on the residuals from the previous correction. For the MHI Biobank, and

the GIANT studies, each anthropometric trait was corrected for sex, age, age-squared and the first ten principal components, and we normalized the resulting residuals using inverse normal transformation. Taking into account the direction of the effect, we combined results across studies using a weighted Z-score meta-analysis method implemented in the software METAL, where the weight is the sample size of the corresponding study (22). To estimate statistical power of our approach, we modeled the effect of a recessive LoF variant on a normally distributed quantitative trait, as previously described (30). This is a simplistic model as we ignore the presence of additional LoF variants in the same gene, which are considered in our method because they can lead to additional individuals that have a predicted gene KO. We assume that the variant has a MAF=5%, explains 1% of the genetic variance, and used a sample size of $N=4,500$ (corresponding to ESP), $\alpha=2 \times 10^{-5}$ (Bonferroni correction for the number of genes with KOs), and 5,000 simulations to perform power calculations. Under this scenario, our gene KO approach would have 95% power to detect the association. Alternatively, testing the association while assuming that the variant has an additive effect would result in only 3% power. Using the same assumptions, we estimated 64% and 1% power for a variant that explains 0.5% of the variance when tested using our gene KO methodology or a simple additive model, respectively.

Candidate-gene enrichment analyses

We explored whether prioritizing gene KOs into different categories could increase the chance to reveal an association. First, we investigated whether the gene was an OMIM disease-causing gene, as defined elsewhere (26). Next, we considered whether the genes were LoF intolerant by either having a Residual Variation Intolerance Score (RVIS) < 15% of the RVIS scores for all genes in the human genome (release 0.3) or a probability of being LoF intolerant (pLI) score > 0.9 (20, 26). We looked for enrichment by overlapping the QQ-plots of genes

belonging to these different categories separately on the QQ-plot containing all genes. We also created subsets of genes based on 30 phenotype categories from the Mouse Genome Informatics (MGI) Database (31). We tested the enrichment using Fisher's exact test.

ACKNOWLEDGMENTS

We thank all participants involved in this project, and Ekat Kritikou for comments on the manuscript. PBM and MC acknowledge this work forms part of the research program of the NIHR Barts Cardiovascular Biomedical Research Unit. MC is a senior National Institute for Health Research Investigator. Sequencing of the MHI Biobank samples was performed at the McGill University and Génome Québec Innovation Centre. The authors wish to acknowledge the support of the National Heart, Lung, and Blood Institute (NHLBI) and the contributions of the research institutions, study investigators, field staff and study participants in creating this resource for biomedical research. SL is funded by a Canadian Institutes of Health research Banting doctoral scholarship. GL is funded by Genome Canada and Génome Québec; the Canada Research Chair program; and the Montreal Heart Institute Foundation. CML is supported by Wellcome Trust [grant numbers 086596/Z/08/Z, 086596/Z/08/A]; and the Li Ka Shing Foundation. NS is funded by National Institutes of Health [grant numbers HL088456, HL111089, HL116747]. The Mount Sinai BioMe Biobank Program is supported by the Andrea and Charles Bronfman Philanthropies. GO ESP is supported by NHLBI [RC2 HL-103010 to HeartGO, RC2 HL-102923 to LungGO, RC2 HL-102924 to WHISP]. The ESP exome sequencing was performed through NHLBI [RC2 HL-102925 to BroadGO, RC2 HL-102926 to SeattleGO]. EGCUT work was supported through the Estonian Genome Center of University of Tartu by the Targeted Financing from the Estonian Ministry of Science and Education [grant number

SF0180142s08]; the Development Fund of the University of Tartu [grant number SP1GVARENG]; the European Regional Development Fund to the Centre of Excellence in Genomics (EXCEGEN) [grant number 3.2.0304.11-0312]; and through FP7 [grant number 313010]. EGCUT were further supported by the US National Institute of Health [grant number R01DK075787].

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflicts of interest.

REFERENCES

- 1 Alkuraya, F.S. (2015) Human knockout research: new horizons and opportunities. *Trends in genetics : TIG*, **31**, 108-115.
- 2 Kamb, A., Harper, S. and Stefansson, K. (2013) Human genetics as a foundation for innovative drug development. *Nature biotechnology*, **31**, 975-978.
- 3 Antonarakis, S.E. and Beckmann, J.S. (2006) Mendelian disorders deserve more attention. *Nat Rev Genet*, **7**, 277-282.
- 4 Amselem, S., Duquesnoy, P., Attree, O., Novelli, G., Bousnina, S., Postel-Vinay, M.C. and Goossens, M. (1989) Laron dwarfism and mutations of the growth hormone-receptor gene. *The New England journal of medicine*, **321**, 989-995.
- 5 Clement, K., Vaisse, C., Lahlou, N., Cabrol, S., Pelloux, V., Cassuto, D., Goumelen, M., Dina, C., Chambaz, J., Lacorte, J.M. *et al.* (1998) A mutation in the human leptin receptor gene causes obesity and pituitary dysfunction. *Nature*, **392**, 398-401.
- 6 Jackson, R.S., Creemers, J.W., Farooqi, I.S., Raffin-Sanson, M.L., Varro, A., Dockray, G.J., Holst, J.J., Brubaker, P.L., Corvol, P., Polonsky, K.S. *et al.* (2003) Small-intestinal dysfunction accompanies the complex endocrinopathy of human proprotein convertase 1 deficiency. *The Journal of clinical investigation*, **112**, 1550-1560.
- 7 1000 Genomes Project Consortium. (2010) A map of human genome variation from population-scale sequencing. *Nature*, **467**, 1061-1073.
- 8 1000 Genomes Project Consortium. (2012) An integrated map of genetic variation from 1,092 human genomes. *Nature*, **491**, 56-65.

- 9 Fu, W., O'Connor, T.D., Jun, G., Kang, H.M., Abecasis, G., Leal, S.M., Gabriel, S., Rieder, M.J., Altshuler, D., Shendure, J. *et al.* (2013) Analysis of 6,515 exomes reveals the recent origin of most human protein-coding variants. *Nature*, **493**, 216-220.
- 10 Tennessen, J.A., Bigham, A.W., O'Connor, T.D., Fu, W., Kenny, E.E., Gravel, S., McGee, S., Do, R., Liu, X., Jun, G. *et al.* (2012) Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science*, **337**, 64-69.
- 11 Alsalem, A.B., Halees, A.S., Anazi, S., Alshamekh, S. and Alkuraya, F.S. (2013) Autozygome sequencing expands the horizon of human knockout research and provides novel insights into human phenotypic variation. *PLoS genetics*, **9**, e1004030.
- 12 MacArthur, D.G., Balasubramanian, S., Frankish, A., Huang, N., Morris, J., Walter, K., Jostins, L., Habegger, L., Pickrell, J.K., Montgomery, S.B. *et al.* (2012) A systematic survey of loss-of-function variants in human protein-coding genes. *Science*, **335**, 823-828.
- 13 Gorlov, I.P., Gorlova, O.Y., Sunyaev, S.R., Spitz, M.R. and Amos, C.I. (2008) Shifting paradigm of association studies: value of rare single-nucleotide polymorphisms. *American journal of human genetics*, **82**, 100-112.
- 14 Sulem, P., Helgason, H., Oddson, A., Stefansson, H., Gudjonsson, S.A., Zink, F., Hjartarson, E., Sigurdsson, G.T., Jonasdottir, A., Jonasdottir, A. *et al.* (2015) Identification of a large set of rare complete human knockouts. *Nature genetics*, **47**, 448-452.
- 15 Li, A.H., Morrison, A.C., Kovar, C., Cupples, L.A., Brody, J.A., Polfus, L.M., Yu, B., Metcalf, G., Muzny, D., Veeraraghavan, N. *et al.* (2015) Analysis of loss-of-function variants and 20 risk factor phenotypes in 8,554 individuals identifies loci influencing chronic disease. *Nat Genet*, in press.

- 16 Lim, E.T., Wurtz, P., Havulinna, A.S., Palta, P., Tukiainen, T., Rehnstrom, K., Esko, T., Magi, R., Inouye, M., Lappalainen, T. *et al.* (2014) Distribution and medical impact of loss-of-function variants in the Finnish founder population. *PLoS genetics*, **10**, e1004494.
- 17 Kaiser, V.B., Svinti, V., Prendergast, J.G., Chau, Y.Y., Campbell, A., Patarcic, I., Barroso, I., Joshi, P.K., Hastie, N.D., Miljkovic, A. *et al.* (2015) Homozygous loss-of-function variants in European cosmopolitan and isolate populations. *Human molecular genetics*, in press.
- 18 Lim, E.T., Raychaudhuri, S., Sanders, S.J., Stevens, C., Sabo, A., MacArthur, D.G., Neale, B.M., Kirby, A., Ruderfer, D.M., Fromer, M. *et al.* (2013) Rare complete knockouts in humans: population distribution and significant role in autism spectrum disorders. *Neuron*, **77**, 235-242.
- 19 Browning, S.R. and Browning, B.L. (2007) Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *American journal of human genetics*, **81**, 1084-1097.
- 20 Lek, M., Karczewski, K., Minikel, E., Samocha, K., Banks, E., Fennell, T., O'Donnell-Luria, A., Ware, J., Hill, A., Cummings, B. *et al.* (2015) Analysis of protein-coding genetic variation in 60,706 humans. *bioRxiv*, in press.
- 21 Hogan, M.C., Griffin, M.D., Rossetti, S., Torres, V.E., Ward, C.J. and Harris, P.C. (2003) PKHDL1, a homolog of the autosomal recessive polycystic kidney disease gene, encodes a receptor with inducible T lymphocyte expression. *Human molecular genetics*, **12**, 685-698.
- 22 Willer, C.J., Li, Y. and Abecasis, G.R. (2010) METAL: Fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*, **26**, 2190-2191.
- 23 Cramer, S.D., Ferree, P.M., Lin, K., Milliner, D.S. and Holmes, R.P. (1999) The gene encoding hydroxypyruvate reductase (GRHPR) is mutated in patients with primary hyperoxaluria type II. *Human molecular genetics*, **8**, 2063-2069.

- 24 Nissel, R., Latta, K., Gagnadoux, M.F., Kelly, D., Hulton, S., Kemper, M.J., Ruder, H., Soderdahl, G., Otte, J.B., Cochat, P. *et al.* (2006) Body growth after combined liver-kidney transplantation in children with primary hyperoxaluria type 1. *Transplantation*, **82**, 48-54.
- 25 Dent, C.E. and Stamp, T.C. (1970) Treatment of primary hyperoxaluria. *Arch Dis Child*, **45**, 735-745.
- 26 Petrovski, S., Wang, Q., Heinzen, E.L., Allen, A.S. and Goldstein, D.B. (2013) Genic intolerance to functional variation and the interpretation of personal genomes. *PLoS Genet*, **9**, e1003709.
- 27 Majithia, A.R., Flannick, J., Shahinian, P., Guo, M., Bray, M.A., Fontanillas, P., Gabriel, S.B., Go, T.D.C., Project, N.J.F.A.S., Consortium, S.T.D. *et al.* (2014) Rare variants in PPARG with decreased activity in adipocyte differentiation are associated with increased risk of type 2 diabetes. *Proc Natl Acad Sci U S A*, **111**, 13127-13132.
- 28 Thormaehlen, A.S., Schuberth, C., Won, H.H., Blattmann, P., Joggerst-Thomalla, B., Theiss, S., Asselta, R., Duga, S., Merlini, P.A., Ardissino, D. *et al.* (2015) Systematic cell-based phenotyping of missense alleles empowers rare variant association studies: a case for LDLR and myocardial infarction. *PLoS Genet*, **11**, e1004855.
- 29 Liu, X., Jian, X. and Boerwinkle, E. (2013) dbNSFP v2.0: a database of human non-synonymous SNVs and their functional predictions and annotations. *Hum Mutat*, **34**, E2393-2402.
- 30 Lettre, G., Lange, C. and Hirschhorn, J.N. (2007) Genetic model testing and statistical power in population-based association studies of quantitative traits. *Genet Epidemiol*, **31**, 358-362.
- 31 Eppig, J.T., Blake, J.A., Bult, C.J., Kadin, J.A., Richardson, J.E. and Mouse Genome Database, G. (2015) The Mouse Genome Database (MGD): facilitating mouse as a model for human biology and disease. *Nucleic Acids Res*, **43**, D726-736.

FIGURE LEGENDS

Figure 1. Distributions of the number of NHLBI Exome Sequence Project (ESP) participants with predicted gene knockouts (KOs). We present distributions in African Americans (**A** and **B**) and European Americans (**C** and **D**). We include all loss-of-function (LoF: nonsense, stop-loss, splice site, frameshift indel) variants in **A** and **C**, whereas only rare/low-frequency LoF variants (minor allele frequency <5%) are included in **B** and **D**. Homo., gene KO due to homozygosity; Comp. het., gene KO due to compound heterozygosity; Both, genes with homozygous and compound heterozygous LoF variants.

Figure 2. Schematic representation of the method to detect association between gene knockouts (KOs) and human quantitative variation. This example depicts a fictive gene with three exons (*GENE1*) that contains several SNPs. Our analytical framework only considers loss-of-function (LoF) variants (shown in red). *GENE1* KOs are individuals who are either compound heterozygous or homozygous for LoF variants (individual 1 and 2). The histogram shows the distribution of a normalized human quantitative trait. Our method tests whether individuals that are KOs for a given gene (red arrows) have on average more extreme phenotypes than the rest of the individuals.

Figure 3. Quantile-quantile (QQ) plots of association results between predicted gene knockouts (KOs) and anthropometric traits in the (**A-C**) NHLBI Exome Sequence Project (ESP) and (**D-F**) GIANT ExomeChip datasets. In these datasets, we only considered loss of function (LoF) variants (nonsense, stop-loss, splice site, frameshift indels (ESP only)) with a minor allele frequency (MAF) <5%. We analyzed three anthropometric traits: (**A**) body mass index (BMI)

($N_{\text{participants}}=4,475$), (**B**) height ($N_{\text{participants}}=4,423$), and (**C**) waist-to-hip ratio (WHR) ($N_{\text{participants}}=2,973$). We performed these analyses stratified by ethnicity, and then combined the European American and African American results using meta-analysis methodology. We analyzed the same traits in the GIANT dataset: (**D**) BMI ($N_{\text{participants}}=103,838$), (**E**) height ($N_{\text{participants}}=102,775$), and (**F**) WHR ($N_{\text{participants}}=62,355$). Results are not corrected for the genomic inflation factor. The dash lines correspond to the 95% confidence interval. λ_{GC} , genomic inflation factor; N_{gene} , number of genes with at least one participant that carries two LoF alleles.

Figure 4. Quantile-quantile (QQ) plots of association results between predicted gene knockouts (KOs) in candidate-genes and anthropometric traits. We restricted these analyses to OMIM disease-causing genes (green), genes with Residual Variation Intolerance Score (RVIS) score < 15% of RVIS scores for all genes in the human genome (red), or genes with a probability of being loss-of-function intolerant (pLI) score > 0.9 (blue). We report results for three anthropometric traits in the NHLBI Exome Sequence Project (ESP): (**A**) body mass index (BMI) ($N_{\text{participants}}=4,475$), (**B**) height ($N_{\text{participants}}=4,423$), and (**C**) waist-to-hip ratio (WHR) ($N_{\text{participants}}=2,973$). We also report results for the same traits in the GIANT ExomeChip datasets: (**D**) BMI ($N_{\text{participants}}=103,838$), (**E**) height ($N_{\text{participants}}=102,775$), and (**F**) WHR ($N_{\text{participants}}=62,355$). Results are not corrected for the genomic inflation factor. The dash lines correspond to the 95% confidence interval. λ_{GC} , genomic inflation factor; N_{gene} , number of genes with at least one participant that carries two LoF alleles.

TABLES

Table 1. Number and frequency of predicted gene knockouts (KO) in 1,727 African Americans and 2,772 European Americans from the NHLBI Exome Sequence Project

(ESP). For this loss-of-function (LoF) variant analysis, we consider autosomal nonsense, stop-loss and splice site variants, as well as frameshift insertion-deletions (indels). Rare LoF variants have a minor allele frequency <5%. In the absence of phasing information, we assume that rare LoF are inherited in trans. As expected, considering phased genotype information significantly impacts the number of gene KOs that we can detect due to compound heterozygosity.

			Variants /individuals	Variants /gene	NOT PHASED		PHASED	
					Gene KOs /individuals	Number of KO genes	Gene KOs /individuals	Number of KO genes
African Americans	All LoF (N=18,137)		237	0.92	33.7	2,530	25.9	2,429
		Homozygotes			23.2	2,384	23.2	2,384
		Compound heterozygotes			10.4	601	2.6	334
	Rare LoF (N=17,446)		65	0.89	4.2	2,174	2.5	2,071
		Homozygotes			2.3	2,028	2.3	2,028
		Compound heterozygotes			1.9	381	0.2	155
European Americans	All LoF (N=21,935)		197	1.12	28.8	1,844	23.2	1,741
		Homozygotes			21.3	1,694	21.3	1,694
		Compound heterozygotes			7.6	487	1.9	247
	Rare LoF		39	1.09	1.8	1,538	1.1	1,433

	(N=21,351)							
		Homozygotes			1	1,390	1.01	1,390
		Compound heterozygotes			0.8	318	0.09	124

Table 2. Association of gene knockouts (KOs) with anthropometric traits in the Exome Sequence Project (ESP) and Montreal Heart Institute (MHI) Biobank DNA sequencing datasets. We attempted to replicate gene KO associations from the ESP whole-exome DNA sequencing dataset in the MHI Biobank whole-genome DNA sequencing dataset. We tested for replication genes with $P < 0.05$ and at least two KO individuals in the ESP dataset. We report genes with a combined $P < 0.005$. We provide the mean gene KO effect size in standard deviation (SD) and metric units, assuming that 1 SD corresponds to 4.5 kg/m², 6.4 cm, and 0.07 for BMI, height, and WHR respectively. N_{KO}: number of individuals that are KO for the given gene. EA: European-ancestry; AA: African-ancestry.

Trait	Gene	ESP					MHI			Combined	
		Mean EA (real units)	N _{KO} EA	Mean AA (real units)	N _{KO} AA	P	Mean (real units)	N _{KO}	P	Weighted average (real units)	P
BMI	<i>PKHD1L1</i>	0.7 (+3.2 kg/m ²)	11	0.5 (+2.3 kg/m ²)	6	0.009	1.6 (+7.2 kg/m ²)	3	0.009	0.8 (+3.6 kg/m ²)	0.0002
	<i>PLIN4</i>	2.7 (+12.2 kg/m ²)	1	3.1 (+14.0 kg/m ²)	1	5x10 ⁻⁵	-0.2 (-0.9 kg/m ²)	2	0.67	1.4 (+6.3 kg/m ²)	0.002
Height	<i>RMDN2</i>	NA	0	-1.1 (-7.0 cm)	4	0.03	-1.6 (-10.2 cm)	2	0.02	-1.3 (-8.3 cm)	0.002
	<i>ASIC4</i>	3.6 (23.0 cm)	1	1.5 (9.6 cm)	2	5x10 ⁻⁵	-0.4 (-2.6 cm)	2	0.56	1.2 (+7.7 cm)	0.002
	<i>SH2B2</i>	-1.6	2	NA	0	0.02	-1.9	1	0.06	-1.7	0.003

		(-10.2 cm)					(-12.2 cm)			(-10.9 cm)	
WHR	<i>CIQTNF5</i>	0.6 (+0.04)	1	1.8 (+0.13)	2	0.04	1.5 (+0.11)	2	0.03	1.4 (0.10)	0.003

Table 3. Top association results between anthropometric traits and predicted gene knockouts (KOs) identified using ExomeChip data from 22 studies participating in the GIANT Consortium. We only report genes with $P < 0.005$ and at least two KO individuals. The weighted mean corresponds to the average phenotype (in standard deviation units) of individuals that are KO for this gene. N_{KO} : number of individuals with a KO gene; N_{study} : number of studies with at least one KO individual for a given gene.

Trait	Gene	N_{KO}	N_{study}	Weighted mean (SD)	P
BMI	<i>CYP20A1</i>	100	15	-0.35	0.001
	<i>ME2</i>	2	2	-1.90	0.002
	<i>KIAA1024</i>	7	5	-0.75	0.002
	<i>TBC1D5</i>	4	3	1.05	0.003
	<i>LRR39</i>	147	16	0.23	0.003
	<i>TAS1R1</i>	191	6	0.15	0.004
	<i>LAMA3</i>	9	2	-1.02	0.004
	<i>KIAA0391</i>	3	2	-1.64	0.004
	<i>TAS2R60</i>	2	2	-2.04	0.005
Height	<i>GRHPR</i>	2	2	-2.28	0.0001
	<i>ABCB7</i>	365	10	-0.12	0.0003
	<i>ZDHHC14</i>	3	2	-2.01	0.0003
	<i>ZFPM1</i>	21	3	-0.60	0.0008
	<i>DHX57</i>	2	2	-2.04	0.0009
	<i>CD8A</i>	2	2	2.35	0.001
	<i>CDC42BPA</i>	4	2	1.70	0.001
	<i>NSUN4</i>	13	3	-0.78	0.002

	<i>ARPC5L</i>	6	2	-1.25	0.002
	<i>CCDC125</i>	45	9	0.35	0.002
	<i>BOK</i>	27	4	0.61	0.003
	<i>NSRP1</i>	9	1	-1.00	0.003
	<i>TEX13A</i>	2	1	2.00	0.004
	<i>RPGRIP1</i>	10	4	-0.72	0.004
	<i>SCGN</i>	6	5	-0.96	0.005
WHR	<i>C18orf56</i>	7	1	1.39	0.0002
	<i>AARS2</i>	3	2	-1.78	0.001
	<i>C18orf34</i>	6	3	1.27	0.002
	<i>CCDC68</i>	13	1	0.83	0.002
	<i>HRG</i>	3	2	-1.52	0.004
	<i>SPTA1</i>	2	2	1.86	0.004
	<i>SPTBN5</i>	191	11	0.15	0.005

ABBREVIATIONS

BMI: Body mass index

ESP: Exome Sequence Project

GIANT: Genetic Investigation of ANthropometric Traits

KO: Knockout

LoF: Loss-of-function

MAF: Minor allele frequency

MHI: Montreal Heart Institute

WES: Whole-exome DNA sequencing

WGS: Whole-genome DNA sequencing

WHR: Waist-to-hip ratio







