## SUPPLEMENTAL FIGURE LEGENDS

**Figure S1.** NM-HA does not form visible aggregates in N2a cells as detected by anti-Sup35p-N rat monoclonal antibody 7H5. N2a cells were transfected with a NM-HA construct and studied 48 h post transfection. NM-HA was detected using rat monoclonal antibody 7H5. *A*) N2a cells expressing NM-HA were fixed and stained for NM-HA (red). Nuclei were visualized using Hoechst staining (blue). No visible foci were observed. Untransfected cells were not stained by the antibody. Scale bar: 10 μm. *B*) Sedimentation assay of NM-HA. Lysate fractions were subjected to SDS gel electrophoresis. The fractions of insoluble (pellet = P) and soluble recombinant proteins (supernatant = S) are shown. As a control, untransfected cells were lysed and lysates fractioned into soluble and insoluble fractions. NM-HA remains exclusively in the supernatant. *C*) Proteinase K digestion of NM-HA. Cell lysates were subjected to PK treatment using the indicated amounts of PK. NM-HA was completely digested upon treatment with 5 μg/ml PK for 30 min.

**Figure S2.** PrP<sub>cyto</sub>, N-PrP and NM-PrP aggregates are not located in intracellular organelles or vesicles. N2a cells expressing the constructs were fixed and stained for PrP<sub>cyto</sub>, N-PrP and NM-PrP with mAb 3F4 48 h post transfection. Nuclei were visualized using Hoechst staining. Merged images demonstrate missing co-localization of aggregates with intracellular compartments. *A*) Co-staining for ER marker Calnexin. *B*) Transfected N2a cells were co-stained for Rab6. *C*) N2a cells were transfected with constructs and Rab4-GFP, a marker for early endosomes. *D*) N2a cells were co-transfected with the constructs and Rab11-GFP that visualizes late and recycling endosomes. Results were independent of the cell line. Scale bars: 10 μm.

**Figure S3.** Analysis of recombinantly produced PrP-M, NM and NM-PrP by SDS-PAGE and subsequent protein staining with Coomassie Blue. Samples of the respective proteins (2 μg of protein per lane) were subjected to SDS-PAGE after 0, 5, 21 and 28 days of incubation. LMW = Low Molecular Weight protein standard (BIORAD). Numbers of the lanes refer to days of incubation.

Fig. S1




Fig. S2




Fig. S2




Fig. S3

