

RESEARCH HIGHLIGHT

Functional consequences of genetic polymorphisms in the NKG2D receptor signaling pathway and putative gene interactions

Antje Isernhagen¹, Dörthe Malzahn², Sebastian Monecke^{1,3}, Daniela Schilling^{4,5}, Pranali Shah¹, Gabriele Multhoff^{4,5}, Gerald Wulf⁶, Dieter Kube⁶, Heike Bickeböller², Ralf Dressel^{1,3}

¹*Institute of Cellular and Molecular Immunology, University Medical Center Göttingen, Göttingen, 37099, Germany*

²*Institute of Genetic Epidemiology, University Medical Center Göttingen, Göttingen, 37099, Germany*

³*DZHK (German Center for Cardiovascular Research), Partner site Göttingen, Göttingen, 37099, Germany*

⁴*Department of Radiation Oncology, Klinikum rechts der Isar, Technische Universität München, München, 81675, Germany*

⁵*Institute of Innovative Radiotherapy (iRT), Department of Radiation Sciences (DRS), Helmholtz Zentrum München, München, 85764, Germany*

⁶*Department of Hematology and Medical Oncology, University Medical Center Göttingen, Göttingen, 37099, Germany*

Correspondence: Ralf Dressel

E-mail: rdresse@gwdg.de

Received: March 25, 2016

Published online: May 02, 2016

NKG2D (NK group 2, member D) is an activating natural killer (NK) receptor, which is expressed on NK and CD8⁺ T cells. On NK cells, NKG2D elicits cytotoxicity and release of cytokines. On CD8⁺ T cells, it functions as a co-stimulatory molecule. The receptor recognizes several ligands including the major histocompatibility complex (MHC) class I chain-related molecules A (MICA) and B (MICB) as well as the UL16-binding proteins (ULBP). The diversity of NKG2D ligands is further increased by a high degree of genetic variability of the ligands. Recently, an amino acid exchange from valine to methionine at position 129 in MICA has been found to be associated with the outcome of allogeneic hematopoietic stem cell transplantation (HSCT), and the functional consequences of this specific genetic variation have been elucidated. The clinical associations found after HSCT were explainable by the functional differences of the MICA-129 variants. Herein, we discuss how the genetic polymorphisms of NKG2D ligands and NKG2D itself interact and may affect the outcome of HSCT and the susceptibility to other diseases, which have been associated with polymorphisms in the NKG2D signaling pathway.

Keywords: NKG2D; NKG2D ligands; MICA; hematopoietic stem cell transplantation; graft versus host disease; genetic polymorphism

To cite this article: Antje Isernhagen, et al. Functional consequences of genetic polymorphisms in the NKG2D receptor signaling pathway and putative gene interactions. Receptor Clin Invest 2016; 3: e1269. doi: 10.14800/rci.1269.

Copyright: © 2016 The Authors. Licensed under a *Creative Commons Attribution 4.0 International License* which allows users including authors of articles to copy and redistribute the material in any medium or format, in addition to remix, transform, and build upon the material for any purpose, even commercially, as long as the author and original source are properly cited or credited.

Introduction

NKG2D (NK group 2, member D) is an activating natural killer (NK) receptor, which is encoded by the *KLRK1* gene in the human NK gene complex on chromosome 12 [1]. The expression of NKG2D at the plasma membrane requires its association with the adaptor protein DNAX-activation protein (DAP)10 [2]. Notably, a truncated isoform of NKG2D, i.e. NKG2D^{TR}, lacking the extracellular domain can be generated by alternative splicing. NKG2D^{TR} can compete with the full-length protein for DAP10 resulting in a reduced expression of the functional NKG2D receptor at the cell surface [3]. NKG2D is expressed on almost all human NK cells, resting CD8⁺ αβ T cells, many γδ T cells and iNKT cells as well as a small subset of effector or memory CD4⁺ T cells [2, 4, 5]. The expression of NKG2D is up-regulated by interleukin (IL)-2, IL-7, IL-12, and IL-15, but down-regulated by transforming growth factor (TGF)-β, interferon (IFN)-β1, and IL-21 [2]. On NK cells, NKG2D signaling elicits degranulation and killing of target cells [6] as well as secretion of cytokines such as IFN-γ [7]. On CD8⁺ αβ T cells, NKG2D functions as a co-stimulatory molecule and provides, in addition to the first signal received by the T cell receptor, a second signal to elicit proliferation and differentiation into effector cytotoxic T lymphocytes (CTL) [8, 9]. NKG2D has been demonstrated to be important for the elimination of tumor cells [10] and for defense against pathogens [11, 12].

The NKG2D receptor recognizes several ligands (Figure 1A) including the major histocompatibility complex (MHC) class I chain-related molecules A (MICA) and B (MICB). The *MICA* and *MICB* genes are located within the human leukocyte antigen (HLA) complex in close proximity to *HLA-B* [13, 14]. The protein structure of MICA and MICB is similar to classical class I molecules with three extracellular domains (α1, α2, and α3), a transmembrane segment, and a carboxy-terminal cytoplasmic tail. However, in contrast to classical MHC class I molecules, MICA and MICB are not associated with β2-microglobulin and do not present peptides. NKG2D binds also to a second family of ligands, the UL16-binding proteins (ULBP) encoded by the *RAET1* gene family, which is also localized on chromosome 6 but outside the HLA complex [15]. Six loci encode functional proteins, i.e. *RAET1I* (ULBP1), *RAET1H* (ULBP2), *RAET1N* (ULBP3), *RAET1E* (ULBP4), *RAET1G* (ULBP5), and *RAET1L* (ULBP6) [16].

NKG2D ligands are constitutively expressed only on a few cell types [17]. They can be expressed during embryonic development and on pluripotent stem cells [18] but in general, the proteins are absent on healthy adult cells. However,

almost all cells exposed to cellular or genotoxic stress can express NKG2D ligands [17, 19]. Therefore, NKG2D ligands are frequently found on malignant or virally infected cells and mark these cells for recognition by NK cells and other lymphocytes expressing NKG2D [20]. Several mechanisms have been described that permit tumors or virally infected cells to escape the NKG2D-mediated immune surveillance. One mechanism is that MICA and other NKG2D ligands can undergo a proteolytic shedding that results in an immune escape mediated not only by the loss of ligands on target cells but in addition by the generation of immunosuppressive soluble NKG2D ligands [16, 21, 22]. Several clinical studies showed an association between tumor-associated or soluble NKG2D ligands and disease progression or poor prognosis in different malignant diseases [23].

The diversity of NKG2D ligands is further increased by genetic polymorphisms (Figure 1B). *MICA* is the most polymorphic non-classical class I gene and currently 100 alleles are known that encode for 79 protein variants (<http://www.ebi.ac.uk/imgt/hla/>, release 3.17.0). *MICB* is also very polymorphic with 40 alleles encoding 26 protein variants. The *RAET1* gene cluster is less polymorphic and the variation is found mainly within the *RAET1E* (ULBP4), *RAET1L* (ULBP6), and *RAET1N* (ULBP3) genes [24, 25]. The importance particularly of *MICA* polymorphisms for cancer has been widely studied [26, 27].

A single nucleotide polymorphism (SNP) (rs1051792) at nucleotide position 454 (G/A) of *MICA* causing a valine (Val) to methionine (Met) exchange at amino acid position 129 in the α2 domain of the protein has gained specific interest because the *MICA* alleles can be separated into two groups with respect to this polymorphism. *MICA* variants containing a methionine at position 129 bind NKG2D with high avidity, whereas those with a valine bind NKG2D with low avidity [28]. The potential relevance of this difference is highlighted by several disease associations described for this SNP. The *MICA*-129 dimorphism has been associated with the risks for nasopharyngeal carcinoma [29] and hepatitis B virus-induced hepatocellular carcinoma [30]. Moreover, it has been associated with several autoimmune diseases, including ankylosing spondylitis [31], rheumatoid arthritis [32], inflammatory bowel disease [33, 34], lupus erythematosus [35], type I diabetes [36], and psoriatic disease [37]. Recently, an association with the severity of chronic Chagas heart disease has been described [38]. Notably, this SNP has also been associated with changes in the serum level of soluble MICA (sMICA). In patients with ulcerative colitis, carriers of the *MICA*-129Val/Val genotype had higher sMICA serum levels [34]. The *MICA*-129Met allele on the other hand has been associated with lower sMICA serum levels in healthy




Figure 1. Summary of the effects of variation in the NKG2D receptor and NKG2D ligand pathway. (A) The NKG2D receptor on lymphocytes can interact with several NKG2D ligands on target cells, which may differ in their efficacy to elicit NKG2D signaling. (B) NKG2D ligands are polymorphic and isoforms of the same ligand, e.g. MICA, vary in their cell surface expression and their capacity to elicit NKG2D signaling. (C) NKG2D vary in the intensity of cell surface expression due to genetic polymorphisms. The functional consequences of polymorphisms in NKG2D and NKG2D ligands may be cooperative or counteracting. The interaction of the variants could be highly important for the outcome of NKG2D signaling and disease associations of the NKG2D signaling pathway.

controls and hepatitis B virus-induced hepatocellular carcinoma patients^[30].

We became interested in the MICA-129 dimorphism in the context of hematopoietic stem cell transplantation (HSCT). HSCT is a potentially curative therapy for several hematological diseases. However, the success is limited by post-transplant complications including graft versus host disease (GVHD), relapse of malignancy, and infections^[39]. The MICA-129 dimorphism has been associated with the incidence of chronic GVHD and relapse of malignancy after HSCT^[40]. However, the finding that high avidity MICA-129Met variants were associated with an increased risk of relapse, whereas the low avidity MICA-129Val variants were associated with an increased risk of chronic GVHD^[40] appeared to be counterintuitive in view of the functions of NKG2D. Thus, we analyzed another cohort of 452 patients undergoing HSCT and found that the *MICA-129Met* allele was associated with an improved survival and a reduced risk to die from acute GVHD. Nonetheless, carriers of the *MICA-129Met/Met* genotype had an increased risk to experience acute GVHD^[41].

To better understand the associations of the MICA-129 dimorphism with the outcomes of HSCT, we determined functional differences of the MICA-129Met and MICA-129Val isoforms^[41, 42]. Binding of the MICA-129Met isoform to NKG2D stimulated a stronger phosphorylation of

SRC family kinases in NK cells than binding of the MICA-129Val isoform. Subsequently, the MICA-129Met ligand triggered more degranulation and IFN- γ production of NK cells than the MICA-129Val ligand. Notably, the extent of degranulation and IFN- γ secretion correlated clearly with the MICA expression intensity on target cells for the MICA-129Val isoform. The expression intensity of the MICA-129Met isoform, in contrast, had either none or even a negative effect on the extent of degranulation, target cell killing, and IFN- γ release. On CD8⁺ T cells, the MICA-129Met isoform induced an earlier co-stimulatory activation than the MICA-129Val isoform. Importantly, the MICA-129Met ligand induced also a stronger down-regulation of NKG2D on both NK and CD8⁺ T cells than the MICA-129Val ligand. This down-regulation of NKG2D impaired the capability of NK and CD8⁺ T cells to receive signals via NKG2D. Thus, MICA-129Met ligands, which elicit strong NKG2D responses, stimulate in parallel a robust negative feedback signal by down-regulation of NKG2D and this appears to limit the initially stronger effects of MICA-129Met ligands^[41]. In target cells, more of the MICA-129Met isoform was retained in intracellular compartments and when transported to the cell surface, it was more susceptible to shedding than the MICA-129Val isoform^[42]. Both processes appear to limit the expression of the high avidity MICA-129Met isoform at the plasma membrane.

These results show how the MICA-129 dimorphism affects the function of NK cells and CTL. In HSCT recipients, carriers of two *MICA-129Met* alleles had an increased risk to experience acute GVHD, which could be the result of rapid and strong effects of the MICA-129Met isoform on NKG2D signaling. A faster co-stimulatory activation of CD8⁺ T cells by the MICA-129Met isoform could be essential for this difference. On the other site, having at least one *MICA-129Met* allele conferred a lower probability of death due to acute GVHD. This finding is explainable by a rapid down-regulation of NKG2D on allo-reactive CD8⁺ T cells mediated by engagement of the high avidity MICA-129Met isoform, which limits the NKG2D-mediated co-stimulation of allo-reactive donor CD8⁺ T cells. In heterozygous recipients, also the risk of occurrence of acute GVHD was reduced suggesting that the effect of the MICA-129Met isoform on the cell surface expression of NKG2D was decisive for this outcome. Consequently, we found an increase in survival after HSCT for recipients carrying a *MICA-129Met* allele. Consistently, recipients having two *MICA-129Val* alleles were at hazard to develop a fatal acute GVHD and they appeared to mainly profit from treatment with anti-thymocyte globulin (ATG), which depletes T cells. This is explainable by a failure to efficiently down-regulate NKG2D on allo-reactive CD8⁺ T cells when having only low avidity MICA-129Val variants. Therefore, this finding is of potential therapeutic relevance for recipients having two *MICA-129Val* alleles. These patients may specifically profit from ATG or other T cell depleting therapies, which are used in some treatment protocols to reduce the risk of acute GVHD.

It has previously been reported that the risk of chronic GVHD was increased for patients having the *MICA-129Val/Val* genotype, whereas the *MICA-129Met/Met* genotype was associated with an increased risk of relapse^[40]. These associations, although not found in our cohort^[41], are also explainable by functional effects of the MICA variants on NKG2D. Sustained NKG2D-mediated activation of allo-reactive CD8⁺ T cells is likely if only a MICA-129Val isoform is present in a patient and this could increase the risk of chronic GVHD. On the other hand, sustained NKG2D-mediated activation of anti-leukemic CD8⁺ T cells and NK cells would be predicted to reduce the risk of relapse.

We demonstrated that the MICA-129Met isoform triggers more NKG2D signaling at low expression intensities. The MICA-129Val isoform, in contrast, produces more NKG2D effects at high expression intensity, at which the MICA-129Met isoform already down-regulates NKG2D leading to a diminished function^[41]. Since MICA expression intensities can vary for certain *MICA* alleles^[43], a functional

interaction of several SNPs within the *MICA* gene can be postulated. The principal relevance of NKG2D signaling for the outcome of allogeneic HSCT has been further demonstrated in mouse models, which developed less GVHD when receiving NKG2D-deficient T cells after transplantation^[44]. The importance of the pathway is also emphasized by human studies demonstrating effects of the genotypes of *RAET1L*^[45] encoding the NKG2D ligand ULBP6 and the NKG2D-encoding gene *KLRK1*^[46] on overall survival of patients after HSCT. Therefore, also gene-gene interactions within the NKG2D signaling pathway presumably affect the outcome of HSCT and other diseases, which have been described to be associated with SNPs in NKG2D ligands or NKG2D itself.

The *KLRK1* gene has a limited degree of variation with only one SNP leading to an amino acid substitution (rs2255336, A>G, Thr72Ala). Despite this limited variation at the protein level, *KLRK1* haplotype alleles constructed from five or three SNPs have been associated with low and high NK cell cytotoxicity and increased or decreased risk of cancer development^[47] and an interaction of the *KLRK1* genotype and lifestyle risk factors for cancer was observed^[48]. It was the *NKG2D-HNK1* haplotype, a haplotype expected to induce greater NK cell activity, that has been associated with significantly improved overall survival after HSCT^[46].

SNPs within the *KLRK1* gene have been associated also with the risk of cholangiocarcinoma in patients with primary sclerosing cholangitis^[49]. The rs2255336 G genotype encoding the NKG2D-72Ala variant was found to be associated with an increased risk of cervical carcinoma and progression to advanced stages of the disease^[50]. For homozygous carriers of this variant, an increased risk for systemic lupus erythematosus (SLE) was observed^[51]. In accordance with these data, presence of the NKG2D-72Thr variant appeared to confer protection against SLE^[52]. The G/G genotype was also associated with an increased risk of early symptomatic infection in cases with congenital cytomegalovirus infection^[53]. The SNP rs2617160 T/T genotype was associated with susceptibility to chronic hepatitis B^[54].

The functional consequences of the *KLRK1* polymorphisms are only partly understood. Notably, the *KLRK1* haplotypes associated with low and high NK cell cytotoxicity^[47], respectively, were described to be associated with low and high expression intensity of NKG2D on NK and CD8⁺ T cells^[55] (Figure 1C). In homozygous carriers of the NKG2D-72Thr variant, a trend was observed towards a higher proliferation in response to stimulation with antibodies against CD3 and NKG2D suggesting a functional

effect also of this polymorphism^[51].

In conclusion, we have shown that the MICA-129 dimorphism affects the strength and kinetics of NKG2D signaling as well as MICA cell surface expression and shedding^[41, 42]. This results in differences in the NK cell cytotoxicity and cytokine secretion as well as CD8⁺ T cell co-stimulation and it affects the NKG2D expression on NK and CD8⁺ T cells by a counter regulatory mechanism. The MICA-129 dimorphism has a significant impact on autoimmune diseases, infections, malignancies and the outcome of HCST. In the future, it might be worthwhile to investigate the interaction of gene variants in the NKG2D signaling pathway. Specific combinations of polymorphisms in NKG2D and NKG2D ligands or downstream signaling molecules may have a higher impact on the outcome of associated diseases than single SNPs in one of the respective genes.

Conflicting interests

The authors have declared that no conflict of interests exists.

Acknowledgments

The work of the authors was supported by the Deutsche Forschungsgemeinschaft (GRK 1034 and SFB 1002 TP C05), and the European Union grant FP7-PEOPLE-2012-ITN-315963 (CELLEUROPE).

Author contributions

RD drafted the manuscript, which AI, DM, SM, DS, PS, GM, GW, DK, and HB edited. All authors approved the final manuscript.

Abbreviations

ATG: anti-thymocyte globulin; CTL: cytotoxic T lymphocytes; DAP10: DNAX-activation protein 10; GVHD: graft versus host disease; HLA: human leukocyte antigen; HSCT: hematopoietic stem cell transplantation; IFN: interferon; IL: interleukin; MHC: major histocompatibility complex; MICA: MHC class I chain-related molecules A; NK: natural killer; NKG2D: NK group 2, member D; NKG2D-L: NKG2D ligand; Smica: soluble MICA; SNP: single nucleotide polymorphism; TGF: transforming growth factor; ULBP: UL16-binding protein.

References

1. Glienke J, Sobanov Y, Brostjan C, Steffens C, Nguyen C, Lehrach H, et al. The genomic organization of NKG2C, E, F, and D

receptor genes in the human natural killer gene complex. *Immunogenetics* 1998; 48:163-173.

2. Lanier LL. NKG2D receptor and its ligands in host defense. *Cancer Immunol Res* 2015; 3:575-582.
3. Karimi MA, Aguilar OA, Zou B, Bachmann MH, Carlyle JR, Baldwin CL, et al. A truncated human NKG2D splice isoform negatively regulates NKG2D-mediated function. *J Immunol* 2014; 193:2764-2771.
4. Raulat DH, Gasser S, Gowen BG, Deng W, Jung H. Regulation of ligands for the NKG2D activating receptor. *Annu Rev Immunol* 2013; 31:413-441.
5. Zafirova B, Wensveen FM, Gulin M, Polic B. Regulation of immune cell function and differentiation by the NKG2D receptor. *Cell Mol Life Sci* 2011; 68:3519-3529.
6. Billadeau DD, Upshaw JL, Schoon RA, Dick CJ, Leibson PJ. NKG2D-DAP10 triggers human NK cell-mediated killing via a Syk-independent regulatory pathway. *Nat Immunol* 2003; 4:557-564.
7. Andre P, Castriconi R, Espeli M, Anfossi N, Juarez T, Hue S, et al. Comparative analysis of human NK cell activation induced by NKG2D and natural cytotoxicity receptors. *Eur J Immunol* 2004; 34:961-971.
8. Groh V, Rhinehart R, Randolph-Habecker J, Topp MS, Riddell SR, Spies T. Costimulation of CD8 α T cells by NKG2D via engagement by MIC induced on virus-infected cells. *Nat Immunol* 2001; 2:255-260.
9. Spear P, Wu MR, Sentman ML, Sentman CL. NKG2D ligands as therapeutic targets. *Cancer Immun* 2013; 13:8.
10. Guerra N, Tan YX, Joncker NT, Choy A, Gallardo F, Xiong N, et al. NKG2D-deficient mice are defective in tumor surveillance in models of spontaneous malignancy. *Immunity* 2008; 28:571-580.
11. Wesselkamper SC, Eppert BL, Motz GT, Lau GW, Hassett DJ, Borchers MT. NKG2D is critical for NK cell activation in host defense against *Pseudomonas aeruginosa* respiratory infection. *J Immunol* 2008; 181:5481-5489.
12. Fang M, Lanier LL, Sigal LJ. A role for NKG2D in NK cell-mediated resistance to poxvirus disease. *PLoS Pathog* 2008; 4:e30.
13. Bahram S, Bresnahan M, Geraghty DE, Spies T. A second lineage of mammalian major histocompatibility complex class I genes. *Proc Natl Acad Sci U S A* 1994; 91:6259-6263.
14. Leelayuwat C, Townend DC, Degli-Esposti MA, Abraham LJ, Dawkins RL. A new polymorphic and multicopy MHC gene family related to nonmammalian class I. *Immunogenetics* 1994; 40:339-351.
15. Radosavljevic M, Cuillerier B, Wilson MJ, Clement O, Wicker S, Gilfillan S, et al. A cluster of ten novel MHC class I related genes on human chromosome 6q24.2-q25.3. *Genomics* 2002; 79:114-123.
16. Chitadze G, Bhat J, Lettau M, Janssen O, Kabelitz D. Generation of Soluble NKG2D Ligands: Proteolytic Cleavage, Exosome Secretion and Functional Implications. *Scand J Immunol* 2013; 78:120-129.
17. Groh V, Bahram S, Bauer S, Herman A, Beauchamp M, Spies T. Cell stress-regulated human major histocompatibility complex class I gene expressed in gastrointestinal epithelium. *Proc Natl*

- Acad Sci U S A 1996; 93:12445-12450.
18. Kruse V, Hamann C, Monecke S, Cyganek L, Elsner L, Hübscher D, *et al.* Human induced pluripotent stem cells are targets for allogeneic and autologous natural killer (NK) cells and killing is partly mediated by the activating NK receptor DNAM-1. *PLoS One* 2015; 10:e0125544.
 19. Gasser S, Orsulic S, Brown EJ, Raulet DH. The DNA damage pathway regulates innate immune system ligands of the NKG2D receptor. *Nature* 2005; 436:1186-1190.
 20. Ullrich E, Koch J, Cerwenka A, Steinle A. New prospects on the NKG2D/NKG2DL system for oncology. *Oncoimmunology* 2013; 2:e26097.
 21. Groh V, Wu J, Yee C, Spies T. Tumour-derived soluble MIC ligands impair expression of NKG2D and T-cell activation. *Nature* 2002; 419:734-738.
 22. Salih HR, Rammensee HG, Steinle A. Cutting edge: down-regulation of MICA on human tumors by proteolytic shedding. *J Immunol* 2002; 169:4098-4102.
 23. El-Gazzar A, Groh V, Spies T. Immunobiology and conflicting roles of the human NKG2D lymphocyte receptor and its ligands in cancer. *J Immunol* 2013; 191:1509-1515.
 24. Antoun A, Jobson S, Cook M, O'Callaghan CA, Moss P, Briggs DC. Single nucleotide polymorphism analysis of the NKG2D ligand cluster on the long arm of chromosome 6: Extensive polymorphisms and evidence of diversity between human populations. *Hum Immunol* 2010; 71:610-620.
 25. Cox ST, Arrieta-Bolanos E, Poeso S, Vullo C, Madrigal JA, Saudemont A. RAET1/ULBP alleles and haplotypes among Kolla South American Indians. *Hum Immunol* 2013; 74:775-782.
 26. Chen D, Gyllenstein U. MICA polymorphism: biology and importance in cancer. *Carcinogenesis* 2014; 35:2633-2642.
 27. Goto K, Kato N. MICA SNPs and the NKG2D system in virus-induced HCC. *J Gastroenterol* 2015; 50:261-272.
 28. Steinle A, Li P, Morris DL, Groh V, Lanier LL, Strong RK, *et al.* Interactions of human NKG2D with its ligands MICA, MICB, and homologs of the mouse RAE-1 protein family. *Immunogenetics* 2001; 53:279-287.
 29. Douik H, Ben Chaaben A, Attia Romdhane N, Romdhane HB, Mamoghli T, Fortier C, *et al.* Association of MICA-129 polymorphism with nasopharyngeal cancer risk in a Tunisian population. *Hum Immunol* 2009; 70:45-48.
 30. Tong HV, Toan NL, Song LH, Bock CT, Kremsner PG, Velavan TP. Hepatitis B virus-induced hepatocellular carcinoma: functional roles of MICA variants. *J Viral Hepat* 2013; 20:687-698.
 31. Amroun H, Djoudi H, Busson M, Allat R, El Sherbini SM, Sloma I, *et al.* Early-onset ankylosing spondylitis is associated with a functional MICA polymorphism. *Hum Immunol* 2005; 66:1057-1061.
 32. Kirsten H, Petit-Teixeira E, Scholz M, Hasenclever D, Hantmann H, Heider D, *et al.* Association of MICA with rheumatoid arthritis independent of known HLA-DRB1 risk alleles in a family-based and a case control study. *Arthritis Res Ther* 2009; 11:R60.
 33. Lopez-Hernandez R, Valdes M, Lucas D, Campillo JA, Martinez-Garcia P, Salama H, *et al.* Association analysis of MICA gene polymorphism and MICA-129 dimorphism with inflammatory bowel disease susceptibility in a Spanish population. *Hum Immunol* 2010; 71:512-514.
 34. Zhao J, Jiang Y, Lei Y, Zou K, Wang C, Huang S, *et al.* Functional MICA-129 polymorphism and serum levels of soluble MICA are correlated with ulcerative colitis in Chinese patients. *J Gastroenterol Hepatol* 2011; 26:593-598.
 35. Yoshida K, Komai K, Shiozawa K, Mashida A, Horiuchi T, Tanaka Y, *et al.* Role of the MICA polymorphism in systemic lupus erythematosus. *Arthritis Rheum* 2011; 63:3058-3066.
 36. Raache R, Belanteur K, Amroun H, Benyahia A, Heniche A, Azzouz M, *et al.* Association of major histocompatibility complex class I chain-related gene a dimorphism with type 1 diabetes and latent autoimmune diabetes in adults in the Algerian population. *Clin Vaccine Immunol* 2012; 19:557-561.
 37. Pollock RA, Chandran V, Pellett FJ, Thavaneswaran A, Eder L, Barrett J, *et al.* The functional MICA-129 polymorphism is associated with skin but not joint manifestations of psoriatic disease independently of HLA-B and HLA-C. *Tissue Antigens* 2013; 82:43-47.
 38. Ayo CM, Oliveira AP, Camargo AV, Mattos CC, Bestetti RB, Mattos LC. Association of the functional MICA-129 polymorphism with the severity of chronic Chagas heart disease. *Clin Infect Dis* 2015; 61:1310-1313.
 39. Dickinson AM. Non-HLA genetics and predicting outcome in HSCT. *Int J Immunogenet* 2008; 35:375-380.
 40. Boukouaci W, Busson M, Peffault de Latour R, Rocha V, Suberbielle C, Bengoufa D, *et al.* MICA-129 genotype, soluble MICA, and anti-MICA antibodies as biomarkers of chronic graft-versus-host disease. *Blood* 2009; 114:5216-5224.
 41. Isernhagen A, Malzahn D, Viktorova E, Elsner L, Monecke S, von Bonin F, *et al.* The MICA-129 dimorphism affects NKG2D signaling and outcome of hematopoietic stem cell transplantation. *EMBO Mol Med* 2015; 7:1480-1502.
 42. Isernhagen A, Schilling D, Monecke S, Shah P, Elsner L, Walter L, *et al.* The MICA-129Met/Val dimorphism affects plasma membrane expression and shedding of the NKG2D ligand MICA. *Immunogenetics* 2016; 68:109-123.
 43. Shafi S, Vantourout P, Wallace G, Antoun A, Vaughan R, Stanford M, *et al.* An NKG2D-mediated human lymphoid stress surveillance response with high interindividual variation. *Sci Transl Med* 2011; 3:113ra124.
 44. Karimi MA, Bryson JL, Richman LP, Fesnak AD, Lechner TM, Satake A, *et al.* NKG2D expression by CD8+ T cells contributes to GVHD and GVT effects in a murine model of allogeneic HSCT. *Blood* 2015; 125:3655-3663.
 45. Antoun A, Vekaria D, Salama RA, Pratt G, Jobson S, Cook M, *et al.* The genotype of RAET1L (ULBP6), a ligand for human NKG2D (KLRK1), markedly influences the clinical outcome of allogeneic stem cell transplantation. *Br J Haematol* 2012; 159:589-598.
 46. Espinoza JL, Takami A, Onizuka M, Sao H, Akiyama H, Miyamura K, *et al.* NKG2D gene polymorphism has a significant impact on transplant outcomes after HLA-fully-matched unrelated bone marrow transplantation for standard risk hematologic malignancies. *Haematologica* 2009; 94:1427-1434.
 47. Hayashi T, Imai K, Morishita Y, Hayashi I, Kusunoki Y, Nakachi

- K. Identification of the NKG2D haplotypes associated with natural cytotoxic activity of peripheral blood lymphocytes and cancer immunosurveillance. *Cancer Res* 2006; 66:563-570.
48. Furue H, Kumimoto H, Matsuo K, Suzuki T, Hasegawa Y, Shinoda M, *et al.* Opposite impact of NKG2D genotype by lifestyle exposure to risk of aerodigestive tract cancer among Japanese. *Int J Cancer* 2008; 123:181-186.
49. Melum E, Karlsen TH, Schrumpf E, Bergquist A, Thorsby E, Boberg KM, *et al.* Cholangiocarcinoma in primary sclerosing cholangitis is associated with NKG2D polymorphisms. *Hepatology* 2008; 47:90-96.
50. Roszak A, Lianeri M, Jagodzinski PP. Prevalence of the NKG2D Thr72Ala polymorphism in patients with cervical carcinoma. *Genet Test Mol Biomarkers* 2012; 16:841-845.
51. Kabalak G, Thomas RM, Martin J, Ortego-Centeno N, Jimenez-Alonso J, de Ramon E, *et al.* Association of an NKG2D gene variant with systemic lupus erythematosus in two populations. *Hum Immunol* 2010; 71:74-78.
52. Piotrowski P, Lianeri M, Olesinska M, Jagodzinski PP. Prevalence of the NKG2D Thr72Ala polymorphism in patients with systemic lupus erythematosus. *Mol Biol Rep* 2012; 39:1343-1347.
53. Taniguchi R, Koyano S, Suzutani T, Goishi K, Ito Y, Morioka I, *et al.* A Thr72Ala polymorphism in the NKG2D gene is associated with early symptomatic congenital cytomegalovirus disease. *Infection* 2015; 43:353-359.
54. Ma J, Guo X, Wu X, Li J, Zhu X, Li Z, *et al.* Association of NKG2D genetic polymorphism with susceptibility to chronic hepatitis B in a Han Chinese population. *J Med Virol* 2010; 82:1501-1507.
55. Imai K, Hayashi T, Yamaoka M, Kajimura J, Yoshida K, Kusunoki Y, *et al.* Effects of NKG2D haplotypes on the cell-surface expression of NKG2D protein on natural killer and CD8 T cells of peripheral blood among atomic-bomb survivors. *Hum Immunol* 2012; 73:686-691.