

## **Thioredoxin reductase 1 suppresses adipocyte differentiation and insulin responsiveness**

Xiaoxiao Peng<sup>1§</sup>, Alfredo Giménez-Cassina<sup>1</sup>, Paul Petrus<sup>2</sup>, Marcus Conrad<sup>3</sup>, Mikael Rydén<sup>2</sup> and Elias S.J. Arnér<sup>\*,1</sup>



<sup>1</sup> Division of Biochemistry, Department of Medical Biochemistry and Biophysics, Karolinska Institutet, SE-171 77 Stockholm, Sweden;

<sup>2</sup> Clinical Research Center, and the Department of Medicine, Huddinge University Hospital, Karolinska Institutet, SE-141 86 Stockholm, Sweden;


<sup>3</sup> Helmholtz Zentrum München, Institute of Developmental Genetics, Ingolstädter Landstrasse 1, 85764 Neuherberg, Germany

<sup>§</sup> Present address: CVMD Translational Medicine Unit, Early Clinical Development, AstraZeneca, Mölndal, Sweden.


**Supplementary material:** Original non-cropped images of western blots



**Non-cropped blots for the experiments shown in Fig. 1D of the main article.**




**Non-cropped blot for the experiment shown in Fig. 3B of the main article.**




**Non-cropped blots for the experiments shown in Fig. 3C of the main article.**





Non-cropped blot for the experiment shown in Fig. 5F of the main article.



**Non-cropped blots for the experiments shown in Fig. 5H of the main article.**