

Growth inhibition, cell-cycle arrest and apoptosis in human T-cell leukemia by the isothiocyanate sulforaphane

Carmela Fimognari¹, Michael Nüsse², Rossano Cesari¹, Renato Iori³, Giorgio Cantelli-Forti¹ and Patrizia Hrelia^{1,4}

¹Dipartimento di Farmacologia, Università di Bologna, via Imerio 48, 40126 Bologna, Italy, ²GSF-Flow Cytometry Group, Neuherberg, Germany and

³Istituto Sperimentale per le Colture Industriali, MiPA, Bologna, Italy

⁴To whom correspondence should be addressed

Email: phrelia@biocfarm.unibo.it

Glucosinolates (GL) can inhibit, retard or reverse experimental multistage carcinogenesis. When brassica plant tissue is broken, GLs are hydrolyzed by the endogenous enzyme myrosinase (Myr), releasing many products including isothiocyanates (ITC). Synthetic ITCs like sulforaphane exert chemopreventive effects against chemically induced tumors in animals, modulating enzymes required for carcinogens' activation/detoxification and/or the induction of cell-cycle arrest and apoptosis in tumor cell lines. To investigate the chemopreventive potential of ITCs while reproducing the circumstances of dietary contact with sulforaphane, we studied proliferation, apoptosis induction and p53, bcl-2 and bax protein expression in Jurkat T-leukemia cells by sulforaphane, the ITC generated *in situ* in a quantitative manner by Myr starting from glucoraphanin (GRA). Jurkat cells were treated with different doses of GRA–Myr mixture. Effects on cell growth or survival were evaluated by counting trypan blue-excluding cells. Cell-cycle progression, apoptosis and expression of p53, bax and bcl-2 proteins were analyzed by flow cytometry. Results were analyzed by two-sided Fisher's exact test. Sulforaphane, but not GRA, caused G₂/M-phase arrest ($P = 0.028$) and increase of apoptotic cell fraction ($P < 0.0001$) in a time- and dose-dependent manner. Necrosis was observed after prolonged exposure to elevated sulforaphane doses. Moreover, it markedly increased p53 and bax protein expression, and slightly affected bcl-2 expression. These findings indicate that sulforaphane but not the native GL GRA can exert both protective and toxic effects inhibiting leukemic cell growth. Sulforaphane therefore deserves study as a potential chemopreventive/chemotherapeutic antileukemic agent.

Introduction

Cancer chemoprevention involves preventing, delaying or reversing carcinogenesis by administering non-toxic synthetic chemicals or chemicals from natural substances before malignancy (1). A number of agents have been reported to be effective against chemical carcinogenesis in several organs, and some are candidates for the inhibition of human cancer development (2–4). The multifactorial nature of the carcino-

genic process provides numerous pathways that could be manipulated by these agents in order to inhibit or delay tumor development (5). As a result of the wide variety of possible chemopreventive agents, short-term screening assays are being developed to help identify potentially effective compounds (6).

Edible brassicas (e.g. broccoli and cauliflower) contain substantial quantities of biosynthesized secondary metabolites known as glucosinolates (GLs), as well as myrosinase (Myr) (thioglucoside glucohydrolase; EC 3.2.3.1). When the vegetables are crushed by chopping or chewing, Myr catalyzes a rapid hydrolytic reaction in which the GLs are then converted into isothiocyanates (ITCs) and other products, depending on the chemical structure of GL side chains (7). Many synthetic ITCs exert versatile chemopreventive effects against tumors induced by chemical carcinogens in various animal organs, suggesting that common mechanisms underlie their chemopreventive properties (8,9). Two sets of putative mechanisms have been identified that help to explain the effects of ITCs. The first involves modulation of enzymes that are required for the activation or detoxification of many carcinogens (10,11). In particular, ITCs have been shown to induce the activity of phase II enzymes (e.g. glutathione *S*-transferase, quinone reductase and glucuronosyltransferases) and/or inhibit phase I enzymes (12–14). Recent studies in humans also revealed that a high consumption of brassicas led to an increase in glutathione *S*-transferase activity, supporting the potential role of ITCs in the prevention of human cancer (15). The second set of putative mechanisms involves suppression of tumor development following the initiation of pre-cancerous cells (16). Possible mechanisms of suppression include the deletion of initiated cells from genetically damaged tissue by apoptosis, so that clonal expansion of the lesion is aborted (17). Some constituents of brassicas have already been shown to induce apoptosis or block proliferation in cancer cells *in vitro*. In particular, Huang *et al.* (18) showed that phenylethyl ITC induces p53 transactivation in mouse epidermal cells in a dose- and time-dependent fashion, while Gamet-Payraastre *et al.* demonstrated that sulforaphane induces cell-cycle arrest and apoptosis in HT29 human colon cancer cells (19).

Most of these studies were performed using synthetic ITCs, which are commonly assumed to correspond to major natural constituents of cruciferous vegetables. However, this is not literally so, as sulforaphane only appears as the product of enzymatic degradation of its naturally occurring parent GL, namely glucoraphanin (GRA). Therefore, dietary exposure to sulforaphane, and indeed other ITCs, actually occurs as a result of the hydrolytic breakdown of precursor GLs on ingestion (e.g. by chewing raw vegetables in salads). Although cooking degrades the Myr present in broccoli, sulforaphane may also be produced in the human intestine, where microflora seem to promote the hydrolysis of GLs into ITCs (20).

In order to mimic more closely the circumstances of dietary contact with sulforaphane following consumption of brassicas and to expand our knowledge on the chemopreventive potential

Abbreviations: EB, ethidium bromide; FITC, fluorescein isothiocyanate; FSC, forward scatter; GL, glucosinolate; GRA, glucoraphanin; GSH, Glutathione; ITC, isothiocyanate; Myr, myrosinase; PI, propidium iodide; SSC, side scatter.

of ITCs, we decided to study the effects of the mixture of GRA and Myr to produce sulforaphane on cell growth, cell-cycle progression and apoptosis induction in Jurkat cells (i.e. human T-cell leukemia). We also assessed the apoptotic response of Jurkat cells in terms of p53, bcl-2 and bax expression.

Materials and methods

GRA was obtained by a recently developed procedure (21) whereby a quantitative yield is produced starting from glucorucin. Glucorucin was isolated with a high purity grade from the ripe seeds of rocket (*Eruca sativa* Miller), according to a well-defined protocol (22). The transformation of glucorucin into GRA is based on the oxidation reaction of sulfides into their corresponding sulfoxides. The semi-synthetic GRA produced was purified according to the method reported by Visentin *et al.* (22). Purity was assessed by HPLC analysis of the desulfo-derivative according to the ISO 9167-1 method (23). The Myr used in the present study was isolated from ripe seeds of white mustards (*Sinapis alba* L.), as reported by Pessina *et al.* (24). The Myr stock solution had a specific activity of ~60 U/mg of soluble protein. One Myr unit was defined as the amount of enzyme able to hydrolyze 1 μ mol of sinigrin/min at pH 6.5 and 37°C. Myr solution was stored at 4°C in sterile distilled water until use.

Jurkat T-leukemia cells were grown in suspension and propagated in RPMI 1640 supplemented with 10% heat-inactivated bovine serum, 1% antibiotics (all obtained from Sigma, St Louis, MO). To maintain exponential growth, the cultures were divided every third day by dilution to a concentration of 1×10^5 cells/ml.

A 4 mM stock solution of GRA was prepared in 0.9% NaCl and stored at 4°C. Samples (0.23 U) of Myr were directly added to 5 ml of the complete cell-culture medium containing increasing concentrations of GRA (3, 10 and 30 μ M), prepared by diluting the stock solution with 0.9% NaCl. Exponentially growing Jurkat cells at a concentration of 400 000 cells/ml were subsequently added. The activity of native GRA or Myr was tested by treating the cell cultures with 30 μ M GRA in absence of the enzyme or with Myr in absence of GRA, respectively. NaCl (0.9%) was used as control. For each experiment, treatments were done in duplicate and separate cultures were set up for each treatment.

In separate experiments, the GRA–Myr mixture was added at various concentrations to exponentially growing cells. Cell concentrations were measured as a function of time by counting trypan blue-excluding cells on cell aliquots removed from culture at the designated times (2, 4, 8, 12, 24 and 48 h). Results were calculated as viable cells in GRA–Myr mixture-treated cultures relative to control. ID₅₀, the drug concentration causing cell toxicity by 50%

Flow cytometry was performed using a FACStar⁺ flow cytometer (Becton Dickinson, Sunnyvale, CA) equipped with an argon laser (Innova 90; Coherent Radiation, Palo Alto, CA) operating at 488 nm (500 mW) for excitation of the ethidium bromide (EB).

The preparation of samples for measurements of the cell-cycle distribution of nuclei by DNA content was performed according to a two-step method reported elsewhere (27,28). Briefly, cultures were centrifuged for 5 min at 800 *g* and treated with 1 ml of solution I (584 mg/l NaCl, 1000 mg/l Na-citrate, 25 mg/l EB, 10 mg/l RNase and 0.3 ml/l Nonidet P-40). Then, after ~1 h, 1 ml of solution II (15 g/l citric acid, 0.25 M sucrose and 40 mg/l EB) was added and the samples were briefly vortexed. EB fluorescence, forward scatter (FSC) and side scatter (SSC) of nuclei in suspension, measured using a combination of KV550 and OG590 longpass filters (Schott, Mainz, Germany), were recorded for cell nuclei in list mode. For each sample, 10 000 events were registered. The fraction of cells in the different compartments of cell cycle was calculated as described by Schreiber *et al.* (29).

It has been shown that loss of phospholipid asymmetry of the plasma membrane is an early event of apoptosis (30,31). The annexin V binds to negatively charged phospholipids, like phosphatidylserine. During apoptosis, the cells react to annexin V once chromatin condenses but before the plasma membrane loses its ability to exclude propidium iodide (PI). Hence, by staining cells with a combination of fluorescein isothiocyanate (FITC) annexin V and PI it is possible to detect non-apoptotic live cells, early apoptotic cells and late apoptotic or necrotic cells (32,33).

Determination by flow cytometry of different proteins involved in the apoptotic process offers rapid and objective quantification of protein expression (34–37). Under the same exposure conditions described above, 1×10^6 cells were fixed and permeabilized by a commercially available kit, namely Leucoperm solution A and B (Serotec, Oxford, UK). They were then incubated with 10 μ l of each antibody, i.e. FITC p53 (35 μ g/ml; Novocastra, Newcastle, UK) against phosphorylated and non-phosphorylated protein, FITC bcl-2 (200 μ g/ml; Serotec) and bax (200 μ g/ml; Santa Cruz Biotechnology, Santa Cruz, CA) or isotype-matched negative control (Serotec). The cells were washed and incubated with 10 μ l of the FITC-labeled secondary antibody for the bax-stained cells (5 μ g/ml; Serotec). Finally, the cells were washed and resuspended in 0.2 ml of 1% paraformaldehyde. The cells were then analyzed to quantify FITC binding by flow cytometry. From each sample 10 000 cells were analyzed and non-specific binding was excluded by gating around those cells which were labeled by the isotype negative control antibodies.

All data are the mean $\pm$ SD of two experiments. Fisher's exact test was adopted for statistical evaluation of the results. A *t*-test was used for the analysis of protein level measurements. All *P* values are two-sided. Precise *P* values are given. All statistical analyses were performed in GraphPad InStat version 3.00 for Windows 95, GraphPad Software, San Diego, CA.

Figure 2 shows the growth curve of Jurkat cells at different time points and after treatment with different concentrations




Fig. 2. The effect of sulforaphane on cell viability. The GRA–Myr mixture was added at the appropriate concentrations to exponentially growing cultures of Jurkat cells. Aliquots were removed from drug-treated and untreated control cultures after different time and stained with trypan blue. Data are means of duplicate experiments. When they do not appear, deviation bars are smaller than the symbol size.

of sulforaphane. When the harvested cells were counted, the numbers of control cells continued to increase, whereas the numbers of treated cells decreased in a time- and dose-dependent manner. The maximum effect was registered after 48 h of treatment and at a concentration of 30 μM sulforaphane. In fact, the number of cells decreased from ~ 6 to 1×10^6 . These changes appear to be irreversible, persisting when the treated cells were recultured in sulforaphane-free medium (data not shown). Only a slight decrease in the number of viable cells in cultures treated with Myr or GRA 30 μM was observed. The ID_{50} , as measured by the number of viable cells in cultures 24 h after the addition of GRA plus Myr, was seen at a concentration of $\sim 15 \mu\text{M}$.

Sulforaphane but not GRA induces cell-cycle arrest

To elucidate the rate at which sulforaphane altered cell-cycle progression, cell-cycle progression was determined at several time points. Table I reports the effects of sulforaphane (and of GRA and Myr) on the cell-cycle distribution of Jurkat cells. The effect of sulforaphane was time dependent. The immediate effects (8 h), observed at drug concentrations of 10–30 μM , appeared primarily as an increase in the proportion of cells in the G_2/M phase of the cell cycle (from ~ 17 to 24%) accompanied by a slight compensatory decrease in G_1 phase cells. Longer exposure (24 h) to sulforaphane concentrations of 3, 10 and 30 μM led to a further decrease in the proportion of G_1 cells, while the percentage of cells in G_2/M phase increased from 18 to 30%. Prolonged (48 h) exposure of Jurkat cells to sulforaphane appeared as a decrease in G_1 -phase cells (from 44 to 33%), a loss of S-phase cells (from 44 to 38%) and a marked increase in the proportion of G_2/M -phase cells (from 12 to 25%). All these effects were particularly apparent following treatment with sulforaphane at concentrations of 10 and 30 μM .

No effect on cell-cycle progression was detected in cell cultures treated with enzyme only. A very slight effect was observed with GRA after 48 h of treatment.

Sulforaphane but not GRA induces apoptosis in a time- and dose-dependent fashion

Apoptosis was observed at the highest dose of sulforaphane and at late stages during the treatment (at 24 and 48 h). It manifested by appearance of cells with decreased DNA content, identified in the DNA content distributions as cells with DNA

Table I. Effects of sulforaphane (and of GRA and Myr) on cell-cycle progression with respect to the control^a

Treatment	8 h				24 h				48 h			
	G_0/G_1 (mean $\pm$ SD)	S (mean $\pm$ SD)	G_2/M (mean $\pm$ SD)	G_0/G_1 (mean $\pm$ SD)	S (mean $\pm$ SD)	G_2/M (mean $\pm$ SD)	G_0/G_1 (mean $\pm$ SD)	S (mean $\pm$ SD)	G_2/M (mean $\pm$ SD)	G_0/G_1 (mean $\pm$ SD)	S (mean $\pm$ SD)	G_2/M (mean $\pm$ SD)
Control	37.24 $\pm$ 2.30	45.91 $\pm$ 3.67	16.84 $\pm$ 2.60	38.22 $\pm$ 2.30	44.12 $\pm$ 2.02	17.65 $\pm$ 1.26	44.26 $\pm$ 2.0	44.17 $\pm$ 1.26	11.56 $\pm$ 3.21	44.26 $\pm$ 2.0	44.17 $\pm$ 1.26	11.56 $\pm$ 3.21
Sulforaphane 3 μM	39.33 $\pm$ 1.36	42.36 $\pm$ 2.44	18.31 $\pm$ 2.07	37.28 $\pm$ 0.78	43.18 $\pm$ 0.01	19.52 $\pm$ 0.78	39.66 $\pm$ 2.38	46.03 $\pm$ 4.96	14.32 $\pm$ 0.78	39.66 $\pm$ 2.38	46.03 $\pm$ 4.96	14.32 $\pm$ 0.78
Sulforaphane 10 μM	33.49 $\pm$ 3.22	42.33 $\pm$ 3.67	24.18 $\pm$ 1.07	26.08 $\pm$ 3.67	43.99 $\pm$ 0.42	29.92 $\pm$ 4.09	42.09 $\pm$ 2.44	37.76 $\pm$ 2.38	20.15 $\pm$ 1.98	42.09 $\pm$ 2.44	37.76 $\pm$ 2.38	20.15 $\pm$ 1.98
Sulforaphane 30 μM	33.56 $\pm$ 4.09	43.28 $\pm$ 1.32	23.16 $\pm$ 0.23	31.87 $\pm$ 4.96	42.90 $\pm$ 2.60	25.23 $\pm$ 2.38	33.33 $\pm$ 2.44	41.50 $\pm$ 2.01	25.17 $\pm$ 1.30 ^b	33.33 $\pm$ 2.44	41.50 $\pm$ 2.01	25.17 $\pm$ 1.30 ^b
GRA 30 μM	36.29 $\pm$ 2.07	44.39 $\pm$ 0.44	19.32 $\pm$ 2.38	40.38 $\pm$ 2.07	43.62 $\pm$ 0.86	15.99 $\pm$ 2.93	51.66 $\pm$ 2.91	37.96 $\pm$ 0.86	10.38 $\pm$ 0.01	51.66 $\pm$ 2.91	37.96 $\pm$ 0.86	10.38 $\pm$ 0.01
Myr	36.32 $\pm$ 0.11	47.64 $\pm$ 1.07	17.44 $\pm$ 0.79	38.86 $\pm$ 3.30	44.34 $\pm$ 2.22	16.80 $\pm$ 1.07	45.11 $\pm$ 0.18	43.84 $\pm$ 0.67	10.99 $\pm$ 0.92	45.11 $\pm$ 0.18	43.84 $\pm$ 0.67	10.99 $\pm$ 0.92

^aAt the indicated time, distribution of the cells in G_0/G_1 , S and G_2/M phase was analyzed by flow cytometry as described in Materials and methods. Results are expressed as the percentage of total cells. Data represent means of duplicate experiments.

^b $p = 0.028$ with respect to the control.

Table II. Incidence of apoptosis as concurrently detected by two methods in Jurkat cells treated with different doses and at different times with sulforaphane^a

Time (h)	Concentrations of sulforaphane (μM)	% Apoptosis	
		Sub-G ₁ DNA (mean ± SD)	Annexin (mean ± SD)
8	Control	1.00 ± 0.14	1.00 ± 0.00
	3	1.50 ± 0.71	1.00 ± 0.00
	10	1.85 ± 0.21	1.50 ± 0.71
	30	2.00 ± 0.00	5.50 ± 2.12
24	Control	1.15 ± 0.21	2.00 ± 0.00
	3	1.00 ± 0.00	3.00 ± 1.41
	10	2.85 ± 0.21	4.00 ± 1.41
	30	8.50 ± 0.71 ^b	16.00 ± 4.24 ^c
48	Control	1.00 ± 0.00	3.50 ± 0.71
	3	1.00 ± 0.00	2.00 ± 0.00
	10	2.5 ± 0.71	4.00 ± 0.00
	30	38.00 ± 0.00 ^d	32.00 ± 0.71 ^d

^aAt the indicated time, total cells were treated as described in Materials and methods. The DNA content (sub-G₁ peak) and the binding of annexin V-FITC to the plasma membrane were analyzed by flow cytometry.

^b*P* = 0.035, ^c*P* = 0.001, ^d*P* < 0.0001 with respect to the control.

values below that of G₁ cells ('sub-G₁ cells'). Thus, as an example, after 8 h of treatment with 30 μM sulforaphane, only 2% of apoptotic cells were found. This fraction significantly increased to 8 and 38% after 24 and 48 h of treatment, respectively (Table II).

The proportion of cells reactive with the annexin V-FITC conjugate indicates that target cells start to die as late as 24 h after effector and target mixing by an apoptosis-related mechanism. Thus, after 8 h of treatment of Jurkat cells with 30 μM sulforaphane, the incidence of annexin V^{pos} and PI^{neg} cells—i.e. apoptotic cells—was 5% (vs 1% in the control). They increased to 16% after 24 h and 32% after 48 h (Table II). Membrane permeabilization, revealed by PI uptake (annexin V^{pos} – PI^{pos}), presented the same kinetics (data not shown). There was only 3% of annexin V^{pos}/PI^{pos} cells after 8 h of treatment. This fraction increased to 14 and 30% after 24 and 48 h of treatment, respectively. No effects on apoptotic event induction were detected in cell cultures treated with GRA or with Myr (data not shown).

Sulforaphane modulates p53 and bax protein expression, but does not affect bcl-2 protein expression

To examine the effect of sulforaphane on levels of p53, bcl-2 and bax, cells were treated for 48 h with sulforaphane 30 μM. Apoptotic cells were gated, based on their changed FSC and SSC distribution, and p53, bcl-2 and bax proteins were then quantified. Figure 3 shows histograms of p53, bcl-2 and bax protein levels following incubation without compound or sulforaphane. p53 and bax expression were found to be increased in treated cells relative to the untreated cells. Typically, p53 and bax protein levels showed ~7- and 3-fold increase with regard to the control, respectively. bcl-2 protein level was only slightly affected.

Discussion

What this study adds to our knowledge is the first demonstration that sulforaphane, generated from GRA by Myr hydrolysis but not its precursor GL, is able to induce cell-cycle arrest and apoptosis or necrosis in a leukemic cell line. The bioactivity




Fig. 3. Representative histograms showing p53, bcl-2 and bax protein level following 48 h culture in the presence of no compound and sulforaphane (30 μM), respectively. When they do not appear, deviation bars are smaller than the symbol size.

of GRA was observed only in the presence of Myr, thereby demonstrating that the hydrolytic product sulforaphane is the biologically active agent, according to the evidence that ITCs are ~1000-fold more potent than the precursors (38).

Treatment of Jurkat cells with generated sulforaphane resulted in a potent inhibition of cell growth *in vitro*. Flow cytometric DNA analysis of treated cultures revealed that complete derangement of cell-cycle progression was responsible for the inhibition of tumor-cell growth. Examination of the Jurkat cells after a 48 h incubation with various doses of sulforaphane revealed a marked increase in the percentage of cells in the G₂/M cell-cycle phase. Such behavior is in line with the recent finding that synthetic sulforaphane is capable of causing cell-cycle arrest in the G₂/M phase in the HT29 colon cancer cell line (19).

Moreover, simultaneous analysis of cell-cycle progression (sub-G₁ peak) and loss of phospholipid asymmetry of the plasma membrane in sulforaphane-treated cultures revealed a strictly dose- and time-dependent increase in the percentage of Jurkat cells bearing the flow cytometric characteristics of apoptosis. The latter became apparent after 24 h of incubation. Taken together, these observations imply that the growth inhibition of Jurkat cells produced by sulforaphane (as generated *in situ* from the GRA-Myr mixture) results from a combination of apoptosis and of cell-cycle derangements in which G₂/M arrest is a key event. The simultaneous appearance of G₂/M arrest and apoptosis clearly indicates that cell death is a primary direct effect due to the hydrolytic product of GRA, and not to activation of apoptotic pathways as a consequence of cells' inability to overcome growth arrest and proceed through the cell cycle.

Interactions between p53, bax, bcl-2 and its associated proteins have been postulated to form one pathway of apoptosis (39,40). We used flow cytometric analysis to evaluate the levels of p53, bcl-2 and bax proteins, after treatment with GRA plus Myr. We found that the generated sulforaphane increased p53 and bax protein levels, whereas bcl-2 levels remained unchanged. These observations support the hypothesis that induction of p53 expression may be an important step during the sulforaphane-induced apoptosis. Apoptosis is activated by many oxidative agents (41), and Payen *et al.* (42) indicated a sulforaphane-mediated production of reactive

oxygen species. Moreover, Zhang (43) demonstrated that ITCs, and sulforaphane among these, are taken by cells predominantly through GSH conjugation reactions in cells. Sulforaphane may induce oxidative stress by reacting with and depleting the intracellular GSH pool, which may then induce apoptosis. The molecular mechanism may be mediated by the c-Jun N-terminal kinase, as demonstrated for other ITCs (44) and, therefore, by p53 known to be a substrate of this kinase (45). Otherwise, Bonnesen *et al.* (46) demonstrated that sulforaphane is not associated with DNA damage in colon cell lines, as assessed by comet assay. *bax* has been found to be up-regulated during apoptosis in studies on several other cell types (47,48). *p53* is a direct transcriptional activator of the *bax* gene (48). So in the presence of increased *p53* expression, it is scarcely surprising that *bax* expression is also affected. The induction of apoptosis recorded in this study seems to be dependent on p53 gene. This is contrary to a recent report (19) where sulforaphane-induced HT29 cell death is not associated with a change in p53 protein expression but is accompanied by an over expression of *bax*. On the other hand, Huang *et al.* (18) demonstrated an essential role of p53 in the induction of apoptosis by phenylethyl ITC in a mouse epidermal cell line. However, it should be borne in mind that many of the genes and proteins thought to be involved in the apoptotic process may have cell- and injury-type-dependent roles (36).

The present study also demonstrates for the first time that high doses of sulforaphane are capable of inducing necrosis. In the Jurkat cells studied by us, sulforaphane induced necrosis after prolonged exposure (48 h) to the highest concentration (30 μ M) investigated. This suggests that very high levels of exposure to produced sulforaphane can cause some form of intensive cell toxicity.

We conclude that the sulforaphane produced *in situ* is able to exert G₂/M phase cell-cycle arrest and produce apoptotic death (and at very high doses even necrosis) of Jurkat tumor cells. The end result of these effects is potent inhibition of the leukemic cell growth. We propose that sulforaphane could be tested as a potential chemopreventive and chemotherapeutic agent, also in the light of its effects on xenobiotic metabolism that were detected in a range of concentrations similar to that used in this study. In particular, sulforaphane is able to induce *in vitro* phase II detoxifying enzymes at 5 μ M, with a maximal activity at 50 μ M and a decrease of activity at 100 μ M (49). Experimental investigations into ways in which sulforaphane could be employed in selective chemoprotective strategies for subjects affected by pre-neoplastic lesions could also be worthwhile. Further in-depth *in vitro* studies have to be performed to compare the activity of sulforaphane on different cell types. *In vivo* studies may then lead to clinical trials to investigate the therapeutic potential of sulforaphane.

Acknowledgements

The authors are grateful to Robin M.T.Cooke for editing. This work was supported by the University of Bologna Special Grant to the project 'Development of fluorescent probes'.

References

- Morse, M.A. and Stoner, G.D. (1993) Cancer chemoprevention. Principles and prospects. *Carcinogenesis*, **14**, 1737–1746.
- Kelloff, G.J., Boone, C.W., Malone, W.F. and Steele, V.E. (1992) Recent results in preclinical and clinical drug development of chemopreventive agents at the National Cancer Institute. In Wattenberg, L., Lipkin, M., Boone, C.W. and Kelloff, G.J. (eds) *Cancer Chemoprevention*. CRC Press, Boca Raton, FL, pp. 41–56.
- Hsieh, T. and Wu, J.M. (1999) Differential effects on growth, cell cycle arrest and induction of apoptosis by resveratrol in human prostate cancer cell lines. *Exp. Cell Res.*, **249**, 109–115.
- Kelloff, G.J., Boone, C.W., Crowell, J.A. and Steele, V.E. (1996) New agents for cancer chemoprevention. *J. Cell. Biochem.*, **26**, 1–28.
- Surh, Y. (1999) Molecular mechanisms of chemopreventive effects of selected dietary and medicinal phenolic substances. *Mutat. Res.*, **428**, 305–327.
- Sharma, S., Stutzman, J.D., Kelloff, G.J. and Steele, V.E. (1994) Screening of potential chemopreventive agents using biochemical markers of carcinogenesis. *Cancer Res.*, **54**, 5848–5855.
- Leoni, O., Iori, R. and Palmieri, S. (2000) Hydrolysis of glucosinolates using nylon-immobilized myrosinase to produce pure bioactive molecules. *Biotechnol. Bioeng.*, **68**, 660–664.
- Zhang, Y. and Talalay, P. (1994) Anticarcinogenic activities of organic isothiocyanates: chemistry and mechanisms. *Cancer Res.*, **54** (suppl.), 1976s–1981s.
- Hecht, S.S. (1995) Chemoprevention by isothiocyanates. *J. Cell. Biochem.*, **22**, 195–209.
- Barcelo, S., Gardiner, J.M., Gescher, A. and Chipman, J.K. (1996) CYP2E1-mediated mechanism of anti-genotoxicity of the broccoli constituent sulforaphane. *Carcinogenesis*, **17**, 277–282.
- Kall, M.A., Vang, O. and Clausen, J. (1996) Effect of dietary broccoli on human *in vivo* drug metabolizing enzymes: evaluation of caffeine, oestrone and chlorzoxazone metabolism. *Carcinogenesis*, **17**, 793–799.
- Zhang, Y., Talalay, P., Cho, C.-G. and Posner, G.H. (1992) A major inducer of anticarcinogenic protective enzymes from broccoli: isolation and elucidation of structure. *Proc. Natl Acad. Sci. USA*, **89**, 2399–2403.
- Tawfiq, N., Heaney, R.K., Plumb, J.A., Fenwick, G.R., Musk, S.R.R. and Williamson, G. (1995) Dietary glucosinolates as blocking agents against carcinogenesis: glucosinolate breakdown products assessed by induction of quinone reductase activity in murine hepatic cells. *Carcinogenesis*, **16**, 1191–1194.
- Morse, M.A., Amin, S.G., Hecht, S.S. and Chung, F.L. (1989) Effects of aromatic isothiocyanates on tumorigenicity, O⁶-methylguanine formation and metabolism of the tobacco-specific nitrosamine-4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone in A/J mouse lung cancer. *Cancer Res.*, **49**, 2894–2897.
- Bogaards, J.J.P., Verhagen, H., Willems, M.I., van Poppel, G. and van Bladeren, P.J. (1994) Consumption of Brussels sprouts results in elevated a-class glutathione S-transferase levels in human blood plasma. *Carcinogenesis*, **15**, 1073–1075.
- Wattenberg, L.W. (1990) Inhibition of carcinogenesis by minor nutrient constituents of the diet. *Proc. Nutr. Soc.*, **49**, 173–183.
- Thompson, C.B. (1995) Apoptosis in the pathogenesis and treatment of disease. *Science*, **267**, 1456–1462.
- Huang, C., Ma, W.Y., Li, J., Hecht, S.S. and Dong, Z. (1998) Essential role of p53 in phenethyl isothiocyanate-induced apoptosis. *Cancer Res.*, **58**, 4102–4106.
- Gamet-Payraastre, L., Li, P., Lumeau, S., Cassar, G., Dupont, M.-A., Chevrolleau, S., Gasc, N., Tulliez, J. and Tercé, F. (2000) Sulforaphane, a naturally occurring isothiocyanate, induces cell cycle arrest and apoptosis in HT29 human colon cancer cells. *Cancer Res.*, **60**, 1426–1433.
- Getahun, S.M. and Chung, F.L. (1999) Conversion of glucosinolates to isothiocyanates in humans after ingestion of cooked watercress. *Cancer Epidemiol. Biomarkers Prev.*, **8**, 447–451.
- Iori, R., Bernardi, R., Gueyrard, D., Rollin, P. and Palmieri, S. (1999) Formation of glucoraphanin by chemoselective oxidation of natural glucoerucin: a chemoenzymatic route to sulforaphane. *Bioorg. Med. Chem. Lett.*, **9**, 1047–1048.
- Visentin, M., Tava, A., Iori, R. and Palmieri, S. (1991) Isolation and identification of *trans*-4-(methylthio)-3-butenyl glucosinolate from radish roots (*Raphanus sativus* L.). *J. Agric. Food Chem.*, **40**, 1687–1691.
- EEC Regulation No. 1864/90 (1990) Enclosure VIII. *Offic. J. Eur. Commun.*, **L170**, 27–34.
- Pessina, A., Thomas, R.M., Palmieri, S. and Luisi, P.L. (1990) An improved method for the purification of myrosinase and its physicochemical characterization. *Arch. Biochem.*, **280**, 383–389.
- Leoni, O., Iori, R., Palmieri, S., Esposito, E., Menegatti, E., Cortesi, R. and Nastruzzi, C. (1997) Myrosinase-generated isothiocyanate from glucosinolates: isolation, characterization and *in vitro* antiproliferative studies. *Bioorgan. Med. Chem.*, **5**, 1799–1806.
- Sun, S.-Y., Yue, P., Shroot, B., Hong, W.K. and Lotan, R. (1997) Induction of apoptosis in human non-small cell lung carcinoma cells by the novel synthetic retinoid CD437. *J. Cell Physiol.*, **173**, 279–284.
- Nüsse, M., Beisker, W., Kramer, J., Miller, B.M., Schreiber, G.A., Viaggi, S., Weller, E.M. and Wessels, J.M. (1994) Measurement of micronuclei by flow

- cytometry. In Darzynkiewicz,Z. and Crissman,H.A. (eds) *Methods in Cell Biology*. Academic Press, San Diego, CA, Vol. 42, pp. 149–58.
28. Fimognari,C., Nüsse,M. and Hrelia,P. (1999) Flow cytometric analysis of genetic damage, effect on cell cycle progression and apoptosis by Thiophanate-methyl in human lymphocytes. *Environ. Mol. Mutagen.*, **33**, 220–223.
 29. Schreiber,G.A., Beisker,W., Braselmann,H., Bauchinger,M., Bögl,K.W. and Nüsse,M. (1992) Technical report: an automated flow cytometric micronucleus assay for human lymphocytes. *Int. J. Radiat. Biol.*, **62**, 695–709.
 30. Fadok,V.A., Voelker,D.R., Campbell,P.A., Cohen,J.J., Bratton,D.L. and Henson,P.M. (1992) Exposure of phosphatidylserine on the surface of apoptotic lymphocytes triggers specific recognition and removal by macrophages. *J. Immunol.*, **148**, 2207–2216.
 31. Koopman,G., Reutelingsperger,C.P.M., Kuijten,G.A.M., Keehnen,R.M., Pals,S.T. and van Oers,M.H. (1994) Annexin V for flow cytometric detection of phosphatidylserine expression of B cells undergoing apoptosis. *Blood*, **84**, 1415–1420.
 32. Vermes,I., Haanen,C., Stefens Nakken,H. and Reutelingsperger,C. (1995) A novel assay for apoptosis. Flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labelled Annexin V. *J. Immunol. Methods*, **184**, 39–51.
 33. Del Bino,G., Darzynkiewicz,Z., Degraef,C., Mosselmans,R., Fokan,D. and Galand,P. (1999) Comparison of methods based on annexin-V binding, DNA content or TUNEL for evaluating cell death in HL-60 and adherent MCF-7 cells. *Cell Prolif.*, **32**, 25–37.
 34. Benini,E., Costa,A., Abolafio,G. and Silvestrini,R. (1998) p53 expression in human carcinomas: could flow cytometry be an alternative to immunohistochemistry? *J. Histochem. Cytochem.*, **46**, 41–47.
 35. Filippini,G., Griffin,S., Uhr,M., Eppenberger,H., Bonilla,J., Cavalli,F. and Soldati,G. (1998) A novel flow cytometric method for the quantification of p53 gene expression. *Cytometry*, **31**, 180–186.
 36. Liu,X. and Zhu,X.-Z. (1999) Roles of p53, c-Myc, Bcl-2, Bax and caspases in glutamate-induced neuronal apoptosis and the possible neuroprotective mechanism of basic fibroblast growth factor. *Mol. Brain Res.*, **71**, 210–216.
 37. Chiu,L.C.M. and Wan,J.M.F. (1999) Induction of apoptosis in HL-60 cells by eicosapentaenoic acid (EPA) is associated with downregulation of bcl-2 expression. *Cancer Lett.*, **145**, 17–27.
 38. Musk,S.R., Smith,T.K. and Johnson,I.T. (1995). On the cytotoxicity and genotoxicity of allyl and phenethyl isothiocyanates and their parent glucosinolates sinigrin and gluconasturtiin. *Mutat. Res.*, **348**, 19–23.
 39. Adams,J.M. and Cory,S. (1998) The Bcl-2 protein family: arbiters of cells survival? *Science*, **281**, 322–326.
 40. Staunton,M.J. and Gaffney,E.F. (1998) Apoptosis: basic concepts and potential significance in human cancer. *Arch. Pathol. Lab. Med.*, **122**, 310–319.
 41. Jacobson,M.D. (1996) Reactive oxygen species and programmed cell death. *Trends Biochem. Sci.*, **21**, 83–86.
 42. Payen,L., Courtois,A., Loewert,M., Guillouzo,A. and Fardel,O. (2001) Reactive oxygen species-related induction of multidrug resistance-associated protein 2 expression in primary hepatocytes exposed to sulforaphane. *Biochem. Biophys. Res. Commun.*, **282**, 257–263.
 43. Zhang,Y. (2001) Molecular mechanism of rapid cellular accumulation of anticarcinogenic isothiocyanates. *Carcinogenesis*, **22**, 425–431.
 44. Chen,Y.-R., Wang,W., Kong,A.-N.T. and Tan,T.-H. (1998) Molecular mechanisms of c-Jun N-terminal kinase-mediated apoptosis induced by anticarcinogenic isothiocyanates. *J. Biol. Chem.*, **273**, 1769–1775.
 45. Milne,D.M., Campbell,L.E. and Meek,D.W. (1995) p53 is phosphorylated *in vitro* and *in vivo* by an ultraviolet radiation-induced protein kinase characteristic of the c-Jun kinase, JNK1. *J. Biol. Chem.*, **270**, 5511–5518.
 46. Bonnesen,C., Eggleston,I.M. and Hayes,J.D. (2001) Dietary indoles and isothiocyanates that are generated from cruciferous vegetables can both stimulate apoptosis and confer protection against DNA damage in human colon cell lines. *Cancer Res.*, **61**, 6120–6130.
 47. Selakumaran,M., Lin,H.-K., Miyashita,T., Wang,H.G., Krajewski,S., Reed,J.C., Hoffman,B. and Liebermann,D. (1994) Immediate early up-regulation of bax expression by p53 but not TGF- β , a paradigm for distinct apoptotic pathways. *Oncogene*, **9**, 1791–1798.
 48. Miyashita,T. and Reed,J.C. (1995) Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. *Cell*, **80**, 293–299.
 49. Yu,R., Lei,W., Mandlekar,S., Webers,M.J., Der,C.J., Wu,J. and Kong,A.-N.T. (1999) Role of a mitogen-activated protein kinase pathway in the induction of phase II detoxifying enzymes by chemicals. *J. Biol. Chem.*, **274**, 27545–27552.

Received June 20, 2001; revised December 3, 2001;
accepted December 4, 2001