

Lieferschein

Bayerische Staatsbibliothek Muenchen

- Dokumentlieferung –
Ludwigstr. 16

D-80539 Muenchen

Tel.: ++49-89-28638-2643
Fax: ++49-89-280-9284
Email: doklief@bsb-muenchen.de

Empfänger

Helmholtz Zentrum Muenchen GmbH

Zentralbibliothek / Fernleihe

D-85764 Neuherberg

Ingolstaedter Landstr. 1

Angaben zur Bestellung:

Bestelldatum: 2017-04-04 12:38:12
Bestellnummer: SUBITO:VE17040400699
Name des Bestellers: Helmholtz Zentrum Muenchen GmbH
Benutzerkennung: SLS02X00668

Lieferdatum: 2017-04-04 16:15:07
Lieferpriorität: NORMAL
Aktueller Lieferweg: Email
E-Mail Adresse: library@helmholtz-muenchen.de

Bemerkungen zur Auslieferung:

Angaben zum Dokument:

Signatur: 4 Z 92.144 Hbzs 752-17 b = Neueste Hefte
Autor:
Titel: Journal of the American Society of Nephrology
Jahr: 2016
Band / Jahrgang: 28/3
Seiten: 981-994
Aufsatzautor: Li, M
Aufsatztitel: SOS2 and ACP1 Loci Identified through Large-Scale Exome Chip Analysis Regulate Kidney Development
ISSN:
ISBN: 1046-6673
CODEN:

Ihre Bemerkung zur Bestellung: Paulini

subito Urheberrechtshinweis



Die Bestellung und Nutzung der über subito gelieferten Aufsatzkopien unterliegen den urheberrechtlichen Bestimmungen. Mit der Registrierung bei subito verpflichten Sie sich, diese einzuhalten, d.h. insbesondere, dass die Kopien ausschließlich zum eigenen Gebrauch bestimmt sind und nicht an Dritte weitergegeben werden dürfen. Sie dürfen ohne Genehmigung des Verlags nicht zum Wiederverkauf, Wiederabdruck, zu systematischer Verteilung, Emailversand, Webhosting eingeschlossen institutionelle Repositorien/Archive oder jedweden anderen kommerziellen Zweck verwendet werden.

Sofern Sie eine Lieferung per Email oder FTP erhalten, dürfen Sie die Kopie nur einmal ausdrucken und müssen diese anschließend dauerhaft löschen.

Die Kopien sind mit einem Wasserzeichen versehen, welches ein Urheberrechtsvermerk enthält. Das von subito e.V. angebrachte Wasserzeichen darf nicht entfernt werden.

FTP

Bestelldatum: 2017-04-04 12:38:12

BSB Bayerische
Staatsbibliothek

NORMAL

Kopie

SUBITO-VE17040400699



Helmholtz Zentrum Muenchen GmbH
Zentralbibliothek / Fernleihe
Herr Rasso Ranzinger
Ingolstaedter Landstr. 1
85764 Neuherberg

Ben.-Gruppe: USER-GROUP-4
Tel: +49 89 31872343
Mail: docdel@subito-doc.de

Fax: +49 89 31873391

Subito-Kundennummer:
SLS02X00668
Subito-Bestellnummer:
SUBITO-VE17040400699

4 Z 92.144 Hbzs 752-17 b = Neueste Hefte

Jahrgang: 2016

Band/Heft: 28/3

Seiten: 981-994

Verfasser: Li, M

**Titel: SOS2 and ACP1 Loci Identified through Large-Scale
Exome Chip Analysis Regulate Kidney Development an**

Journal of the American Society of Nephrology
ISSN: 1046-6673

Bemerkung: Paulini

Beschreibung:

Die Abrechnung dieser Lieferung erfolgt über die subito-Zentralregulierung

Bei Rückfragen wenden Sie sich bitte innerhalb von 10 Tagen an die Bayerische Staatsbibliothek, Direktlieferdienste
Tel. ++49 89 28 638-26 43, doklief@bsb-muenchen.de

Wir weisen den Empfänger darauf hin, dass Sie nach geltendem Urheberrecht die von uns übersandten Vervielfältigungsstücke ausschließlich zu Ihrem privaten oder sonstigen Gebrauch verwenden und weder entgeltlich noch unentgeltlich in Papierform oder als elektronische Kopien verbreiten dürfen.

SOS2 and ACP1 Loci Identified through Large-Scale Exome Chip Analysis Regulate Kidney Development and Function

CKDGen Consortium

Due to the number of contributing authors, the authors and affiliations are listed at the end of this article.

ABSTRACT

Genome-wide association studies have identified >50 common variants associated with kidney function, but these variants do not fully explain the variation in eGFR. We performed a two-stage meta-analysis of associations between genotypes from the Illumina exome array and eGFR on the basis of serum creatinine (eGFR_{crea}) among participants of European ancestry from the CKDGen Consortium (n_{Stage1} : 111,666; n_{Stage2} : 48,343). In single-variant analyses, we identified single nucleotide polymorphisms at seven new loci associated with eGFR_{crea} (*PPM1J*, *EDEM3*, *ACP1*, *SPEG*, *EYA4*, *CYP1A1*, and *ATXN2L*; $P_{\text{Stage1}} < 3.7 \times 10^{-7}$), of which most were common and annotated as nonsynonymous variants. Gene-based analysis identified associations of functional rare variants in three genes with eGFR_{crea}, including a novel association with the SOS Ras/Rho guanine nucleotide exchange factor 2 gene, *SOS2* ($P = 5.4 \times 10^{-8}$ by sequence kernel association test). Experimental follow-up in zebrafish embryos revealed changes in glomerular gene expression and renal tubule morphology in the embryonic kidney of *acp1*- and *sos2*-knockdowns. These developmental abnormalities associated with altered blood clearance rate and heightened prevalence of edema. This study expands the number of loci associated with kidney function and identifies novel genes with potential roles in kidney formation.

J Am Soc Nephrol 28: 981–994, 2017. doi: 10.1681/ASN.2016020131

CKD is considered a complex phenotype with a genetic predisposition.¹ Previous genome-wide association studies (GWAS) have successfully identified multiple common genetic risk variants associated with the CKD-defining measures of eGFR and urinary albumin-to-creatinine ratio (UACR).^{2–5} Together, these variants explain only a small proportion of the variation in eGFR and UACR.⁶ To comprehensively interrogate protein-coding regions and assess the effects of rare variants (minor allele frequency [MAF] < 1%), we carried out a two-stage meta-analysis of the association between eGFR on the basis of serum creatinine (eGFR_{crea}) and variants genotyped on the Illumina HumanExome chip (http://genome.sph.umich.edu/wiki/Exome_Chip_Design) among 111,666 European ancestry (EA) participants from the CKDGen Consortium and assessed the role of genes significantly associated with eGFR_{crea} in kidney development using embryonic zebrafish models. In secondary analyses, we examined associations with

eGFR_{crea} stratified by diabetes status, and in a smaller subset of EA participants, we also tested eGFR on the basis of cystatin C (eGFR_{cys}) and UACR. An additional 9624 participants of African ancestry (AA) were also used in an independent exome-chip discovery meta-analysis.

Received February 3, 2016. Accepted August 22, 2016.

M. Li, Y. Li, and O.W. contributed equally to this work.

W.G., A. Köttgen, and A.Y.C. jointly oversaw this work.

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Audrey Y. Chu, NHLBI's Framingham Heart Study, 73 Mt Wayte Ave Suite #2, Framingham, MA 01702, or Dr. Anna Köttgen, University Medical Center Freiburg, Berliner Allee 29, 79110 Freiburg, Germany. Email: audrey.chu@nih.gov or anna.koettgen@uniklinik-freiburg.de

Copyright © 2017 by the American Society of Nephrology

RESULTS

Up to 120,357 participants from 27 studies of EA and up to 11,386 participants from seven studies of AA were included in stage 1 analyses for eGFR_{crea}, eGFR_{cys}, or UACR. An additional 48,343 participants from 12 studies of EA were included in stage 2 analysis of eGFR_{crea}. All participants provided informed consent and each of the studies was approved by its governing ethics committee or Institutional Review Board. Sample characteristics and genotyping information for each study are summarized in Supplemental Tables 1 and 2.

In stage 1 single-variant EA analyses, we identified 33 loci associated with eGFR_{crea} (Supplemental Table 3, Table 1) that met our *a priori* chip-wide significance threshold of $P < 3.7 \times 10^{-7}$ (0.05/134299 variants). Of these, eight had not been identified in association with eGFR_{crea} from previous GWAS analyses; six were missense variants: rs34611728 (*PPM1J*), rs78444298 (*EDEM3*), rs11553746 (*ACP1*), rs2307394 (*ORC4*), rs55760516 (*SPEG*), and rs9493627 (*EYA4*); and two were GWAS tag single nucleotide polymorphisms (SNPs) included on the exome chip due to prior associations in the National Human Genome Research Institute (NHGRI) GWAS Catalog for caffeine/coffee intake:^{7,8} rs2472297 (intergenic, near *CYP1A1*) and inflammatory bowel disease⁹ rs8049439 (intronic, *ATXN2L*); all were common variants ($MAF > 1\%$). A Manhattan plot displaying all 33 chip-wide significant loci is shown in Figure 1. Quantile-quantile plots indicated no inflation of the overall *P* value distribution (Supplemental Figure 1). Regional association plots show the associations of other variants within the 500 Mb region of the index variant of the eight newly identified loci in Supplemental Figure 2.

In stage 2 analyses, we followed up the eight newly identified eGFR_{crea} loci meeting our significance threshold among an additional 48,343 EA participants from 12 studies (sample characteristics and genotyping information are summarized in Supplemental Tables 2 and 4). All loci but *ORC4* met criteria for replication (a direction of effect consistent with stage 1 analysis; a $P_{1-sided} < 0.05$ from stage 2 analysis; and $P < 3.7 \times 10^{-7}$ from a combined stage 1 and stage 2 analysis; Table 1). Because diabetes mellitus is a major risk factor for CKD, we assessed the genetic associations at these loci in the presence or absence of diabetes to obtain additional insights into potential mechanistic pathways. In total, 11,040 and 94,677 participants were included in the diabetes- and nondiabetes-specific analyses, respectively. In analyses stratified by diabetes status, the β -coefficients were directionally consistent and of similar magnitude between the strata for six out of seven newly identified loci (Supplemental Table 5).

To identify additional novel loci associated with alternative measures of kidney function, we tested the association of single variants with eGFR_{cys} ($n = 32,861$ EA participants) and UACR ($n = 31,164$ EA participants). All observed associations achieving the significance threshold ($P < 3.7 \times 10^{-7}$; Supplemental

Table 6) have been previously reported in association with eGFR_{crea}, eGFR_{cys}, or UACR.^{3,4,6}

To further investigate the role of rare variants in kidney function, we performed gene-based tests for 9990 autosomal genes that contained at least two nonsynonymous or splice-site variants with $MAF < 1\%$. No evidence of inflation was observed in quantile-quantile plots for gene-based tests (Supplemental Figure 3). Three genes, *SOS2*, *SLC47A1*, and *LRP2*, met the experiment-wide threshold for gene-based significance ($P < 2.5 \times 10^{-6}$; 0.05/19,922 tests [9961 genes \times 2 tests]; Table 2; Supplemental Figure 4). The association for *SLC47A1*, a renal solute transporter, was driven by the presence of a single variant, rs111653425, with MAF approximately 1% (Supplemental Table 7). Common variants at *SLC47A1* and *LRP2* have been previously reported in association with eGFR_{crea},^{4,6} thus implicating both rare and common variants at both loci.^{4,6} Conversely, the association with *SOS2* was novel ($P_{SKAT} = 5.38 \times 10^{-8}$; $P_{TI} = 3.25 \times 10^{-6}$). No genes reached the threshold for chip-wide significance in gene-based associations for eGFR_{crea} stratified by diabetes status, eGFR_{cys}, or UACR.

To identify genes that may play a role during kidney development, we used knockdown zebrafish embryos generated by injecting morpholino oligonucleotides (MOs) into single-cell stage embryos. Morpholinos are a commonly used tool in zebrafish screens because they enable an efficient identification of genes that may have a role in developmental processes. The morpholinos targeted genes with nonsynonymous variants (*ppm1j*, *acp1*, *eya4*, *speg*, and *edem3*) and the novel gene-based finding, *sos2* (Supplemental Figure 5A and Supplemental Table 8). General defects in the pronephros or embryonic kidney structure (marked by expanded *pax2a* expression) were observed in *acp1* ATG MO- and *sos2* ATG MO-injected embryos compared with controls (Figure 2, A, E, and I; $P < 0.001$ for both). Both *acp1* ATG- and *sos2* ATG-knockdowns showed elongated proximal tubules (increased *slc20a1a* expression) compared with the control group (Figure 2, B, F, and J; mean difference in proximal tubule length: *sos2* = 81.7 μm ; $P < 0.001$; and *acp1* = 74.7 μm ; $P < 0.001$; Figure 2M). The increase in proximal tubule length was not a consequence of increased embryo length, as both *sos2* ATG and *acp1* ATG-morphants had significantly reduced body length relative to controls (Supplemental Figure 5B). Additionally, *acp1* ATG-knockdowns showed shorter distal tubule length (*slc12a3* expression), which may be a consequence of reduced body length (Figure 2, C, G, and K; $P < 0.001$). No abnormalities were observed for podocytes (*wt1a* expression) for *sos2* or *acp1* compared with controls (Figure 2, D, H, and I; $P > 0.05$).

Because of the potential off-target effects of morpholinos, these developmental findings were validated with secondary splice-site morpholinos designed to target the *sos2* and *acp1* pre-mRNA (Supplemental Figure 5A). The *sos2* and *acp1* splice morpholino-injected embryos had significantly increased proximal tubule length relative to controls (Supplemental Figure 5, C–E). Furthermore, the *acp1* splice-site morpholino-injected embryos had shortened distal tubules (*slc12a3* expression)

consistent with the phenotype induced by the ATG morpholino (Supplemental Figure 5F). The reproducibility of the proximal tubule developmental defects with ATG and splice-site morpholinos adds confidence to the specificity of the morpholino-induced tubule phenotype.

Follow-up *in situ* hybridization experiments to determine expression patterns of *sos2* and *acp1* during zebrafish development did not reveal kidney-specific expression of *sos2* or *acp1*; however, *sos2* and *acp1* were broadly expressed throughout embryogenesis at key stages of kidney development and may be acting to control kidney development (Supplemental Figure 6). In humans, both *sos2* and *acp1* protein are detected in adult renal tubules.¹⁰ No significant developmental abnormalities were observed among MO knockdowns for the remaining genes (Supplemental Figure 6C). These findings suggest that both *SOS2* and *ACPI* may influence embryonic renal development, and that genetic influences on kidney development may contribute to variation in kidney function.

We next sought to determine whether *sos2* and *acp1*-mediated developmental alterations led to abnormalities in kidney function. In zebrafish, edema is a common sign of kidney failure.¹¹ We first performed an edema prevalence study and identified incidence rates of pericardial and global edema in *sos2* and *acp1* morphant larva. Both *sos2* and *acp1* ATG morpholino-injected embryos had a heightened incidence of pericardial edema beginning at 72 hours post fertilization (hpf) (Figure 3, A–C). The *sos2* morphants developed severe global edema by 120 hpf, whereas the *acp1* morphants presented with only pericardial edema (Figure 3, A–C). Both *sos2* and *acp1* morphants showed indications of embryonic lethality by 120–144 hpf (Figure 3, A–C).

An additional metric for kidney function is the assessment of glomerular filtration and fluid flow by fluorescent dextran clearance.¹² Control, *sos2*, or *acp1* morphant embryos were injected with equal volumes of rhodamine-labeled 70 kDa molecular mass dextran in the cardiac sinus venosus at 72 hpf. Dextran clearance rate was assessed by the quantification of rhodamine fluorescence intensity in a standardized area of the cardiac region at 2, 24, and 48 hours post injection (hpi). *sos2* and *acp1* morphant embryos exhibited decreased dextran clearance at both 24 and 48 hpi relative to controls (Figure 3, D–I, and M). Furthermore, renal tubules marked by fluorescent dextran in *sos2* and *acp1* morphant larva displayed an abnormal morphology (Figure 3, J–L). Specifically, the proximal convoluted tubules were reduced in size and lacked coiling depth. Failed clearance of 70 kDa molecular mass dextran and abnormal tubular structure suggests that morphants may have defects in tubular fluid flow and glomerular filtration.

We also evaluated heart rate in the *sos2* and *acp1* morphants relative to controls because compromised cardiovascular function could contribute to defects in dextran clearance. At 96 hpf (24 hours after dextran injection), the *sos2* (145.5 ± 3.854 bpm; $P=0.003$) and *acp1* (151.2 ± 2.653 bpm; $P<0.001$) morphants had an elevated mean heart rate relative to controls (127.8 ± 3.353 bpm) (Supplemental Figure 2G). Although altered

Table 1. Novel variants associated with eGFR_{crea} in EA participants from single-variant analysis in stage 1 meeting chip-wide significance ($P<3.7 \times 10^{-7}$) and associations in stage 2 and combined analysis

Locus ^a	dbSNPID	Chr	Position ^b	Variation (Substitution)	Stage 1 ^c			Stage 2 ^d			Combined		
					A1/A2 (A1 AF)	β (SEM)	P Value	r ²	β (SEM)	1-sided P	P Value	β (SEM)	P Value
PPM1J	rs34611728	1	113255456	c.639G>T (L213F)	A/C (0.13)	-0.0103 (0.0013)	1.2E-14	13.2	-0.0059 (0.0023)	4.7E-03	-0.0092 (0.0011)	3.3E-16	0.05
EDEM3	rs78444298	1	184672098	c.2236C>T (P746S)	A/G (0.02)	-0.0183 (0.0034)	5.2E-08	15.3	-0.0225 (0.0055)	1.8E-05	-0.0195 (0.0029)	1.5E-11	0.03
ACP1	rs11553746	2	272203	c.129C>T (T95I)	T/C (0.35)	-0.0049 (0.0009)	2.0E-07	20.7	-0.0032 (0.0016)	2.2E-02	-0.0045 (0.0008)	1.0E-08	0.02
ORC4 ^e	rs2307394	2	148716428	c.233A>G (N78S)	C/T (0.32)	-0.0058 (0.0010)	6.8E-09	14.3	-0.0025 (0.0016)	6.0E-02	-0.0049 (0.0009)	8.4E-09	0.03
SPEG	rs55760516	2	220354108	c.8191A>G (R2731G)	G/A (0.33)	0.0059 (0.0009)	4.8E-10	0.5	0.0054 (0.0016)	3.7E-04	0.0058 (0.0008)	1.7E-13	0.04
EYA4	rs9493627	6	133789728	c.829G>A (G223S)	A/G (0.31)	0.0061 (0.0010)	2.3E-10	0.0	0.0049 (0.0016)	1.4E-03	0.0058 (0.0009)	1.4E-11	0.04
CYP1A1	rs2472297	15	75027880	intergenic	T/C (0.24)	0.0057 (0.0010)	7.0E-08	0.0	0.0059 (0.0017)	3.2E-04	0.0058 (0.0009)	3.0E-11	0.03
ATXN2L	rs8049439	16	28837515	intronic	C/T (0.40)	0.0048 (0.0009)	1.3E-07	7.1	0.0045 (0.0016)	1.8E-03	0.0047 (0.0008)	1.2E-09	0.03

A1, effect allele; A2, non-effect allele; A1 AF, effect allele frequency; Chr, chromosome; Prop Var Exp, proportion of variance in ln(eGFR_{crea}) explained.

^aLoci are named according to the closest gene on the basis of the position of the lead SNP for new loci.

^bPosition is reported in UCSC Genome Browser/build hg19.

^cSample size for stage 1 analysis: $n=111,666$.

^dSample size for stage 2 analysis: $n=48,343$.

^eThis variant reached chip-wide significance ($P<3.7 \times 10^{-7}$) in the stage 1 samples but did not meet validation criteria in stage 2.




Figure 1. Manhattan plot for single-variant analysis in eGFRcrea among 111,666 EA participants. Newly identified variants are in dark orange. The gene, *ORC4*, not successfully replicated, is in orange. Known loci are in blue.

kidney development, abnormal tubular structure, edema, and decreased dextran clearance all support the conclusion that both *sos2* and *acp1* are regulators of kidney development and function, it remains possible that altered hemodynamics contribute to the edema and dextran clearance phenotypes observed.

In separate single variant analyses among 9624 AA participants, we identified three loci in association with eGFRcrea at chip-wide significance (Supplemental Table 9; $P < 3.7 \times 10^{-7}$). These variants were rare and the limited availability of AA cohorts prevented replication of these findings. The *APOL1* G1 variant, rs73885319, that was a known risk factor for kidney function decline and ESRD in AA populations, was included on the exome array. However, it was not associated with eGFRcrea using an additive genetics model in the AA participants here ($P = 0.70$).

To investigate if the newly identified eGFRcrea loci were also associated with diabetes mellitus and arterial hypertension, major risk factors for CKD, we tested for associations between the seven validated eGFRcrea loci with BP and type 2 diabetes (T2D) among EA participants in collaborations with the Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium (CHARGE) Blood Pressure¹³ and CHARGE Glycemia-T2D¹⁴ Working Groups. Consistent with prior observations, the majority of variants were not associated with BP traits or T2D (Supplemental Table 10). The exception was rs2472297 at the *CYP11A1* locus, a GWAS tag SNP previously associated with coffee/caffeine intake.^{7,8} This SNP was also associated with systolic and diastolic BP ($P = 7.4 \times 10^{-7}$, and 4.6×10^{-11} , respectively).¹³

DISCUSSION

Our main findings are four-fold. First, we identified and validated seven loci associated with eGFRcrea through genotyping on the exome-chip among 160,009 participants of EA in a two-stage study design. Second, the majority of the newly uncovered

associations were for common variants with modest effect sizes, which argues against the presence of rare protein-coding variants with large effect sizes represented on the exome chip. Third, we identified one novel association for *SOS2* through gene-based testing. Fourth, we demonstrated altered kidney development and function in zebrafish *sos2* and *acp1* knockdowns.

Our study emphasizes the continued success of efforts combining population-based genetics and model organisms to identify genes underlying kidney function.^{4,15} The zebrafish knockdowns provide a systematic model to examine the consequences of gene perturbations in the embryonic renal system of the fish. Our zebrafish morpholino experiments revealed a potential role for *sos2* and *acp1* in kidney development and function. Mutations in the *SOS* gene family (*SOS1* and *SOS2*) lead to Noonan syndrome, a congenital RASopathy (syndromes caused by germline mutations controlling signal transduction pathways) that can feature mild kidney

Table 2. Genes associated with eGFRcrea in EA participants from gene-based analyses meeting chip-wide significance thresholds ($P < 2.5 \times 10^{-6}$)

Gene ^a	Chr	cMAF	N Variants ^b	T1 ^c			SKAT ^d
				β	SEM	P Value	P Value
<i>LRP2</i>	2	0.070	38	0.003	0.002	6.7E-02	3.5E-7 ^e
<i>SLC47A1</i> ^f	17	0.033	4	-0.033	0.004	7.8E-15 ^e	3.4E-12 ^e
<i>SOS2</i> ^g	14	0.040	8	0.020	0.004	3.3E-06	5.4E-08 ^e

Chr, chromosome; cMAF, cumulative MAF used in analysis; SKAT, sequence kernel association test.

^aGene name.

^bNumber of variants used in analysis.

^cThe standard burden test collapses the variants with MAF < 1% into a single variable and tests the association between this variable with a phenotype.³⁰

^dThe SKAT aggregates individual variant score test statistics.³²

^eMeets chip-wide significant threshold, $P < 2.5 \times 10^{-6}$.

^fGene-based association results driven by one variant.

^gNovel gene.




Figure 2. *sos2* and *acp1* knockdowns result in defective kidney development. (A–D) Whole mount *in situ* hybridization in control embryos demonstrates normal expression of kidney markers, including *pax2a* (global kidney, A), *slc20a1a* (proximal tubules, B), and *slc12a3* (distal tubules, C) at 48 hpf, and *wt1a* (podocytes, D) at 24 hpf. (E–L) *sos2* and *acp1* ATG morpholino (MO) knockdown embryos develop glomerular gene expression defects (E, I, arrowheads) and display elongated proximal tubules (F, J). Knockdown of *acp1* shortened the distal tubules, whereas *sos2* knockdown left distal tubule *slc12a3* expression unaffected (G, K). No abnormalities in podocyte marker *wt1a* were observed for *sos2* ATG- and *acp1* ATG-MOs (H, L). (M) Quantitative assessment of proximal tubule length (*slc20a1a* expression) shows that proximal tubules are elongated in *sos2* ATG- and *acp1* ATG-MO injected embryos. *t* test used to calculate *P* values. (N) Table of observed abnormal embryos and total number examined by kidney markers *pax2a*, *wt1a*, and *slc12a3*, and MO-injected or control status. Fisher exact test used to calculate *P* values.

dysfunction¹⁶ and renal anomalies.^{17,18} The *SOS2* protein is expressed in the glomeruli and tubules of kidneys from adult humans.¹⁰ Together, these findings provide further evidence for a potential role for *SOS* proteins in kidney development and function. There are no prior reports of an association between kidney function and *ACPI*, a gene that encodes for an acid phosphatase involved in the immune response and found in erythrocytes.¹⁹ Our observations of kidney abnormalities in *sos2*- and *acp1*-knockdowns provide genes and target tissues for prioritization in future studies of more extensive functional follow-up, diagnostic screening, and potentially drug development.

Although morpholinos are an efficient tool for the rapid evaluation of GWAS hits, they have the potential for off-target effects, and morphant phenotypes are not always recapitulated in genetic mutant models. In this study, we evaluated *sos2* and *acp1* kidney development phenotypes using two independent morpholinos, which we believe adds confidence in the specificity of our phenotypes. Furthermore, we provided evidence that kidney developmental changes correlate with edema and

reduced dextran clearance from the blood. We believe that our morpholino screen has allowed us to clarify promising candidates for further study. Simultaneously, we acknowledge that future studies in genetic mutants will enhance and clarify these findings. Because genetic knockout techniques do not necessarily recapitulate exact features of the identified human variants, future experiments are also needed to evaluate the effect of specific variants on kidney development and function.

The majority of identified novel variants were common. The strongest single-variant association with eGFR_{crea} to date is the common variant rs13329952 at the *UMOD* locus with an effect size of 0.016 ln(ml/min per 1.73 m²) (MAF=0.19).¹⁵ Given our large discovery sample, our study was adequately powered (>80%) to detect effect sizes of 0.11–0.008 ln(ml/min per 1.73 m²) for very rare variants to more common variants (0.0005>MAF>0.10) (details of the power calculation can be found in the Supplemental Material). Although the selection of nonsynonymous content was expected to enrich for functional variants with large effect sizes, a possible reason for the lack of these findings might be the design-based,




Figure 3. *sos2* and *acp1* knockdowns result in altered kidney function. (A) *sos2* and *acp1* morphants develop edema, which is a sign of kidney failure in zebrafish. *sos2* morphants display severe global edema at 120 hpf, with fluid accumulation in the pericardium (black arrow) and intestinal tract (black star). At 96 hpf, *acp1* morphants have severe pericardial edema (black arrow). (B–C) Incidence of edema and embryonic lethality in *acp1* and *sos2* morphants with Fisher exact test. *Indicates P value <0.05 ; **indicates P value <0.01 ; ***indicates P value <0.001 . (D–L) Embryos were injected with control, *sos2*, or *acp1* morpholino at the single cell stage and subsequently injected with 70 kDa molecular mass fluorescent rhodamine dextran at 72 hpf. Dextran-injected embryos show equal loading at 2 hpi. Compared with control embryos, *sos2* and *acp1* MO injected embryos have reduced dextran clearance in the cardiovascular region over time. (J–L) Morphant embryos have altered convoluted tubule morphology at 120 hpf (48 hpi) (white arrows). (M) Dextran fluorescence intensity over time as normalized to starting fluorescence intensity.

limited coverage of rare variants on the exome chip. The exome chip was primarily designed to assess nonsynonymous rare variants that had been observed among approximately 12,000 sequenced participants that were not selected on the basis of kidney disease (http://genome.sph.umich.edu/wiki/Exome_Chip_Design), and thus, we would not expect kidney disease-causing mutations to be well represented on the exome chip. Although the convenience and low cost of chip-based genotyping arrays focused on exonic variants were

influential in facilitating a large number of cohorts to participate, as is necessary for a well powered study, the limited coverage of the exome affected the ability of both single-variant and gene-based tests to assess all exonic variation in association with kidney function. Thus, we cannot rule out *bona fide* rare variant associations with eGFR_{crea}. Large-scale whole exome or whole genome sequencing will be able to adequately address the unresolved issue of the contribution of rare variants to the variation of kidney function in the general population. Because of limited

sample size in the diabetes stratum compared with the nondiabetes stratum, we did not have adequate power to assess unique genetic associations in the diabetes stratified analysis. We did not implement a random effects meta-analysis as an alternative screening procedure for novel loci because the between-study heterogeneity was small. The analysis module that we used cannot correctly account for the different allele dosage between women and men and therefore we were unable to implement association analysis for chromosome X variants. Although our study supports the observation that gene-based analyses aggregating rare nonsynonymous variants are able to identify new loci in association with common complex phenotypes^{14,20–22} and can additionally uncover new rare missense variants within known loci, the number of loci identified through gene-based methods remains a small minority of the findings compared with single-variant results. Finally, the zebrafish knockdowns helped us to screen novel loci that appear to have a role in kidney development or function. However, extrapolation of the relationship between the novel loci among patients with more advanced CKD remains to be determined; follow up studies are needed to assess the association of these loci with ESRD and incident CKD.

In summary, we identified eight novel loci (seven common single variants and one gene with multiple rare coding variants) associated with kidney function. Functional experiments in zebrafish highlighted potential roles for *SOS2* and *ACPI* in embryonic kidney development. Future whole exome and whole genome sequencing studies will be needed to assess the full spectrum of rare genetic variants on kidney function.

CONCISE METHODS

Study Participants

Across all traits analyzed, a total of 120,357 participants from 27 studies of EA were included in stage 1 of this study. An additional 48,343 participants from 12 studies of EA were included in stage 2. A total of 11,386 participants from seven studies of AA were also included in separate analyses. Study-specific characteristics are summarized in Supplemental Tables 1, 2, and 4. All participants provided informed consent and each study was approved by its governing ethics committee or Institutional Review Board.

Phenotype Definitions

Serum creatinine was measured in each study as described in the Study-Specific Methods section in the Supplemental Material, and calibrated to the National Health and Nutrition Examination Study data to account for between-laboratory variation.^{23,24} eGFR_{crea} was estimated using the four-variable Modification of Diet in Renal Disease (MDRD) Study Equation.²⁵ Cystatin C, an alternative biomarker of kidney function, was measured in a sub-set of participating studies. eGFR_{cys} was estimated as $76.7 \times (\text{serum cystatin C})$.²⁵ All eGFR_{crea} and eGFR_{cys} values <15 ml/min per 1.73m^2 were set to 15, and those >200 ml/min per 1.73m^2 were set to 200 to avoid undue influence from outliers. UACR was defined as urinary albumin (mg/L)/urinary creatinine (mg/dl) $\times 100$. All analyzed traits (eGFR_{crea},

eGFR_{cys}, and UACR) were natural log (ln)-transformed. Diabetes was defined as fasting glucose ≥ 126 mg/dl, pharmacologic treatment for diabetes, or by self-report. Hypertension was defined as systolic BP ≥ 140 mmHg, diastolic BP ≥ 90 mmHg, or pharmacologic treatment for hypertension.¹⁵

Genotypes

Genotyping was conducted in each study using the Illumina Human Exome BeadChip (http://genome.sph.umich.edu/wiki/Exome_Chip_Design). This genotyping array containing 247,870 markers focuses on exonic variants discovered through exome sequencing of approximately 12,000 individuals. Illumina's GenTrain version 2.0 clustering algorithm in GenomeStudio, zCall,²⁶ or a combination of both procedures were used to call genotypes. To improve genotype calling of low frequency and rare variants, genotypes from eight of the contributing cohorts (Atherosclerosis Risk in Communities [ARIC], Age, Gene/Environment Susceptibility-Reykjavik Study [AGES], Cardiovascular Health Study [CHS], Framingham Heart Study [FHS], Rotterdam Study [RS], Health, Aging and Body Composition Study [Health ABC], Family Heart Study [FamHS], and Jackson Heart Study [JHS]) were jointly clustered and called *via* the CHARGE Consortium algorithm.²⁷ Other participating cohorts were called individually. Among them, CROATIA-Korcula and Generation Scotland (GS) applied the cluster file from the CHARGE Consortium for genotype calling. Details regarding genotyping and quality control within each study are summarized in Supplemental Table 2.

Statistical Methods for Stage 1

By following a centralized analysis plan, each study performed two sets of analyses: single-variant analysis and gene-based analysis. The primary meta-analyses were focused on the EA population and a secondary set of meta-analyses were focused on the AA population due to a substantially smaller sample size. Where not specified otherwise, R software was used for data management, statistical analyses, and graphing.²⁸

Single-Variant Analysis

Each study performed association analyses of the following phenotypes and models: (1) ln-transformed eGFR_{crea}, (2) ln-transformed eGFR_{cys}, (3) ln-transformed UACR, and (4) ln-transformed eGFR_{crea} stratified by diabetes status. These association analyses were based on linear regression models adjusting for age, sex, study site (if applicable), family structure (if applicable), and the first ten principal components to control for population stratification. All analyses were performed assuming an additive genetic effect and all analyses were stratified by ancestry.

For single-variant meta-analysis, study-specific results were combined for each trait in a fixed-effects model using METAL.²⁹ We restricted single-variant meta-analyses to (1) autosomal variants, (2) polymorphic variants, (3) variants existing in the joint calling effort within the CHARGE consortium²⁷, and (4) with minor allele count ≥ 20 across all cohorts in each ancestry group. Bonferroni correction for the number of variants tested within each ancestry was used to set the significance threshold for each analysis (chip-wide significance), corresponding to $P < 3.7 \times 10^{-7}$ (0.05/134299 variants) for EA analysis and $P < 5.5 \times 10^{-7}$ (0.05/91187 variants) for AA analysis.

Gene-Based Analysis

Single-variant analysis methods have limited power to detect association for rare variants. Recently developed gene-based methods, where defined variants contained in a gene region are collapsed into one unit for analysis, provide additional statistical power³⁰ to investigate the role of rare variants on kidney function traits.

For gene-based meta-analysis, study-specific results were combined using the seqMeta package for R.³¹ The same phenotypes and covariates were used as for single-variant testing and analyses were again stratified by ancestry. We used two gene-based tests for aggregated analysis of rare single nucleotide variants (SNVs): (1) T1,³⁰ which is more powerful when all variants within the gene region affect the phenotype in the same direction; and (2) the sequence kernel association test,³² which allows for bidirectional effects and is more powerful when there are both protective and deleterious variants within the same gene. Both gene-based tests were restricted to variants with MAF < 1% and variants likely to exert a major effect on the gene product (stop gain/loss, nonsynonymous, or splice-site variants on the basis of annotation with dbNSFP [v.2.0]).³³ Genes containing at least two variants with a cumulative minor allele count ≥ 20 ²⁰ were included in the analysis. In total, we tested 9990 autosomal genes meeting our established thresholds and filters. Bonferroni correction for the number of genes and tests performed was used to set the significance threshold for the gene-based analysis corresponding to $P < 2.5 \times 10^{-6}$ (0.05/19,922 tests [996 genes \times 2 tests]) in EA analysis and $P < 2 \times 10^{-6}$ (25,378 tests [12,689 genes \times 2 tests]) in AA analysis.

Statistical Methods for Stage 2

Chip-wide significant results from single-variant meta-analyses were brought forward for testing in stage 2 where each new study followed the same methodology as stage 1. Details regarding the genotyping and population characteristics of each cohort can be found in Supplemental Tables 2 and 4.

Study-specific results from stage 2 cohorts were combined using the same meta-analysis approach and software as in stage 1 (fixed-effects model in METAL²⁹). Criteria for validation were: (1) a direction of effect consistent with stage 1 analysis, (2) a one-sided P value < 0.05 from stage 2 analysis, and (3) $P < 3.7 \times 10^{-7}$ from combined stage 1 and stage 2 analysis.

The percentage of phenotypic variance explained by each novel locus was estimated as $R^2 = \beta^2 \text{var}(\text{SNP}) / \text{var}(\ln[\text{eGFR}_{\text{crea}}])$, where β is the estimated effect of the SNP on $\ln(\text{eGFR}_{\text{crea}})$, and $\text{var}(\text{SNP}) = 2 * \text{MAF}_{\text{SNP}} * (1 - \text{MAF}_{\text{SNP}})$. $\text{Var}(\ln[\text{eGFR}_{\text{crea}}])$ was estimated in the ARIC study. All loci were assumed to have independent effects on the phenotype.

Associations with Other Traits

To examine potential associations of the novel eGFR-associated variants with other correlated traits and conditions, we performed external look ups for systolic and diastolic BP in collaboration with the CHARGE Blood Pressure Working Group ($n = 145,872$)¹³ and for T2D in collaboration with the CHARGE Glycemia-T2D Working Group ($n = 10,240$ T2D cases and 63,105 controls)²⁰ among EA participants.

NHGRI GWAS Catalog and PubMed Queries

For all newly identified and validated variants, we interrogated the NHGRI GWAS Catalog (<https://www.ebi.ac.uk/gwas/>; 10/12/15) for the lead SNP at each locus and for SNPs in linkage disequilibrium

with the lead SNP (within 1 Mbp and $r^2 > 0.5$ from 1000 Genomes Pilot in CEU; <http://www.broadinstitute.org/mpg/snp/>) to assess association with other traits (see Supplemental Table 11 for full listing of GWAS Catalog associations). Additionally, for each locus we searched PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>; 10/12/15) for publications on kidney/renal function or CKD.

Functional Studies in Zebrafish

To investigate a potential role of the newly identified genes during kidney development, we assessed the functional consequences of gene knockdown in zebrafish embryos. We used antisense MO technology to knock down genes identified on the basis of validated associations for nonsynonymous variants and for novel gene-based loci. Two independent MO probes were used for each gene. Zebrafish were maintained in accordance with established Institutional Animal Care and Use Committee protocols.

Morpholinos (Gene Tools) were designed against zebrafish target genes. Morpholino sequences are as follows: *sos2* (5' GCACCGGGAA-CAACCACACAACCTTT 3'), *sos2* (exon 2) (5' CCTGCACCTATAAACA-CAGAATAGA 3'), *ppm1j* (5' AATTTGTGACATCAGCGGCACGGTA 3'), *acp1* ATG (5' TCCGCTGGAAGCCGCCATATTGGTC 3'), *acp1* (exon 1) (5' TATAGCATTTTCTTACCCAAGCACAC 3'), *spg* ATG (5' TCTTCTCTTCAGTAACCTTTCTCAT 3'), *edem3* (exon 1) (5' AGTCCTCACACAGACACATACCTCA 3'), and *eya4* ATG (5' CAGATCCTGTGTATTCTCCATCAGT 3'). Zebrafish embryos were injected with various concentrations of MO (*sos2* ATG–150 μM , *sos2* (exon 2)–400 μM , *ppm1j* ATG, *acp1* ATG, and *acp1* (exon 1)–400 μM , *edem3* ATG–300 μM , *eya4* ATG–400 μM , *spg* ATG–100 μM) at the one-cell stage. We fixed embryos in 4% paraformaldehyde at relevant developmental stages for analysis by *in situ* hybridization (<http://zfin.org/ZFIN/Methods/ThisseProtocol.html>). Distinct pronephros (embryonic kidney) structures were visualized using a series of established markers: *pax2a* (global), *wt1a* (podocyte), *slc20a1a* (proximal tubule), and *slc12a3* (distal tubule). *acp1* probe was generated from zebrafish cDNA using the following primers: 5' TGGAGAATAGACAGTGCCGC 3' (forward 1) and 5' TTITCAGCTGCTTGCCITC 3' (reverse 1), and 5' GTGGAGAATA-GACAGTGCCG 3' (forward 2) and 5' CAGGAAGGCTTTGCATC 3' (reverse 2). *sos2* probe was generated from zebrafish cDNA using the following primers: 5' GTGTTCGAGGAAGGAGCACA 3' (forward) and 5' TGATGTTCCACCCACTGACG 3' (reverse).

Abnormal gene expression patterns were identified by direct comparison to control embryos that were injected with a standard control MO designed by GeneTools (SynGene, Cambridge, UK). Developmental phenotypes were scored by two independent researchers. Fisher exact tests were used to test for normal and abnormal embryonic phenotypes for the *pax2a*, *wt1a*, and *slc12a3* markers and t test was used to test for differences in proximal tubule length for the *slc20a1a* marker; $P < 0.05$ was set as a threshold for statistical significance. To evaluate proximal tubule length, the distance between the most anterior and the most posterior tip of the right proximal tubule (from standardized dorsal-view images) was measured using the imageJ measurement tool. Fisher exact test was used to test for differences in edema prevalence; $P < 0.05$ was set as a threshold for statistical significance.

Dextran clearance experiments were performed following a previously described protocol.¹² Seventy-two hours after morpholino

injection, embryos were anesthetized in a 1:20 dilution of 4 mg/ml Tricaine in embryo water and placed on a 2% agarose injection mold. An equal volume of tetramethylrhodamine dextran (70 kDa molecular mass; Invitrogen, Carlsbad, CA) was injected into the cardiac sinus venosus of each embryo, and individual embryos were sorted into designated wells for timelapse imaging. Fluorescent microscopy images were taken at 2 hpi (74 hpf), 24 hpi (96 hpf), and 48 hpi (120 hpf) for each sorted embryo to assess loading fluorescence and the dextran clearance over time. Fluorescence intensity in the cardiac region was measured as the mean grayscale value using Image J as previously described.³⁴ Remaining fluorescence intensity at each time point was normalized to the starting intensity and plotted as a percentage of the initial fluorescence intensity. To evaluate edema, morpholinos were injected into the single-cell stage embryo and embryos were examined every 24 hours for evidence of edema. The number of affected embryos was recorded as a fraction of total number of injected embryos.

Power Calculation

Power for association was evaluated for eGFR_{crea} assuming a mean of 4.5 ln(ml/min per 1.73 m²) with standard deviation of 0.2 ln(ml/min per 1.73 m²), estimates from the EA samples in the ARIC study, using QUANTO power calculator, version 1.2.4 (<http://biostats.usc.edu/Quanto.html>) at the significance level of 3.7×10^{-7} for a variant with MAF of 0.1, 0.05, 0.03, 0.01, 0.005, 0.001, or 0.0005. In stage 1 with sample size $n=111,666$, there was at least 80% power to detect effect sizes of 0.008, 0.011, 0.015, 0.026, 0.036, 0.08, or 0.11 ln(ml/min/1.73 m²), respectively. For stage 2, using one-sided tests, there was 92% power to detect an effect size of 0.018 for a variant with MAF of 0.02 (minimum MAF among all eight SNPs tested) on the basis of 48,343 samples in stage 2 and correcting for the eight SNPs tested.

ACKNOWLEDGMENTS

The views expressed in this manuscript are those of the authors and do not necessarily represent the views of the National Heart, Lung, and Blood Institute, the National Institutes of Health, or the US Department of Health and Human Services. Acknowledgments and funding information for all contributing studies are presented in the Supplemental Material. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The full list of consortium members is provided in the supplemental material.

DISCLOSURES

Caroline S. Fox and Audrey Y. Chu are employed by Merck Research Laboratories as of December 14, 2015 and July 18, 2016, respectively.

REFERENCES

1. Satko SG, Sedor JR, Iyengar SK, Freedman BI: Familial clustering of chronic kidney disease. *Semin Dial* 20: 229–236, 2007

- Böger CA, Chen MH, Tin A, Olden M, Köttgen A, de Boer IH, Fuchsberger C, O'Seaghdha CM, Pattaro C, Teumer A, Liu CT, Glazer NL, Li M, O'Connell JR, Tanaka T, Peralta CA, Kutalik Z, Luan J, Zhao JH, Hwang SJ, Akyzbekova E, Kramer H, van der Harst P, Smith AV, Lohman K, de Andrade M, Hayward C, Kollerits B, Tönjes A, Aspelund T, Ingelsson E, Eiriksdóttir G, Launer LJ, Harris TB, Shuldiner AR, Mitchell BD, Arking DE, Franceschini N, Boerwinkle E, Egan J, Hernandez D, Reilly M, Townsend RR, Lumley T, Siscovick DS, Psaty BM, Kestenbaum B, Haritunians T, Bergmann S, Vollenweider P, Waeber G, Mooser V, Waterworth D, Johnson AD, Florez JC, Meigs JB, Lu X, Turner ST, Atkinson EJ, Leak TS, Aasærd K, Skorpen F, Syvänen AC, Illig T, Baumert J, Koenig W, Krämer BK, Devuyst O, Mychaleckyj JC, Minelli C, Bakker SJ, Kedenko L, Paulweber B, Coassin S, Endlich K, Kroemer HK, Biffar R, Stracke S, Völzke H, Stumvoll M, Mägi R, Campbell H, Vitart V, Hastie ND, Gudnason V, Kardina SL, Liu Y, Polasek O, Curhan G, Kronenberg F, Prokopenko I, Rudan I, Arnlöv J, Hallan S, Navis G, Parsa A, Ferrucci L, Coresh J, Shlipak MG, Bull SB, Paterson NJ, Wichmann HE, Wareham NJ, Loos RJ, Rotter JI, Pramstaller PP, Cupples LA, Beckmann JS, Yang Q, Heid IM, Rettig R, Dreisbach AW, Bochud M, Fox CS, Kao WH, Kao WH: CKDGen Consortium: CUBN is a gene locus for albuminuria. *J Am Soc Nephrol* 22: 555–570, 2011
- Köttgen A, Pattaro C, Böger CA, Fuchsberger C, Olden M, Glazer NL, Parsa A, Gao X, Yang Q, Smith AV, O'Connell JR, Li M, Schmidt H, Tanaka T, Isaacs A, Ketkar S, Hwang SJ, Johnson AD, Dehghan A, Teumer A, Paré G, Atkinson EJ, Zeller T, Lohman K, Cornelis MC, Probst-Hensch NM, Kronenberg F, Tönjes A, Hayward C, Aspelund T, Eiriksdóttir G, Launer LJ, Harris TB, Rumpusaud E, Mitchell BD, Arking DE, Boerwinkle E, Struchalin M, Cavalieri M, Singleton A, Giallauria F, Metter J, de Boer IH, Haritunians T, Lumley T, Siscovick D, Psaty BM, Zillikens MC, Oostra BA, Feitosa M, Province M, de Andrade M, Turner ST, Schillert A, Ziegler A, Wild PS, Schnabel RB, Wilde S, Munzel TF, Leak TS, Illig T, Klopp N, Meisinger C, Wichmann HE, Koenig W, Zgaga L, Zemunik T, Kolcic I, Minelli C, Hu FB, Johansson A, Igl W, Zabolj G, Wild SH, Wright AF, Campbell H, Ellinghaus D, Schreiber S, Aulchenko YS, Felix JF, Rivadeneira F, Uitterlinden AG, Hofman A, Imboden M, Nitsch D, Brandstätter A, Kollerits B, Kedenko L, Mägi R, Stumvoll M, Kovacs P, Boban M, Campbell S, Endlich K, Völzke H, Kroemer HK, Nauck M, Völker U, Polasek O, Vitart V, Badola S, Parker AN, Ridker PM, Kardina SL, Blankenberg S, Liu Y, Curhan GC, Franke A, Roach T, Paulweber B, Prokopenko I, Wang W, Gudnason V, Shuldiner AR, Coresh J, Schmidt R, Ferrucci L, Shlipak MG, van Duijn CM, Borecki I, Krämer BK, Rudan I, Gyllenstein U, Wilson JF, Witteman JC, Pramstaller PP, Rettig R, Hastie N, Chasman DI, Kao WH, Heid IM, Fox CS: New loci associated with kidney function and chronic kidney disease. *Nat Genet* 42: 376–384, 2010
- Pattaro C, Köttgen A, Teumer A, Garnaas M, Böger CA, Fuchsberger C, Olden M, Chen MH, Tin A, Taliun D, Li M, Gao X, Gorski M, Yang Q, Hundertmark C, Foster MC, O'Seaghdha CM, Glazer N, Isaacs A, Liu CT, Smith AV, O'Connell JR, Struchalin M, Tanaka T, Li G, Johnson AD, Gierman HJ, Feitosa M, Hwang SJ, Atkinson EJ, Lohman K, Cornelis MC, Johansson Å, Tönjes A, Dehghan A, Chouraki V, Holliday EG, Sorice R, Kutalik Z, Lehtimäki T, Esko T, Deshmukh H, Ulivi S, Chu AY, Murgja F, Trompet S, Imboden M, Kollerits B, Pistis G, Harris TB, Launer LJ, Aspelund T, Eiriksdóttir G, Mitchell BD, Boerwinkle E, Schmidt H, Cavalieri M, Rao M, Hu FB, Demirkan A, Oostra BA, de Andrade M, Turner ST, Ding J, Andrews JS, Freedman BI, Koenig W, Illig T, Döring A, Wichmann HE, Kolcic I, Zemunik T, Boban M, Minelli C, Wheeler HE, Igl W, Zabolj G, Wild SH, Wright AF, Campbell H, Ellinghaus D, Nöthlings U, Jacobs G, Biffar R, Endlich K, Ernst F, Homuth G, Kroemer HK, Nauck M, Stracke S, Völker U, Völzke H, Kovacs P, Stumvoll M, Mägi R, Hofman A, Uitterlinden AG, Rivadeneira F, Aulchenko YS, Polasek O, Hastie N, Vitart V, Helmer C, Wang JJ, Ruggiero D, Bergmann S, Kähönen M, Viikari J, Nikopoulou T, Province M, Ketkar S, Colhoun H, Doney A, Robino A, Giulianini F, Krämer BK, Portas L,

- Ford I, Buckley BM, Adam M, Thun GA, Paulweber B, Haun M, Sala C, Metzger M, Mitchell P, Ciullo M, Kim SK, Vollenweider P, Raitakari O, Metspalu A, Palmer C, Gasparini P, Pirastu M, Jukema JW, Probst-Hensch NM, Kronenberg F, Toniolo D, Gudnason V, Shuldiner AR, Coresh J, Schmidt R, Ferrucci L, Siscovick DS, van Duijn CM, Borecki I, Kardia SL, Liu Y, Curhan GC, Rudan I, Gyllensten U, Wilson JF, Franke A, Pramstaller PP, Rettig R, Prokopenko I, Witteman JC, Hayward C, Ridker P, Parsa A, Bochud M, Heid IM, Goessling W, Chasman DI, Kao WH, Fox CS, Goessling W, Chasman DI, Kao WH, Fox CS; CARDIoGRAM Consortium; ICBP Consortium; CARE Consortium; Wellcome Trust Case Control Consortium 2 (WTCCC2): Genome-wide association and functional follow-up reveals new loci for kidney function. *PLoS Genet* 8: e1002584, 2012
5. Okada Y, Sim X, Go MJ, Wu JY, Gu D, Takeuchi F, Takahashi A, Maeda S, Tsunoda T, Chen P, Lim SC, Wong TY, Liu J, Young TL, Aung T, Seielstad M, Teo YY, Kim YJ, Lee JY, Han BG, Kang D, Chen CH, Tsai FJ, Chang LC, Fann SJ, Mei H, Rao DC, Hixson JE, Chen S, Katsuya T, Isono M, Ogihara T, Chambers JC, Zhang W, Kooner JS, Albrecht E, Yamamoto K, Kubo M, Nakamura Y, Kamatani N, Kato N, He J, Chen YT, Cho YS, Tai ES, Tanaka T; KidneyGen Consortium; CKDGen Consortium; GUGC consortium: Meta-analysis identifies multiple loci associated with kidney function-related traits in east Asian populations. *Nat Genet* 44: 904–909, 2012
 6. Pattaro C, Teumer A, Gorski M, Chu AY, Li M, Mijatovic V, Garnaas M, Tin A, Sorice R, Li Y, Taliun D, Olden M, Foster M, Yang Q, Chen M-H, Pers TH, Johnson AD, Ko Y-A, Fuchsberger C, Tayo B, Nalls M, Feitosa MF, Isaacs A, Dehghan A, d'Adamo P, Adeyemo A, Dieffenbach AK, Zonderman AB, Nolte IM, van der Most, PJ, Wright, AF, Shuldiner, AR, Morrison, AC, Hofman, A, Smith, AV, Dreisbach, AW, Franke, A, Uitterlinden, AG, Metspalu, A, Tonjes, A, Lupo, A, Robino, A, Johansson, A, Demirkan, A, Kollerits, B, Freedman, BI, Ponte, B, Oostra, BA, Paulweber, B, Kramer, BK, Mitchell, BD, Buckley, BM, Peralta, CA, Hayward, C, Helmer, C, Rotimi, CN, Shaffer, CM, Muller, C, Sala, C, van Duijn, CM, Saint-Pierre, A, Ackermann, D, Shriner, D, Ruggiero, D, Toniolo, D, Lu, Y, Cusi, D, Czamara, D, Ellinghaus, D, Siscovick, DS, Ruderfer, D, Gieger, C, Grallert, H, Rohtchina, E, Atkinson, EJ, Holliday, EG, Boerwinkle, E, Salvi, E, Bottinger, EP, Murgia, F, Rivadeneira, F, Ernst, F, Kronenberg, F, Hu, FB, Navis, GJ, Curhan, GC, Ehret, GB, Homuth, G, Coassin, S, Thun, G-A, Pistis, G, Gambaro, G, Malerba, G, Montgomery, GW, Eiriksdottir, G, Jacobs, G, Li, G, Wichmann, HE, Campbell, H, Schmidt, H, Wallaschofski, H, Volzke, H, Brenner, H, Kroemer, HK, Kramer, H, Lin, H, Leach, IM, Ford, I, Guessous, I, Rudan, I, Prokopenko, I, Borecki, I, Heid, IM, Kolcic, I, Persico, I, Jukema, JW, Wilson, JF, Felix, JF, Divers, J, Lambert, J-C, Stafford, JM, Gaspoz, J-M, Smith, JA, Faul, JD, Wang, JJ, Ding, J, Hirschhorn, JN, Attia, J, Whitfield, JB, Chalmers, J, Viikari, J, Coresh, J, Denny, JC, Karjalainen, J, Fernandes, JK, Endlich, K, Butterbach, K, Keene, KL, Lohman, K, Portas, L, Launer, LJ, Lytikainen, L-P, Yengo, L, Franke, L, Ferrucci, L, Rose, LM, Kedenko, L, Rao, M, Struchalin, M, Kleber, ME, Cavalieri, M, Haun, M, Cornelis, MC, Ciullo, M, Pirastu, M, de Andrade, M, McEvoy, MA, Woodward, M, Adam, M, Cocca, M, Nauck, M, Imboden, M, Waldenberger, M, Pruijm, M, Metzger, M, Stumvoll, M, Evans, MK, Sale, MM, Kahonen, M, Boban, M, Bochud, M, Rheinberger, M, Verweij, N, Bouatia-Naji, N, Martin, NG, Hastie, N, Probst-Hensch, N, Soranzo, N, Devuyst, O, Raitakari, O, Gottesman, O, Franco, OH, Polasek, O, Gasparini, P, Munroe, PB, Ridker, PM, Mitchell, P, Muntner, P, Meisinger, C, Smit, JH, Consortium, I, Consortium, A, Cardiogram, Group, CH-HF, Consortium, EC, Kovacs, P, Wild, PS, Froguel, P, Rettig, R, Magi, R, Biffar, R, Schmidt, R, Middelberg, RPS, Carroll, RJ, Penninx, BW, Scott, RJ, Katz, R, Sedaghat, S, Wild, SH, Kardia, SLR, Ulivi, S, Hwang, S-J, Enroth, S, Kloiber, S, Trompet, S, Stengel, B, Hancock, SJ, Turner, ST, Rosas, SE, Stracke, S, Harris, TB, Zeller, T, Zemunik, T, Lehtimäki, T, Illig, T, Aspelund, T, Nikopoulou, T, Esko, T, Tanaka, T, Gyllensten, U, Volker, U, Emilsson, V, Vitart, V, Aalto, V, Gudnason, V, Chouraki, V, Chen, W-M, Igl, W, Marz, W, Koenig, W, Lieb, W, Loos, RJF, Liu, Y, Snieder, H, Pramstaller, PP, Parsa, A, O'Connell, JR, Susztak, K, Hamet, P, Tremblay, J, de Boer, IH, Boger, CA, Goessling, W, Chasman, DI, Kottgen, A, Kao, WH, Fox, CS: Genetic associations at 53 loci highlight cell types and biological pathways relevant for kidney function. *Nat Commun* 7: 10023, 2016
 7. Cornelis MC, Monda KL, Yu K, Paynter N, Azzato EM, Bennett SN, Berndt SI, Boerwinkle E, Chanock S, Chatterjee N, Couper D, Curhan G, Heiss G, Hu FB, Hunter DJ, Jacobs K, Jensen MK, Kraft P, Landi MT, Nettleton JA, Purdue MP, Rajaraman P, Rimm EB, Rose LM, Rothman N, Silverman D, Stolzenberg-Solomon R, Subar A, Yeager M, Chasman DI, van Dam RM, Caporaso NE: Genome-Wide Meta-Analysis Identifies Regions on 7p21 AHR and 15q24 CYP1A2 As Determinants of Habitual Caffeine Consumption. *PLoS Genet* 7: e1002033, 2011
 8. Sulem P, Gudbjartsson DF, Geller F, Prokopenko I, Feenstra B, Aben KKH, Franke B, den Heijer M, Kovacs P, Stumvoll M, Mägi R, Yanek LR, Becker LC, Boyd HA, Stacey SN, Walters GB, Jonasdottir A, Thorleifsson G, Holm H, Gudjonsson SA, Rafnar T, Björnsson G, Becker DM, Melbye M, Kong A, Tönjes A, Thorgeirsson T, Thorsteinsdottir U, Kiemenev LA, Stefansson K: Sequence variants at CYP1A1-CYP1A2 and AHR associate with coffee consumption. *Hum Mol Genet* 20: 2071–2077, 2011
 9. Imielinski M, Baldassano RN, Griffiths A, Russell RK, Annese V, Dubinsky M, Kugathasan S, Bradfield JP, Walters TD, Sleiman P, Kim CE, Muise A, Wang K, Glessner JT, Saeed S, Zhang H, Frackelton EC, Hou C, Flory JH, Otieno G, Chiavacci RM, Grundmeier R, Castro M, Latiano A, Dallapiccola B, Stempak J, Abrams DJ, Taylor K, McGovern D, Silber G, Wrobel I, Quiros A, Barrett JC, Hansoul S, Nicolae DL, Cho JH, Duerr RH, Rioux JD, Brant SR, Silverberg MS, Taylor KD, Barnuda MM, Bitton A, Dassopoulos T, Datta LW, Green T, Griffiths AM, Kistner EO, Murtha MT, Regueiro MD, Rotter JJ, Schumm LP, Steinhard AH, Targor SR, Xavier RJ, Libioulle C, Sandor C, Lathrop M, Belaiche J, Dewit O, Gut I, Heath S, Laukens D, Mni M, Rutgeerts P, Van Gossum A, Zelenika D, Franchimont D, Hugot JP, de Vos M, Vermeire S, Louis E, Cardon LR, Anderson CA, Drummond H, Nimmo E, Ahmad T, Prescott NJ, Onnie CM, Fisher SA, Marchini J, Ghori J, Bumpstead S, Gwillam R, Tremelling M, Delukas P, Mansfield J, Jewell D, Satsangi J, Mathew CG, Parkes M, Georges M, Daly MJ, Heyman MB, Ferry GD, Kirschner B, Lee J, Essers J, Grand R, Stephens M, Levine A, Piccoli D, Van Limbergen J, Cucchiara S, Monos DS, Guthery SL, Denson L, Wilson DC, Grant SF, Daly M, Silverberg MS, Satsangi J, Hakonarson H; Western Regional Alliance for Pediatric IBD; International IBD Genetics Consortium; NIDDK IBD Genetics Consortium; Belgian-French IBD Consortium; Wellcome Trust Case Control Consortium: Common variants at five new loci associated with early-onset inflammatory bowel disease. *Nat Genet* 41: 1335–1340, 2009
 10. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, Olsson I, Edlund K, Lundberg E, Navani S, Szgyarto CA, Odeberg J, Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist PH, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P, Schwenk JM, Hamsten M, von Feilitzen K, Forsberg M, Persson L, Johansson F, Zwahlen M, von Heijne G, Nielsen J, Pontén F: Proteomics. Tissue-based map of the human proteome. *Science* 347: 1260419, 2015
 11. Hanke N, Staggs L, Schroder P, Litteral J, Fleig S, Kaufeld J, Pauli C, Haller H, Schiffer M: "Zebrafishing" for novel genes relevant to the glomerular filtration barrier. *BioMed Res Int* 2013: 658270, 2013
 12. Hentschel DM, Mengel M, Boehme L, Liebsch F, Albertin C, Bonventre JV, Haller H, Schiffer M: Rapid screening of glomerular slit diaphragm integrity in larval zebrafish. *Am J Physiol Renal Physiol* 293: F1746–F1750, 2007
 13. Liu C, Kraja AT, Smith JA, Brody JA, Franceschini N, Bis JC, Rice K, Morrison AC, Lu Y, Weiss S, Guo X, Palmas W, Martin LW, Chen YD, Surendran P, Drenos F, Cook JP, Auer PL, Chu AY, Giri A, Zhao W, Jakobsdottir J, Lin LA, Stafford JM, Amin N, Mei H, Yao J, Voorman A, Larson MG, Grove ML, Smith AV, Hwang SJ, Chen H, Huan T, Kosova G, Stitzel NO, Kathiresan S, Samani N, Schunkert H, Deloukas P, Li M,

- Fuchsberger C, Pattaro C, Gorski M, Kooperberg C, Papanicolaou GJ, Rossouw JE, Faul JD, Kardina SL, Bouchard C, Raffel LJ, Uitterlinden AG, Franco OH, Vasani RS, O'Donnell CJ, Taylor KD, Liu K, Bottinger EP, Gottesman O, Daw EW, Giulianini F, Ganesh S, Salfati E, Harris TB, Launer LJ, Dorr M, Felix SB, Rettig R, Volzke H, Kim E, Lee WJ, Lee IT, Sheu WH, Tsoie KS, Edwards DR, Liu Y, Correa A, Weir DR, Volker U, Ridker PM, Boerwinkle E, Gudnason V, Reiner AP, van Duijn CM, Borecki IB, Edwards TL, Chakravarti A, Rotter JJ, Psaty BM, Loos RJ, Fornage M, Ehret GB, Newton-Cheh C, Levy D, Chasman DI: Meta-analysis identifies common and rare variants influencing blood pressure and overlapping with metabolic trait loci. *Nat Genet* 48: 1162–1170, 2016
14. Wessel J, Chu AY, Willems SM, Wang S, Yaghoobkar H, Brody JA, Dauriz M, Hivert M-F, Raghavan S, Lipovich L, Hidalgo B, Fox K, Huffman JE, An P, Lu Y, Rasmussen-Torvik LJ, Grarup N, Ehm MG, Li L, Baldridge AS, Stancakova A, Abrol R, Besse C, Boland A, Bork-Jensen J, Fornage M, Freitag DF, Garcia ME, Guo X, Hara K, Isaacs A, Jakobsdottir J, Lange LA, Layton JC, Li M, Zhao H: J, Meidtner, K, Morrison, AC, Nalls, MA, Peters, MJ, Sabater-Lleal, M, Schurmann, C, Silveira, A, Smith, AV, Southam, L, Stoiber, MH, Strawbridge, RJ, Taylor, KD, Varga, TV, Allin, KH, Amin, N, Aponte, JL, Aung, T, Barbieri, C, Bihlmeyer, NA, Boehnke, M, Bombieri, C, Bowden, DW, Burns, SM, Chen, Y, Chen, Y-D, Cheng, C-Y, Correa, A, Czajkowski, J, Dehghan, A, Ehret, GB, Eiriksdottir, G, Escher, SA, Farmaki, A-E, Franberg, M, Gambaro, G, Giulianini, F, Goddard, WA, Goel, A, Gottesman, O, Grove, ML, Gustafsson, S, Hai, Y, Hallmans, G, Heo, J, Hoffmann, P, Ikram, MK, Jensen, RA, Jorgensen, ME, Jorgensen, T, Karaleftheri, M, Khor, CC, Kirkpatrick, A, Kraja, AT, Kuusisto, J, Lange, EM, Lee, JT, Lee, W-J, Leong, A, Liao, J, Liu, C, Liu, Y, Lindgren, CM, Linneberg, A, Malerba, G, Mamakou, V, Marouli, E, Maruthur, NM, Matchan, A, McKean-Cowdin, R, McLeod, O, Metcalf, GA, Mohlke, KL, Muzny, DM, Ntalla, I, Palmer, ND, Pasko, D, Peter, A, Rayner, NW, Renstrom, F, Rice, K, Sala, CF, Sennblad, B, Serafetinidis, I, Smith, JA, Soranzo, N, Speliotes, EK, Stahl, EA, Stirrups, K, Tentolouris, N, Thanopoulou, A, Torres, M, Traglia, M, Tsaftantakis, E, Javad, S, Yanek, LR, Zengini, E, Becker, DM, Bis, JC, Brown, JB, Adrienne Cupples, L, Hansen, T, Ingelsson, E, Karter, AJ, Lorenzo, C, Mathias, RA, Norris, JM, Peloso, GM, Sheu, WHH, Toniolo, D, Vaidya, D, Varma, R, Wagenknecht, LE, Boeing, H, Bottinger, EP, Dedoussis, G, Deloukas, P, Ferrannini, E, Franco, OH, Franks, PW, Gibbs, RA, Gudnason, V, Hamsten, A, Harris, TB, Hattersley, AT, Hayward, C, Hofman, A, Jansson, J-H, Langenberg, C, Launer, LJ, Levy, D, Oostra, BA, O'Donnell, CJ, O'Rahilly, S, Padmanabhan, S, Pankow, JS, Polasek, O, Province, MA, Rich, SS, Ridker, PM, Rudan, I, Schulze, MB, Smith, BH, Uitterlinden, AG, Walker, M, Watkins, H, Wong, TY, Zeggini, E, The, E-IC, Laakso, M, Borecki, IB, Chasman, DI, Pedersen, O, Psaty, BM, Shyong Tai, E, van Duijn, CM, Wareham, NJ, Waterworth, DM, Boerwinkle, E, Linda Kao, WH, Florez, JC, Loos, RJF, Wilson, JG, Frayling, TM, Siscovick, DS, Dupuis, J, Rotter, JJ, Meigs, JB, Scott, RA, Goodarzi, MO: Low-frequency and rare exome chip variants associate with fasting glucose and type 2 diabetes susceptibility. *Nat Commun* 6: 5897, 2015
15. Pattaro C, Teumer A, Gorski M, Chu AY, Li M, Mijatovic V, Gamaas M, Tin A, Sorice R, Li Y, Taliun D, Olden M, Foster M, Yang Q, Chen MH, Pers TH, Johnson AD, Ko YA, Fuchsberger C, Tayo B, Nalls M, Feitosa MF, Isaacs A, Dehghan A, d'Adamo P, Adeyemo A, Dieffenbach AK, Zonderman AB, Nolte IM, van der Most PJ, Wright AF, Shuldiner AR, Morrison AC, Hofman A, Smith AV, Dreisbach AW, Franke A, Uitterlinden AG, Metspalu A, Tonjes A, Lupo A, Robino A, Johansson A, Demirkan A, Kollerits B, Freedman BI, Ponte B, Oostra BA, Paulweber B, Krämer BK, Mitchell BD, Buckley BM, Peralta CA, Hayward C, Helmer C, Rotimi CN, Shaffer CM, Müller C, Sala C, van Duijn CM, Saint-Pierre A, Ackermann D, Shriner D, Ruggiero G, Toniolo D, Lu Y, Cusi D, Czamara D, Ellinghaus D, Siscovick DS, Ruderfer D, Gieger C, Grallert H, Rohtchchina E, Atkinson EJ, Holliday EG, Boerwinkle E, Salvi E, Bottinger EP, Murgia F, Rivadeneira F, Ernst F, Kronenberg F, Hu FB, Navis GJ, Curhan GC, Ehret GB, Homuth G, Coassin S, Thun GA, Pistis G, Gambaro G, Malerba G, Montgomery GW, Eiriksdottir G, Jacobs G, Li G, Wichmann HE, Campbell H, Schmidt H, Wallaschofski H, Völzke H, Brenner H, Kroemer HK, Kramer H, Lin H, Leach IM, Ford I, Guessous I, Rudan I, Prokopenko I, Borecki I, Heid IM, Kolcic I, Persico I, Jukema JW, Wilson JF, Felix JF, Divers J, Lambert JC, Stafford JM, Gaspoz JM, Smith JA, Faul JD, Wang JJ, Ding J, Hirschhorn JN, Attia J, Whitfield JB, Chalmers J, Viikari J, Coresh J, Denny JC, Karjalainen J, Fernandes JK, Endlich K, Butterbach K, Keene KL, Lohman K, Portas L, Launer LJ, Lyttikäinen LP, Yengo L, Franke L, Ferrucci L, Rose LM, Kedenko L, Rao M, Struchalin M, Kleber ME, Cavalieri M, Haun M, Cornelis MC, Ciullo M, Pirastu M, de Andrade M, McEvoy MA, Woodward M, Adam M, Cocca M, Nauck M, Imboden M, Waldenberger M, Pruijm M, Metzger M, Stumvoll M, Evans MK, Sale MM, Kähönen M, Boban M, Bochud M, Rheinberger M, Verweij N, Bouatia-Naji N, Martin NG, Hastie N, Probst-Hensch N, Soranzo N, Devuyst O, Raitakari O, Gottesman O, Franco OH, Polasek O, Gasparini P, Munroe PB, Ridker PM, Mitchell P, Muntner P, Meisinger C, Smit JH, Kovacs P, Wild PS, Froguel P, Rettig R, Mägi R, Biffar R, Schmidt R, Middelberg RP, Carroll RJ, Penninx BW, Scott RJ, Katz R, Sedaghat S, Wild SH, Kardina SL, Ulivi S, Hwang SJ, Enroth S, Kloiber S, Trompet S, Stengel B, Hancock SJ, Turner ST, Rosas SE, Stracke S, Harris TB, Zeller T, Zemunik T, Lehtimäki T, Illig T, Aspelund T, Nikopensius T, Esko T, Tanaka T, Gyllenstein U, Völker U, Emilsson V, Vitart V, Aalto V, Gudnason V, Chouraki V, Chen WM, Igl W, März W, Koenig W, Lieb W, Loos RJ, Liu Y, Snieder H, Pramstaller PP, Parsa A, O'Connell JR, Susztak K, Hamet P, Tremblay J, de Boer IH, Böger CA, Goessling W, Chasman DI, Köttgen A, Kao WH, Fox CS; ICBP Consortium; AGEN Consortium; CARDIOGRAM; CHARGE-Heart Failure Group; ECHOGen Consortium: Genetic associations at 53 loci highlight cell types and biological pathways relevant for kidney function. *Nat Commun* 7: 10023, 2016
16. Yamamoto GL, Aguena M, Gos M, Hung C, Pilch J, Fahiminiya S, Abramowicz A, Cristian I, Buscarilli M, Naslavsky MS, Malaquias AC, Zatz M, Bodamer O, Majewski J, Jorge AAL, Pereira AC, Kim CA, Passos-Bueno MR, Bertola DR: Rare variants in *SOS2* and *LZTR1* are associated with Noonan syndrome. *J Med Genet* 52: 413–421, 2015
17. Romano AA, Allanson JE, Dahlgren J, Gelb BD, Hall B, Pierpont ME, Roberts AE, Robinson W, Takemoto CM, Noonan JA: Noonan syndrome: clinical features, diagnosis, and management guidelines. *Pediatrics* 126: 746–759, 2010
18. George CD, Patton MA, el Sawi M, Sharland M, Adam EJ: Abdominal ultrasound in Noonan syndrome: a study of 44 patients. *Pediatr Radiol* 23: 316–318, 1993
19. Mancini F, Rigacci S, Berti A, Balduini C, Torti M: The low-molecular-weight phosphotyrosine phosphatase is a negative regulator of FcγRIIA-mediated cell activation. *Blood* 110: 1871–1878, 2007
20. Peloso GM, Auer PL, Bis JC, Voorman A, Morrison AC, Stitzel NO, Brody JA, Khetarpal SA, Crosby JR, Fornage M, Isaacs A, Jakobsdottir J, Feitosa MF, Davies G, Huffman JE, Manichaikul A, Davis B, Lohman K, Joon AY, Smith AV, Grove ML, Zanon P, Redon V, Demissie S, Lawson K, Peters U, Carlson C, Jackson RD, Ryckman KK, Mackey RH, Robinson JG, Siscovick DS, Schreiner PJ, Mychaleckyj JC, Pankow JS, Hofman A, Uitterlinden AG, Harris TB, Taylor KD, Stafford JM, Reynolds LM, Marioni RE, Dehghan A, Franco OH, Patel AP, Lu Y, Hindy G, Gottesman O, Bottinger EP, Melander O, Orho-Melander M, Loos RJ, Duga S, Merlini PA, Farrall M, Goel A, Asselta R, Girelli D, Martinelli N, Shah SH, Kraus WE, Li M, Rader DJ, Reilly MP, McPherson R, Watkins H, Ardicino D, Zhang Q, Wang J, Tsai MY, Taylor HA, Correa A, Griswold ME, Lange LA, Starr JM, Rudan I, Eiriksdottir G, Launer LJ, Ordovas JM, Levy D, Chen YD, Reiner AP, Hayward C, Polasek O, Deary IJ, Borecki IB, Liu Y, Gudnason V, Wilson JG, van Duijn CM, Kooperberg C, Rich SS, Psaty BM, Rotter JJ, O'Donnell CJ, Rice K, Boerwinkle E, Kathiresan S, Cupples LA; NHLBI GO Exome Sequencing Project: Association of low-frequency and rare coding-sequence variants with blood lipids and coronary

Fernando Rivadeneira,¹ Jacques E. Rossouw,² Frank Schmidt,³ David Siscovick,⁴ Nicole Soranzo,⁵ Konstantin Strauch,⁶ Daniela Toniolo,⁷ Stephen T. Turner,⁸ André G. Uitterlinden,⁹ Sheila Ulivi,¹⁰ Dinesh Velayutham,¹¹ Uwe Völker,¹² Henry Völzke,¹³ Melanie Waldenberger,¹⁴ Jie Jin Wang,¹⁵ David R. Weir,¹⁶ Daniel Witte,¹⁷ Helena Kuivaniemi,¹⁸ Caroline S. Fox,¹⁹ Nora Franceschini,²⁰ Wolfram Goessling,²¹ Anna Köttgen,²² and Audrey Y. Chu²³

¹Department of Epidemiology, Johns Hopkins University, Baltimore, Maryland; ²Division of Nephrology and Department of Human Genetics, University of Utah, Salt Lake City, Utah; ³Division of Genetic Epidemiology, Medical Center – University of Freiburg, Freiburg, Germany; ⁴Genetics Division, ⁵Division of Preventive Medicine, ⁶Cardiovascular Medicine Division, and ⁷Gastroenterology Division, Brigham and Women’s Hospital, Harvard Medical School, Boston, Massachusetts; ⁸Department of Life and Reproduction Sciences, and ⁹Section of Biology and Genetics, Department of Life and Reproduction Sciences, University of Verona, Verona, Italy; ¹⁰Institute for Community Medicine, ¹¹Department of Anatomy and Cell Biology, ¹²Institute of Clinical Chemistry and Laboratory Medicine, ¹³Institute of Physiology, and ¹⁴Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald, Greifswald, Germany; ¹⁵National Heart, Lung, and Blood Institute’s Framingham Heart Study and the Center for Population Studies, Framingham, Massachusetts; ¹⁶Center for Genomic & Experimental Medicine, Institute of Genetics and Molecular Medicine, ¹⁷Medical Research Council Human Genetics, Institute of Genetics and Molecular Medicine, and ¹⁸Center for Population Health Sciences, University of Edinburgh, Edinburgh, Scotland, United Kingdom; ¹⁹The Sigfried and Janet Weis Center for Research, Geisinger Health System, Danville, Pennsylvania; ²⁰Division of Molecular Biology and Human Genetics, Department of Biomedical Sciences, Faculty of Medicine and Health Sciences, Stellenbosch University, Tygerberg, South Africa; ²¹Center for Biomedicine, European Academy of Bolzano/Bozen, affiliated to the University of Lübeck, Bolzano/Bozen, Italy; ²²Department of Biostatistics, Center for Statistical Genetics, ²³Survey Research Center, Institute for Social Research, and ²⁴Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, Michigan; ²⁵Department of Genetic Epidemiology, Institute of Epidemiology and Preventive Medicine, University of Regensburg, Regensburg, Germany; ²⁶Department of Nephrology, University Hospital Regensburg, Regensburg, Germany; ²⁷Department of Clinical Chemistry, Fimlab Laboratories and University of Tampere School of Medicine, Tampere, Finland; ²⁸Institute of Genetics and Biophysics “Adriano Buzzati-Traverso”, Napoli, Italy; ²⁹Department of Epidemiology, and ³⁰Department of Internal Medicine, Erasmus Medical Center, Rotterdam, The Netherlands; ³¹Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts; ³²Novo Nordisk Foundation Center for Basic Metabolic Research, Section of Metabolic Genetics, and ³³Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark; ³⁴Steno Diabetes Center, Gentofte, Denmark; ³⁵McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland; ³⁶Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, Tennessee; ³⁷Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, Massachusetts; ³⁸Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, Illinois; ³⁹Department of Genetics, School of Medicine, Washington University School of Medicine, St. Louis, Missouri; ⁴⁰Division of Nephrology, Gemelli Foundation University Hospital, Catholic University, Rome, Italy; ⁴¹Institute for Maternal and Child Health, IRCCS “Burlo Garofolo”, University of Trieste, Trieste, Italy; ⁴²Institute of Genetic Epidemiology and ⁴³Research Unit of Molecular Epidemiology, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany; ⁴⁴Institute of Hematology, Rui Jin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China; ⁴⁵State Key Laboratory of Medical Genomics, Shanghai, China; ⁴⁶Swiss Tropical and Public Health Institute, Basel, Switzerland; ⁴⁷The Charles Bronfman Institute for Personalized Medicine, and ⁴⁸Division of Psychiatric Genomics, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York; ⁴⁹Kidney Research Institute, ⁵⁰Cardiovascular Health Research Unit, Department of Medicine, and ⁵¹Departments of Epidemiology and Health Services, University of Washington, Seattle, Washington; ⁵²Center for Vision Research, Department of Ophthalmology and Westmead Millennium Institute for Medical Research, University of Sydney, Westmead, Sydney, New South Wales, Australia; ⁵³Department of Medicine, University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland; ⁵⁴Department of Epidemiology, and ⁵⁵Department of Epidemiology, Leiden University Medical Center, Leiden, The Netherlands; ⁵⁶Division of Public Health Sciences, Biostatistical Sciences, and ⁵⁷Epidemiology and Prevention, Division of Public Health Sciences, Wake Forest University School of Medicine, Winston-Salem, North Carolina; ⁵⁸Estonian Genome Center of University of Tartu, Tartu, Estonia; ⁵⁹Department of Genetics, and ⁶⁰Department of Epidemiology, University of North Carolina, Chapel Hill, North Carolina; ⁶¹Epidemiology Section, Department of Biostatistics Epidemiology & Scientific Computing, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia; ⁶²Institute for Maternal and Child Health – IRCCS “Burlo Garofolo”, Trieste, Italy; ⁶³Department of Health Science, University of Milano, Milano, Italy; ⁶⁴Fred Hutchinson Cancer Research Center, Seattle, Washington; ⁶⁵Department of Clinical Science, Lund University, Malmö, Sweden; ⁶⁶Icelandic Heart Association, Kopavogur, Iceland; ⁶⁷Faculty of Medicine, University of Iceland, Reykjavik, Iceland; ⁶⁸San Raffaele Scientific Institute, Milano, Italy; ⁶⁹Program in Personalized and Genomic Medicine and Department of Medicine, School of Medicine, University of Maryland, Baltimore, Maryland; ⁷⁰Department of Medicine and Department of Biomedical Sciences, and ⁷¹Division of Endocrinology, Diabetes and Metabolism, Cedars-Sinai Medical Center, Los Angeles, California; ⁷²Division of Statistical Genomics, Department of Genetics and Center for Genome Sciences and Systems Biology, Washington University, St. Louis, Missouri; ⁷³Department of Epidemiology, Fairbanks School of Public Health, Indianapolis, Indiana; ⁷⁴Department of Medicine, Indiana University School of Medicine, Indianapolis, Indiana; ⁷⁵The Human Genetics Center and Institute of Molecular Medicine, University of Texas Health Science Center, Houston, Texas; ⁷⁶Department of Clinical Biochemistry, and

Department of Internal Medicine and Endocrinology, Vejle Hospital, Vejle, Denmark; Department of Medicine, University of South Alabama College of Medicine, Mobile, Alabama; Renal Division and Channing Laboratory, Brigham and Women's Hospital, Boston, Massachusetts; Division of Nephrology, San Paolo Hospital, Milan, Italy; Division of Nephrology, Department of Medicine, Leiden University Medical Center and Leiden University, The Netherlands; University of Zurich, Institute of Physiology, Zurich Center for Integrative Human Physiology, Mechanisms of Inherited Kidney Disorders Group, Zürich, Switzerland; Department of Medicine, University of Mississippi, Jackson, Mississippi; Division of Endocrinology, Children's Hospital Boston, Boston, Massachusetts; Department of Genetics, Harvard Medical School, Boston, Massachusetts; Department of Medical Genetics and Molecular Biochemistry, Temple University School of Medicine, Philadelphia, Pennsylvania; Research Center for Prevention and Health, The Capital Region of Denmark, Glostrup University Hospital, Glostrup, Denmark; National Institutes on Aging, Bethesda, Maryland; Division of Applied Health Sciences, University of Aberdeen, Aberdeen, Scotland, United Kingdom; Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts; Department of Internal Medicine, The Ohio State University, Columbia, Ohio; Department of Internal Medicine II, Cardiology, University of Ulm Medical Center, Ulm, Germany; The German Heart Center Munich, Technical University of Munich, Munich, Germany; German Center for Cardiovascular Research, partner site Munich Heart Alliance, Munich, Germany; Department of General Practice, Aarhus University, Aarhus, Denmark; Department of Clinical Experimental Research, Rigshospitalet, Denmark; Renal Unit, Department of Medicine, University Hospital of Verona, Verona, Italy; German Center for Cardiovascular Research, partner site Greifswald, Greifswald, Germany; Cologne Center for Genomics, University of Cologne, Cologne, Germany; Division of Nephrology, Department of Medicine, University of Maryland School of Medicine, Baltimore, Maryland; German Center for Cardiovascular Disease Research, Munich, Germany; Group Health Research Institute, Group Health Cooperative, Seattle, Washington; Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, and Research Center of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland; National Heart, Lung, and Blood Institute, Bethesda, Maryland; Institute of Medical Informatics, Biometry and Epidemiology, Chair of Genetic Epidemiology, Ludwig-Maximilians-Universität, Munich, Germany; Division of Nephrology and Hypertension, Mayo Clinic, Rochester, Minnesota; German Center for Diabetes Research, partner site Greifswald, Greifswald, Germany; Department of Public Health, Aarhus University, Aarhus, Denmark; Harvard Stem Cell Institute, Cambridge, Massachusetts; and Dana-Farber Cancer Institute, Boston, Massachusetts