

REVIEW PAPER

Nitric oxide production in plants: an update

Jeremy Astier, Inonge Gross* and Jörg Durner

Helmholtz Zentrum München, Department of Environmental Science, Institute of Biochemical Plant Pathology D-85764 Neuherberg, Germany

* Correspondence: inonge.gross@helmholtz-muenchen.de

Received 1 August 2017; Editorial decision 30 October 2017; Accepted 2 November 2017

Editor: Ismail Turkan, Ege University, Turkey

Abstract

Nitric oxide (NO) is a key signaling molecule in plant physiology. However, its production in photosynthetic organisms remains partially unresolved. The best characterized NO production route involves the reduction of nitrite to NO via different non-enzymatic or enzymatic mechanisms. Nitrate reductases (NRs), the mitochondrial electron transport chain, and the new complex between NR and NOFNiR (nitric oxide-forming nitrite reductase) described in *Chlamydomonas reinhardtii* are the main enzymatic systems that perform this reductive NO production in plants. Apart from this reductive route, several reports acknowledge the possible existence of an oxidative NO production in an arginine-dependent pathway, similar to the nitric oxide synthase (NOS) activity present in animals. However, no NOS homologs have been found in the genome of embryophytes and, despite an increasing amount of evidence attesting to the existence of NOS-like activity in plants, the involved proteins remain to be identified. Here we review NO production in plants with emphasis on the presentation and discussion of recent data obtained in this field.

Keywords: Arginase, arginine, copper amine oxidase, nitrate reductase, nitric oxide, nitric oxide production, plant.

Introduction

Nitric oxide (NO) is a small molecule that possesses a wide range of physiological functions in living organisms. Although this radical gas was originally described as an air pollutant, it has since been shown to be involved in signaling functions in all living organisms. In plants, NO has been shown to be involved in many physiological processes, such as germination, flowering, or leaf senescence, and in response to environmental stresses (Mur *et al.*, 2013).

The mechanism of action of NO in plants has been investigated for more than two decades. In plants, NO donors or endogenous NO are able to modulate the expression of several genes involved in hormonal signaling, primary

metabolism, or stress responses (Besson-Bard *et al.*, 2009; Grün *et al.*, 2006). Acting as a molecular messenger, NO has been shown to interact with the signaling pathways dependent on cGMP, Ca²⁺, and notably with reactive oxygen species (ROS) (Astier *et al.*, 2010; Mur *et al.*, 2013). Several lines of evidence demonstrated that NO impacts the signaling of many phytohormones involved in developmental as well as in defense processes (Freschi, 2013). At the molecular level, NO has been shown to be responsible for three main specific post-translational modifications (PTMs) of proteins. It can reversibly bind the thiol group of cysteinyl residues leading to their S-nitrosation in a so-called S-nitrosylation process,

Abbreviations: ADC, arginine decarboxylase; AO, aldehyde oxidase; ARC, amidoxime reducing component; AS, argininosuccinate; AtNOA1, nitric oxide associated 1; BH₄, tetrahydrobiopterin; CaM, calmodulin; eNOS, endothelial NOS; GSNOR, S-nitrosoglutathione reductase; iNOS, inducible NOS; mETC, mitochondrial electron transport chain; Moco, molybdenum cofactor; Ni-NR activity, nitrite:NO reductase activity; nNOS, neuronal NOS; NO, nitric oxide; NOFNiR, nitric oxide-forming nitrite reductase; NOS, nitric oxide synthase; NR, nitrate reductase; PA, polyamine; PTM, post-translational modification; ROS, reactive oxidative species; SO, sulfite oxidase; TH₄, tetrahydrofolate; THB1, truncated hemoglobin 1; XO, xanthine oxidase.

© The Author(s) 2017. Published by Oxford University Press on behalf of the Society for Experimental Biology. All rights reserved. For permissions, please email: journals.permissions@oup.com

impacting the conformation, the activity, or the localization of the target protein. NO can also nitrosylate tyrosine residues in a so-called tyrosine nitration reaction, forming 3-nitrotyrosine, a process which is mainly irreversible. Finally, NO can also interact reversibly with the heme center of metalloproteins, generally resulting in conformational changes that will impact their activity (Astier and Lindermayr, 2012). Recently, the nitration of fatty acids by NO has also been demonstrated to be an important part of NO signaling in plants (Mata-Pérez *et al.*, 2017). Similarly to any signaling, the NO pathway in plants is tightly controlled through the action of specific enzymes. The turnover of NO messaging is, for example, dependent on the action of non-symbiotic hemoglobins that can oxidize it to nitrate, as well as on the activity of *S*-nitrosoglutathione reductase (GSNOR) that control the GSNO content in the cell, a major reservoir of NO (Mur *et al.*, 2013).

Despite intensive studies that revealed its variety of functions and reactivity in photosynthetic organisms, NO production in plants is still not fully understood and remains one of the most challenging issues of the field. NO synthesis in plants can be schematically achieved via two main routes defined by their chemical properties, one reductive and one oxidative. The reductive pathway is based on the reduction of nitrites to NO, while the oxidative route relies on the oxidation of aminated molecules.

In this review, we provide an update about NO production in plants in general, reviewing the overall and recent data available concerning this question.

Reduction of nitrite

NO can originate from different routes and substrates in living organisms. One of the synthesis routes concerns the reduction of nitrite. To date, this reductive route is the most firmly described and evidenced synthesis pathway for NO in plants.

Non-enzymatic reduction of nitrite

The reduction of nitrites to NO can occur non-enzymatically in particular conditions, such as low pH or highly reducing environments, when high concentrations of nitrate are present. These specific situations can happen, for example, in the apoplast of barley aleurone layers. Although such conditions are rarely encountered, they can lead to an efficient and rapid production of NO from nitrite (Bethke *et al.*, 2004).

Nitrate reductase

In addition to this non-enzymatic reduction, several proteins have been described to catalyze the production of NO from nitrites.

Nitrate reductase (NR) is a multifunctional cytoplasmic enzyme involved in nitrogen assimilation and metabolism. It is responsible for the first rate-limiting step of nitrate assimilation by catalyzing the reduction of nitrate to nitrite using

NADH as an electron donor. The active enzymatic homodimeric complex requires the presence of molybdopterin, heme, and FAD as cofactors (Campbell, 2001). Interestingly, in addition to this primary NR activity, this enzyme has also been described to possess a nitrite:NO reductase activity (Ni-NR activity; Yamasaki *et al.*, 1999; Yamasaki and Sakihama, 2000; Rockel *et al.*, 2002). This second reaction is relatively low and represents only 1% of the nitrate-reducing capacity of NR in normal conditions. Whereas the NR K_m for nitrite is notably higher than that for nitrate (100 μM compared with 10 μM), the Ni-NR activity requires a nitrite accumulation to occur and is efficiently inhibited by nitrates (K_i 50 μM). However, the reaction can be promoted by specific conditions such as anoxic or acidic environments, which leads to a substantial production of NO, from 2 nmol gFW⁻¹ h⁻¹ to 200 nmol gFW⁻¹ h⁻¹ *in vivo* (Rockel *et al.*, 2002; Meyer *et al.*, 2005). Despite these specific requirements, the importance of NO production by NR in plant physiology has been clearly demonstrated using both pharmacological and genetic approaches (Wilson *et al.*, 2008; Mur *et al.*, 2013). For example, the use of *Arabidopsis thaliana* NR-impaired lines *nial*, *nia2*, or the double mutant *nial1/nia2*, revealed the crucial role of NR-dependent NO production in various processes, such as stomatal movements (Desikan *et al.*, 2002; Hao *et al.*, 2010), hormone responses (Kolbert and Erdei, 2008; Hao *et al.*, 2010), salt, osmotic, or cold stress responses (Zhao *et al.*, 2009; Kolbert *et al.*, 2010; Xie *et al.*, 2013), and floral or root development (Seligman *et al.*, 2008; Méndez-Bravo *et al.*, 2010; Lombardo and Lamattina, 2012).

The new nitric oxide-forming nitrite reductase

The crucial role of NR in NO signaling in plants has been reinforced by recent works. Indeed, another NO-producing mechanism by NR has been unraveled in *Chlamydomonas reinhardtii*. In this unicellular alga, it has been shown that NR can interact with the partner protein NOFNiR (nitric oxide-forming nitrite reductase) to produce NO from nitrite. NOFNiR belongs to the amidoxime reducing component (ARC) protein family. Although ARC proteins were initially described from the animal field for reducing amidoxime prodrugs to their corresponding amino form *in vitro*, their exact physiological role *in vivo* is not fully understood (Havemeyer *et al.*, 2006). In *C. reinhardtii*, it has been demonstrated that NOFNiR can reduce nitrite to NO in an NAD(P)H reaction using electrons provided by the diaphorase activity of NR. Interestingly, this NO-producing activity occurs in normoxia and is not inhibited by nitrate, in contrast to the Ni-NR activity. In addition, the gene expression patterns and enzymatic activity of the two components of this system have been shown to correlate (Chamizo-Ampudia *et al.*, 2016, 2017). Interestingly, the *A. thaliana* genome contains two genes for ARC protein (Chamizo-Ampudia *et al.*, 2016), one of them presenting an NO-producing activity *in vitro* (Yang *et al.*, 2015). The determination of the existence of an NR:NOFNiR system in higher plants, similar to what is found in *C. reinhardtii*, would provide some information about the ubiquity

of this system in the green lineage and could better explain the crucial role of NR observed in plant NO production.

In parallel, the same team demonstrated that in addition to NOFNiR, NR could also associate with the truncated hemoglobin 1 (THB1) of *C. reinhardtii*. NR reduces THB1 through its diaphorase activity, which becomes active and can efficiently convert NO to nitrate in the presence of oxygen (Sanz-Luque *et al.*, 2015). This apparent contradictory role of NR in NO signaling, participating both in the production and the turnover of this signaling molecule, is actually coherent when considered in the light of the complex regulation of the nitrate cycle of the alga (reviewed by Calatrava *et al.*, 2017). However, this complexity highlights the importance of defining further the precise physiological role of NO produced by this specific system, considering its potential involvement in developmental or defense processes.

Plasma membrane-bound NR

In addition to the involvement of the cytoplasmic NR, the participation of a membrane-bound nitrite reductase (Ni:NOR) in the production of NO in plants has been reported. Using membrane fractions from tobacco roots, nitrite-dependent NO production was measured and attributed to a putative Ni:NOR that is yet to be identified (Stöhr *et al.*, 2001). This membrane-bound protein would be exclusively found in roots, and produces NO from nitrite in the apoplast of the cells, using NAD(P)H as electron donor. Its activity is dependent on low oxygen pressure and it would function together with an apoplastic membrane-bound NR that would provide nitrite from nitrate (Stöhr and Ullrich, 2002; Stöhr and Stremlau, 2006). Further work suggested a role for this Ni-NOR-produced NO in the mycorrhizal colonization of tobacco roots (Moche *et al.*, 2010).

Role of other molybdoenzymes

Both NR and NOFNiR display the presence of a molybdenum cofactor (Moco) in their structural features. In plants, other Moco-containing enzymes exist, namely xanthine oxidases (XOs), aldehyde oxidases (AOs), and sulfite oxidases (SOs), and they have been shown potentially to possess an NO-producing activity from nitrite.

XO is a highly conserved enzyme described initially in mammals as being responsible for purine catabolism, and hydroxylating hypoxanthine to xanthine and xanthine to urea. In plants, two XOs have been shown to contribute to ROS homeostasis during biotic stress, by generating both superoxide anions that contribute to the ROS burst and ureic acid involved in H₂O₂ removal in chloroplast (Yesbergenova *et al.*, 2005; Ma *et al.*, 2016). The potential nitrite reduction capacity of mammalian isoforms under anaerobic conditions has been documented *in vitro* for several years (Maia and Moura, 2015). In white lupine roots, a pharmacological approach using allopurinol, an inhibitor of XO, resulted in an inhibition of NO accumulation during development, suggesting a potential role for XO in this mechanism *in vivo* (Wang *et al.*, 2010). However, the data concerning a potential *in vivo*

role for XO in NO production in plants are scarce, and no NO emission from recombinant protein could be evidenced *in vitro* (Planchet *et al.*, 2005).

A structurally close enzyme related to XO is AO. AOs are cytoplasmic enzymes that generally catalyze the oxidation of aldehydes to carboxylates, producing superoxide anions. Its nitrite reduction activity has also been confirmed under anaerobiosis for several mammalian homologs *in vitro* (Maia and Moura, 2015). In plants, AOs participate in the synthesis of phytohormones such as abscisic acid (ABA) or indole-3-acetic acid (IAA), and contribute to ROS production (Zarepour *et al.*, 2012; Yergaliyev *et al.*, 2016), therefore being important for developmental processes and defense responses. However, no information is available about their NO-producing capacity *in vivo* in plants.

A last member of the Moco-containing enzyme family in plants is SO. SO is also a conserved enzyme, found in the peroxisomes, that catalyzes the oxidation of sulfite to sulfate, by an O₂-dependent mechanism (Eilers *et al.*, 2001). Its capability to reduce nitrite to NO has been quite recently demonstrated *in vitro* for the human isoform under anoxia, but this reaction requires more specific conditions and is less potent than the one observed for mammalian XO and AO (Wang *et al.*, 2015). In plants, the role of SOs is mainly assumed to concern the removal of toxic sulfite in the cell (Yarmolinsky *et al.*, 2013). Similarly to AOs, its involvement in nitrite reduction *in planta* has not been addressed yet.

Mitochondrial electron transport chain

In addition to the mechanisms described above, NO can be produced from nitrite through the action of the mitochondrial electron transport chain (mETC) in plants. After pioneer works demonstrating that nitrite-dependent NO formation could be prevented by mETC inhibitors in algae and tobacco (Tischner *et al.*, 2004; Planchet *et al.*, 2005), mETC-dependent NO production has been demonstrated in various species of plants such as pea, tobacco, and barley (Gupta *et al.*, 2005; Gupta and Kaiser, 2010). This reaction was located to the membrane of the mitochondria, involving mainly complex III and IV. It is determined by the availability in nitrite (K_m of 175 μ M) and requires anaerobic conditions, as oxygen can readily inhibit the reaction (K_i of 0.6 μ M). This reaction therefore is restricted to tissues exposed to hypoxia such as roots, and its occurrence can be explained by the requirement for an electron acceptor to preserve respiration, when oxygen is lacking (Gupta and Igamberdiev, 2011). In addition to metabolism preservation and allowing a correct functioning of mitochondria, this mETC-dependent NO production has also been suggested to be involved in signaling regulation processes (Palmieri *et al.*, 2010).

NOS-like activity in plants

In addition to the reductive pathway from nitrite, several lines of evidence demonstrate the existence of an oxidative route for NO production in plants, similar to the main pathway described in animals.

In mammals, although a reductive route by molybdenum-containing enzymes or non-enzymatically has recently been highlighted under acidic/reducing environments (Maia and Moura, 2015), the production of NO is principally achieved through the enzymatic activity of specialized enzymes: the nitric oxide synthases (NOSs).

These enzymes catalyze the formation of L-citrulline and NO from L-arginine through double mono-oxygenation. They work as homodimers, and contain schematically two main parts, the N-terminal domain possessing an oxidative activity and the C-terminal domain presenting a reducing activity. These two domains are linked to a calmodulin (CaM)-binding site. The binding of CaM triggers a structural change of the homodimer required for the enzymatic activity. This activity also needs the presence of several cofactors: FMN, FAD, and tetrahydrobiopterin (BH₄). Additionally, the reaction requires electrons from NADPH and the presence of oxygen (Förstermann and Sessa, 2012).

Several NOS isoforms have been characterized in animals, which possess specific features. In humans, three main isoforms have been studied. Two of them are constitutive, the endothelial NOS (eNOS) and the neuronal NOS (nNOS), with the other one being inducible (iNOS). The eNOS and nNOS activity requires the presence of a Ca²⁺-loaded CaM. Their activation leads to a quick, short, and relatively small release of NO (pmol min⁻¹ mg⁻¹ NOS), classically associated with NO signaling-dependent cellular processes. The iNOS does not require the presence of a Ca²⁺-loaded CaM for activation, and leads to a stronger and long-lasting release of NO (nmol min⁻¹ mg⁻¹ NOS). Its activation is involved generally in immune responses or pathology, where NO acts as a cytotoxic agent (Förstermann and Sessa, 2012).

NOS in plants

With the identification of NO as a crucial mediator of physiological processes in plants in the late 1990s, several studies sought to determine NO sources in the plant kingdom, primarily aiming to identify and characterize NOS homologs. Two main candidates have been described. The biochemical purification of a NOS-like activity from kilograms of tobacco leaves led to the identification of a P variant of the glycine decarboxylase complex. Unfortunately, further studies demonstrated that this protein does not produce NO; consequently, the corresponding articles were retracted. Another approach undertaken in *A. thaliana* led to the identification of a candidate that presented homology to an enzyme implicated in NO synthesis in the snail *Helix pomatia* (Guo *et al.*, 2003). Investigations demonstrated that the corresponding mutant displays an impaired NO content. However, the enzyme, initially named AtNOS1 (nitric oxide synthase 1), was further characterized as a functional small GTPase and therefore renamed AtNOA1 (nitric oxide associated 1; Moreau *et al.*, 2008). It is noteworthy that even if the mechanism underlying its impaired NO production is unclear, the *Atnoa1* mutant is used as a general NOS-like impaired tool, leading to an increasing amount of data referring to the study of a NOS-like activity in plant. These two unsuccessful examples of

plant NOS identification drove the community to question seriously their existence in the green lineage (Zemojtel *et al.*, 2006; Fröhlich and Durner, 2011).

However, in the beginning of 2010, the first NOS from the plant kingdom was characterized in the green algae *Ostreococcus tauri* (Foresi *et al.*, 2010). This enzyme was identified by sequence homology to the human NOS, with ~43% similarity to the eNOS sequence. The cofactor-binding sites for FAD, FMN, BH₄, and CaM are present, as well as L-arginine and an NADPH-binding site. This enzyme was shown to produce NO from L-arginine similarly to the animal NOSs. Its importance in light irradiance stress responses was also demonstrated in the *O. tauri* model (Foresi *et al.*, 2010). These results demonstrated the possibility of the presence of an endogenous and functional NOS in plants, with an actual role in plant physiology.

The first description of a canonical NOS from the plant kingdom was recently completed with an extensive analysis of the transcriptomes and genomes of >1300 species of plants, looking for the presence of NOS homologs (Jeandroz *et al.*, 2016). These authors screened the 1000 Plants (1KP) international multidisciplinary consortium's transcriptome database and the publicly available algal genome sequences, using the OtNOS and nNOS from human as templates. They could highlight 15 complete sequences presenting enough similarity with templates to be identified as NOS, all belonging to algal species. The identified sequences contain the key features of NOS, and the binding sites for NOS cofactors are conserved. The oxidative domain, especially in its N-terminal part, presents some diversity in the different candidates identified that could impact the dimerization of the enzyme or the binding of BH₄. This hypothesis is reinforced by the fact that OtNOS uses tetrahydrofolate (TH₄) instead of BH₄ to accomplish the enzymatic reaction *in vitro* and *in vivo* (Foresi *et al.*, 2010, 2015), and that the screen of 1KP database reveals the absence of the enzymes responsible for BH₄ synthesis in plants. First structural and phylogenetic analyses of these plant NOS candidates show that the activity is likely to be achieved independently of Ca²⁺, and demonstrate the presence of a diversity of structures that may result in a variety of functions (Jeandroz *et al.*, 2016; Santolini *et al.*, 2017). These observations raise the question of their role in algal physiology and constitute a promising new aspect of research to better understand the role of NO in the plant kingdom in general.

If these recent data confirmed the existence of NOS in several photosynthetic organisms, they also show that no homologs of NOS sequence can be found in any of the >1000 transcriptomes of land plants screened (Jeandroz *et al.*, 2016). These results, together with the unsuccessful attempts to purify candidates, tend to demonstrate that canonical NOSs probably do not exist in embryophytes. According to the phylogeny, it is likely that the NOS gene was transmitted from a common ancestor before the formation of the eukaryotic supergroup, and was later lost in land plants, the NOS from algae being the remaining testimony of these events (Jeandroz *et al.*, 2016; Santolini *et al.*, 2017).

The confirmation of the absence of canonical NOS in land plants raises the question of the relevance of a NOS activity in plants. Indeed, several studies carried out in plants suggested the existence of an oxidative route for NO production, so-called NOS-like activity.

Measurements of NOS-like activity in plants

Originally, the assumption that NOS would be present in plants comes from the measurement of NOS-like activity in plant tissues. Pioneer studies carried out in the mid-/late 1990s attested to the presence of this NOS-like activity in several plant models, such as maize, pea, tobacco, and lupine (Cueto *et al.*, 1996; Ninnemann and Maier, 1996; Durner *et al.*, 1998; Barroso *et al.*, 1999; Ribeiro *et al.*, 1999). It is noteworthy that these original works use the same technique to measure NOS activity: the citrulline-based assay. The principle of this technique is to follow the conversion of radiolabeled arginine provided as a substrate to radiolabeled citrulline (Bredt and Snyder, 1989). The reaction mixture generally contains all the common NOS cofactors and, after incubation, is applied to a cation exchange chromatography column that will retain the positively charged arginine but not citrulline. The radioactivity in the flowthrough is then assumed to refer to the converted citrulline, and its count theoretically directly correlates with the NOS activity present in the sample. However, this assay does not identify citrulline as a product, and its relevance to

follow NOS-like activity in plants was seriously questioned (Tischner *et al.*, 2007). It was actually demonstrated that the arginine-dependent activity measured from *A. thaliana* leaf extracts using the citrulline assay in normal conditions was mainly producing argininosuccinate (AS) rather than citrulline. Indeed, primary metabolism in plants differs from that in animals, and arginine can be metabolized in several different pathways, including through the action of AS lyase resulting in the measurement of AS formation (Fig. 1). These results highlight the caution needed in the transposition and interpretation of techniques used from other fields.

Nevertheless, the presence of NOS-like activity was later confirmed in plants using other techniques that directly measure the production of NO, such as chemiluminescence assay (Corpas *et al.*, 2004, 2006; Valderrama *et al.*, 2007; Chaki *et al.*, 2009) or EPR (Caro and Puntarulo, 1999; Pagnussat *et al.*, 2002; Dordas *et al.*, 2004; Simontacchi *et al.*, 2004; Jasid *et al.*, 2006, 2008). These activities were referred to as NOS-like activity as they were reported to be strictly dependent on the presence of arginine and NADPH, and several NOS co-factors. The localization of this NOS-like activity has been proposed, such as in chloroplasts or peroxisomes (Barroso *et al.*, 1999; Jasid *et al.*, 2006; Corpas and Barroso, 2014), but a clear picture is yet to be obtained regarding the enzymatic activity. More importantly, the corresponding enzymes remain to be identified.




Fig. 1. Schematic representation of the principal arginine metabolism pathways in plants. Arginine can be the substrate for several enzymes. Argininosuccinate lyase can generate argininosuccinate from arginine and fumarate. The guanidyl group of arginine can also be processed by arginase to ornithine and urea, or possibly by a NOS-like activity to citrulline. Arginine decarboxylase is another enzyme using arginine as a substrate, metabolizing it to agmatine. Arginine is the precursor of the principal polyamines in plants, such as putrescine, spermidine, and spermine. SPDS, Spermidine synthase; SPMS, Spermine synthase.

Table 1. Summary of principal reports using NOS inhibitors in plant systems

Organism	Tissue/cell type	Inhibitors applied (concentration μM)	NO production inhibition (%)	Technique used for NO production monitoring	Reference
<i>Arabidopsis thaliana</i>	Root	L-NAME (100)	75	DAF	Tossi et al. (2013)
<i>Arabidopsis thaliana</i>	Leaf	L-NAME (25)	75	DAF	Hao et al. (2010)
<i>Arabidopsis thaliana</i>	Leaf/root	L-NAME (200–1000)	50–80	Citrulline assay/DAF	Guo et al. (2003)
<i>Arabidopsis thaliana</i>	Root	L-NAME (5000); AG (2000)	Strong (not quantified)	DAR-4MAM	Corpas et al. (2009)
<i>Arabidopsis thaliana</i>	Leaf	L-NAME (3000); L-NNA (300)	100; 100	DAF	Ji et al. (2016)
<i>Arabidopsis thaliana</i>	Leaf	L-NAME (300); L-NNA (300)	100; 100	DAF	Zhao et al. (2009)
<i>Arabidopsis thaliana</i>	Root	L-NMMA (1000)	0	DAF	Kolbert et al. (2010)
<i>Arabidopsis thaliana</i>	Leaf	L-NNA (10000)	85	Citrulline assay/DAF	Zhao et al. (2007)
<i>Brassica rapa</i>	Root	L-NMMA (200)	90		Chen et al. (2014)
<i>Chorisporea bungeana</i>	Cell suspension	L-NAME (300)	100	Oxyhemoglobin assay/ Greiss assay	Liu et al. (2010)
<i>Cucurbita maxima</i> × <i>C. moschata</i>	Seedling	L-NAME (200)	80	DAF	Li et al. (2017)
<i>Elymus nutans</i>	Leaf	L-NNA (150)	100		Fu et al. (2015)
<i>Glycine max</i>	Chloroplast	L-NAME (5000); L-NNA (5000)	100; 100	EPR	Jasid et al. (2006)
<i>Glycine max</i>	Cotyledon	L-NIL (3000)	30	Citrulline assay	Modolo et al. (2002)
<i>Helianthus annuus</i>	Hypocotyl	AG (5000); L-NMMA (1000)	100; 100	Ozone chemiluminescence	Chaki et al. (2011)
<i>Hibiscus moscheutos</i>	Root	L-NNA (10000)	40	Citrulline assay/DAF	Tian et al. (2007)
<i>Lupinus albus</i>	Root	L-NAME (1000); L-NMMA (1000)	50; 50	Citrulline assay	Cueto et al. (1996)
<i>Lycopersicon esculentum</i>	Seedling	L-NAME (200)	70	DAF	Diao et al. (2016)
<i>Malus domestica</i>	seed	L-NAME (300)	100	Oxyhemoglobin assay/DAF	Krasuska et al. (2016)
<i>Nicotiana benthamiana</i>	Leaf	L-NAME (200)	50	DAF	Deng et al. (2016)
<i>Nicotiana tabacum</i>	Leaf	L-NAME (5000)	100	DAF	Zhang et al. (2011)
<i>Nicotiana tabacum</i>	Cell suspension	L-NAME (10000)	55	DAF	Lamotte et al. (2004)
<i>Nicotiana tabacum</i>	Cell suspension.	L-NMMA (1000)	37	Citrulline assay	Durner et al. (1998)
<i>Xanthi</i>	Extracts				
<i>Nicotiana tabacum</i>	Leaf	L-NMMA (na)	50	DAF	Foissner et al. (2000)
<i>Olea europaea</i>	Leaf	AG (1000)	100	Ozone chemiluminescence	Valderrama et al. (2007)
<i>Paulownia tomentosa</i>	Pollen tube	L-NAME (50)	100		He et al. (2007)
<i>Pennisetum glaucum</i>	Seedling	L-NAME (10000)	50	DAF	Manjunatha et al. (2009)
<i>Pinus bungeana</i>	Pollen tube	L-NNA (45)	40	DAF	Wang et al. (2009)
<i>Pisum sativum</i>	Leaf extract	AG (1000)	70	Ozone chemiluminescence	Corpas et al. (2008)
<i>Pisum sativum</i>	Plant	L-NAME (1000); AG (2000)	55; 85	DAF/EPR	Corpas et al. (2004)
<i>Pisum sativum</i>	Extract	L-NAME (1000); AG (1000); L-NMMA (1000); L-NIL (1000)	90; 100; 88; 59	Citrulline assay	Barroso et al. (1999)
<i>Scutellaria baicalensis</i>	Cell suspension	L-NNA (100)	100	DAF	Zhang et al. (2014)
<i>Solanum lycopersicum</i>	Root	L-NAME (20)	50	DAF	Negi et al. (2010)
<i>Solanum lycocarpum</i>	Root	L-NAME (500)	90		Jin et al. (2011)
<i>Vicia faba</i>	Leaf	L-NAME (1000)	100		Garcia-Mata and Lamattina (2007)
<i>Vicia faba</i>	Leaf	L-NAME (25)		DAF	Yan et al. (2007)
<i>Zea Mays</i>	Seedling	L-NAME (100)	35	DAF	Tossi et al. (2009a)
<i>Zea mays</i>	Leaf	L-NAME (200)	80	DAF	Sang et al. (2008b)
<i>Zea mays</i>	Leaf	L-NAME (na)	70		Tossi et al. (2009b)
<i>Zea mays</i>	Leaf/root	L-NAME (3000); AG (3000)	30; 30	Citrulline assay	Ribeiro et al., 1999)
<i>Zea mays</i>	Leaf	L-NAME (200)	71		Sang et al. (2008a)

L-NAME, *N*^ω-nitro-L-arginine methyl ester; L-NNA, L-*N*^ω-nitroarginine; L-NMMA, *N*^G-monomethyl- L-arginine; L-NIL, *N*⁶-(1-iminoethyl)- L-lysine; AG, aminoguanidine; DAF, diamino fluorescein; DAR, daminorhodamine.

Pharmacological approaches: use of inhibitors

Another substantial part of the work providing evidence for the presence of a NOS-like activity in plants comes from the analyses of NOS inhibitor effects in plant systems. NOS inhibitors are mainly arginine analogs, which compete for the active

site of the enzyme. For the last two decades, NOS inhibitors were used in various conditions on various plant models. The compilation of the data available concerning the use of NOS inhibitors in plants reveals a very strong variation in the effect observed (Table 1). The discrepancy between the different

concentrations used (from 25 μM to up to 10 mM) and the effectiveness of the inhibition recorded (from 0 to 100% inhibition, sometimes considering the same model) highlights the complexity of studying the NO-producing system in plants.

Another concern to be raised in the use of these NOS inhibitors is regarding their specificity. It is well defined that analogs of arginine can impact the activity of several enzymes (Víteček *et al.*, 2012). As an example, aminoguanidine can efficiently inhibit amine oxidase enzymes in plants (Planas-Portell *et al.*, 2013). Considering also that the main results characterizing the NOS-like activity in plants are obtained using complex systems such as crude extract, it cannot be excluded that the observed NO production impairment comes from an indirect enzymatic mechanism. Moreover, in the absence of their target(s), inhibitors can display enhanced off-target effects. The determination of the precise inhibitor target(s) is therefore a prerequisite for the correct interpretation of their observed effects. For all these reasons, these pharmacological approaches must be interpreted with caution.

Nevertheless, these pharmacological approaches share a consistency in the potential of a NOS inhibitor to prevent NO production in general, in different plant species and plant cell types. They also show that using different NO-monitoring techniques favors the existence of NOS-like activity in plants.

Heterologous expression of NOS and hydroxylamine oxidation in plants

In addition to the direct measurement of NOS-like activity in plants, additional approaches provide hints confirming its existence.

Genetic constructs aiming to express NOS in plants have been generated. The expression of recombinant nNOS from rat resulted in higher NO content, observed in *A. thaliana*, tobacco, and rice, correlated with higher resistance to biotic and abiotic stresses (Chun *et al.*, 2012; Shi *et al.*, 2012; Cai *et al.*, 2015). Similarly, expression of the OtNOS for algae in *A. thaliana* resulted in a functional enzyme producing NO *in planta*, correlated with a better germination and a tolerance to salt, oxidative, and drought stresses (Foresi *et al.*, 2015). Taken together, these approaches demonstrate that the cofactors and conditions required for a functional NOS activity are present in plants, arguing in favor of the existence of the oxidative NO production route.

In the same direction, *in vitro* experiments conducted on tobacco cell suspensions demonstrated that plants possess the ability to oxidize hydroxylamines to NO (Rümer *et al.*, 2009). However, the occurrence of this substrate in the natural physiology of plants is questionable and the involved enzyme has yet to be identified.

Arginine metabolism and NO oxidative production route in plant

Measurements of NOS-like activity and the use of NOS inhibitors have suggested an arginine-dependent NO production pathway in plants. Several other works have also linked arginine-dependent metabolism with NO signaling

in photosynthetic organisms. As an example, the commonly used mutant *Atnox1*, impaired in the expression of a chloroplast phosphoenolpyruvate/phosphate translocator, presents elevated levels of arginine correlated with a constitutive overproduction of NO (He *et al.*, 2004; Frungillo *et al.*, 2014). In plants, the arginine pool depends on the activity of several enzymes that use it as a substrate (Fig. 1).

Arginases are enzymes that catalyze the conversion of arginine into ornithine and urea reacting with the guanidyl group of the amino acid. Two isoforms are found in *A. thaliana*; both have been localized to the mitochondria (Flores *et al.*, 2008). Interestingly, genetic approaches demonstrated that *A. thaliana* mutants impaired in arginase expression display an increased NO content correlated with higher putrescine and spermine levels (Flores *et al.*, 2008; Shi *et al.*, 2013). Conversely, overexpression of arginase led to a decreased NO production and putrescine and spermine levels, which correlated to a susceptibility to abiotic stresses (Shi *et al.*, 2013). Recently, the involvement of a higher arginase activity impacting the arginine pool was found to be responsible for the impaired NO production and developmental phenotype observed in the *A. thaliana* mutant for the copper amine oxidase 8 (CuAO8), an enzyme involved in polyamine (PA) catabolism (Groß *et al.*, 2017). Similar results were obtained in cotton where an increased arginase activity due to the overexpression of the rice arginase gene resulted in decreased NO production that correlated with the developmental phenotype in roots (Meng *et al.*, 2015).

Arginine decarboxylases (ADCs) are enzymes responsible for the formation of agmatine through the decarboxylation of arginine. This constitutes the first step of the unique PA synthetic route in *A. thaliana*. Two isoforms are also found in *A. thaliana*, both being chloroplastic (Borrell *et al.*, 1995). Interestingly, transient overexpression of the pepper ADC1 resulted in an increased NO accumulation in tobacco cells, together with an accumulation of PAs (Kim *et al.*, 2013). Accordingly, the *A. thaliana* mutant *adc2.1* was impaired in melatonin- or iron deficiency-induced NO accumulation, correlated with PA accumulation deficiency (Zhou *et al.*, 2016).

PAs are found in all living kingdoms. The most common PAs found in plants are putrescine, spermine, and spermidine containing two, three, and four amine groups, respectively. These molecules have been shown to be involved in a wide range of physiological mechanisms in plants, from development to stress responses (Tiburcio *et al.*, 2014; Liu *et al.*, 2015). Over the last 15 years, several works reported that exogenous application of PAs results in NO production in several plant models (Tun *et al.*, 2006; Yang *et al.*, 2014; Diao *et al.*, 2016; Zhou *et al.*, 2016). In agreement with these data, *A. thaliana* mutants impaired in the expression of two different enzymes regulating PA catabolism, CuAO1 and CuAO8, presented an altered NO production (Wimalasekera *et al.*, 2011a, b; Groß *et al.*, 2017). It is important to note that arginine is the precursor of PA synthesis, connecting their metabolism (Fig. 1).

Taken together, these data strengthen the link existing between arginine metabolism and NO in plants, favoring

the existence of an oxidative NO production route in higher plants, even if the enzymes responsible for this potential activity remain to be identified.

Concluding remarks

The determination of NO sources in plants has clearly been and remains a challenging issue of the field. The intensive studies carried out over the last decade depict the emergence of a complex system where several players are involved. It is now apparent that nitrite reduction is the main source of NO. The recent findings concerning the nitrite reduction through the association of NR and NOFNiR proteins in *C. reinhardtii* open up an interesting aspect of research to determine if this mechanism is also present in higher plants. More generally, the in-depth characterization of the other Moco-containing proteins could provide information on their role in the reductive NO production route.

The amount of data regarding the oxidative NO production pathway in plants has also accumulated in recent years. The identification of a dozen NOSs restricted to the algal genome is surprising and interrogative. The characterization of their activity and the corresponding impact on algal metabolism could help to better define the physiological role of NO in these models. On the other hand, it is now clear that no canonical NOSs are present in embryophyte transcriptomes. However, several pieces of evidence reported in this review are in favor of the existence of NOS-like activity. This activity is dependent on arginine, or at least the arginine metabolic pathways. The apparent contradiction between the measurement of NOS-like activity and the absence of NOS in higher plants could be explained by the requirement for protein complex formation to bring different polypeptides required to reconstitute a full NOS activity into close proximity, similarly to what is observed for NR:NOFNiR, as recently suggested (Corpas and Barroso, 2017). The identification and characterization of the proteins involved and the precise substrate/cofactors needed are a prerequisite for a better understanding of NO formation in plants.

Taken together, the data available on NO production in plants reveal a deep complexity and diversity. The specificity of each source needs clarification as well as a better determination of the enzymes involved in its production. These constitute an important and promising aspect of research for a better comprehension of NO physiological function in plants.

References

Astier J, Besson-Bard A, Wawer I, Parent C, Rasul S, Jeandroz S, Dat J, Wendehenne D. 2010. Nitric oxide signalling in plants: cross-talk with Ca²⁺, protein kinases and reactive oxygen species. *Annual Plant Reviews* **42**, 147–170.

Astier J, Lindermayr C. 2012. Nitric oxide-dependent posttranslational modification in plants: an update. *International Journal of Molecular Sciences* **13**, 15193–15208.

Barroso JB, Corpas FJ, Carreras A, Sandalio LM, Valderrama R, Palma JM, Lupiáñez JA, del Río LA. 1999. Localization of nitric-oxide synthase in plant peroxisomes. *Journal of Biological Chemistry* **274**, 36729–36733.

Besson-Bard A, Astier J, Rasul S, Wawer I, Dubreuil-Maurizi C, Jeandroz S, Wendehenne D. 2009. Current view of nitric oxide-responsive genes in plants. *Plant Science* **177**, 302–309.

Bethke PC, Badger MR, Jones RL. 2004. Apoplastic synthesis of nitric oxide by plant tissues. *The Plant Cell* **16**, 332–341.

Borrell A, Culiñez-Macia FA, Altabella T, Besford RT, Flores D, Tiburcio AF. 1995. Arginine decarboxylase is localized in chloroplasts. *Plant Physiology* **109**, 771–776.

Bredt DS, Snyder SH. 1989. Nitric oxide mediates glutamate-linked enhancement of cGMP levels in the cerebellum. *Proceedings of the National Academy of Sciences, USA* **86**, 9030–9033.

Cai W, Liu W, Wang WS, Fu ZW, Han TT, Lu YT. 2015. Overexpression of rat neurons nitric oxide synthase in rice enhances drought and salt tolerance. *PLoS One* **10**, e0131599.

Calatrava V, Chamizo-Ampudia A, Sanz-Luque E, Ocaña-Calahorra F, Llamas A, Fernandez E, Galvan A. 2017. How Chlamydomonas handles nitrate and the nitric oxide cycle. *Journal of Experimental Botany* **68**, 2593–2602.

Campbell WH. 2001. Structure and function of eukaryotic NAD(P)H:nitrate reductase. *Cellular and Molecular Life Sciences* **58**, 194–204.

Caro A, Puntarulo S. 1999. Nitric oxide generation by soybean embryonic axes. Possible effect on mitochondrial function. *Free Radical Research* **31**(Suppl), S205–S212.

Chaki M, Fernández-Ocaña AM, Valderrama R, et al. 2009. Involvement of reactive nitrogen and oxygen species (RNS and ROS) in sunflower–mildew interaction. *Plant and Cell Physiology* **50**, 265–279.

Chaki M, Valderrama R, Fernández-Ocaña AM, et al. 2011. High temperature triggers the metabolism of S-nitrosothiols in sunflower mediating a process of nitrosative stress which provokes the inhibition of ferredoxin-NADP reductase by tyrosine nitration. *Plant, Cell and Environment* **34**, 1803–1818.

Chamizo-Ampudia A, Sanz-Luque E, Llamas A, Galvan A, Fernandez E. 2017. Nitrate reductase regulates plant nitric oxide homeostasis. *Trends in Plant Science* **22**, 163–174.

Chamizo-Ampudia A, Sanz-Luque E, Llamas A, Ocaña-Calahorra F, Mariscal V, Carreras A, Barroso JB, Galván A, Fernández E. 2016. A dual system formed by the ARC and NR molybdoenzymes mediates nitrite-dependent NO production in Chlamydomonas. *Plant, Cell and Environment* **39**, 2097–2107.

Chen Y, Mo HZ, Hu LB, Li YQ, Chen J, Yang LF. 2014. The endogenous nitric oxide mediates selenium-induced phytotoxicity by promoting ROS generation in *Brassica rapa*. *PLoS One* **9**, e110901.

Chun HJ, Park HC, Koo SC, et al. 2012. Constitutive expression of mammalian nitric oxide synthase in tobacco plants triggers disease resistance to pathogens. *Molecules and Cells* **34**, 463–471.

Corpas FJ, Barroso JB. 2014. Peroxisomal plant nitric oxide synthase (NOS) protein is imported by peroxisomal targeting signal type 2 (PTS2) in a process that depends on the cytosolic receptor PEX7 and calmodulin. *FEBS Letters* **588**, 2049–2054.

Corpas FJ, Barroso JB. 2017. Nitric oxide synthase-like activity in higher plants. *Nitric Oxide* **68**, 5–6.

Corpas FJ, Barroso JB, Carreras A, et al. 2004. Cellular and subcellular localization of endogenous nitric oxide in young and senescent pea plants. *Plant Physiology* **136**, 2722–2733.

Corpas FJ, Barroso JB, Carreras A, Valderrama R, Palma JM, León AM, Sandalio LM, del Río LA. 2006. Constitutive arginine-dependent nitric oxide synthase activity in different organs of pea seedlings during plant development. *Planta* **224**, 246–254.

Corpas FJ, Chaki M, Fernández-Ocaña A, et al. 2008. Metabolism of reactive nitrogen species in pea plants under abiotic stress conditions. *Plant and Cell Physiology* **49**, 1711–1722.

Corpas FJ, Hayashi M, Mano S, Nishimura M, Barroso JB. 2009. Peroxisomes are required for in vivo nitric oxide accumulation in the cytosol following salinity stress of Arabidopsis plants. *Plant Physiology* **151**, 2083–2094.

Cueto M, Hernández-Perera O, Martín R, Bentura ML, Rodrigo J, Lamas S, Golvano MP. 1996. Presence of nitric oxide synthase activity in roots and nodules of *Lupinus albus*. *FEBS Letters* **398**, 159–164.

Deng XG, Zhu T, Zou LJ, Han XY, Zhou X, Xi DH, Zhang DW, Lin HH. 2016. Orchestration of hydrogen peroxide and nitric oxide

in brassinosteroid-mediated systemic virus resistance in *Nicotiana benthamiana*. *The Plant Journal* **85**, 478–493.

Desikan R, Griffiths R, Hancock J, Neill S. 2002. A new role for an old enzyme: nitrate reductase-mediated nitric oxide generation is required for abscisic acid-induced stomatal closure in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **99**, 16314–16318.

Diao QN, Song YJ, Shi DM, Qi HY. 2016. Nitric oxide induced by polyamines involves antioxidant systems against chilling stress in tomato (*Lycopersicon esculentum* Mill.) seedling. *Journal of Zhejiang University. Science. B* **17**, 916–930.

Dordas C, Hasinoff BB, Rivoal J, Hill RD. 2004. Class-1 hemoglobins, nitrate and NO levels in anoxic maize cell-suspension cultures. *Planta* **219**, 66–72.

Durner J, Wendehenne D, Klessig DF. 1998. Defense gene induction in tobacco by nitric oxide, cyclic GMP, and cyclic ADP-ribose. *Proceedings of the National Academy of Sciences, USA* **95**, 10328–10333.

Eilers T, Schwarz G, Brinkmann H, et al. 2001. Identification and biochemical characterization of *Arabidopsis thaliana* sulfite oxidase. A new player in plant sulfur metabolism. *Journal of Biological Chemistry* **276**, 46989–46994.

Flores T, Todd CD, Tovar-Mendez A, et al. 2008. Arginase-negative mutants of *Arabidopsis* exhibit increased nitric oxide signaling in root development. *Plant Physiology* **147**, 1936–1946.

Foissner I, Wendehenne D, Langebartels C, Durner J. 2000. In vivo imaging of an elicitor-induced nitric oxide burst in tobacco. *The Plant Journal* **23**, 817–824.

Foresi N, Correa-Aragunde N, Parisi G, Caló G, Salerno G, Lamattina L. 2010. Characterization of a nitric oxide synthase from the plant kingdom: NO generation from the green alga *Ostreococcus tauri* is light irradiance and growth phase dependent. *The Plant Cell* **22**, 3816–3830.

Foresi N, Mayta ML, Lodeyro AF, Scuffi D, Correa-Aragunde N, García-Mata C, Casalougué C, Carrillo N, Lamattina L. 2015. Expression of the tetrahydrofolate-dependent nitric oxide synthase from the green alga *Ostreococcus tauri* increases tolerance to abiotic stresses and influences stomatal development in *Arabidopsis*. *The Plant Journal* **82**, 806–821.

Förstermann U, Sessa WC. 2012. Nitric oxide synthases: regulation and function. *European Heart Journal* **33**, 829–37, 837a.

Freschi L. 2013. Nitric oxide and phytohormone interactions: current status and perspectives. *Frontiers in Plant Science* **4**, 398.

Fröhlich A, Durner J. 2011. The hunt for plant nitric oxide synthase (NOS): is one really needed? *Plant Science* **181**, 401–404.

Frungillo L, Skelly MJ, Loake GJ, Spoel SH, Salgado I. 2014. S-Nitrosothiols regulate nitric oxide production and storage in plants through the nitrogen assimilation pathway. *Nature Communications* **5**, 5401.

Fu J, Chu X, Sun Y, Miao Y, Xu Y, Hu T. 2015. Nitric oxide mediates 5-aminolevulinic acid-induced antioxidant defense in leaves of *Elymus nutans* griseb. exposed to chilling stress. *PLoS One* **10**, e0130367.

García-Mata C, Lamattina L. 2007. Abscisic acid (ABA) inhibits light-induced stomatal opening through calcium- and nitric oxide-mediated signaling pathways. *Nitric Oxide* **17**, 143–151.

Groß F, Rudolf EE, Thiele B, Durner J, Astier J. 2017. Copper amine oxidase 8 regulates arginine-dependent nitric oxide production in *Arabidopsis thaliana*. *Journal of Experimental Botany* **68**, 2149–2162.

Grün S, Lindermayr C, Sell S, Durner J. 2006. Nitric oxide and gene regulation in plants. *Journal of Experimental Botany* **57**, 507–516.

Guo FQ, Okamoto M, Crawford NM. 2003. Identification of a plant nitric oxide synthase gene involved in hormonal signaling. *Science* **302**, 100–103.

Gupta KJ, Igamberdiev AU. 2011. The anoxic plant mitochondrion as a nitrite:NO reductase. *Mitochondrion* **11**, 537–543.

Gupta KJ, Kaiser WM. 2010. Production and scavenging of nitric oxide by barley root mitochondria. *Plant and Cell Physiology* **51**, 576–584.

Gupta KJ, Stoimenova M, Kaiser WM. 2005. In higher plants, only root mitochondria, but not leaf mitochondria reduce nitrite to NO, *in vitro* and *in situ*. *Journal of Experimental Botany* **56**, 2601–2609.

Hao F, Zhao S, Dong H, Zhang H, Sun L, Miao C. 2010. Nia1 and Nia2 are involved in exogenous salicylic acid-induced nitric oxide generation and stomatal closure in *Arabidopsis*. *Journal of Integrative Plant Biology* **52**, 298–307.

Havemeyer A, Bittner F, Wollers S, Mendel R, Kunze T, Clement B. 2006. Identification of the missing component in the mitochondrial benzamidoxime prodrug-converting system as a novel molybdenum enzyme. *Journal of Biological Chemistry* **281**, 34796–34802.

He JM, Bai XL, Wang RB, Cao B, She XP. 2007. The involvement of nitric oxide in ultraviolet-B-inhibited pollen germination and tube growth of *Paulownia tomentosa* in vitro. *Physiologia Plantarum* **131**, 273–282.

He Y, Tang RH, Hao Y, et al. 2004. Nitric oxide represses the *Arabidopsis* floral transition. *Science* **305**, 1968–1971.

Jasid S, Simontacchi M, Bartoli CG, Puntarulo S. 2006. Chloroplasts as a nitric oxide cellular source. Effect of reactive nitrogen species on chloroplastic lipids and proteins. *Plant Physiology* **142**, 1246–1255.

Jasid S, Simontacchi M, Puntarulo S. 2008. Exposure to nitric oxide protects against oxidative damage but increases the labile iron pool in sorghum embryonic axes. *Journal of Experimental Botany* **59**, 3953–3962.

Jeandroz S, Wipf D, Stuehr DJ, et al. 2016. Occurrence, structure, and evolution of nitric oxide synthase-like proteins in the plant kingdom. *Science Signaling* **9**, re2.

Ji Y, Liu J, Xing D. 2016. Low concentrations of salicylic acid delay methyl jasmonate-induced leaf senescence by up-regulating nitric oxide synthase activity. *Journal of Experimental Botany* **67**, 5233–5245.

Jin CW, Du ST, Shamsi IH, Luo BF, Lin XY. 2011. NO synthase-generated NO acts downstream of auxin in regulating Fe-deficiency-induced root branching that enhances Fe-deficiency tolerance in tomato plants. *Journal of Experimental Botany* **62**, 3875–3884.

Kim NH, Kim BS, Hwang BK. 2013. Pepper arginine decarboxylase is required for polyamine and γ -aminobutyric acid signaling in cell death and defense response. *Plant Physiology* **162**, 2067–2083.

Kolbert Z, Erdei L. 2008. Involvement of nitrate reductase in auxin-induced NO synthesis. *Plant Signaling and Behavior* **3**, 972–973.

Kolbert Z, Ortega L, Erdei L. 2010. Involvement of nitrate reductase (NR) in osmotic stress-induced NO generation of *Arabidopsis thaliana* L. roots. *Journal of Plant Physiology* **167**, 77–80.

Krasuska U, Ciacka K, Orzechowski S, Fettke J, Bogatek R, Gniazdowska A. 2016. Modification of the endogenous NO level influences apple embryos dormancy by alterations of nitrated and biotinylated protein patterns. *Planta* **244**, 877–891.

Lamotte O, Gould K, Lecourieux D, Sequeira-Legrand A, Lebrun-Garcia A, Durner J, Pugin A, Wendehenne D. 2004. Analysis of nitric oxide signaling functions in tobacco cells challenged by the elicitor cryptogein. *Plant Physiology* **135**, 516–529.

Li L, Shu S, Xu Q, An YH, Sun J, Guo SR. 2017. NO accumulation alleviates H₂O₂-dependent oxidative damage induced by Ca(NO₃)₂ stress in the leaves of pumpkin-grafted cucumber seedlings. *Physiologia Plantarum* **160**, 34–45.

Liu JH, Wang W, Wu H, Gong X, Moriguchi T. 2015. Polyamines function in stress tolerance: from synthesis to regulation. *Frontiers in Plant Science* **6**, 827.

Liu Y, Jiang H, Zhao Z, An L. 2010. Nitric oxide synthase like activity-dependent nitric oxide production protects against chilling-induced oxidative damage in *Chorisporea bungeana* suspension cultured cells. *Plant Physiology and Biochemistry* **48**, 936–944.

Lombardo MC, Lamattina L. 2012. Nitric oxide is essential for vesicle formation and trafficking in *Arabidopsis* root hair growth. *Journal of Experimental Botany* **63**, 4875–4885.

Ma X, Wang W, Bittner F, et al. 2016. Dual and opposing roles of xanthine dehydrogenase in defense-associated reactive oxygen species metabolism in *Arabidopsis*. *The Plant Cell* **28**, 1108–1126.

Maia LB, Moura JJ. 2015. Nitrite reduction by molybdoenzymes: a new class of nitric oxide-forming nitrite reductases. *Journal of Biological Inorganic Chemistry* **20**, 403–433.

Manjunatha G, Niranjana-Raj S, Prashanth GN, Deepak S, Amruthesh KN, Shetty HS. 2009. Nitric oxide is involved in chitosan-induced systemic resistance in pearl millet against downy mildew disease. *Pest Management Science* **65**, 737–743.

- Mata-Pérez C, Sánchez-Calvo B, Padilla MN, Begara-Morales JC, Valderrama R, Corpas FJ, Barroso JB.** 2017. Nitro-fatty acids in plant signaling: new key mediators of nitric oxide metabolism. *Redox Biology* **11**, 554–561.
- Méndez-Bravo A, Raya-González J, Herrera-Estrella L, López-Bucio J.** 2010. Nitric oxide is involved in alkamide-induced lateral root development in *Arabidopsis*. *Plant and Cell Physiology* **51**, 1612–1626.
- Meng Z, Meng Z, Zhang R, Liang C, Wan J, Wang Y, Zhai H, Guo S.** 2015. Expression of the rice arginase gene OsARG in cotton influences the morphology and nitrogen transition of seedlings. *PLoS One* **10**, e0141530.
- Meyer C, Lea US, Provan F, Kaiser WM, Lillo C.** 2005. Is nitrate reductase a major player in the plant NO (nitric oxide) game? *Photosynthesis Research* **83**, 181–189.
- Moche M, Stremlau S, Hecht L, Göbel C, Feussner I, Stöhr C.** 2010. Effect of nitrate supply and mycorrhizal inoculation on characteristics of tobacco root plasma membrane vesicles. *Planta* **231**, 425–436.
- Modolo LV, Cunha FQ, Braga MR, Salgado I.** 2002. Nitric oxide synthase-mediated phytoalexin accumulation in soybean cotyledons in response to the *Diaporthe phaseolorum* f. sp. *meridionalis elicitor*. *Plant Physiology* **130**, 1288–1297.
- Moreau M, Lee GI, Wang Y, Crane BR, Klessig DF.** 2008. AtNOS/AtNOA1 is a functional *Arabidopsis thaliana* cGTPase and not a nitric-oxide synthase. *Journal of Biological Chemistry* **283**, 32957–32967.
- Mur LA, Mandon J, Persijn S, et al.** 2013. Nitric oxide in plants: an assessment of the current state of knowledge. *AoB Plants* **5**, pls052.
- Negi S, Santisree P, Kharshiing EV, Sharma R.** 2010. Inhibition of the ubiquitin–proteasome pathway alters cellular levels of nitric oxide in tomato seedlings. *Molecular Plant* **3**, 854–869.
- Ninnemann H, Maier J.** 1996. Indications for the occurrence of nitric oxide synthases in fungi and plants and the involvement in photocondiation of *Neurospora crassa*. *Photochemistry and Photobiology* **64**, 393–398.
- Pagnussat GC, Simontacchi M, Puntarulo S, Lamattina L.** 2002. Nitric oxide is required for root organogenesis. *Plant Physiology* **129**, 954–956.
- Palmieri MC, Lindermayr C, Bauwe H, Steinhauser C, Durner J.** 2010. Regulation of plant glycine decarboxylase by S-nitrosylation and glutathionylation. *Plant Physiology* **152**, 1514–1528.
- Planas-Portell J, Gallart M, Tiburcio AF, Altabella T.** 2013. Copper-containing amine oxidases contribute to terminal polyamine oxidation in peroxisomes and apoplast of *Arabidopsis thaliana*. *BMC Plant Biology* **13**, 109.
- Planchet E, Jagadis Gupta K, Sonoda M, Kaiser WM.** 2005. Nitric oxide emission from tobacco leaves and cell suspensions: rate limiting factors and evidence for the involvement of mitochondrial electron transport. *The Plant Journal* **41**, 732–743.
- Ribeiro EA Jr, Cunha FQ, Tamashiro WM, Martins IS.** 1999. Growth phase-dependent subcellular localization of nitric oxide synthase in maize cells. *FEBS Letters* **445**, 283–286.
- Rockel P, Strube F, Rockel A, Wildt J, Kaiser WM.** 2002. Regulation of nitric oxide (NO) production by plant nitrate reductase in vivo and in vitro. *Journal of Experimental Botany* **53**, 103–110.
- Rümer S, Gupta KJ, Kaiser WM.** 2009. Plant cells oxidize hydroxylamines to NO. *Journal of Experimental Botany* **60**, 2065–2072.
- Sang J, Jiang M, Lin F, Xu S, Zhang A, Tan M.** 2008a. Nitric oxide reduces hydrogen peroxide accumulation involved in water stress-induced subcellular anti-oxidant defense in maize plants. *Journal of Integrative Plant Biology* **50**, 231–243.
- Sang J, Zhang A, Lin F, Tan M, Jiang M.** 2008b. Cross-talk between calcium–calmodulin and nitric oxide in abscisic acid signaling in leaves of maize plants. *Cell Research* **18**, 577–588.
- Santolini J, André F, Jeandroz S, Wendehenne D.** 2017. Nitric oxide synthase in plants: where do we stand? *Nitric Oxide* **63**, 30–38.
- Sanz-Luque E, Ocaña-Calahorra F, de Montaigu A, Chamizo-Ampudia A, Llamas Á, Galván A, Fernández E.** 2015. THB1, a truncated hemoglobin, modulates nitric oxide levels and nitrate reductase activity. *The Plant Journal* **81**, 467–479.
- Seligman K, Saviani EE, Oliveira HC, Pinto-Maglio CA, Salgado I.** 2008. Floral transition and nitric oxide emission during flower development in *Arabidopsis thaliana* is affected in nitrate reductase-deficient plants. *Plant and Cell Physiology* **49**, 1112–1121.
- Simontacchi M, Jasid S, Puntarulo S.** 2004. Nitric oxide generation during early germination of sorghum seeds. *Plant Science* **167**, 839–847.
- Shi H, Ye T, Chen F, Cheng Z, Wang Y, Yang P, Zhang Y, Chan Z.** 2013. Manipulation of arginase expression modulates abiotic stress tolerance in *Arabidopsis*: effect on arginine metabolism and ROS accumulation. *Journal of Experimental Botany* **64**, 1367–1379.
- Shi HT, Li RJ, Cai W, Liu W, Wang CL, Lu YT.** 2012. Increasing nitric oxide content in *Arabidopsis thaliana* by expressing rat neuronal nitric oxide synthase resulted in enhanced stress tolerance. *Plant and Cell Physiology* **53**, 344–357.
- Stöhr C, Stremlau S.** 2006. Formation and possible roles of nitric oxide in plant roots. *Journal of Experimental Botany* **57**, 463–470.
- Stöhr C, Strube F, Marx G, Ullrich WR, Rockel P.** 2001. A plasma membrane-bound enzyme of tobacco roots catalyses the formation of nitric oxide from nitrite. *Planta* **212**, 835–841.
- Stöhr C, Ullrich WR.** 2002. Generation and possible roles of NO in plant roots and their apoplastic space. *Journal of Experimental Botany* **53**, 2293–2303.
- Tian QY, Sun DH, Zhao MG, Zhang WH.** 2007. Inhibition of nitric oxide synthase (NOS) underlies aluminum-induced inhibition of root elongation in *Hibiscus moscheutos*. *New Phytologist* **174**, 322–331.
- Tiburcio AF, Altabella T, Bitrián M, Alcázar R.** 2014. The roles of polyamines during the lifespan of plants: from development to stress. *Planta* **240**, 1–18.
- Tischner R, Galli M, Heimer YM, Bielefeld S, Okamoto M, Mack A, Crawford NM.** 2007. Interference with the citrulline-based nitric oxide synthase assay by argininosuccinate lyase activity in *Arabidopsis* extracts. *FEBS Journal* **274**, 4238–4245.
- Tischner R, Planchet E, Kaiser WM.** 2004. Mitochondrial electron transport as a source for nitric oxide in the unicellular green alga *Chlorella sorokiniana*. *FEBS Letters* **576**, 151–155.
- Tossi V, Cassia R, Lamattina L.** 2009a. Apocynin-induced nitric oxide production confers antioxidant protection in maize leaves. *Journal of Plant Physiology* **166**, 1336–1341.
- Tossi V, Lamattina L, Cassia R.** 2009b. An increase in the concentration of abscisic acid is critical for nitric oxide-mediated plant adaptive responses to UV-B irradiation. *New Phytologist* **181**, 871–879.
- Tossi V, Lamattina L, Cassia R.** 2013. Pharmacological and genetical evidence supporting nitric oxide requirement for 2,4-epibrassinolide regulation of root architecture in *Arabidopsis thaliana*. *Plant Signaling and Behavior* **8**, e24712.
- Tun NN, Santa-Catarina C, Begum T, Silveira V, Handro W, Floh EI, Scherer GF.** 2006. Polyamines induce rapid biosynthesis of nitric oxide (NO) in *Arabidopsis thaliana* seedlings. *Plant and Cell Physiology* **47**, 346–354.
- Valderrama R, Corpas FJ, Carreras A, et al.** 2007. Nitrosative stress in plants. *FEBS Letters* **581**, 453–461.
- Víteček J, Lojek A, Valacchi G, Kubala L.** 2012. Arginine-based inhibitors of nitric oxide synthase: therapeutic potential and challenges. *Mediators of Inflammation* **2012**, 318087.
- Wang BL, Tang XY, Cheng LY, et al.** 2010. Nitric oxide is involved in phosphorus deficiency-induced cluster-root development and citrate exudation in white lupin. *New Phytologist* **187**, 1112–1123.
- Wang J, Krizowski S, Fischer-Schrader K, et al.** 2015. Sulfite oxidase catalyzes single-electron transfer at molybdenum domain to reduce nitrite to nitric oxide. *Antioxidants and Redox Signaling* **23**, 283–294.
- Wang Y, Chen T, Zhang C, Hao H, Liu P, Zheng M, Baluška F, Šamaj J, Lin J.** 2009. Nitric oxide modulates the influx of extracellular Ca²⁺ and actin filament organization during cell wall construction in *Pinus bungeana* pollen tubes. *New Phytologist* **182**, 851–862.
- Wilson ID, Neill SJ, Hancock JT.** 2008. Nitric oxide synthesis and signalling in plants. *Plant, Cell and Environment* **31**, 622–631.
- Wimalasekera R, Tebartz F, Scherer GF.** 2011a. Polyamines, polyamine oxidases and nitric oxide in development, abiotic and biotic stresses. *Plant Science* **181**, 593–603.
- Wimalasekera R, Villar C, Begum T, Scherer GF.** 2011b. COPPER AMINE OXIDASE1 (CuAO1) of *Arabidopsis thaliana* contributes to abscisic

acid- and polyamine-induced nitric oxide biosynthesis and abscisic acid signal transduction. *Molecular Plant* **4**, 663–678.

Xie Y, Mao Y, Lai D, Zhang W, Zheng T, Shen W. 2013. Roles of NIA/ NR/NOA1-dependent nitric oxide production and HY1 expression in the modulation of *Arabidopsis* salt tolerance. *Journal of Experimental Botany* **64**, 3045–3060.

Yamasaki H, Sakihama Y. 2000. Simultaneous production of nitric oxide and peroxyxynitrite by plant nitrate reductase: in vitro evidence for the NR-dependent formation of active nitrogen species. *FEBS Letters* **468**, 89–92.

Yamasaki H, Sakihama Y, Takahashi S. 1999. An alternative pathway for nitric oxide production in plants: new features of an old enzyme. *Trends in Plant Science* **4**, 128–129.

Yan J, Tsuichihara N, Etoh T, Iwai S. 2007. Reactive oxygen species and nitric oxide are involved in ABA inhibition of stomatal opening. *Plant, Cell and Environment* **30**, 1320–1325.

Yang B, Wu J, Gao F, Wang J, Su G. 2014. Polyamine-induced nitric oxide generation and its potential requirement for peroxide in suspension cells of soybean cotyledon node callus. *Plant Physiology and Biochemistry* **79**, 41–47.

Yang J, Giles LJ, Ruppelt C, Mendel RR, Bittner F, Kirk ML. 2015. Oxy and hydroxyl radical transfer in mitochondrial amidoxime reducing component-catalyzed nitrite reduction. *Journal of the American Chemical Society* **137**, 5276–5279.

Yarmolinsky D, Brychkova G, Fluhr R, Sagi M. 2013. Sulfite reductase protects plants against sulfite toxicity. *Plant Physiology* **161**, 725–743.

Yergaliyev TM, Nurbekova Z, Mukiyanova G, et al. 2016. The involvement of ROS producing aldehyde oxidase in plant response to Tombusvirus infection. *Plant Physiology and Biochemistry* **109**, 36–44.

Yesbergenova Z, Yang G, Oron E, Soffer D, Fluhr R, Sagi M. 2005. The plant Mo-hydroxylases aldehyde oxidase and xanthine dehydrogenase have distinct reactive oxygen species signatures and are induced by drought and abscisic acid. *The Plant Journal* **42**, 862–876.

Zarepour M, Simon K, Wilch M, Nieländer U, Koshiba T, Seo M, Lindel T, Bittner F. 2012. Identification of superoxide production by *Arabidopsis thaliana* aldehyde oxidases AAO1 and AAO3. *Plant Molecular Biology* **80**, 659–671.

Zemojtel T, Fröhlich A, Palmieri MC, et al. 2006. Plant nitric oxide synthase: a never-ending story? *Trends in Plant Science* **11**, 524–525; author reply 526.

Zhang H, Zhao X, Yang J, Yin H, Wang W, Lu H, Du Y. 2011. Nitric oxide production and its functional link with OIPK in tobacco defense response elicited by chitoooligosaccharide. *Plant Cell Reports* **30**, 1153–1162.

Zhang JJ, Li XQ, Sun JW, Jin SH. 2014. Nitric oxide functions as a signal in ultraviolet-B-induced baicalin accumulation in *Scutellaria baicalensis* suspension cultures. *International Journal of Molecular Sciences* **15**, 4733–4746.

Zhao MG, Chen L, Zhang LL, Zhang WH. 2009. Nitric reductase-dependent nitric oxide production is involved in cold acclimation and freezing tolerance in *Arabidopsis*. *Plant Physiology* **151**, 755–767.

Zhao MG, Tian QY, Zhang WH. 2007. Nitric oxide synthase-dependent nitric oxide production is associated with salt tolerance in *Arabidopsis*. *Plant Physiology* **144**, 206–217.

Zhou C, Liu Z, Zhu L, Ma Z, Wang J, Zhu J. 2016. Exogenous melatonin improves plant iron deficiency tolerance via increased accumulation of polyamine-mediated nitric oxide. *International Journal of Molecular Sciences* **17**, 1777.