

Clinical Research in Cardiology

Impact of generalized anxiety disorder (GAD) on prehospital delay of acute myocardial infarction patients. Findings from the multicenter MEDEA Study

--Manuscript Draft--

Manuscript Number:	CRCD-D-17-00648R2	
Full Title:	Impact of generalized anxiety disorder (GAD) on prehospital delay of acute myocardial infarction patients. Findings from the multicenter MEDEA Study	
Article Type:	Original Paper	
Keywords:	Generalized anxiety disorder; behavior response; decision time; prehospital delay	
Corresponding Author:	Karl-Heinz Ladwig Helmholtz Zentrum Munchen Deutsches Forschungszentrum fur Umwelt und Gesundheit GERMANY	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Helmholtz Zentrum Munchen Deutsches Forschungszentrum fur Umwelt und Gesundheit	
Corresponding Author's Secondary Institution:		
First Author:	Xiaoyan Fang	
First Author Secondary Information:		
Order of Authors:	Xiaoyan Fang Derek Spieler Loai Albarqouni Joram Ronel Karl-Heinz Ladwig	
Order of Authors Secondary Information:		
Funding Information:	Deutsche Herzstiftung (8810002296)	Prof. Karl-Heinz Ladwig
Abstract:	<p>Background: Anxiety has been identified as a cardiac risk factor. However, less is known about the impact of generalized anxiety disorder (GAD) on prehospital delay during an acute myocardial infarction (AMI). This study assessed the impact of GAD on prehospital delay and delay related cognition and behavior.</p> <p>Methods: Data were from the cross sectional Munich Examination of Delay in Patients Experiencing Acute Myocardial Infarction (MEDEA) study with a total of 619 ST-elevated myocardial infarction (STEMI) patients. Data on sociology-demographic, clinical and psycho-behavioral characteristics were collected at bedside. The outcome was assessed with the Generalized Anxiety Disorder scale (GAD-7). A GAD-7 score greater than or equal to 10 indicates general anxiety disorder.</p> <p>Results: A total of 11.47% (n=71) MI patients suffered from GAD. GAD was associated with increased odds of early arrival (OR: 1.72, 95%CI 1.04-2.88), which was more significant in women (112 vs. 238 mins, p= 0.02) than in men (150 vs. 198 mins, p=0.38). GAD was highly correlated with acute anxiety (p=0.004) and fear of death (p=0.005). Nevertheless, the effect remained significant after controlling for these two covariates. GAD patients were more likely to perceive a higher cardiovascular risk (OR: 2.56, 95%CI 1.37-4.76) in six months before MI, which leading to the higher likelihood of making self-decision to go to hospital (OR: 2.68, 95%CI 1.48-4.85) in the acute phase. However, GAD was also highly associated with impaired psychological well-being, stress and fatigue (p<0.0001).</p> <p>Conclusions: In AMI patients, GAD was independently associated with less prehospital delay, but led to an impaired psychological state.</p>	

Response to Reviewers:

see attachment

Impact of generalized anxiety disorder (GAD) on prehospital delay of acute myocardial infarction patients

Findings from the multicenter MEDEA Study

Fang XY ^{1,2)}, Spieler D, MD ^{1,2)*}, Albarqouni L, MD ¹⁾, Ronel J, MD ²⁾, Ladwig K-H, PhD, MD ^{1,2,3)*}

¹ Institute of Epidemiology II, Mental Health Research Unit, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

² Department of Psychosomatic Medicine and Psychotherapy, Klinikum rechts der Isar, Technische Universität München, Munich, Germany

³ Deutsches Zentrum für Herz-Kreislauf-Forschung (DZHK), Partnersite Munich, Germany

Abstract Word Count: 249

Manuscript Word Count: 3080

(Text body)

Number of Tables: 3

Number of Figures: 1

Number of References: 35

* Shared the first authorship

*Corresponding author:

Prof. Dr. K.H. Ladwig

Institute of Epidemiology II

HelmholtzZentrum München

German Research Center for Environmental Health

Ingolstädter Landstr. 1

85764 Neuherberg, Germany

Phone ++49-89-3187-3623

Fax ++49-89-3187-3667

Email: ladwig@helmholtz-muenchen.de

Website: <http://www.helmholtz-muenchen.de/epi2/research/research-group-3-mental-health-epidemiology/objectives/index.html>

Abstract

Background: Anxiety has been identified as a cardiac risk factor. However, less is known about the impact of generalized anxiety disorder (GAD) on prehospital delay during an acute myocardial infarction (AMI). This study assessed the impact of GAD on prehospital delay and delay related cognition and behaviour.

Methods: Data were from the cross sectional *Munich Examination of Delay in Patients Experiencing Acute Myocardial Infarction (MEDEA)* study with a total of 619 ST-elevated myocardial infarction (STEMI) patients. Data on sociodemographic, clinical and psycho-behavioural characteristics were collected at bedside. The outcome was assessed with the Generalized Anxiety Disorder scale (GAD-7). A GAD-7 score greater than or equal to 10 indicates general anxiety disorder.

Results: A total of 11.47% (n=71) MI patients suffered from GAD. GAD was associated with decreased odds of delay compared to patients without GAD (OR: 0.58, 95%CI 0.35-0.96), which was more significant in women (112 vs. 238 mins, p= 0.02) than in men (150 vs. 198 mins, p=0.38). GAD was highly correlated with acute anxiety (p=0.004) and fear of death (p=0.005). Nevertheless, the effect remained significant after controlling for these two covariates. GAD patients were more likely to perceive a higher cardiovascular risk (OR: 2.56, 95%CI 1.37-4.76) in six months before MI, which leading to the higher likelihood of making self-decision to go to hospital (OR: 2.68, 95%CI 1.48-4.85) in the acute phase. However, GAD was also highly associated with impaired psychological well-being, stress and fatigue (p<0.0001).

Conclusions: In AMI patients, GAD was independently associated with less prehospital delay, but led to an impaired psychological state.

Abstract Word Count: 249

Keywords: Generalized anxiety disorder; behaviour response; decision time; prehospital delay.

Abbreviations:

AMI: Acute Myocardial Infarction, **STEMI:** ST segment Elevation Myocardial Infarction, **PHD:** Prehospital Delay, **MEDEA:** Munich Examination of Delay in Patients Experiencing Acute Myocardial Infarction, **CHD** Coronary Heart Disease, **MACE** Major Adverse Cardiac Event

Introduction

1
2 Anxiety and fear are closely related basic emotions. They comprise anticipatory affective, cognitive, and
3 behavioral changes executed to avoid or reduce the impact of a potential threat or a danger [1]. A key difference
4 between fear and anxiety rests in the certainty or uncertainty of the threat. Fear is the response to a rather certain
5 and objective threat while anxiety is the response to rather uncertain perceived subjective threat. Recent research
6 has provided persuasive neurochemical and neuroanatomical evidence for this psychological distinction [2].
7

8
9 Once these anticipatory processes to uncertainty become maladaptive by being executed disproportionately to
10 the likelihood or severity of the threat, pathological forms of anxiety develop, which can severely interfere with
11 normal live [3,4]. Anxiety disorders have been classified into several distinct disorders described in the DSM-
12 5/ICD-10, one of which is referred to as generalized anxiety disorder (GAD) [5,6]. With GAD, patients present
13 with unfocused worry and anxiety that is not connected to recent stressful events. It is characterized by feelings
14 of threat, restlessness, irritability, insomnia, tension, and physical symptoms such as palpitations, dry mouth, or
15 sweating, lasting six month or longer. Due to the relapsing course of GAD, the disorder is often associated with
16 seriously impaired social and occupational functioning. GAD is a common condition, with life time prevalence
17 rates of 4-7% in the general population [7], women being twice as much affected [8]. In coronary heart disease
18 (CHD) patients, its prevalence is even higher, ranging from 5.42% to 11.57% [9,10].
19
20
21
22
23
24
25

26 Studies examining the impact of GAD on cardiovascular prognosis have yielded conflicting results: On one
27 hand, GAD has been identified as an etiological risk factor of adverse cardiovascular events [11] such as
28 ischemic stroke [12], myocardial infarction [13,9]. On the other hand, recently several large scale studies show
29 that GAD patients had a better prognosis following a cardiac event [14-17]. A probable reason for this positive
30 effect of GAD might be due to higher alertness and increased health promoting behavior [15].
31
32
33

34 Time to treatment is a crucial determinant of survival in patients who have suffered an acute myocardial
35 infarction (AMI) [18,19]: the earlier interventional or thrombolytic therapy is given, the greater the reduction of
36 infarct size and subsequent disability and mortality. Among numerous somatic and psychological factors which
37 have the potential to influence delay time, it is already well established that acute fear and anxiety during AMI
38 onset reduce the decision delay to seek medical help [10, 11]. However, no study has been conducted so far to
39 investigate the role of GAD on prehospital delay during AMI.
40
41
42
43

44 Thus, the objectives of our study are: 1) to assess the impact of GAD on prehospital delay and 2) to test whether
45 a putative effect of GAD remains even after controlling for acute anxiety conditions, 3) to assess the impact of
46 GAD on patient's behavioral responses to the symptoms during the acute phase of an AMI and 4) to further
47 explore the impact of GAD on the post-acute course of AMI.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Methods

The multicenter, retrospective cross-sectional MEDEA study (*Munich Examination of Delay in Patients Experiencing Acute Myocardial Infarction*) was conceived with the aim to evaluate prehospital delay of STEMI patients, and the factors which may contribute to prolonged delay.

Study design

The patients were recruited from eight different university or municipal hospitals with coronary care units, belonging to the Munich emergency system network clinics. The MEDEA study was approved by the Ethic Commission of the Faculty of Medicine of the Technische Universität München (TUM) on 10.12.2007 and the consent of the Munich Institut für klinische Forschung (IKF) for the participating four municipal hospitals (9.4.2008). The main inclusion criterion was diagnosis of STEMI as evidenced by typical clinical symptoms, ECG changes and myocardial biomarkers levels. Exclusion criteria were: In-hospital STEMI, resuscitation at AMI-onset and language barriers or cognitive impairment impeding patients to answer the questionnaires properly. There were no age restrictions.

Standardized operation procedures (SOPs) were implemented to ensure the consecutive referral of eligible patients into the study. All patients were informed of the aim and procedures of the study and also that taking part in the study would have no effect on their treatment. All patients were required to sign a declaration of consent. Bed-side interviews and self-administered questionnaires were conducted in the hospital ward within 24 h after referral from intensive care.

Sample

From December, 12. 2007 until May, 31. 2012, a total of 755 patients were screened for eligibility. In 619 patients, a diagnosis of STEMI was confirmed. Approximately 18% of patients were excluded: 4% due to not meeting inclusion criteria and 14% due to absence of consent.

Data collection

The data collection process was divided into three sections. Firstly, a structured bedside interview was conducted with trained personnel. Secondly, a self-administered questionnaire was filled by the patient without supervision. Thirdly, data were collected from the hospitals' patient charts.

Measures

Prehospital Delay (PHD)

Patients were asked to recall at what time acute symptoms began. The time difference between symptom onset and first ECG at hospital entry constitutes "prehospital delay" (PHD), measured in minutes. PHD was thus available as a continuous variable which was heavily left-skewed and did not approximate a normal distribution after log-transformations.

Generalized anxiety disorder

Anxiety was assessed with the German version of Generalized Anxiety Disorder scale (GAD-7). It is composed of 7 items, rated on a 4-point Likert scale from not present to very high, leading to an overall score ranging from

0 to 21. A suspected diagnosis of GAD is defined by a GAD-7 score greater than or equal to 10. Using the threshold score of 10, the GAD-7 has a sensitivity of 82% for GAD. [20]

Psychological measures

Depression was assessed with the Major Depression Inventory (MDI) - a self-report mood questionnaire able to generate an ICD-10 or DSM-IV diagnosis of clinical depression. The MDI contains 12 items. According to the DSM-IV definition, patients who had at least five symptoms in the MDI scale, of which at least one must be a 'core' symptom, were diagnosed with major depression [21].

Well-being was evaluated through the WHO-Five Well-Being index. It contains five items on a 6-point scale that range from 0 to 25. Thereafter, the raw scores are transformed into a scale that range from 0 to 100. [22]. WHO-5 score less than or equal to 50 indicates suboptimal well-being[23]. Effectiveness of the index has been supported in evaluation of emotional well-being in patients with cardiovascular diseases.

Vital exhaustion was assessed using a 4-item index on a 5 point Likert Scale that range from 0 to 16. Two items are from The Maastricht Questionnaire ("Do you often feel tired?" and "Do you often feel weak all over?"). The other two were obtained from the CES-D ("I felt that everything I did was an effort" and "I could not get going"). In present study, we applied the median split as a cut-off point, leading to an exhausted (>7) and non-exhausted (<=7) group. The predictive validity of the exhaustion index has been reported elsewhere 3.18 and the internal consistency (Cronbach's) of this scale were 0.55 [24].

Psychological stress was assessed with three single-item questions relating to stress at work, at home and financial stress, rated on a 4 point Likert scale, ranging from 3 (never) to 12 (permanent stress). Stress was defined as feeling irritable, filled with anxiety, or as having sleeping difficulties as a result of conditions at work or at home. In present study, we applied the median split as a cut-off point, leading to a stressed (>5) and non-stressed (<=5) group.

Patient behavioral responses to STEMI

A German version of the Response to Symptoms Questionnaire was applied [25], which assesses the behavior and subsequent reactions of both the patient as well as witnesses in the following areas: social context in which symptoms occurred and bystanders responses, behavioral responses to the symptoms, cognitive responses to the symptoms and emotional responses to the symptoms. The instrument also includes one item on symptom expectation, which measures the congruence between symptom expectation and perception (11 items, 5 point Likert scale, >3 rated was used as cut-off to define a high level).

Data analysis

Differences between dichotomous variables were assessed using the Chi-square test. When comparing ordinal variables with more than two outcomes, the Mantel-Haenszel Chi-square test was used. Differences in age were assessed using the t test. The nonparametric Wilcoxon test was used for assessing differences in median prehospital delay times. Multivariate Logistic regression model was applied to assess the association between GAD and patients' responses to the symptom onset. In addition, the additional effect of stress and exhaustion on patients' responses was also assessed by logistic regression model. Because anxiety level is highly correlated

1 with other psychometric factors, logistic regression with different grades of adjustments for psychological
2 factors was applied to assess the association between GAD and the chance of longer delay. Patients who delayed
3 more than two hours are defined as delayed group. Adjustments were made for fear of death, acute anxiety
4 during the symptom onset (model 2), and additionally for stress (model 3), exhaustion (model 4) and depression
5 (model 5).The relative risk for longer delay is presented as odds ratio (OR) with 95% confidence interval (95%
6 CI).
7
8

9
10 All statistical analyses were run in SAS (Version 9.3, SAS-Institute Inc., Cary, NC, USA). The significance
11 level was set at $p < 0.05$. The analysis and description in this paper follow the STROBE guidelines for cross-
12 sectional studies [26].
13
14
15
16
17
18

19 **Results**

20
21 A total of 619 patients were included in the present study with 162 (26.17%) women and 457 (73.83%) men
22 aged between 30 and 93 years (mean age 62.50 years, $SD = 12.15$). In the total sample, the median delay time
23 was 200 (100-652) minutes.
24
25

26 **Prevalence and distribution of GAD in patients with STEMI**

27
28 The GAD-7 score was right skewed with a mean of 5.98 ± 4.40 and a median of 5, leading to a total of 71 (11%)
29 patients with GAD ($GAD-7 \geq 10$). We identified a similar prevalence in women (11.11%) and in men
30 (11.60%) ($p = 0.87$). As shown in **Table 1**, patients with GAD were more likely to be younger ($p = 0.05$) but did
31 not show differences with respect to social demographic characteristics (education levels, employment status
32 and living situation).
33
34
35

36 **Characteristics of patients with GAD during the six months prior to STEMI**

37
38 As displayed in **Table 1**, patients with GAD were more likely to report stress ($p < 0.0001$), vital exhaustion (p
39 < 0.0001), suboptimal well-being ($p < 0.0001$) and depression ($p < 0.0001$).
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Sociodemographic and clinical characteristics of the study population stratified by with GAD (n=71) and without GAD (n=548).

	GAD (n=619)			P value
	Missing	With GAD	Without GAD	Overall
All Patients	-	71 (11.47%)	548 (88.53%)	
Socio-demographic Factors				
Age(>65)	-	58.71 ±11.91	62.98±12.86	0.005
Sex (Female)		18 (11.11%)	144 (88.89%)	0.87
Sex (Male)	-	53 (11.60%)	404 (88.40%)	
Living alone	-	27 (38.03%)	155 (28.28%)	0.09
Employed	-	37 (52.11%)	267 (48.72%)	0.59
Education (secondary school and above)	-	25 (35.21%)	231 (42.15%)	0.26
AMI symptoms				
Chest pain		63 (88.73%)	489 (89.23%)	0.90
Dyspnea		29 (40.85%)	165 (30.16%)	0.07
Racing heart		10 (14.08%)	44 (8.03%)	0.22
Sweating		48 (67.61%)	305 (55.76%)	0.06
Faint		5 (7.04%)	30 (5.48%)	0.59
Exhaustion		16 (22.54%)	74 (13.53%)	0.04
Vomiting		11 (15.49%)	78 (14.26%)	0.78
Nausea		29 (40.85%)	212 (38.76%)	0.73
Heart burn		6 (8.45%)	32 (5.85%)	0.39
Stomachache		9 (12.68%)	37 (6.76%)	0.07
Psychological factors				
Perceived stress		56 (24.89%)	10 (3.47)	<0.0001
Vital exhaustion		53 (23.66%)	18 (4.56)	<0.0001
Fear of death		20 (28.99%)	68 (13.00%)	0.0005
Acute anxiety		31 (43.66%)	148(27.21%)	0.004
Depression		25 (35.21%)	13(2.82%)	<0.0001
Optimal well-being		21 (29.58%)	367 (66.97%)	<0.0001

Values are n (%). Bold means significant p values at <0.05 level.

Impact of GAD on patients' symptom perception, behaviour responses during STEMI

In the acute phase of STEMI, patients with GAD were more likely to perceive exhaustion (p=0.04), fear of death (p=0.0005) and a higher level of acute anxiety (p=0.004). As can be seen in **Table 2**, patients with GAD perceived a higher subjective cardiovascular risk as compared to patients without GAD (OR: 0.39, 95%CI 0.21-0.73) and were more likely to make self-decisions to go to the hospital (OR: 2.68, 95%CI 1.48-4.85). The associations remained significant in GAD patients who additionally suffered from stress or vital exhaustion.

Table 2. The impact of GAD, further stratified for GAD population with stress (n=56) and exhaustions (n=53)

	GAD vs. No GAD (71 vs. 548) OR (95% CI)	GAD with stress vs. Others (56 vs 457) OR (95% CI)	GAD with exhaustion vs. Others (53 vs 566) OR (95% CI)
Cognitive responses			
Heart misattribution	1.00 0.61-1.65	1.01 0.58-1.75	0.97 0.55-1.71
Failed to recognize the symptoms as MI	1.32 0.81-2.17	1.51 0.87-2.63	1.47 0.83-2.59
Insufficient risk perception	0.39 0.21-0.73	0.32 0.16-0.61	0.36 0.18-0.72
Behavioral responses			
Take medicine	0.86 0.52-1.44	1.06 0.52-2.17	1.29 0.59-2.81
Wait until the symptom resolve	0.70 0.43-1.16	0.69 0.39-1.19	0.90 0.50-1.61
Continue doing the activity on going	1.01 0.54-1.88	1.03 0.52-2.06	1.06 0.52-2.18
Try to relax	1.25 0.75-2.10	1.05 0.60-1.85	1.45 0.79-2.63
Call someone for help	2.32 0.55-9.88	1.80 0.42-7.68	1.65 0.39-7.07
Call general physician	0.95 0.41-2.17	1.41 0.49-4.05	1.29 0.45-3.72
Call emergency doctor	1.24 0.75-2.05	1.21 0.69-2.12	1.52 0.86-2.69
Used ambulance to get to the hospital	0.86 0.52-1.44	0.90 0.51-1.59	0.76 0.42-1.38
Drive themselves to the hospital	1.28 0.68-2.41	1.16 0.58-2.31	1.63 0.75-3.55
Made self-decision to go to the hospital	2.68 1.48-4.85	2.89 1.46-5.70	2.67 1.35-5.29
Post-acute course			
With complication	0.44 0.20-0.99	0.60 0.26-1.35	0.43 0.17-1.10*
Cardiac arrest	2.11 0.76-5.84	2.81 1.01-7.83	0.97 0.00-4.24
Intensive care \geq 3 days	0.91 0.54-1.52	0.95 0.53-1.68	0.89 0.50-1.59

Bold means significant p values at <0.05 level. *p=0.08

Impact of GAD on prehospital delay

The median delay time in patients with GAD tended to be shorter than in patients without GAD (median delay time 134 vs. 213 mins, $p=0.059$).

GAD was associated with decreased odds of delay (delay time $>$ or \leq 2hrs) compared to patients without GAD (OR: 0.58, 95%CI 0.35-0.96). As can be seen in **Table 3**, the effects were independent from the acute anxiety at onset of symptoms and even fear of death (**model 2**) and remained significant after stepwise adjustment for stress, exhaustion and depression (**model 3-5**).

As can be seen in **Figure 1**, sex stratified analysis illustrated that the effect of GAD on prehospital delay in women (112 vs. 238 mins, $p=0.02$) is more significant than in men (150 vs. 198 mins, $p=0.38$). Likewise, GAD was associated with decreased odds of delay longer than 2 hours in women (OR: 0.30, 95%CI 0.11-0.85, $p=0.02$) but not in men (OR: 0.71, 95%CI 0.40-1.30, $p=0.26$).

Table 3. Association of GAD and pre-hospital delay assessed by logistic regression, adjusted by fear of death, acute anxiety, stress, exhaustion and depression

Delay > 2hrs vs. Delay \leq 2hrs (426 vs. 193)					
OR (95% CI)					
Emotional factors	model 1	model 2	model 3	model 4	model 5
GAD	0.58 0.35-0.96	0.60 0.35-0.99	0.48 0.27-0.84	0.49 0.27-0.89	0.50 0.26-0.97
Fear of death		0.64 0.33-1.24	0.77 0.35-1.67	0.77 0.35-1.67	0.78 0.36-1.71
Acute anxiety		0.96 0.91-1.01	0.96 0.90-1.02	0.96 0.90-1.02	0.96 0.90-1.01
Stress			1.05 0.94-1.18	1.05 0.94-1.18	1.04 0.93-1.12
Exhaustion				1.01 0.67-1.52	0.98 0.67-1.50
Depression					1.00 0.44-2.26

Bold means significant p values at <0.05 level.

All the model were adjusted for sex and age

Model 1: The crude model

Model 2: Adjusted with acute anxiety condition (including fear of death and acute anxiety)

Model 3: Further adjusted with self-perceived burden of daily stress

Model 4: Further adjusted with vital exhaustion

Model 5: Further adjusted with depression

The post-acute course of patients with GAD

In the post-acute infarction phase during ICU stay, patients with GAD were less likely to have complications (OR: 0.44, 95%CI 0.22-0.99). The GAD patients additionally suffering from stress were more likely to experience in-hospital cardiac arrest, but did not show differences regarding complication and ICU stay compared to their counterparts. GAD patients suffering additionally from vital exhaustion tended to experience less cardiac complications (OR: 0.43, 95%CI 0.17-1.10).

Discussion

To the best of our knowledge, this is the first comprehensive evaluation of the impact of GAD on prehospital delay in patients facing an AMI. The major finding of the present study is that GAD had a favorable effect on reducing prehospital delay during AMI. This effect of GAD on prehospital delay was significant in women while in men we identified solely a non-significant trend. Moreover, GAD was associated with better prognosis in the post-acute phase of AMI.

Patients suffered from GAD also presented a comorbidity pattern of impaired mental health, meaning the patients with GAD were also significantly more likely to suffer from acute anxiety, depression, vital exhaustion and perceived stress. It has been well documented that pronounced acute anxiety/fear owing to the sudden onset of the life-threatening AMI leads to a shorter delay time, hereby favoring a good prognosis [27-29].

Of note, the beneficial effects of GAD on prehospital delay and prognosis found in our homogeneous STEMI sample remained significant even after we controlled for acute fear of death [30], depression, exhaustion and perceived stress. This finding underscores that GAD is a powerful and independent protective factor on its own in patients facing an AMI.

This is a remarkable finding, which points to a specific alertness of GAD patients more likely to be present at the time long before the onset of AMI. This assumption is supported by our finding showing that GAD patients had a higher self-perceived MI risk than non-GAD patients. In that line, GAD has been found to be a ‘driver’ for individuals to address their health needs more regularly and conscientiously and seek help at the early signs of the disease. Dubayova et al. [31] reported in a systemic review including 15 studies that being ‘anxious’ has a significant positive effect on decision making in help seeking behavior. Parker et al. [14] found that GAD patients received more medical test and tended to take part more often in post-AMI rehabilitation programs. Interestingly, GAD patients did not experience a different pattern of acute symptoms compared to non-GAD patients. This is noteworthy because it is unlikely that the GAD patients sought help faster because of more severe symptoms.

Moreover, the study reveals the association of GAD patients with a better prognosis in the post-acute phase of AMI. It is not unlikely that this is a consequence of the reduced delay time in GAD patients as well, based on the earlier treatment and hereby improved course with less symptoms, since every minute of delay to treatment for STEMI has previously been shown to affect the 1-year mortality [32]. Yet, the post-acute outcome was not favorable anymore, if GAD was accompanied by stress or exhaustion (**Table 2**).

Contrary to expectation, we found no sex difference of GAD prevalence in our clinical sample. This is remarkable because in general population, women are twice as much affected with GAD than men [8]. The analysis shows a sex specific impact of anxiety on delay time though. In women, the difference of delay time was highly significant, whereas in men, there was only a trend towards a reduced delay. Currently, we have no possible reasons to explain the differences.

Although this study identified favorable effects in patients meeting GAD criteria having shorter time to treatment and fewer complications, it seems to be essential to balance this ‘advantage’ with the disease burden

1 of GAD itself: GAD patients were more likely to suffer from higher levels of negative emotions (including
2 depression, exhaustion and perceived stress and thus impaired psychological well-being). This is in line with the
3 observation showing that anxiety and depression frequently co-occur [33,34].
4

5 To our knowledge, this is the first study investigating the impact of generalized anxiety disorder on prehospital
6 delay in a strictly defined population of STEMI patients. There are a few study limitations that are worth
7 considering. First, all data were collected at bedside within a very narrow time frame (< 24 hours after referral
8 from intensive care) after STEMI, nevertheless we cannot fully exclude the possibility of recall bias. We had
9 relatively small numbers of women, so replications of these results in larger datasets are warranted. Furthermore,
10 selection bias could have resulted from excluding STEMI patients who died before reaching the hospital. Finally,
11 GAD diagnosis was based on GAD-7 questionnaire data which provides a sensitivity of 82% for GAD [35]
12 using a threshold score of 10.
13
14
15
16

17 **Conclusion**

18
19
20 Our study demonstrates that in patients facing an AMI, GAD is associated with an increased chance of early
21 arrival and thus had fewer complications, despite its known adverse effects on psychological well-being. The
22 higher perceived MI risk and the higher chance of making self-decision to seek medical help in GAD patients
23 suggests that GAD patients are particularly sensitive to early sign of the disease, ultimately resulting in shorter
24 time to treatment and better prognosis. The shorter delay time and appropriate behavioral responses during AMI
25 indicated the protective effect of GAD on patients' acute situation. However, our study does not provide
26 information regarding long term effect of GAD on patients' cardiac outcome. Further investigation will be
27 necessary to reveal whether the impaired psychological well-being caused by GAD will affect long term
28 prognosis, in order to provide clinically implication for the appropriate timing to intervene GAD in CHD
29 patients.
30
31
32
33
34
35
36
37
38
39
40
41

42 **Funding**

43
44 Our work was supported by a research grant from the Deutsche Herzstiftung (8810002296) (to Prof. Dr. KH
45 Ladwig). This investigation was realized under the umbrella of the Munich Heart Alliance (MHA).
46
47

48 **Conflict of interest**

49
50 None to declare.
51

52 **Acknowledgements**

53
54 Cooperating clinics in the city of Munich (Germany): Klinikum-Augustinum (Prof. Dr. Michael Block),
55 Klinikum-Bogenhausen (Prof.Dr. Ellen Hoffmann), Deutsche-Herz-Zentrum München (Prof. Dr.Heribert
56 Schunkert), Klinikum-Harlaching (Prof. Dr. Harald Köhl), Universitäts-Klinikum der LMU-Innenstadt (Prof. Dr.
57
58
59
60
61
62
63
64
65

1 Axil Bauer), Klinikum-Neuperlach (Prof. Dr. Harald Mudra), Universitäts-Klinikum Rechts der Isar-der-TUM
2 (Prof. Dr. Karl-Ludwig Laugwitz) and Klinikum-Schwabing (Prof. Dr. Stefan Sack).
3
4
5
6
7

8 **Ethic statements**

9
10 All patients were informed of the aim and procedures of the study and also that taking part in the study would
11 have no effect on their treatment. All patients were required to sign a declaration of consent. Details that might
12 disclose the identity of the subjects under study has been omitted.
13
14
15
16
17

18 **Figure Captions**

19
20 **Fig 1.** Non-parametric test for comparing median delay time (in min) for all patients with and without GAD and
21 stratified for women and men.
22
23
24
25
26
27

28 **References**

- 29
30 1. Grupe DW, Nitschke JB (2013) Uncertainty and anticipation in anxiety: an integrated neurobiological and
31 psychological perspective. *Nat Rev Neurosci* 14 (7):488-501. doi:10.1038/nrn3524
32
33 2. Tovote P, Fadok JP, Luthi A (2015) Neuronal circuits for fear and anxiety. *Nat Rev Neurosci* 16 (6):317-331.
34 doi:10.1038/nrn3945
35
36 3. Moser DK (2007) "The rust of life": impact of anxiety on cardiac patients. *Am J Crit Care* 16 (4):361-369
37
38 4. Celano CM, Millstein RA, Bedoya CA, Healy BC, Roest AM, Huffman JC (2015) Association between
39 anxiety and mortality in patients with coronary artery disease: A meta-analysis. *Am Heart J* 170 (6):1105-1115.
40 doi:10.1016/j.ahj.2015.09.013
41
42 5. APA (May 18, 2013) *Diagnostic and Statistical Manual of Mental Disorders*. 5th edn., Washington, DC
43
44 6. WHO (1992) *The ICD-10 classification of mental and behavioural disorders: Clinical descriptions and*
45 *diagnostic guidelines*. Geneva: World Health Organization.
46
47 7. Hoge EA, Ivkovic A, Fricchione GL (2012) Generalized anxiety disorder: diagnosis and treatment. *BMJ*
48 345:e7500. doi:10.1136/bmj.e7500
49
50 8. Wittchen HU (2002) Generalized anxiety disorder: prevalence, burden, and cost to society. *Depress Anxiety*
51 16 (4):162-171. doi:10.1002/da.10065
52
53 9. Tully PJ, Cosh SM, Baumeister H (2014) The anxious heart in whose mind? A systematic review and meta-
54 regression of factors associated with anxiety disorder diagnosis, treatment and morbidity risk in coronary heart
55 disease. *J Psychosom Res* 77 (6):439-448. doi:10.1016/j.jpsychores.2014.10.001
56
57 10. Blumenthal JA, Feger BJ, Smith PJ, Watkins LL, Jiang W, Davidson J, Hoffman BM, Ashworth M, Mabe
58 SK, Babyak MA, Kraus WE, Hinderliter A, Sherwood A (2016) Treatment of anxiety in patients with coronary
59
60
61
62
63
64
65

heart disease: Rationale and design of the UNderstanding the benefits of exercise and escitalopram in anxious patients With coroNary heart Disease (UNWIND) randomized clinical trial. *Am Heart J* 176:53-62.

doi:10.1016/j.ahj.2016.03.003

11. Emdin CA, Oduyayo A, Wong CX, Tran J, Hsiao AJ, Hunn BH (2016) Meta-Analysis of Anxiety as a Risk Factor for Cardiovascular Disease. *Am J Cardiol* 118 (4):511-519. doi:10.1016/j.amjcard.2016.05.041

12. Perez-Pinar M, Ayerbe L, Gonzalez E, Mathur R, Foguet-Boreu Q, Ayis S (2017) Anxiety disorders and risk of stroke: A systematic review and meta-analysis. *Eur Psychiatry* 41:102-108. doi:10.1016/j.eurpsy.2016.11.004

13. Martens EJ, de Jonge P, Na B, Cohen BE, Lett H, Whooley MA (2010) Scared to death? Generalized anxiety disorder and cardiovascular events in patients with stable coronary heart disease:The Heart and Soul Study. *Arch Gen Psychiatry* 67 (7):750-758. doi:10.1001/archgenpsychiatry.2010.74

14. Parker G, Hyett M, Hadzi-Pavlovic D, Brotchie H, Walsh W (2011) GAD is good? Generalized anxiety disorder predicts a superior five-year outcome following an acute coronary syndrome. *Psychiatry Res* 188 (3):383-389. doi:10.1016/j.psychres.2011.05.018

15. Herrmann C, Brand-Driehorst S, Buss U, Ruger U (2000) Effects of anxiety and depression on 5-year mortality in 5,057 patients referred for exercise testing. *J Psychosom Res* 48 (4-5):455-462. doi:
[http://dx.doi.org/10.1016/S0022-3999\(99\)00086-0](http://dx.doi.org/10.1016/S0022-3999(99)00086-0)

16. Meyer T, Herrmann-Lingen C, Chavanon ML, Nolte K, Pasedach CA, Binder L, Pieske B, Hasenfuss G, Wachter R, Edelmann F (2015) Higher plasma levels of MR-pro-atrial natriuretic peptide are linked to less anxiety: results from the observational DIAST-CHF study. *Clin Res Cardiol* 104 (7):574-581. doi:10.1007/s00392-015-0820-9

17. Kobe J, Hucklenbroich K, Geisendorfer N, Bettin M, Frommeyer G, Reinke F, Dechering D, Burgmer M, Eckardt L (2017) Posttraumatic stress and quality of life with the totally subcutaneous compared to conventional cardioverter-defibrillator systems. *Clin Res Cardiol* 106 (5):317-321. doi:10.1007/s00392-016-1055-0

18. O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Jr., Chung MK, de Lemos JA, Ettinger SM, Fang JC, Fesmire FM, Franklin BA, Granger CB, Krumholz HM, Linderbaum JA, Morrow DA, Newby LK, Ornato JP, Ou N, Radford MJ, Tamis-Holland JE, Tommaso JE, Tracy CM, Woo YJ, Zhao DX (2013) 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 127 (4):529-555. doi:10.1161/CIR.0b013e3182742c84

19. Ladwig KH, Meisinger C, Hymer H, Wolf K, Heier M, von Scheidt W, Kuch B (2011) Sex and age specific time patterns and long term time trends of pre-hospital delay of patients presenting with acute ST-segment elevation myocardial infarction. *Int J Cardiol* 152 (3):350-355. doi:10.1016/j.ijcard.2010.08.003

20. Spitzer RL, Kroenke K, Williams JB, Lowe B (2006) A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 166 (10):1092-1097. doi:10.1001/archinte.166.10.1092


21. Olsen LR, Jensen DV, Noerholm V, Martiny K, Bech P (2003) The internal and external validity of the Major Depression Inventory in measuring severity of depressive states. *Psychol Med* 33 (2):351-356

22. Birket-Smith M, Rasmussen A (2008) Screening for mental disorders in cardiology outpatients. *Nord J Psychiatry* 62 (2):147-150. doi:10.1080/08039480801983562

23. Bech P (2004) Measuring the Dimension of Psychological General Well-Being by the WHO-5. . *QoL Newsletter* 32:15-16

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
24. Kop WJ, Gottdiener JS, Tangen CM, Fried LP, McBurnie MA, Walston J, Newman A, Hirsch C, Tracy RP (2002) Inflammation and coagulation factors in persons > 65 years of age with symptoms of depression but without evidence of myocardial ischemia. *Am J Cardiol* 89 (4):419-424. doi: [http://dx.doi.org/10.1016/S0002-9149\(01\)02264-0](http://dx.doi.org/10.1016/S0002-9149(01)02264-0)
25. Burnett RE, Blumenthal JA, Mark DB, Leimberger JD, Califf RM (1995) Distinguishing between early and late responders to symptoms of acute myocardial infarction. *Am J Cardiol* 75 (15):1019-1022
26. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP (2008) The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 61 (4):344-349. doi:10.1016/j.jclinepi.2007.11.008
27. Moser DK, McKinley S, Dracup K, Chung ML (2005) Gender differences in reasons patients delay in seeking treatment for acute myocardial infarction symptoms. *Patient Educ Couns* 56 (1):45-54. doi:10.1016/j.pec.2003.11.011
28. Khraim FM, Carey MG (2009) Predictors of pre-hospital delay among patients with acute myocardial infarction. *Patient Educ Couns* 75 (2):155-161. doi:10.1016/j.pec.2008.09.019
29. Abed MA, Khalil AA, Moser DK (2015) The Contribution of Symptom Incongruence to Prehospital Delay for Acute Myocardial Infarction Symptoms Among Jordanian Patients. *Res Nurs Health* 38 (3):213-221. doi:10.1002/nur.21658
30. Albarqouni L, Rothe AV, Ronel J, Meinertz T, Ladwig KH (2016) Frequency and covariates of fear of death during myocardial infarction and its impact on prehospital delay: findings from the multicentre MEDEA Study. *Clin Res Cardiol* 105 (2):135-144. doi:10.1007/s00392-015-0895-3
31. Dubayova T, van Dijk JP, Nagyova I, Rosenberger J, Havlikova E, Gdovinova Z, Middel B, Groothoff JW (2010) The impact of the intensity of fear on patient's delay regarding health care seeking behavior: a systematic review. *Int J Public Health* 55 (5):459-468. doi:10.1007/s00038-010-0149-0
32. Guerchicoff A, Brener SJ, Maehara A, Witzenbichler B, Fahy M, Xu K, Gersh BJ, Mehran R, Gibson CM, Stone GW (2014) Impact of delay to reperfusion on reperfusion success, infarct size, and clinical outcomes in patients with ST-segment elevation myocardial infarction: the INFUSE-AMI Trial (INFUSE-Anterior Myocardial Infarction). *JACC Cardiovasc Interv* 7 (7):733-740. doi:10.1016/j.jcin.2014.01.166
33. Tyrer P, Baldwin D (2006) Generalised anxiety disorder. *Lancet* 368 (9553):2156-2166. doi:10.1016/s0140-6736(06)69865-6
34. Bunz M, Lenski D, Wedegartner S, Ukena C, Karbach J, Bohm M, Kindermann I (2016) Heart-focused anxiety in patients with chronic heart failure before implantation of an implantable cardioverter defibrillator: baseline findings of the Anxiety-CHF Study. *Clin Res Cardiol* 105 (3):216-224. doi:10.1007/s00392-015-0909-1
35. Kroenke K, Spitzer RL, Williams JB, Monahan PO, Lowe B (2007) Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. *Ann Intern Med* 146 (5):317-325

Delay in Minutes



Corresponding author:

Karl-Heinz Ladwig, PhD, MD, habil.
Helmholtz Zentrum München
German Research Center for Environmental Health
Institute of Epidemiology II
Ingolstädter Landsstraße 1
85764 Neuherberg, Germany
Phone +49-89-3187-3623
Fax +49-89-3187-3667

Dear Prof. M. Böhm, dear Prof.A. Katus

We thank you for your willingness to reconsider an abridged version of our manuscript. Please find enclosed the revised version of our manuscript (Ms. No. CRC-D-17-00648) entitled “**Impact of generalized anxiety disorder (GAD) on prehospital delay of acute myocardial infarction patients**” for consideration in *Clinical Research in Cardiology*.

We would also like to take this opportunity to thank you and the reviewers for the very valuable comments (dated January 19th, 2018). The reviewer’s comments have been carefully considered and changes have been made to the manuscript which we hope have led to a significant improvement. In the second round, reviewer#2 asked for some additional points. Although we are unable to follow the reviewer’s suggestion completely, we have updated our manuscript by providing a comprehensive regression of all psychological conditions as the reviewer asked for and also have discussed this particular issue in the discussion part. Please find changes to the manuscript highlighted in yellow. Our responses to the reviewer are listed below.

All authors have read and approved the final revised manuscript. The results of this paper have not been published elsewhere nor are they under consideration at any other journal.

With kind regards,

Prof. Dr. Karl-Heinz Ladwig

Responses to the reviewer comments:

We appreciate the reviewers are satisfied follow the revision of our manuscript highlighting “**Impact of generalized anxiety disorder (GAD) on prehospital delay of acute myocardial infarction patients**”. There is an additional issue in the second round of review which remains to be addressed.

Reviewer #2: I only have one comment left regarding question 3. The authors stated that GAD was prospectively defined as a Parameter to be analyzed, but they plan to analyze perceived stress, vital exhaustion, acute anxiety, and depression in upcoming manuscripts. As there is a lot of overlap, it may turn out that similar effect as for GAD can be seen for the other parameters. I proposed to do an analysis with these factors instead of table 3 to give the reader a better impression whether e. g. vital exhaustion gives similar information. I think this topic should also be covered in the discussion. It does not help to provide a manuscript saying GAD prolongs hospital stay and come up with another manuscript a year later: We found out that vital exhaustion is doing the same... We want to know which of these factors is the most relevant or do they just carry the same information.

First, we apologize for our misleading answer in the first response to reviewer about this issue. The reviewer is actually right in the last comments that “**if GAD is by far the most relevant influence factor, they should continue with the proposed multivariate model**”. Hereby, we offer the following finding as background information: patients with acute anxiety also delay less than those who without acute anxiety (135 vs. 213mins, $p=0.002$). However, there is no significant difference of delay time between patients with or without stress (197 vs. 207 $p=0.96$), exhaustion (222.5 vs 200 $p=0.42$), depression (157.5 vs 210, $p=0.32$). In addition, as marked in **Table 3** below, the effect of acute anxiety is no longer significant in the regression model adjusted by GAD. Taken together, GAD is exactly the most relevant factor.

Table 3. Association of GAD and pre-hospital delay assessed by logistic regression, adjusted by fear of death, acute anxiety, stress, exhaustion and depression

	Delay > 2hrs vs. Delay <= 2hrs (426 vs. 193)				
	OR (95% CI)				
Emotional factors	model 1	model 2	model 3	model 4	model 5
GAD	0.58 0.35-0.96	0.60 0.35-0.99	0.48 0.27-0.84	0.49 0.27-0.89	0.50 0.26-0.97
Fear of death		0.64 0.33-1.24	0.77 0.35-1.67	0.77 0.35-1.67	0.78 0.36-1.71
Acute anxiety		0.96 0.91-1.01	0.96 0.90-1.02	0.96 0.90-1.02	0.96 0.90-1.01
Stress			1.05 0.94-1.18	1.05 0.94-1.18	1.04 0.93-1.12
Exhaustion				1.01 0.67-1.52	0.98 0.67-1.50
Depression					1.00 0.44-2.26

Bold means significant p values at <0.05 level.

All the model were adjusted for sex and age

Model 1: The crude model

Model 2: Adjusted with acute anxiety condition (including fear of death and acute anxiety)

Model 3: Further adjusted with self-perceived burden of daily stress

Model 4: Further adjusted with vital exhaustion

Model 5: Further adjusted with depression

As has been mentioned by the other reviewer, we have applied the complete adjustment strategy in **Table 3** and added further explanation in the legend. The completely adjusted model in **Table 3** shows that acute anxiety and other mental health factors are not significant when they are adjusted with GAD. Hence, the **Table 3** shows a powerful and independent effect of GAD on delay which is not affected by the other mental health factors.

Following the suggestion of the reviewer, we also add this point to the discussion part; **Page 10, Para 2:**

Patients suffered from GAD also presented a comorbidity pattern of impaired mental health, meaning the patients with GAD were also significantly more likely to suffer from acute anxiety, depression, vital exhaustion and perceived stress. It has been well documented that pronounced acute anxiety/fear owing to the sudden onset of the life-threatening AMI leads to a shorter delay time, hereby favoring a good prognosis [27-29]. Of note, the beneficial effects of GAD on prehospital delay and prognosis found in our homogeneous STEMI sample remained significant even after we controlled for acute fear of death [30], *depression, exhaustion and perceived stress.* *This finding underscores that GAD is a powerful and independent protective factor on its own in patients facing an AMI.*

We thank the reviewer for giving us the comments to further improve our results and discussion part.



Click here to access/download
Supplementary Material
Generalized Anxiety Disorder.pdf

