

Direct pericyte-to-neuron reprogramming via unfolding of a neural stem cell-like program

Marisa Karow^{1,2,13*}, J. Gray Camp^{3,13}, Sven Falk^{2,4}, Tobias Gerber³, Abhijeet Pataskar⁵, Malgorzata Gac-Santel³, Jorge Kageyama³, Agnieszka Brazovskaja³, Angela Garding⁵, Wenqiang Fan¹, Therese Riedemann², Antonella Casamassa^{1,12}, Andrej Smiyakin⁶, Christian Schichor⁷, Magdalena Götz^{2,4}, Vijay K. Tiwari⁵, Barbara Treutlein^{3,8,9,14*} and Benedikt Berninger^{1,10,11,14*}

Ectopic expression of defined transcription factors can force direct cell-fate conversion from one lineage to another in the absence of cell division. Several transcription factor cocktails have enabled successful reprogramming of various somatic cell types into induced neurons (iNs) of distinct neurotransmitter phenotype. However, the nature of the intermediate states that drive the reprogramming trajectory toward distinct iN types is largely unknown. Here we show that successful direct reprogramming of adult human brain pericytes into functional iNs by *Ascl1* and *Sox2* encompasses transient activation of a neural stem cell-like gene expression program that precedes bifurcation into distinct neuronal lineages. During this transient state, key signaling components relevant for neural induction and neural stem cell maintenance are regulated and functionally contribute to iN reprogramming and maturation. Thus, *Ascl1*- and *Sox2*-mediated reprogramming into a broad spectrum of iN types involves the unfolding of a developmental program via neural stem cell-like intermediates.

Direct lineage reprogramming is an emerging strategy for harnessing the cellular plasticity of differentiated cells for lineage conversion into desired target cell types for disease modeling and tissue repair^{1–4}. While direct lineage reprogramming from starting to target-cell type classically occurs without cell division, thereby sharply contrasting with reprogramming toward induced pluripotency⁵, little is known about the intermediate states that bridge the trajectory between start and end points. Two models have been proposed, according to which direct reprogramming is mediated either through direct conversion between fully differentiated states or through reversal to a developmentally immature state⁶. Furthermore, reprogramming efficiency and final differentiation outcomes are highly cellular-context-dependent, for which the underlying reasons are only incompletely understood^{7,8}. Analyses of the transcriptome alterations induced by the reprogramming factors have yielded fundamental insights into the molecular mechanisms of iN conversion^{9–12}. For instance, a single factor, *Ascl1*, can reprogram mouse astrocytes into iNs with high efficiency¹³, while the same factor induces a muscle cell-like fate in mouse embryonic fibroblasts (MEF) alongside neuronal fates^{11,14}. Efficient reprogramming of MEFs into iNs requires co-expression of additional factors (for example, *Brn2*, *Ascl1* and *Myl1l*, collectively referred to as BAM)^{9,11,12,15}. Moreover, *Ascl1* induces a GABAergic neuron identity in mouse astrocytes^{10,13}, while BAM-transduced fibroblasts predominantly adopt a glutamatergic phenotype¹⁵, raising questions of how the respective reprogramming

trajectories translate into distinct iN transmitter and subtype identities.

In the present study, by analyzing transcriptomes at population and single cell level, we aimed to reconstruct the trajectories underlying direct lineage conversion of adult human brain pericytes into iNs by forced expression of *Ascl1* and *Sox2* (AS)¹⁶. This allowed us to scrutinize the contribution of the starting cell population's heterogeneity to the variability in reprogramming success. By identifying cells of distinct reprogramming competence, we were able to reconstruct a trajectory of productive AS-mediated iN generation, allowing us to uncover intermediate states during successful conversion. Unexpectedly, we found that despite the absence of cell division, cells in the productive trajectory passed through a neural stem cell-like state. Transiently induced genes, many of which are core components of signaling pathways, typified this intermediate state, and interference with these signaling pathways demonstrated their functional importance for the reprogramming process. Finally, the productive reprogramming trajectory revealed an unexpected point of bifurcation into lineages whose transcriptomes were dominated by transcription factor families involved in the specification of GABAergic and glutamatergic subclasses of forebrain neurons.

Results

***Ascl1* and *Sox2* synergism in inducing neuronal gene expression in pericytes.** We have recently shown that adult human brain pericytes can be reprogrammed into iNs via forced expression of the

¹Institute of Physiological Chemistry, University Medical Center Johannes Gutenberg University Mainz, Mainz, Germany. ²Physiological Genomics, Biomedical Center, Ludwig Maximilians University Munich, Planegg/Martinsried, Germany. ³Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany. ⁴Institute for Stem Cell Research, Helmholtz Center Munich, German Research Center for Environmental Health, Neuherberg, Germany. ⁵Institute of Molecular Biology (IMB), Mainz, Germany. ⁶Miltenyi Biotec GmbH, Bergisch Gladbach, Germany. ⁷Department of Neurosurgery, Ludwig Maximilians University, Munich, Germany. ⁸Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, Germany. ⁹Technical University Munich, Munich, Germany. ¹⁰Focus Program Translational Neuroscience, Johannes Gutenberg University Mainz, Mainz, Germany. ¹¹Centre for Neurodevelopmental Biology, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK. ¹²Present address: Division of Pharmacology, School of Medicine, "Federico II" University of Naples, Naples, Italy. ¹³These authors contributed equally: Marisa Karow, J.Gray Camp. ¹⁴These authors jointly supervised this work: Barbara Treutlein, Benedikt Berninger. *e-mail: marisa.karow@med.uni-muenchen.de; barbara_treutlein@eva.mpg.de; berningb@uni-mainz.de




Fig. 1 | Ascl1-Sox2 synergism is required for pericyte-to-iN reprogramming. **a**, Schematic of experiments in this figure. Cells expressing Ascl1 and/or Sox2 are fluorescently labeled and isolated by fluorescence-activated cell-sorting (FACS) for bulk and scRNA-seq analyses at indicated timepoints following transduction. **b**, Representative micrographs of cultured human pericytes expressing pericyte markers PDGFRB and SMA, before (left) and after transformation (right) into TUBB3⁺ and GABA⁺ iNs by overexpressing AS at 46 dpi ($n > 30$). Nuclei are stained with DAPI. Scale bar, 50 μ m. **c**, Bulk RNA-seq with pericytes derived from three individual donors was performed at 2 and 7 dpi with Ascl1-only, Sox2-only, or AS. Bar graph shows the number of differentially expressed (DE) genes (adjusted P value (P_{adj}) < 0.01 ; calculated according to Benjamini-Hochberg) in each condition compared to pericytes transduced with a control vector. The Euler diagram shows the overlap of the DE genes at 7 dpi. Note that the majority of DE genes results from AS synergism. **d**, Heatmaps show normalized expression (z-score) of representative DE genes highlighting the induction of mesodermal, neurogenesis-related, and GABAergic signature genes at both 2 and 7 dpi. **e**, scRNA-seq was performed at 2 and 7 dpi on cells transduced with Ascl1-only ($n = 82$ cells at 2 dpi and 64 cells at 7 dpi) and AS ($n = 86$ cells at 2 dpi and 48 cells at 7 dpi). Principal component analysis (PCA; calculated on a total of 280 cells) followed by t-SNE shows that the pericyte signature is diminished in many 7-dpi Ascl1-only and strongly diminished in the majority of 7-dpi AS cells, concomitant with the acquisition of a mesoderm and GABAergic neuron (iGN) signature in Ascl1-only and AS-cells, respectively. Signatures were calculated by summing the expression of the fate-determinants highlighted in **d** (Supplementary Table 5). **f**, The iGN signature is plotted for all Ascl1-only and AS cells relative to the pericyte signature.

transcription factors Ascl1 and Sox2, and time-lapse imaging showed that this conversion occurs in the absence of cell division, qualifying it as direct lineage reprogramming¹⁶. Given that adult human brain pericyte reprogramming into functional iNs requires co-expression of Sox2 alongside Ascl1¹⁶, we first addressed the contribution of each factor individually or in combination with the gene expression programs underlying pericyte-to-neuron conversion (Fig. 1a,b). We performed RNA-seq of early-passage cultured human brain pericytes, obtained from three different adult donors and transduced with retroviruses encoding a reporter for control, Ascl1, Sox2, or AS at early stages (2 d postinfection (dpi) and 7 dpi) of reprogramming

(Fig. 1a). Unexpectedly, Sox2 only induced minor changes in gene expression, at both 2 and 7 dpi (Fig. 1c, Supplementary Fig. 1a,e, and Supplementary Table 1). In contrast, Ascl1 and AS substantially altered gene expression at both stages (Fig. 1c and Supplementary Fig. 1a,e,f). Notably, Ascl1 and AS changed the expression of distinct sets of genes. We noticed that several of the Ascl1-only altered genes are expressed in cells of the mesodermal lineage, indicative of a failure to cross the lineage barrier toward neurogenesis. In sharp contrast, AS resulted in significant induction of genes related to neurogenesis (Fig. 1d, and Supplementary Fig. 1e,f, and Supplementary Table 1). Moreover, we detected upregulation of

several transcription factors and noncoding RNAs playing key roles in forebrain GABAergic neurogenesis^{17,18} (Fig. 1d, Supplementary Fig. 1e,f, and Supplementary Tables 1 and 2). Yet we also observed a significant increase in *NEUROG2* expression (Fig. 1d), which is associated with diverse excitatory neuron identities.

Comparison of the genes upregulated by *Ascl1*-only or by AS with those transactivated by *Ascl1* in mouse neural stem cells¹⁹ revealed a progressive induction of direct *Ascl1* neural stem cell target genes between 2 and 7 dpi (Supplementary Fig. 1b and Supplementary Table 3). However, many of the direct *Ascl1* neural stem cell target genes became induced only upon co-expression of *Sox2* (Supplementary Fig. 1c,d), indicating that the proposed on-target pioneer factor activity of *Ascl1*^{9,19} is highly context-dependent.

To further dissect the differences in the early gene expression programs induced by *Ascl1* or AS, we measured 280 single-cell transcriptomes of *Ascl1*- ($n = 146$) and AS-expressing ($n = 134$) cells by single-cell RNA-seq at 2 and 7 dpi. Principal component analysis followed by *t*-stochastic neighbor embedding (t-SNE) of single-cell transcriptomes revealed an early and progressive separation of *Ascl1*-only and AS-expressing cells (Fig. 1e). Pericyte identity genes (for example, *PDGFRB*, *COL1A1*, and *CAV1*) became downregulated in *Ascl1*- and AS-expressing cells (Supplementary Fig. 1g), but only the latter acquired a GABAergic neuron fate signature (*DLX1/2*, *DLX5/6*, *SATB1*, etc.; Fig. 1e and Supplementary Fig. 1h). In agreement with our bulk RNA-seq data and published data from MEF-to-neuron reprogramming¹¹, *Ascl1*-expressing cells induced myocyte differentiation genes (for example, *MUSTN1*; Supplementary Fig. 1h). Occasionally, individual AS cell transcriptomes clustered with those of *Ascl1*-only cells, suggesting failed AS synergism as a potential mechanism underlying reprogramming failure (Fig. 1f). Overall, these data demonstrate that *Ascl1* alone is unable to induce a neuronal program in adult human brain pericytes but requires synergism with *Sox2*.

Pericyte heterogeneity and reprogramming competence.

To define the competence of adult human brain pericytes for AS-induced reprogramming, we next compared the transcriptomes of control pericytes with those of AS-transduced cells at early and later stages of reprogramming. To our surprise, t-SNE analysis revealed that control cells fell into two discernible clusters (Fig. 2a,b and Supplementary Fig. 2a,b), here referred to as group 1 and group 2 pericytes, with differentially enriched gene ontology terms (Supplementary Fig. 2c and Supplementary Table 4). While both groups highly expressed several classical pericyte genes (i.e., *PDGFRB*, *CAV1*, *DCN*, etc.; Fig. 2a and Supplementary Fig. 2b), other pericyte-associated genes such as *ANGPT1*, *APOE*, and *LEPR* were differentially expressed (Supplementary Fig. 2a,b), and such differential expression could be confirmed on the protein level (Fig. 2c,f and Supplementary Fig. 2d). Notably, transcriptomes of AS-transduced cells exhibited distinct degrees of relatedness to the two pericyte starting populations, with cells undergoing successful reprogramming being more similar to group 2 pericytes (Fig. 2d,e). These data strongly suggest that the two pericyte groups differed markedly in their response to AS. In fact, t-SNE analysis indicated that productive reprogramming toward neurogenesis originated specifically from group 2 pericytes (Fig. 2a,e). In contrast, group 1 pericytes appeared to give rise to a distinct population positive for the hypothalamic neuronal marker *PMCH* but lacking expression of other neuronal genes, which thus precluded identifying these cells as hypothalamic neurons (Fig. 2a). Besides AS-transduced cells clustering differentially with group 1 and group 2 pericytes, we observed two smaller clusters of AS-transduced cells enriched in genes involved in cell-cycle progression (for example, *MKI67*) and potentially an alternative fate marked by the expression of *POU2F3*. To independently corroborate differential neurogenic competence of group 1 and group 2 pericytes, we used fluorescence-activated

cell sorting to purify these populations via antibodies specific to the leptin receptor, encoded by the *LEPR* gene (Fig. 2f,g and Supplementary Fig. 2a,b). Consistent with the observation that iNs may originate from group 2 pericytes, we found that leptin-receptor-negative cells were more prone to undergo AS-induced neurogenesis than leptin-receptor-positive cells (Fig. 2h). These data provide experimental evidence that the two pericyte starter populations display distinct degrees of reprogramming competence.

Transient activation of a neural stem cell-like program. We next reconstructed the transcriptome trajectory of reprogramming-competent pericytes into iNs by pseudotemporal ordering²⁰ (Fig. 3a). Genes that mark pericyte identity, such as *PDGFRB*, *CAV1*, and *CFH*, became gradually downregulated. Conversely, genes associated with the acquisition of a neuronal fate were progressively upregulated with more linear (for example, *CHD7* and *DLX5*) or nonlinear dynamics (for example, *SNAP25*; Fig. 3b,c), possibly reflecting distinct gene expression waves during early and later phases of neuronal differentiation. Notably, we identified a set of genes that became upregulated early during the reprogramming process, but then declined again as neuronal differentiation progressed (Fig. 3b,c). We refer to these genes as 'switch genes'. These include genes involved in the regulation of cell signaling such as *NOG*, *LEFTY2*, *DKK1*, and *NOTCH2*, suggesting that modulation of signaling pathways is important during early phases of productive reprogramming (see below). The conspicuous dynamics of the regulation of these genes urged us to interrogate their expression during mouse embryonic development. Notably, the switch genes were markedly enriched in the germinal zones of the developing CNS containing the neural stem cells (Fig. 3d and Supplementary Fig. 3a). This strongly suggests that cells undergoing productive reprogramming by AS transiently acquire a neural stem cell-like state. This was further corroborated when analyzing the expression levels of these genes in human fetal brain tissue²¹, where higher levels of expression were found in distinct human neural stem cells (i.e., apical and basal radial glia) as compared to neurons (Fig. 3e). Consistent with the upregulation of the switch genes during successful reprogramming, mapping the switch gene signature onto the t-SNE plot shown in Fig. 2a revealed its specific occurrence in the cell population that connects productive group 2 pericytes with iNs (Fig. 3f). In contrast, the switch gene signature was absent from transcriptomes of AS-transduced cells in the immediate neighborhood of group 1 pericytes (Fig. 3f), indicating that acquisition of a neural stem cell like-state is critical for AS-mediated pericyte-to-iN reprogramming. Of note, mapping the same switch gene signature onto previously published single-cell transcriptomes undergoing MEF-to-iN reprogramming¹¹ revealed an unexpectedly high base level of switch gene expression in the MEF starting population, and its expression did not increase at any stage along the MEF-to-neuron axis but was found to be strongly decreased in neurons (Supplementary Fig. 3b). These data are indicative of fundamental differences in the reprogramming trajectories of these two distinct reprogramming pathways. Notably, time-lapse imaging of pericytes during AS reprogramming revealed the occurrence of different cellular morphologies: while at early phases of reprogramming, cells displayed a flat, fibroblast-like morphology, at subsequent phases, processes undergoing dynamic turnover akin to multipolar progenitors appeared (Fig. 3g and Supplementary Video 1). Finally, at the end of the reprogramming process, neuron-like cells dramatically decreased their motility and protruded processes of increased stability. Thus, the cellular behavior and morphology are consistent with the notion of distinct cellular states underlying the reprogramming of pericytes into iNs.

Modulation of signaling pathways. The conspicuous regulation of several components of signaling pathways known to play key




Fig. 2 | Pericyte heterogeneity correlates with distinct reprogramming competence. **a**, PCA (total of 419 cells) followed by t-SNE reveals heterogeneity during iN reprogramming, with genes marking distinct clusters colored on the adjacent t-SNE plots. Control-transduced pericytes (green) segregate into two distinct clusters. One cluster links to cells that express iN fate determinants (productive), whereas the other cluster is unlinked to iN reprogramming (unproductive). Cells expressing *MKI67* and other hallmarks of cycling cells are observed, as well as a group of cells expressing *POU2F3*. **b**, Heatmap shows the expression of genes that correlate with PC1 from PCA on control pericytes only. Hierarchical clustering reveals two distinct groups of pericytes with selected genes indicated below the heatmap. **c**, Left: micrographs showing cultured human brain pericytes stained against the pan-pericyte marker PDGFRB. Right: the same field of view of cultured human brain pericytes stained against pericyte group 1 marker ITGA6 and pericyte group 2 marker CD4. Nuclei are stained with DAPI ($n = 3$ individual pericyte donors; two independent experiments). Scale bar, 50 μm . **d**, Cells from the productive reprogramming cluster in **a** have a higher similarity to group 2 pericytes, while cells from the unproductive reprogramming clusters have a higher similarity to group 1 pericytes. **e**, Lineage network based on pairwise correlations between cells suggests that group 2 pericytes (lighter green population) are more competent to contribute to productive iN reprogramming. **f**, Top: violin blots show the density distribution of RNA expression of *LEPR* in pericyte groups 1 (31 cells) and 2 (44 cells). Bottom: representative flow cytometry plots show *LEPR* expression in cultured human brain pericytes (four independent experiments). **g**, Human brain pericytes were sorted based on *LEPR* expression (group 1 marker), plated, and transduced with AS to induce lineage conversion. Micrographs show AS-transduced pericytes at 35 dpi, with inset showing higher magnification of reprogrammed pericytes that acquired neuronal morphology and GABA immunoreactivity ($n = 4$). Nuclei are stained with DAPI. Scale bars, 50 μm . **h**, Quantification of reprogramming efficiency (dots represent independent individual experiments; $n = 4$; data are represented as boxplots with whiskers; two-tailed unpaired Student's t test; $***P = 0.000593$) reveals that the *LEPR*⁻ pericyte subpopulation is more competent to iN reprogramming using AS, confirming predictions from scRNA-seq. Boxplots show median, quartiles (box), and range (whiskers).




Fig. 3 | A transient neural precursor-like state emerges on the reprogramming path to iNs. **a**, PCA was performed on 211 single-cell transcriptomes from the productive reprogramming cluster shown in Fig. 2a. Monocle2 was used to infer a pseudotemporal ordering of cells. **b**, Scaled expression of representative marker genes in all 211 cells of the productive reprogramming path shown as a function of pseudotime. Shaded gray represents 95% confidence interval. **c**, Heatmap shows the expression of genes identified by PCA and ordered by hierarchical clustering. Cells are ordered according to pseudotime. The reprogramming path is characterized by three phases of gene expression changes. In the first phase, genes associated with pericyte identity are downregulated, concomitant with a transient upregulation of switch genes (for example, *NOG* and *DKK1*). In the second phase, genes associated with iN fate determination are upregulated (for example, *DLX5*). In the third phase, neuron maturation factors (for example, *DLX1* and *SNAP25*) are upregulated alongside phase 2 genes. **d**, Chart shows fraction of pericyte, switch, and neuronal genes (from **c**) expressed in blood vessels and meninges, ventricular zones, or neuron compartments determined from in situ patterns in the developing mouse brain. **e**, Violin plots show the density distribution of the switch gene signature from scRNA-seq of fetal human cortex (dots within the violins represent medians). Cortex cells are grouped based on cell type (41 apical progenitor cells (AP); 19 basal progenitor cells (BP); 42 early-born neurons (Neuron)). **f**, Switch gene signatures projected onto the t-SNE plot from Fig. 2a shows the induction of switch genes at an intermediate stage exclusively within the productive reprogramming path. **g**, Time-lapse imaging of AS-transduced cells over time (three independent experiments) shows the morphological changes during successful pericyte-to-neuron reprogramming (timepoints specified within the images). Arrows indicate a successfully reprogrammed pericyte throughout the different morphological changes mentioned below the images. Scale bar, 50 μm .

roles during neural induction and neural stem cell maintenance²², such as the BMP inhibitor *NOG*, the *ACTIVIN/NODAL* inhibitor *LEFTY2*, and *NOTCH2* and its downstream targets *HEY1* and *ID1*, prompted us to investigate whether these pathways are of functional importance for successful reprogramming. To test the importance of modulation of *NODAL* and *BMP* signaling, we treated pericytes during early phases of reprogramming with recombinant *NODAL* (1 $\mu\text{g}/\text{mL}$) and *BMP4* (30 ng/mL ; Fig. 4a,b). These treatments resulted in a significant reduction of reprogramming as determined by the number of *TUBB3*⁺ cells amongst AS-transduced cells (Fig. 4c). Conversely, inhibition of *BMP*, *ACTIVIN/NODAL*, and *TGF- β* signaling via the small molecules dorsomorphin (1 μM) and SB431542 (10 μM) caused a threefold increase in the number of reprogrammed iNs (Fig. 4d,e). To address the relevance of *NOTCH* signaling in the reprogramming process, we treated AS-transduced pericytes with the γ -secretase inhibitor *N*-[*N*-(3,5-difluorophenacetyl)-*L*-alanyl]-*S*-phenylglycine *t*-butyl ester (*DAPT*). *DAPT* treatment (10 μM) resulted in a marked increase in the number of iNs (Fig. 4d,e). This finding is consistent with the role of *NOTCH* signaling in neurogenesis inhibition and neural stem cell maintenance²³.

Though AS induction leads to productive iN reprogramming, we observed that maturation seemed to stall at 14 dpi, either because cells at later timepoints failed to mature further or because of a technical bias against harvesting healthy iNs at later stages (Fig. 3a). Our data showed that productive reprogramming involved inhibition of *BMP* signaling, and blocking *BMP* signaling appeared to promote maturation, as suggested by increased morphological iN complexity (Fig. 4e). We therefore analyzed the effect of the *BMP* inhibitor dorsomorphin during early phases of the reprogramming process on subsequent neuronal maturation (Fig. 5a). Dorsomorphin-treated AS-induced neurons exhibited markedly increased morphological complexity and soma size (Fig. 5b,c and Supplementary Fig. 4a), as well as increased membrane capacitance and decreased membrane resistance (Fig. 5d and Supplementary Fig. 4b). Single-cell RNA (scRNA)-seq on dorsomorphin-treated AS-induced neurons revealed that genes associated with synapse formation and synaptic function showed increased expression relative to untreated AS-transduced cells (Fig. 5e). In line with enhanced iN maturation, we also noted enhanced *GABA* and *PVALB* immunoreactivity (Supplementary Fig. 4c).




Fig. 4 | Modulation of signaling pathways identified during neural stem cell-like state. **a**, Schematic overview of the experimental timeline for the experiments contained in this figure. Cells were treated with either small molecules (SM) or growth factors (GF). **b**, Schematic overview of switch gene expression associated with signaling pathways and summary of treatments. **c**, Treatment with NODAL or BMP4 decreases the reprogramming efficiency. Reprogramming efficiency was calculated by quantifying TUBB3-immunoreactive cells among reporter-positive transduced cells 3–5 weeks after transduction with AS. Fold-changes were calculated by determining the ratio of growth factor in treated vs. nontreated TUBB3⁺Reporter⁺ cells per total Reporter⁺ cells of the transduced pericytes (dots represent independent individual experiments; untreated, $n=9$; NODAL, $n=4$; BMP, $n=3$; data are represented as boxplots with whiskers; two-tailed unpaired Student's t test; $P=0.01202$ for NODAL; $P=0.01821$ for BMP4; $*P<0.05$). Boxplots show median, quartiles (box), and range (whiskers). **d**, Reprogramming efficiencies (TUBB3⁺Reporter⁺ cells per total Reporter⁺ cells) increase following enhancing treatments with dorsomorphin (dorso) + SB431542 (SB) or DAPT. The fold change was calculated by determining the ratio of small-molecule-treated vs. nontreated TUBB3⁺Reporter⁺ cells per total Reporter⁺ cells of the transduced pericytes (dots represent independent individual experiments; untreated, $n=9$; dorso + SB, $n=4$; DAPT, $n=7$; data are represented as boxplots with whiskers; two-tailed unpaired Student's t test; $*P=0.02133$ for dorso + SB; $**P=0.0050$ for DAPT). Boxplots show median, quartiles (box), and range (whiskers). **e**, Representative micrographs of AS-transduced pericyte cultures (green) and stained with anti-TUBB3 (magenta). Note the increased appearance of reprogrammed pericytes that acquired neuronal morphology following dorso + SB ($n=4$) and DAPT ($n=7$) treatment compared to the untreated cells. Nuclei are stained with DAPI. Scale bars, 50 μ m.

Bifurcation into distinct iN lineages. These data prompted us to have a closer look at the neuronal subtype specification induced by AS with or without dorsomorphin (ASD). We focused on iNs that expressed both *SNAP25* and *MAP2* and analyzed 20 AS and 72 ASD cells. The majority of AS and ASD iNs exhibited a fore-brain GABAergic interneuron program characterized by coordinated expression of multiple members of the *DLX* gene family (Fig. 5f). Notably, we found evidence for further subspecification among the *DLX*-expressing iNs. We observed distinct clusters of *VIP*-expressing neurons, some of which also co-expressed *CCK*. Likewise, we noted two clusters specifically expressing *SST* (Fig. 5f).

However, we also noted a subset of iNs that expressed a transcription factor of the glutamatergic lineage, *NEUROG2*. Notably, these iNs also expressed downstream targets *NEUROD1* and *NEUROD4*, consistent with the expression of a telencephalic glutamatergic neuron program^{10,24}. The fact that this subset also expressed *RELN* (Fig. 5f) may indicate that these iNs acquire a Cajal–Retzius neuron-like program. It is noteworthy that, while the majority of *DLX1*-expressing cells were *NEUROG2*⁻ and many high-*NEUROG2*⁺ cells

were *DLX1*⁻, we observed some outliers expressing both genes (Supplementary Fig. 4d). This may reflect the possibility that the definitive decision between the two major neuron lineages (*GABA* versus *glutamate*) had not yet taken place in these cells. To reveal the developmental trajectory toward a *DLX*- or *NEUROG2*-dominated fate, we employed pseudotemporal ordering of the transcriptomes of AS cells of the productive path and 14-dpi ASD cells. Notably, we observed a bifurcation of the trajectory into *DLX*- or *NEUROG2*-dominated pathways, which preceded neuronal differentiation marked by *SNAP25* expression (Fig. 5g and Supplementary Fig. 4e). Projecting the switch gene signature identified in Fig. 3c onto the reprogramming path revealed that the transient expression of neural stem cell-like genes occurred and ceased before lineage bifurcation (Fig. 5h). Consistent with the increased maturation of ASD cells, the distribution of ASD transcriptomes was shifted farther along the trajectory of pericyte-to-iN reprogramming (Fig. 5i). We corroborated the emergence of *DLX*- or *NEUROG2*-expressing inhibitory and excitatory neuronal lineages using an alternative scRNA-seq method and a second pericyte donor (Supplementary Fig. 5).




Fig. 5 | Pericytes give rise to distinct neuronal subtypes, and targeting BMP signaling promotes maturation. **a**, Schematic overview of experiments in this figure. **b**, Representative examples of untreated and dorsomorphin-treated AS-transduced pericytes showing complex branching of dendritic arbors in dorsomorphin-treated cells. Neuronal morphology was reconstructed using TUBB3 immunoreactivity and used for Sholl analyses. In the Sholl masks (right), warmer hues indicate higher number of intersections (untreated, $n=14$ cells from 3 independent experiments; dorsomorphin-treated, $n=14$ cells from 3 independent experiments). **c**, Single-cell neuromorphology reconstruction by Sholl analysis on dorsomorphin treated AS-transduced cells was compared to untreated cells. Note the increase in complexity of dorsomorphin-treated cells, which display more intersections with concentric shells and an overall increase in the length of the dendrites (see also Supplementary Fig. 4a). Data are represented in a line graph as mean \pm s.e.m. (shaded region) (untreated, $n=14$ cells from 3 independent experiments; dorsomorphin-treated, $n=14$ cells from 3 independent experiments). **d**, Electrophysiological assessment of AS and ASD cells. Membrane capacitance (C_N) is plotted as a function of the membrane resistance (R_N). AS cells have greater R_N but smaller C_N compared to ASD cells. **e**, Violin blots show the density distribution of expression of selected neuronal maturation genes in neuronal (SNAP25⁺, MAP2⁺, PDGFRB⁻) AS cells ($n=20$) and ASD cells ($n=75$). **f**, PCA followed by hierarchical clustering was used to characterize neuronal cells (SNAP25⁺, MAP2⁺, PDGFRB⁻) AS cells ($n=20$) and ASD cells ($n=75$) at 14 dpi according to their expression of neuronal genes. Note the different interneuron clusters marked by the expression of interneuron subtype-specific genes *VIP*, *SST*, and *CCK*. **g**, Monocle2 was used to compare maturation of AS and ASD cells. **h**, Switch gene signature from Fig. 3c projected onto pseudotemporal ordered transcriptomes from **g**. **i**, The density distributions (i.e., normalized cell numbers per timepoint of pseudotime) along the pseudotime of AS transcriptomes (14 dpi) from the productive path and ASD transcriptomes (from **g**) are plotted as a function of the distance from the start point. Note the shift of ASD cell transcriptomes compared to AS transcriptomes. The dotted line indicates the bifurcation point in **g**. **j**, Jitter-boxplot (boxplots show median, quartiles (box), and range (whiskers)) showing the expression of *TUBB3* in all SNAP25⁺, MAP2⁺, PDGFRB⁻ AS and ASD ($n=71$), and AS + overexpressed *NEUROG2* (ASN; $n=21$) cells used for this analysis. Boxplots show the strong increase of *NEUROG2* and decrease in *DLX2* gene expression in ASN cells as compared to AS and ASD cells.

460 To test whether *NEUROG2* indeed suffices to induce a glutamatergic phenotype, we overexpressed *NEUROG2* alongside AS (Supplementary Fig. 4f). *NEUROG2* overexpression resulted in the almost complete suppression of *DLX2* (Fig. 5j), suggesting that *NEUROG2* can divert iNs from adopting predominantly a GABAergic phenotype toward generating mostly glutamatergic neurons. Accordingly, we found that *NEUROG2*-overexpressing AS neurons exhibited vesicular glutamate transporter immunoreactivity (Supplementary Fig. 4g).

470 Discussion

471 Here we have shown that reprogramming success of adult human brain pericytes into iNs by the transcription factors *Ascl1* and *Sox2* (AS) critically depends on cellular context, as revealed by the observation that pericyte heterogeneity is a key determinant for reprogramming competence. Successful reprogramming by AS encompasses the passage through a neural stem cell-like intermediate state, yet it occurs in the absence of cell division. Moreover, regulation of signaling pathways during the neural stem cell-like state was of functional importance for the reprogramming outcome. This data indicates that AS-mediated reprogramming involves the unfolding of developmental programs and argues for the engagement of hierarchical developmental gene-regulatory networks⁶ rather than direct interconversion between two states of terminal differentiation. Finally, we found that, following the transition through a neural stem cell-like state, the reprogramming trajectory eventually bifurcates to give rise to two distinct branches characterized by *DLX*- or *NEUROG*-dominated gene expression and indicative of bifurcation into GABAergic or glutamatergic lineages, respectively. This provides a mechanistic explanation for the common observation that a single reprogramming cocktail can yield neurons of distinct neurotransmitter phenotypes^{15,25}.

492 We observed that the reprogramming competence of adult human brain pericytes is highly variable and a main source for this variability is pericyte heterogeneity. Heterogeneity of pericytes has been described in many tissues and may reflect distinct embryonic origins²⁶. Our scRNA-seq experiments revealed two distinct populations, one of which, characterized by high *LEPR* expression, displayed markedly reduced reprogramming propensity. Notably, a recent study using scRNA-seq showed that several of the heterogeneously expressed genes are also expressed at highly variable levels in acutely isolated human midbrain pericytes²⁷, which might indicate that similar heterogeneity occurs in vivo. However, our study may actually underestimate overall pericyte heterogeneity, as we included in our scRNA-seq analysis only retrovirus-transduced cells, for which ongoing cell division at the time of transduction is required. We would also expect that proliferative pericytes do not perfectly match pericytes under resting conditions, but may be more akin to those found to undergo cell division in response to severe CNS injury²⁸.

510 Revealing the cell-context requirements for reprogramming is of greatest importance if direct lineage reprogramming is to be of therapeutic value. Many reprogramming-factor cocktails that work well with mouse cells (for example, MEFs) are rather inefficient with human cells, and in particular, when the cells are of adult-tissue origin. It will be therefore a fruitful field of investigation to identify, in addition to transcriptome differences, epigenetic disparities between the two subpopulations of pericytes identified here. This may yield potential molecular targets for improved reprogramming strategies that may apply to other adult human somatic cell types.

520 A key finding of our study is the observation that AS-transduced cells pass through a neural stem cell-like state before differentiating into iNs. This neural stem cell-like state is characterized by expression of a battery of genes that are normally expressed in neural stem cells or progenitor cells during forebrain embryonic development, referred to here as switch genes as they are dynamically regulated

during the reprogramming process. While referring to the state characterized by switch gene expression as a neural stem cell-like state, we do not equate it to a bona fide neural stem cell state. This distinction is warranted given the absence of classical markers of neural stem cells such as *MSH1* (Musashi) or *NES* (Nestin) during the switch state, some anomalies in gene regulation such as *DLX5* expression preceding *DLX1* expression in time, and above all the absence of cell division and of a transcriptomic signature of an active cell cycle. We hypothesize that genes induced during the switch state represent a neural stem cell gene-expression module specifically regulated by *Ascl1* and *Sox2* and that other transcription factors may be required to induce other neural stem cell markers. Notably, the AS-induced neural stem cell expression module appears to be sufficient to drive the trajectory toward neuronal differentiation.

Switch genes include components of several signaling pathways, such as the *ACTIVIN/NODAL* (*LEFTY2*), *BMP* (*NOG*), and *NOTCH* (*HES5*, *HEY1*, *ID1*, *NOTCH2*) signaling pathways. By activating or inhibiting the *ACTIVIN/NODAL* and *BMP* pathways during the early phase of reprogramming through recombinant ligands or pharmacologically, we showed that these pathways exert an important influence on reprogramming efficiency. The fact that inhibition of *ACTIVIN/NODAL* and *BMP* signaling is required for reprogramming is consistent with the fact that inhibition of these pathways is important for neural induction during embryonic development²², can be used for driving human pluripotent stem cells toward neural lineages²⁹, and enhances transcription factor-mediated reprogramming^{25,30}. Crucially, we found that inhibition of *NOTCH* signaling promoted reprogramming. This is consistent with the role of *NOTCH* signaling in preventing neuronal differentiation of neural stem cells³¹. The conspicuous induction of the *NOTCH* ligand *DLL1* during reprogramming suggests that *DLL1*⁺ cells exert a differentiation inhibitory effect on other AS-transduced cells, an inhibition that can be relieved pharmacologically. Notably, iNs appear to express *Myt1*, which has recently reported to be induced cell-autonomously by *Ascl1* and to repress Notch signaling³². Likewise, its close relative *Myt1l*, a widely used component of the *BAM* reprogramming cocktail, has been shown to repress Notch signaling¹². This suggests that the *BAM* cocktail exhibits similarities to AS' mechanism of reprogramming, but that the addition of the postmitotic repressor *Myt1l* serves to curtail molecular pathways of the switch state that keep neuronal differentiation in check. Another intriguing aspect of *NOTCH2* expression during reprogramming is the fact that *Notch2* has been recently found to repress cell cycle-related genes and drive neural stem cells to quiescence, which may account in part for the lack of cell division during the switch state³³.

Unexpectedly, we found that human brain pericytes reprogrammed by AS bifurcate into lineages dominated by transcription factors that specify inhibitory and excitatory neuron fates. This bifurcation was corroborated using two distinct scRNA-seq platforms (Fluidigm C1 and 10×Genomics). While the *DLX* gene family-dominated branch was enriched for genes characteristic of GABAergic neuron lineage (for example, *GAD1* and *GAD2*), the *NEUROG*-expressing branch expressed other transcription factors characteristic of the glutamatergic neuron lineage, such as *NEUROD1* and *NEUROD4*. Moreover, cells of the latter lineage also expressed *RELN*, suggesting similarities to the Cajal-Retzius subtype of glutamatergic neurons. The fact that forced expression of *Neurog2* in AS-transduced pericytes suppresses *DLX* gene expression may indicate that lineage bifurcation is driven by mutual cross-repression of *NEUROG* and *DLX* family genes. Our data raise the intriguing possibility that the bipotent neural stem cell-like state observed during AS reprogramming relates to the suggested common precursor generating both glutamatergic and GABAergic neurons in the cerebral cortexes of human and nonhuman primates³⁴.

Overall, our study not only provides new insights into the biology underlying iN reprogramming, but also sheds light on the capacity of two transcription factors, *Ascl1* and *Sox2*, to cooperate in the generation of diverse neuronal subtypes, a cooperation that may be relevant during human brain development. The identification of molecular programs that establish cellular intermediates and lineage bifurcations during iN reprogramming provides avenues for improving lineage conversion of human brain-resident cells toward therapeutically relevant cell types.

Methods

Methods, including statements of data availability and any associated accession codes and references, are available at <https://doi.org/10.1038/s41593-018-0168-3>.

Received: 12 February 2018; Accepted: 17 April 2018;

References

- Heinrich, C., Spagnoli, F. M. & Berninger, B. In vivo reprogramming for tissue repair. *Nat. Cell Biol.* **17**, 204–211 (2015).
- Amamoto, R. & Arlotta, P. Development-inspired reprogramming of the mammalian central nervous system. *Science* **343**, 1239882 (2014).
- Mertens, J., Marchetto, M. C., Bardy, C. & Gage, F. H. Evaluating cell reprogramming, differentiation and conversion technologies in neuroscience. *Nat. Rev. Neurosci.* **17**, 424–437 (2016).
- Srivastava, D. & DeWitt, N. In vivo cellular reprogramming: the next generation. *Cell* **166**, 1386–1396 (2016).
- Ruiz, S. et al. A high proliferation rate is required for cell reprogramming and maintenance of human embryonic stem cell identity. *Curr. Biol.* **21**, 45–52 (2011).
- Morris, S. A. Direct lineage reprogramming via pioneer factors; a detour through developmental gene regulatory networks. *Development* **143**, 2696–2705 (2016).
- Vierbuchen, T. & Wernig, M. Molecular roadblocks for cellular reprogramming. *Mol. Cell* **47**, 827–838 (2012).
- Gascón, S., Masserdotti, G., Russo, G. L. & Götz, M. Direct neuronal reprogramming: achievements, hurdles, and new roads to success. *Cell Stem Cell* **21**, 18–34 (2017).
- Wapinski, O. L. et al. Hierarchical mechanisms for direct reprogramming of fibroblasts to neurons. *Cell* **155**, 621–635 (2013).
- Masserdotti, G. et al. Transcriptional mechanisms of proneural factors and REST in regulating neuronal reprogramming of astrocytes. *Cell Stem Cell* **17**, 74–88 (2015).
- Treutlein, B. et al. Dissecting direct reprogramming from fibroblast to neuron using single-cell RNA-seq. *Nature* **534**, 391–395 (2016).
- Mall, M. et al. Myt1l safeguards neuronal identity by actively repressing many non-neuronal fates. *Nature* **544**, 245–249 (2017).
- Heinrich, C. et al. Directing astroglia from the cerebral cortex into subtype specific functional neurons. *PLoS Biol.* **8**, e1000373 (2010).
- Chanda, S., Marro, S., Wernig, M. & Südhof, T. C. Neurons generated by direct conversion of fibroblasts reproduce synaptic phenotype caused by autism-associated neuroligin-3 mutation. *Proc. Natl Acad. Sci. USA* **110**, 16622–16627 (2013).
- Vierbuchen, T. et al. Direct conversion of fibroblasts to functional neurons by defined factors. *Nature* **463**, 1035–1041 (2010).
- Karow, M. et al. Reprogramming of pericyte-derived cells of the adult human brain into induced neuronal cells. *Cell Stem Cell* **11**, 471–476 (2012).
- Long, J. E. et al. *Dlx1&2* and *Mash1* transcription factors control striatal patterning and differentiation through parallel and overlapping pathways. *J. Comp. Neurol.* **512**, 556–572 (2009).
- Liu, S. J. et al. Single-cell analysis of long non-coding RNAs in the developing human neocortex. *Genome Biol.* **17**, 67 (2016).
- Raposo, A.A. et al. *Ascl1* coordinately regulates gene expression and the chromatin landscape during neurogenesis. *Cell Rep.* <https://doi.org/10.1016/j.celrep.2015.02.025> (2015).
- Trapnell, C. et al. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, 381–386 (2014).
- Camp, J. G. et al. Human cerebral organoids recapitulate gene expression programs of fetal neocortex development. *Proc. Natl Acad. Sci. USA* **112**, 15672–15677 (2015).
- Suzuki, I. K. & Vanderhaeghen, P. Is this a brain which I see before me? Modeling human neural development with pluripotent stem cells. *Development* **142**, 3138–3150 (2015).
- Imayoshi, I., Sakamoto, M., Yamaguchi, M., Mori, K. & Kageyama, R. Essential roles of Notch signaling in maintenance of neural stem cells in developing and adult brains. *J. Neurosci.* **30**, 3489–3498 (2010).
- Guillemot, F. Spatial and temporal specification of neural fates by transcription factor codes. *Development* **134**, 3771–3780 (2007).
- Ladewig, J. et al. Small molecules enable highly efficient neuronal conversion of human fibroblasts. *Nat. Methods* **9**, 575–578 (2012).
- Dias Moura Prazeres, P. H. et al. Pericytes are heterogeneous in their origin within the same tissue. *Dev. Biol.* **427**, 6–11 (2017).
- La Manno, G. et al. Molecular diversity of midbrain development in mouse, human, and stem cells. *Cell* **167**, 566–580.e19 (2016).
- Göriz, C. et al. A pericyte origin of spinal cord scar tissue. *Science* **333**, 238–242 (2011).
- Chambers, S. M. et al. Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat. Biotechnol.* **27**, 275–280 (2009).
- Liu, M. L. et al. Small molecules enable neurogenin 2 to efficiently convert human fibroblasts into cholinergic neurons. *Nat. Commun.* **4**, 2183 (2013).
- Kageyama, R., Ohtsuka, T., Shimojo, H. & Imai, I. Dynamic regulation of Notch signaling in neural progenitor cells. *Curr. Opin. Cell Biol.* **21**, 733–740 (2009).
- Vasconcelos, F. F. et al. MyT1 counteracts the neural progenitor program to promote vertebrate neurogenesis. *Cell Rep.* **17**, 469–483 (2016).
- Engler, A. et al. Notch2 signaling maintains NSC quiescence in the murine ventricular-subventricular zone. *Cell Rep.* **22**, 992–1002 (2018).
- Radonjić, N. V. et al. Diversity of cortical interneurons in primates: the role of the dorsal proliferative niche. *Cell Rep.* **9**, 2139–2151 (2014).

Acknowledgements

We thank M. Wernig (Stanford University) for generously providing us with the *Sox2* coding sequence. We are also very grateful to B. Sutor (BMC, LMU Munich) for help with the electrophysiological experiments, R. Menon (UMC Mainz) for help with the cell culture experiments, A. Bosio (Miltenyi Biotec) for help with the multidimensional fluorescence stainings, and F. Calzolari (UMC Mainz) for comments on the manuscript. We thank B. Höber, A. Weihmann, and J. Kelso of MPI-EVA for sequencing and bioinformatics support with this project. Flow cytometric cell sorting was performed at the “Core Unit Durchflusszytometrie” (CUDZ) of the Center for Infectious Diseases at the College of Veterinary Medicine, University of Leipzig, Leipzig, Germany. S.F. was supported by a fellowship from the Swiss National Science Foundation (PA00P3_139709). W.F. was supported by a Fellowship from the China Research Council. This work was supported by the following grants: advanced ERC ChroNeuroRepair to M.G.; Bavarian State Ministry of Sciences, Research and the Arts to M.K. and B.B. (ForiPS D2-F2412.26); Schram foundation (T287/29577/2017) and Wings For Life (WFL-DE-012/14) to M.K.; Max Planck Society to B.T.; and DFG (INST 161/875-2; BE 4182/8-1), NEURON ERA-NET (01EW1604), and Wellcome Trust (206410/Z/17/Z) to B.B.

Author contributions

M.K., J.G.C., B.T., and B.B. conceived the study and designed experiments; M.K. performed direct reprogramming experiments; M.K., S.F., A.P., and V.K.T. analyzed bulk RNA-seq data; A.B. helped with processing of the 10xGenomics data; A.G. performed RNA isolation for bulk RNA-seq analysis; W.F. performed time-lapse imaging experiments; T.R. performed electrophysiological recordings; A.C. performed Sholl analyses; A.S. performed immunocytochemical analyses; C.S. provided human brain biopsies; M.G. provided material; J.G.C., M.G.-S., and T.G. performed single-cell RNA-seq experiments and sequenced libraries; J.G.C., J.K., and B.T. analyzed single-cell RNA-seq data; all authors discussed the data; and M.K., J.G.C., S.F., B.T., and B.B. wrote the paper.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41593-018-0168-3>.

Reprints and permissions information is available at www.nature.com/reprints.

Correspondence and requests for materials should be addressed to M.K. or B.T. or B.B.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

592 Methods

593 **Culture of primary human pericytes.** Primary pericytes were derived as
594 described previously from adult human brain tissue^{16,35}. Briefly, specimens of
595 cerebral cortex were obtained from standard surgical interventions of patients
596 aged 19–70 years old and of both sexes. The study was approved by the ethical
597 committee of the Medical Faculty of the LMU Munich, and written informed
598 consent was obtained from all patients. Human tissue was enzymatically (TrypLE,
599 Life technologies) and mechanically dissociated, and, following centrifugation at
600 1,000 rpm for 5 min and resuspension in pericyte medium, cells were plated in
601 T75 cell-culture flasks. Pericyte growth medium consisted of DMEM high glucose
602 with Glutamax, 20% FBS, and penicillin/streptomycin. Medium was changed
603 twice per week and subcultivation at a ratio of 1:3 was performed every 10–14 d.
604 Cells were grown under low-oxygen conditions (5% O₂, 5% CO₂; Galaxy 170R,
605 New Brunswick).

606 **Retroviral transduction and treatments of human pericytes.** The retroviral
607 backbone used for lineage conversion of pericytes into iNs allowed for the
608 polycistronic expression of *Ascl1* and *Sox2* (connected via p2A) under the control
609 of an internal chicken β -actin promoter with cytomegalovirus enhancer (CAG)
610 together with either *DsRed* or *GFP* downstream of an internal ribosomal entry site
611 (IRES). For control, cultures were transduced with a virus encoding only *DsRed* or
612 *GFP* behind an IRES site as described previously^{13,16}.

613 Retroviral transduction of primary pericyte cultures was performed 24 h after
614 plating on either poly-D-lysine-coated glass coverslips or in T25 or T75 cell-culture
615 flasks without coating, using VSV-G (vesicular stomatitis virus glycoprotein)-
616 pseudotyped retroviruses encoding neurogenic fate determinants as described
617 previously^{16,35}. Samples (pericyte donors, coverslips in 24-well plates or in T25 or
618 T75 cell-culture flasks) were randomly assigned for transduction with different
619 viruses. Twenty-four hours after transduction, the medium was replaced by a
620 differentiation medium consisting of DMEM high glucose with Glutamax and B27
621 supplement (Gibco). For growth factor or small-molecule treatments, addition
622 was performed 1, 3, and 5 d following transduction. Factors were added to a final
623 concentration of 1 μ M^{30,36} for dorsomorphin (Sigma-Aldrich), 10 μ M for DAPT
624 [*N*-[*N*-(3,5-difluorophenacetyl)-L-alanyl]-*S*-phenylglycine *t*-butyl ester; Stem Cell
625 Technologies], 10 μ M^{37,38} for SB431542 (Stem Cell Technologies), 30 ng/mL^{39,40}
626 for recombinant human BMP4 (Preprotech), and 1 μ g/mL^{41,42} for recombinant
627 human NODAL (RnD Systems). Cells were allowed to differentiate under low-
628 oxygen conditions (5% O₂, 5% CO₂). Reprogramming efficiency was calculated
629 by quantifying TUBB3-immunoreactive cells among reporter-positive transduced
630 cells 3–5 weeks following transduction with retroviruses.

631 **Fluorescence-activated cell sorting (FACS).** For sorting of transduced cells
632 for further culturing, bulk RNA-sequencing, or scRNA-sequencing, primary
633 pericytes were detached from the culture dish using TrypLE for 4–6 min and
634 subsequently resuspended in 500–1,000 μ L pericyte growth medium. Cell sorting
635 was performed by taking advantage of the combined expression of *Ascl1* and *Sox2*
636 with a fluorescent reporter protein (either *DsRed* or *GFP*). Gating was achieved
637 via subtracting the autofluorescence of nontransduced cells; control (*DsRed* or
638 *GFP* only)-transduced cells were used as respective controls. Following sorting,
639 cells were (i) collected in pericyte growth medium and plated on PDL-coated glass
640 cover slips on 24-well plates for further culturing, (ii) directly collected into RLT
641 buffer (Qiagen) and stored at –80 °C until RNA isolation for bulk RNA-seq, or
642 (iii) prepared for single-cell loading onto a C1 Fluidigm chip for scRNA-seq. To
643 separate LEPR⁺ and LEPR⁻ pericyte populations, pericyte cultures were detached
644 from the culture dish using TrypLE for 4–6 min and subsequently 1 \times 10⁵–5 \times 10⁵
645 cells were resuspended in 100 μ L staining solution (PBS plus 0.5% BSA). Primary
646 antibody (Alexa Fluor 647-conjugated CD295 (anti-LEPR; 1:20, BD Pharmingen,
647 cat.no. 564376) was added and cells were incubated for 30 min on ice in the dark.
648 After washing three times in staining solution, cells were resuspended in 500 μ L
649 pericyte growth medium and subjected to cell sorting using a FACS Aria (BD). An
650 Alexa Fluor 647-conjugated isotype control antibody (1:100, BD Pharmingen) was
651 used to gate the proper populations.

652 **Immunohistochemical staining.** Cell cultures were fixed in 4% paraformaldehyde
653 (PFA) in phosphate-buffered saline (PBS) for 15 min at room temperature. Cells
654 were first pretreated in blocking solution consisting of 0.2–0.5% Triton X-100 and
655 10% donkey serum in PBS for 60 min, followed by incubation with the primary
656 antibodies in 100 μ L in the same solution for 1 h at room temperature or overnight
657 at 4 °C. After extensive washing in PBS, cells were incubated in the same solution
658 with appropriate species- or subclass-specific secondary antibodies conjugated
659 to fluorophores. Coverslips were finally mounted onto a glass slide with an anti-
660 fade mounting medium (Aqua Poly/Mount; Polysciences, Warrington, PA). For
661 multidimensional immunofluorescence staining, fixed cell cultures were subjected
662 to sequential immunofluorescence staining/destaining cycles adapted from a
663 technique published by Schubert et al.⁴³.

664 **Microscopy and time-lapse imaging.** Immunocytochemical stainings were
665 first examined with an epifluorescence microscope (BX61, Olympus) equipped
666 with the appropriate filter sets. Stainings were further analyzed with a LSM710

667 laser-scanning confocal microscope (Carl Zeiss.). Digital images were captured
668 using the ZEN software (Carl Zeiss).

669 We performed time-lapse microscopy to follow the reprogramming process
670 of pericytes into iNs. Pericytes were transduced with *Ascl1*-*Sox2*-CAG-GFP
671 retrovirus. Twenty-four hours after transduction, medium was replaced by a
672 differentiation medium consisting of DMEM high glucose with Glutamax and
673 B27 supplement (Gibco). The microwell plate containing these cells was 48 h
674 later placed on a heated microscopic stage with 5% CO₂ and 37 °C and imaged
675 continuously for up to 14 dys. Fluorescent images were taken subsequently once
676 every 4 h and brightfield images once every 5 min. After completion of time-lapse
677 imaging, cells were fixed with 4% PFA, and after imaging ICC was performed
678 to corroborate the results from the imaging. Data analysis was performed using
679 Timm's Tracking Tool (TTT) software.

680 **Sholl analysis.** Sholl analysis was performed by using the ImageJ plugin Sholl
681 Analysis⁴⁴. Confocal images of iNs with immunocytochemical stainings against
682 TUBB3 were used for tracing individual neuronal processes of selected cells in
683 ImageJ (Fiji) software⁴⁵. After assigning the center of each cell soma, a grid with
684 concentric circles with increasing diameter (5 μ m) was superimposed. The data
685 are expressed as the mean \pm s.e.m. of the values obtained in four independent
686 experiments; untreated *n* = 14, dorsomorphin-treated *n* = 14. The investigators
687 carried out blinded analyses.

688 **Neuromorphometry.** Several parameters of cell morphology were examined.
689 Neuronal complexity quantification was conducted with the following
690 measurements: (i) primary branches, i.e., processes emerging directly from the
691 soma per neuron; (ii) dendritic segment, i.e., part of the dendrite between two
692 branching points; (iii) branching point, i.e., the point at the dendrite where a
693 dendrite ramifies into two or more; (iv) maximum dendritic length or ending
694 radius, i.e., the radius of the largest circle of the superimposed Sholl mask; (v)
695 soma size (in μ m²), i.e., cross sectional surface area of the cell body; and (vi) sum
696 of intersections, i.e., the sum of all intersections between the dendritic arbors
697 and the concentric circles radiating from the cell body. The numbers of primary
698 branches, as well as the numbers of dendritic segments and branching points, were
699 counted manually. ImageJ Fiji software was used to measure soma size. The sum
700 of intersections and the ending radius were measured using the Sholl method (see
701 “Sholl analysis” section, above).

702 **Statistics.** To test for statistical significance, two-tailed unpaired Student's *t* tests
703 were used. Asterisks indicate statistically significant differences across the two
704 groups: **P* < 0.05, ***P* < 0.01, ****P* < 0.001. The analyses were done using Prism
705 (GraphPad) or R. Data distribution was assumed to be normal, but this was not
706 formally tested. Throughout the study, boxplots show medians, quartiles (box), and
707 ranges (whiskers). No statistical methods were used to predetermine sample sizes,
708 but our sample sizes are similar to those reported in previous publications^{11,16}. If
709 not indicated otherwise, data collection and analysis were not performed blind to
710 the conditions of the experiments. No data points were excluded from the analysis,
711 except for cells in the scRNA-seq analyses that did not fulfill the required criteria
712 (see below sections on scRNA-seq analyses).

713 **Electrophysiology.** For electrophysiological recordings, coverslips with
714 reprogrammed cells were transferred to a recording chamber mounted on the stage
715 of an upright microscope (Axioscope FS, Zeiss, Germany). Cells were perfused
716 with a bathing solution consisting of (in mM): NaCl 150, KCl 3, CaCl₂ 3, MgCl₂ 2,
717 HEPES 10, and D-glucose 10. The pH of the solution was adjusted to 7.4 (NaOH);
718 the osmolarity ranged from 309 to 313 mOsmol. All recordings were performed at
719 room temperature (23–24 °C). Electrodes for whole-cell patch-clamp recordings
720 were fabricated from borosilicate glass capillaries (OD: 1.5 mm, ID: 0.86 mm; Hugo
721 Sachs Elektronik-Harvard Apparatus) and filled with a solution composed of (in
722 mM): potassium-gluconate 135, KCl 4, NaCl 2, EGTA 0.2, HEPES (potassium salt)
723 10, adenosine-triphosphate (magnesium salt, ATP[Mg]) 4, sodium guanosine-
724 triphosphate (NaGTP) 0.5, and phosphocreatine 10 (pH: 7.25–7.30, osmolarity:
725 288–291 mOsmol). The electrodes (resistance: 5–7 M Ω) were connected to the
726 headstage of a NPI ELC-03XS amplifier (NPI, Tamm, Germany). To visualize the
727 cultured cells, the microscope was equipped with differential interference contrast
728 (DIC) optics and with epifluorescence optics for green and red fluorescence
729 (filter sets: Sachs BP450–490, LP520, Zeiss BP546/12, LP590). Images were taken
730 and displayed using a software-operated CCD microscope camera (ORCA R,
731 Hamamatsu, Germany). Following membrane rupture, the cells were voltage-
732 clamped to a holding potential of –60 mV and kept under this condition until
733 the holding current stabilized (3–5 min). Then the amplifier was switched to
734 current-clamp mode. The recorded signals were amplified (\times 10), filtered at 10 or
735 20 kHz (current clamp) and at 5 kHz (voltage clamp), digitized at a sampling rate
736 of 10 or 20 kHz and stored on a computer for offline analysis. Data acquisition
737 and generation of command pulses was done using a CED 1401 Micro 3 system in
738 conjunction with Signal6 data acquisition software (Cambridge electronic design).
739 Data analysis was performed using Igor Pro 6 (WaveMetrics, Lake Oswego,
740 USA) together with the NeuroMatic IGOR plugin (www.neuromatic.thinkrandom.com).
741 Determination of the input resistance, RN, was performed by measuring the

658 amplitude of a voltage deviation induced by a small hyperpolarizing current
659 pulse (1 s, 2–10 pA). The total membrane capacity CN was estimated
660 using a method described by Zemankovics et al.⁴⁶. The ability of the cells to
661 generate action potentials was tested by injecting depolarizing current pulses
662 (50 ms) with increasing current strengths (ΔI : 2–10 pA) or by depolarizing
663 current ramps (50 ms) from 0–100 pA. Spike discharge was analyzed by injecting a
664 series of depolarizing current pulses (duration: 1 s) with a stepwise increment
665 (ΔI : 2–10 pA).

665 **Bulk RNA sequencing.** Primary pericytes from three different human donors were
666 transduced with Ascl1, Sox2, AS, and/or control retroviruses and purified by FACS
667 at 2 and 7 dpi. RNA was isolated using the RNeasy Micro Kit (Qiagen). Following
668 Ribo-Zero removal, the RNA-seq library was prepared in accordance with
669 Illumina's instructions using oligo-dT primers. The RNA-seq output in FASTQ
670 format was aligned to the human hg38 genome (sourced from UCSC) using
671 TopHat v2.0.8⁴⁷ and only uniquely mapped reads were retained for further analysis.
672 SAMTOOLS v.0.1.19⁴⁸ was used for file format conversions (SAM and BAM). The
673 read counts per gene were calculated using HTSeq v0.5.4p1⁴⁹. The DESeq package⁵⁰
674 was used thereafter for differential expression analysis. P_{adj} values were calculated
675 with the Benjamini–Hochberg procedure.

675 **GO terms analysis of bulk RNA-seq data.** GO enrichment analysis was performed
676 using the Bioconductor package TopGO employing the default algorithm weight01⁵¹.
677 Genes were considered significantly deregulated with $P_{adj} < 0.01$. GO terms were
678 ordered according to their significance as determined by Fisher's exact test.

679 **Capture of single cells and preparation of cDNA.** Transduced human brain
680 pericytes were sorted using FACS and single cells were captured on a medium-
681 sized (10- to 17- μ m cell diameter) microfluidic RNA-seq chip using the Fluidigm
682 C1 system. Cells were loaded onto the chip at a concentration of 350–500 cells per
683 μ L and imaged by phase-contrast to assess number of cells per capture site. Only
684 single cells were included in the analysis. cDNAs were prepared on chip using the
685 SMARTer v4 Ultra Input Low RNA kit for Illumina (Clontech).

686 **RNA-seq library construction and cDNA sequencing.** Size distribution and
687 concentration of single-cell cDNA was assessed on a capillary electrophoresis-
688 based fragment analyzer (Advanced Analytical Technologies), and only single
689 cells with high quality cDNA were further processed. Sequencing libraries
690 were constructed in 96-well plates using the Illumina Nextera XT DNA Sample
691 Preparation kit, using primer sets A and B according to the protocol supplied
692 by Fluidigm and as described previously⁴¹. Libraries were quantified by Agilent
693 Bioanalyzer using a High Sensitivity DNA analysis kit, as well as fluorimetrically
694 using Qubit dsDNA HS Assay kits and a Qubit 2.0 Fluorimeter (Invitrogen,
695 Thermo Fisher Scientific). Up to 192 single-cell libraries were pooled and 100-bp
696 paired-end sequenced on one lane of the Illumina HiSeq 2500 to a depth of at
697 least 500,000 reads per cell. Base calling, adaptor trimming, and de-multiplexing
698 was performed as described^{52,53}. The transcriptomes of a total of 769 cells was
699 measured from the following 12 independent experiments: 2-dpi control (76 cells,
700 1 experiment), 2-dpi Ascl1-only (82 cells, 1 experiment), 7-dpi Ascl1-only (64 cells,
701 1 experiment), 2-dpi AS (86 cells, 1 experiment), 7-dpi AS (48 cells, 1 experiment),
702 14-dpi AS (79 cells, 2 experiments), 21/22-dpi AS (130 cells, 2 experiments), 14-
703 dpi ASD (183 cells, 2 experiments), and 14-dpi ASN (21 cells, 1 experiment). See
704 Supplementary Table 5 for the transcriptome data for all 769 cells with annotations
705 (quantification in log₂(FPKM)).

706 **Processing, analysis, and graphic display of single-cell RNA-seq data.** Reads
707 were aligned to a Bowtie2⁵⁴-indexed human genome (hg38 sourced from Ensembl)
708 supplemented with DNA sequences for *Egfp*, *mCherry*, *DsRed*, mouse *Ascl1*, and
709 mouse *Sox2* using TopHat⁴⁷ with default settings. Transcript levels were quantified
710 as fragments per kilobase of mapped reads (FPKM) generated by Cufflinks⁵⁵
711 using GENCODE protein-coding genes (hg38 Havana). We excluded cells that
712 had less than 100,000 reads, expressed $< 1,000$ genes, or did not express either of
713 two housekeeping genes *ACTB* and *GAPDH*. Transcript levels were converted
714 to the log₂-space by taking the log₂(FPKM). R studio (<https://www.rstudio.com/>)
715 was used to run custom R scripts to perform PCA (FactoMineR package) and
716 hierarchical clustering (stats package) and to construct heatmaps, correlation plots,
717 scatter plots, violin plots, dendrograms, bar graphs, and histograms. Generally,
718 ggplot2 and gplots packages were used to generate data graphs. The Seurat package
719 implemented in R was used to identify cell clusters and perform differential gene
720 expression based on t-SNE⁵⁶. The Monocle2 package⁵⁰ was used to analyze cell
721 lineage relationships. Covariance network analysis and visualizations were done
722 using igraph implemented in R (<http://igraph.org/>). Signatures were calculated
723 by summing the log₂(FPKM) expression values of each gene in a set of genes
724 comprising a signature (Supplementary Table 6).

725 **10× Genomics scRNA-seq experiment.** For the 10× Genomics experiment, cells
726 were transfected with AS, treated with dorsomorphin, and analyzed at 14 dpi.
727 Cells were sorted based on the expression of GFP and used for one encapsulation.
728 10× Genomics sample libraries were sequenced on an Illumina HiSeq 2500 and

base calling, adaptor trimming, and de-multiplexing of single cells were performed
using 10× Genomics Cell Ranger 2.0 software. We performed PCA and t-SNE
analyses using the Seurat v2.0 package for R on 3,419 single cells with 1,000–
7,000 genes detected (Supplementary Table 7). We used genes correlating and
anticorrelating with the first eight principal components to cluster the cells, and
found that clustering patterns were robust across multiple PC inclusions. Neuronal
cluster-specific markers were found using Seurat's implementation of the 'bimod'
likelihood-ratio test for single-cell gene expression data, and the top genes were
selected based on the average log fold-change.

Antibodies. The following antibodies were used: mouse (IgG2b) anti-TUBB3
(Sigma; cat. no. T8660; 1:300), rat IgG2a anti-CD49f-PE (Miltenyi Biotec; cat. no.
130-100-096; 1:11), recombinant human anti-CD4 (Miltenyi; cat. no. 130-109-
537; 1:11), rabbit anti-GABA (Abcam; cat. no. ab17413; 1:1,000), chick anti-GFP
(Aves; cat. no. GFP-1020; 1:500), mouse (IgG1) anti-Pvalb (Swant; cat. no. PV-235;
1:1,000), rabbit anti-Pdgfrb (Cell Signaling; cat. no. 3169S; 1:300), rat anti-RFP
(Chromotek; cat. no. 5F8; 1:500), mouse (IgG2b) anti-SMA (Sigma; cat. no. A5228;
1:500), and rabbit anti-VGLUT1 (Synaptic Systems, cat. no. 135302; 1:500). For
FACS we additionally used mouse (IgG2b) anti-LEPR Al647 (BD Pharmingen;
cat. no. 564376; 1:20) and corresponding isotype control (BD Pharmingen; cat.
no. 557903; 1:20). Antibodies were selected according to the antibody validation
reported by the distributing companies.

Accession codes. GEO: scRNA-seq data, [GSE113036](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE113036).

Reporting Summary. Further information on experimental design is available in
the Nature Research Reporting Summary linked to this article.

Data availability and accession codes. The scRNA-seq data used in this study
have been deposited in the Gene Expression Omnibus (GEO) under accession
number [GSE113036](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE113036). The data that support the findings of this study are available
from the corresponding author upon reasonable request.


References

- Karow, M., Schichor, C., Beckervordersandforth, R. & Berninger, B. Lineage-reprogramming of pericyte-derived cells of the adult human brain into induced neurons. *J. Vis. Exp.* <https://doi.org/10.3791/51433> (2014).
- Yu, P. B. et al. Dorsomorphin inhibits BMP signals required for embryogenesis and iron metabolism. *Nat. Chem. Biol.* **4**, 33–41 (2008).
- Inman, G. J. et al. SB-431542 is a potent and specific inhibitor of transforming growth factor-beta superfamily type I activin receptor-like kinase (ALK) receptors ALK4, ALK5, and ALK7. *Mol. Pharmacol.* **62**, 65–74 (2002).
- Laping, N. J. et al. Inhibition of transforming growth factor (TGF)-beta1-induced extracellular matrix with a novel inhibitor of the TGF-beta type I receptor kinase activity: SB-431542. *Mol. Pharmacol.* **62**, 58–64 (2002).
- Martynoga, B. et al. Epigenomic enhancer annotation reveals a key role for NFIX in neural stem cell quiescence. *Genes Dev.* **27**, 1769–1786 (2013).
- Graham, A. C., Francis-West, P., Brickell, P. & Lumsden, A. The signalling molecule BMP4 mediates apoptosis in the rhombencephalic neural crest. *Nature* **372**, 684–686 (1994).
- Chen, A. E., Borowiak, M., Sherwood, R. I., Kweudjeu, A. & Melton, D. A. Functional evaluation of ES cell-derived endodermal populations reveals differences between Nodal and Activin A-guided differentiation. *Development* **140**, 675–686 (2013).
- Kumar, A. et al. Nodal signaling uses activin and transforming growth factor-beta receptor-regulated Smads. *J. Biol. Chem.* **276**, 656–661 (2001).
- Schubert, W. Multiple antigen-mapping microscopy of human tissue. in *Experta Medica: Advances in Analytical Cellular Pathology* (eds. Burger, G., Oberholzer, M. & Vooijs, G.P.) 97–98 (Elsevier, Amsterdam, 1990).
- Ferreira, T. A. et al. Neuronal morphometry directly from bitmap images. *Nat. Methods* **11**, 982–984 (2014).
- Schindelin, J. et al. Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).
- Zemankovics et al., 2010
- Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* **25**, 1105–1111 (2009).
- Li, H. et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).
- Anders, S., Pyl, P. T. & Huber, W. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).
- Oshlack, A., Robinson, M. D. & Young, M. D. From RNA-seq reads to differential expression results. *Genome Biol.* **11**, 220 (2010).
- Alexa, A., Rahnenführer, J. & Lengauer, T. Improved scoring of functional groups from gene expression data by decorrelating GO graph structure. *Bioinformatics* **22**, 1600–1607 (2006).
- Renaud, G., Kircher, M., Stenzel, U. & Kelso, J. freeBis: an efficient basecaller with calibrated quality scores for Illumina sequencers. *Bioinformatics* **29**, 1208–1209 (2013).

- 724 53. Renaud, G., Stenzel, U., Maricic, T., Wiebe, V. & Kelso, J. deML: robust
725 demultiplexing of Illumina sequences using a likelihood-based approach.
726 *Bioinformatics* **31**, 770–772 (2015).
- 727 54. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2.
728 *Nat. Methods* **9**, 357–359 (2012).
- 729
- 730
- 731
- 732
- 733
- 734
- 735
- 736
- 737
- 738
- 739
- 740
- 741
- 742
- 743
- 744
- 745
- 746
- 747
- 748
- 749
- 750
- 751
- 752
- 753
- 754
- 755
- 756
- 757
- 758
- 759
- 760
- 761
- 762
- 763
- 764
- 765
- 766
- 767
- 768
- 769
- 770
- 771
- 772
- 773
- 774
- 775
- 776
- 777
- 778
- 779
- 780
- 781
- 782
- 783
- 784
- 785
- 786
- 787
- 788
- 789
55. Trapnell, C. et al. Transcript assembly and quantification by RNA-Seq reveals
unannotated transcripts and isoform switching during cell differentiation.
Nat. Biotechnol. **28**, 511–515 (2010).
56. Macosko, E. Z. et al. Highly parallel genome-wide expression profiling of
individual cells using nanoliter droplets. *Cell* **161**, 1202–1214 (2015).

Uncorrected proofs

QUERY FORM

Nature Neuroscience	
Manuscript ID	[Art. Id: 168]
Author	Marisa Karow 

AUTHOR:

The following queries have arisen during the editing of your manuscript. Please answer by making the requisite corrections directly in the e-proofing tool rather than marking them up on the PDF. This will ensure that your corrections are incorporated accurately and that your paper is published as quickly as possible.

Query No.	Nature of Query
Q1:	Your paper has been copyedited. Please review every sentence to ensure that it conveys your intended meaning; if changes are required, please provide further clarification rather than reverting to the original text. Please note that formatting (including hyphenation and reference citations) has been made consistent with our house style.
Q2:	Author surnames have been highlighted - please check these carefully and indicate if the first name or surname have been marked up incorrectly. Please note that this will affect indexing of your article, such as in PubMed.
Q3:	Please note that the eproof should be amended in only one browser window at any one time, otherwise changes will be overwritten.
Q4:	In the sentence beginning “In sharp contrast, AS...” please provide an exact P value to support this assertion of significance, or revise as “substantial” or similar. The P value may also appear in the supplementary material, but it must appear in the main text.
Q5:	Define violin plot elements, i.e., median, interquartile range, 95% confidence interval, etc.
Q6:	In panel F, define RFU and FSC, and define the blue and purple colors.
Q7:	In D, define ISH
Q8:	In panel B, define P.
Q9:	scRNA correct as defined? Or define scRNA.
Q10:	In the sentence beginning “Cells were first pretreated...” please provide a numerical temperature/temperature range for “room temperature.”
Q11:	The citation “Zemankovics et al., 2010” was moved to the reference list as #46. Please provide sufficient information to identify this paper or remove the reference.
Q12:	iGraph URL okay as updated?
Q13:	Please ensure that this accession code record is released within 1–2 weeks of receiving your eProof.
Q14:	Please confirm that you have provided names and affiliations for all researchers who provided materials.
Q15:	Please check that the Competing Interests declaration is correct as stated. If you declare competing interests, please check the full text of the declaration for accuracy and completeness.
Q16:	Fill out this reference with either: author names, journal name, volume and page number(s) or DOI.

QUERY FORM

Nature Neuroscience	
Manuscript ID	[Art. Id: 168]
Author	Marisa Karow

AUTHOR:

The following queries have arisen during the editing of your manuscript. Please answer by making the requisite corrections directly in the e-proofing tool rather than marking them up on the PDF. This will ensure that your corrections are incorporated accurately and that your paper is published as quickly as possible.

<i>Query No.</i>	<i>Nature of Query</i>
Q17:	Define violin plot elements, i.e., median, interquartile range, 95% confidence interval, etc.

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see [Authors & Referees](#) and the [Editorial Policy Checklist](#).

Statistical parameters

When statistical analyses are reported, confirm that the following items are present in the relevant location (e.g. figure legend, table legend, main text, or Methods section).

n/a Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- An indication of whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistics including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated
- Clearly defined error bars
State explicitly what error bars represent (e.g. SD, SE, CI)

Our web collection on [statistics for biologists](#) may be useful.

Software and code

Policy information about [availability of computer code](#)

Data collection

For Sholl analyses images were analyzed using the ImageJ(v1.51a-1.51h) Plugin Sholl Analysis v3.6.4; tracing individual neuronal processes was performed using ImageJ software.

Data analysis

Details about data analysis and visualization are provided in the Online Methods section of the paper. The following versions were used: Bioconductor package TopGO v2.3.1 employing the default algorithm weight01, GraphPad Prism v6.01, Igor Pro6 6.0.3.0; Custom R packages were used: TopHat v2.0.8, SAMTOOLS v.0.1.19, HTSeq v0.5.4p1, DESeq2 v1.3.0, FactoMineR v1.34, Seurat v1.4, Monocle2 v2.6.4, igraph v1.2.1, Seurat v2.1-2.2

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors/reviewers upon request. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

The scRNA-seq data used in this study have been in the Gene Expression Omnibus (GEO) under accession number GSE113036. The data that support the findings of this study are available from the corresponding author upon reasonable request. Correspondence and requests for materials and data should be addressed to M.K. (marisa.karow@med.uni-muenchen.de), and B.T. (barbara_treutlein@eva.mpg.de), and B.B. (berningb@uni-mainz.de).

Field-specific reporting

Please select the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/authors/policies/ReportingSummary-flat.pdf](https://www.nature.com/authors/policies/ReportingSummary-flat.pdf)

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	No statistical methods were used to pre-determine sample sizes but our sample sizes are similar to those reported in previous publications (Karow et al., Cell Stem Cell 2012; Treutlein et al., Nature 2016)
Data exclusions	No data points were excluded for the analysis, except for cells in the scRNA-seq analyses not fulfilling the required criteria. For the fluidigm C1 scRNA-seq data: We excluded cells that had less than 100,000 reads, did not express > 1000 genes, or did not express either of two housekeeping genes ACTB and GAPDH. For the 10x Genomics scRNA-seq data: only single cells with 1,000-7,000 genes were included; cells with lower or higher number of detected genes were excluded.
Replication	For all experiments all replicates are indicated in the figure legends or the methods sections. We have provided all informations to reproduce the experiments. All replications were successful.
Randomization	Samples (pericyte donors, coverslips in 24-well plates, T25 or T75 cell culture flasks) were randomly assigned for transduction with different viruses.
Blinding	scRNA-sequencing analyses were performed unbiasedly and therefore blinding is not applicable. Regarding all other data, if not indicated otherwise (e.g. Sholl analysis), data collection and analysis were not performed blind to the conditions of the experiments. These experiments were performed by a single experimentator (MK)

Reporting for specific materials, systems and methods

Materials & experimental systems

n/a	Included in the study
<input type="checkbox"/>	<input checked="" type="checkbox"/> Unique biological materials
<input type="checkbox"/>	<input checked="" type="checkbox"/> Antibodies
<input type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input type="checkbox"/>	<input type="checkbox"/> Palaeontology
<input type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input type="checkbox"/>	<input checked="" type="checkbox"/> Human research participants

Methods

n/a	Included in the study
<input type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input type="checkbox"/>	<input checked="" type="checkbox"/> Flow cytometry
<input type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

Unique biological materials

Policy information about [availability of materials](#)

Obtaining unique materials All unique material (Ascl1/Sox2 encoding viruses) are available from the authors upon

Obtaining unique materials

Antibodies

Antibodies used

Validation

Eukaryotic cell lines

Policy information about [cell lines](#)

Cell line source(s)

Authentication

Mycoplasma contamination

Commonly misidentified lines (See [ICLAC](#) register)

Palaeontology

Specimen provenance

Specimen deposition

Dating methods

Tick this box to confirm that the raw and calibrated dates are available in the paper or in Supplementary Information.

Animals and other organisms

Policy information about [studies involving animals](#); [ARRIVE guidelines](#) recommended for reporting animal research

Laboratory animals

Wild animals

Field-collected samples

Human research participants

Policy information about [studies involving human research participants](#)

Population characteristics

Recruitment

ChIP-seq

Data deposition

Confirm that both raw and final processed data have been deposited in a public database such as [GEO](#).

Confirm that you have deposited or provided access to graph files (e.g. BED files) for the called peaks.

Data access links
May remain private before publication.

Files in database submission

Genome browser session (e.g. [UCSC](#))

Methodology

Replicates	Describe the experimental replicates, specifying number, type and replicate agreement.
Sequencing depth	Describe the sequencing depth for each experiment, providing the total number of reads, uniquely mapped reads, length of reads and whether they were paired- or single-end.
Antibodies	Describe the antibodies used for the ChIP-seq experiments; as applicable, provide supplier name, catalog number, clone name, and lot number.
Peak calling parameters	Specify the command line program and parameters used for read mapping and peak calling, including the ChIP, control and index files used.
Data quality	Describe the methods used to ensure data quality in full detail, including how many peaks are at FDR 5% and above 5-fold enrichment.
Software	Describe the software used to collect and analyze the ChIP-seq data. For custom code that has been deposited into a community repository, provide accession details.

Flow Cytometry

Plots

Confirm that:

- The axis labels state the marker and fluorochrome used (e.g. CD4-FITC).
- The axis scales are clearly visible. Include numbers along axes only for bottom left plot of group (a 'group' is an analysis of identical markers).
- All plots are contour plots with outliers or pseudocolor plots.
- A numerical value for number of cells or percentage (with statistics) is provided.

Methodology

Sample preparation	For sorting of transduced cells for further i) culturing, ii) bulk RNA-sequencing, iii) scRNA-sequencing, primary pericytes were detached from the culture dish using TrypLE for 4-6 minutes and subsequently resuspended in 500-1000 μ l pericyte growth medium. For the separation of LEPR-positive and -negative pericyte populations, pericyte cultures were detached from the culture dish using TrypLE for 4-6 minutes and subsequently 1-5 x 10 ⁵ cells were resuspended in 100 μ l staining solution (PBS plus 0.5% BSA).
Instrument	FACS Aria (BD)
Software	FACS Diva Software
Cell population abundance	The purity of the fluorescent reporter-positive populations that were used for (bulk- and) scRNA-sequencing was confirmed via quantification of the reporter gene expression and resulted in a purity of more than 92%. Due to limitations of using the directly PE conjugated anti-LepR antibody, the purity of the post-sort fraction of the LepR-positive cell population could not be confirmed by additional post-FACS immunohistochemistry.
Gating strategy	For sort of transduced cells: Gating was achieved via subtracting the autofluorescence of non transduced cells and control (DsRed or GFP only) transduced cells were used as respective controls. For the LEPR-based sort: An Alexa647-conjugated isotype control antibody (1:100, BD Pharmingen) was used to gate the proper populations.
<input checked="" type="checkbox"/> Tick this box to confirm that a figure exemplifying the gating strategy is provided in the Supplementary Information.	

Magnetic resonance imaging

Experimental design

Design type	no MRI imaging used in this study
Design specifications	Specify the number of blocks, trials or experimental units per session and/or subject, and specify the length of each trial or block (if trials are blocked) and interval between trials.
Behavioral performance measures	State number and/or type of variables recorded (e.g. correct button press, response time) and what statistics were used to establish that the subjects were performing the task as expected (e.g. mean, range, and/or standard deviation across subjects).

Acquisition

Imaging type(s)

Field strength

Sequence & imaging parameters

Area of acquisition

Diffusion MRI Used Not used

Preprocessing

Preprocessing software

Normalization

Normalization template

Noise and artifact removal

Volume censoring

Statistical modeling & inference

Model type and settings

Effect(s) tested

Specify type of analysis: Whole brain ROI-based Both

Statistic type for inference (See [Eklund et al. 2016](#))

Correction

Models & analysis

n/a	Involvement in the study
<input type="checkbox"/>	<input type="checkbox"/> Functional and/or effective connectivity
<input type="checkbox"/>	<input type="checkbox"/> Graph analysis
<input type="checkbox"/>	<input type="checkbox"/> Multivariate modeling or predictive analysis

Functional and/or effective connectivity

Graph analysis

Multivariate modeling and predictive analysis