

## The algorithms of adjuvant therapy in gliomas and their effect on survival.

Journal: Journal of Neurosurgical Sciences

Paper code: J Neurosurg Sci-4610

Submission date: October 8, 2018

Article type: Review Article

Files:

1. Manuscript

Version: 1

Description: manoscritto

File format: application/msword

2. Figures 1

Version: 1

Description: figure 1

File format: application/pdf

3. Figures 2

Version: 1

Description: figure 2

File format: application/pdf

4. Figures 3

Version: 1

Description: figure 3

File format: application/pdf

5. Figures 4

Version: 1

Description: figure 4

File format: application/pdf

## The algorithms of adjuvant therapy in gliomas and their effect on survival.

Christoph Straube<sup>1,3</sup>, Friederike Schmidt-Graf<sup>4</sup>, Benedikt Wiestler<sup>5</sup>, Claus Zimmer<sup>5</sup>, Bernhard Meyer<sup>6</sup>,  
Stephanie E. Combs<sup>1,2,3</sup>

1 – Department of Radiation Oncology, Klinikum rechts der Isar, Technical University of Munich (TUM), Munich, Germany

2 – Institut für Innovative Radiotherapie (iRT), Department of Radiation Sciences (DRS), Helmholtz Zentrum München, Munich, Germany

3 – Deutsches Konsortium Translationale Krebsforschung (DKTK), Partner Site Munich, Munich, Germany

4 – Department of Neurology, Klinikum rechts der Isar, Technical University of Munich (TUM), Munich, Germany

5 – Department of Neuroradiology, Klinikum rechts der Isar, Technical University of Munich (TUM), Munich, Germany

6 – Department of Neurosurgery, Klinikum rechts der Isar, Technical University of Munich (TUM), Munich, Germany

### Corresponding author:

Univ.-Prof. Dr. Stephanie E. Combs  
Department of Radiation Oncology  
Technische Universität München  
Ismaninger Str. 22  
81675 München  
Tel.: +49-89-4140-4501  
Fax.: +49-89-4140-4477  
e-mail: Stephanie.Combs@tum.de

1  
2  
3 **Abstract**

4 The treatment of gliomas became more sophisticated during the last decades. As by now, adjuvant  
5 treatment after maximum safe resection is considered an important and effective treatment strategy  
6 in most gliomas, yet the decision is based on several factors. This review summarizes the available  
7 evidence for the current adjuvant treatment algorithms with a focus on the impact on the survival of  
8 glioma patients. The review is based on the current guidelines, but it also includes new insights which  
9 have not yet been included into the official guidelines.

10  
11 **Key-words**

12 Glioma, Glioblastoma, Oligodendroglioma, adjuvant treatment, Radiotherapy  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

PEER REVIEW COPY  
Journal of Neurosurgical Sciences

## Introduction

The adjuvant treatment of gliomas has undergone important changes within the past years [1, 2], which now involves surgical, medical as well as technical therapies side by side. Within the past two centuries, important improvements in all of these therapeutic strategies have improved the outcomes of our patients. Beginning with neurosurgery, microsurgical techniques with the use of intraoperative CT- and MR-imaging, neuro-navigation as well as 5-alpha-aminolevulinat (ALA) enhanced resections, among others, have improved the safety as well as the extent of resection of glioma patients. This development has led to an improvement of the prognosis [3, 4]. Not only the surgical techniques, but also the pathological review has made a large step forward. With the advent and inclusion of molecular findings into the diagnosis of primary brain tumors, the accuracy of the diagnosis has improved and the prognostication has become more accurate [5]. The molecular factors in the new WHO classification are also predictive for the following adjuvant treatments [6, 7]. Systemic treatments are now a standard of care in most primary brain tumors, where the choice of the treatment regimen is largely influenced by the molecular pattern of the disease [1]. Radiotherapy has undergone fundamental changes within the past years, too. Modern linear accelerators, enhanced by modern computer based planning algorithms, allow highly conformal dose distributions to the tumor with simultaneous protection of adjacent organs at risk. Furthermore, the precision of the treatment delivery has reached a millimeter or even sub-millimeter level with the use of stereotactic or image-guidance techniques. Besides this, the advent of particle therapies now allows further dose escalations. All of these techniques went hand-in-hand with improvements in neuroimaging, which evolved from anatomic imaging towards functional and metabolic visualisation which now allows to capture the heterogeneity within subvolumes of the tumor [8].

All of these improvements have led us to a plethora of trials, many of them comparing different adjuvant treatments. In this review, we present the milestone trials and their results and generate an algorithm for the adjuvant treatment of gliomas. The algorithm takes into account the 2017 EANO guidelines as well as the 2018 NCCN guidelines [1, 9]. However, we modified the recommendation, where appropriate, to include novel results. The target volume recommendations for radiation therapy are based on the recommendation from the EORTC as well as the ESTRO[10].

## Low-Grade Glioma

The treatment of low-grade gliomas (LGG) is currently shifting towards a more intense treatment [11]. After maximum safe resection, national as well as international guidelines recommend adjuvant treatment in all patients who fulfill a set of risk factors [1][9]. The definition of these risk factors has changed over the past two decades, thereby focussing on the age of  $\geq 40$  years and a subtotal resection as risk factors. Additional clinical features, such as a tumor size of  $> 6$ cm or the presence of neurologic deficits, which were suggested earlier by Pignatti et al. [12], have currently lost their influence. Currently, molecular factors have gained more attention which is discussed in detail later in this article.

The regimen with the highest level of evidence (IB) in the adjuvant situation in LGG is the RTOG 9802 regimen. The trial included patients with diffuse gliomas: Patients with low-risk features as defined above were observed, while patients with high-risk-features were randomized to receive either a mono-radiotherapy or radiotherapy (RT) followed by six cycles of combination chemotherapy (ChT). The radiotherapy dose was 54 Gy in 30 fractions. The adjuvant combination chemotherapy included procarbazine (60 mg/m<sup>2</sup>/d8-21, q56), CCNU (110 mg/m<sup>2</sup>/d1,q56) and Vincristine (1.4mg/m<sup>2</sup>/d8+29,q56)(PCV). As anticipated, the combination regimen was associated with an increased rate of hematologic and serologic side effects, and the average number of cycles was 3 for procarbazine and 4 for CCNU and vincristine. Besides this, the two regimens were equally tolerated. The long-term results of RTOG 9802 were published in 2016 and showed a significant improvement in the long term-survival of LGG patients. The median overall survival (mOS) was increased from 7.8 to 13.3 years, with an even more pronounced effect in patients with oligodendroglial histology [13]. Accordingly, the adjuvant treatment with an RT to 54 Gy in 30 fractions, followed by six cycles of PCV is the best-established standard for high-risk LGGs. The RT is prescribed to the resection cavity as

well as mass like hyperintensities in the fluid-attenuated inversion recovery (FLAIR) sequence plus an anatomically adapted margin of 1.0 to 1.5 cm (clinical target volume, CTV), a margin of 0.3 to 0.5 mm is added to account for positioning uncertainties (planning target volume, PTV).

As PCV is associated with an increased rate of hematologic as well as liver toxicities, especially when combined with temozolomide (TMZ), many centers use TMZ instead of PCV for the adjuvant treatment of diffuse astrocytomas [11]. The evidence for this regimen is based on three arguments. Firstly, the phase II study RTOG 0424 showed a favorable outcome in patients receiving RT with concomitant and adjuvant TMZ as compared to historical controls from EORTC 22844 and 22845 [12, 14]. The 3-year survival in this regimen was 73.1%. A recent secondary analysis further validated the prognostic impact of O6-Methylguanin-DNA-methyltransferase (MGMT) promoter methylation status as well as IDH1/2 mutations in LGGs [15]. Secondly, EORTC 22033-26033 showed the general efficacy of TMZ as a stand-alone treatment. The randomized trial patients to receive either RT to 50.4 Gy in 28 fractions or 12 cycles of TMZ (150-200mg/m<sup>2</sup>/d1-5,q28). The trial was designed to detect a 13% increase in the progression-free survival (PFS) at five years of follow up in favor of TMZ. As the trial was formally negative, a mono-therapy with TMZ is currently not recommended. None the less, TMZ was at least not significantly inferior to a mono- RT, which argues in favor of its efficacy [16]. Thirdly, the results from NOA-04 are frequently quoted, although the trial only included patients with high-grade gliomas (HGG). The trial randomized HGG patients to receive either a mono RT with 59.4 Gy in 33 fractions or chemotherapy. The chemotherapy-arm was divided in either TMZ or PCV. The important strength of this trial was the inclusion of a molecular analysis which was able to stratify an increased efficacy for PCV to patients with a co-deletion of 1p19q (loss of heterogeneity, LOH) [7]. As the less toxic TMZ was not inferior to PCV in patients without a co-deletion of 1p19q, TMZ is considered to be equally effective to PCV in diffuse astrocytomas. The obvious limitation of this argumentation is that NOA-04 only included HGGs.

Currently, the results from the CODEL-trial are pending (NCT00887146). This originally four-armed trial randomized patients with either grade II or III oligodendrogliomas, as defined by a co-deletion of 1p19q, to receive either RT alone, TMZ alone, RT with concomitant TMZ followed by 12 cycles of TMZ or RT followed by six cycles of PCV. The mono-TMZ arm was closed early after an interim analysis due to a worse PFS and OS; the mono-RT arm was closed after the results from RTOG 9402/EORTC 26951 were published [17].

While the general indication for high-risk LGGs is relatively clear, the evidence for the adjuvant treatment of low-risk LGGs is less established. Prospective data from the observation arm of RTOG 9802 supplies the best evidence for this situation. As reported by an abstract in 2006, the 111 patients in the observation arm, the 2- and five years OS was 99% and 94%; the PFS was 82% and 50% [18]. While the OS times thus were superior to the RT+PCV-Arm of the trial, the PFS was inferior to the RT-PCV arm at least at five years. Noteworthy, patients in the RT-PCV were considered to have an unfavorable prognosis as defined by the inclusion criteria [13].

Noteworthy, newer data stress the importance of the molecular pattern of LGGs as risk factors, even beyond the clinical features [6]. While some of these molecular features have already been implemented into the new WHO classification of brain malignancies [5], other markers seem to have an impact on the prognosis of LGG patients, too. The MGMT promoter methylation status is among these markers [15]. As IDH wild-type and nonmethylated MGMT promoters seem to be negative prognostic markers, these patients should undergo active treatment, even despite the presence of clinical low-risk features [1, 11].

Based on these considerations, we propose an adjuvant treatment algorithm as depicted in Figure 1.

INSERT FIGURE 1

### Anaplastic Gliomas

Comparable to LGGs, anaplastic gliomas are now preferably treated with a multimodal adjuvant approach. After maximum safe resection, usually, RT is prescribed. The RT is 59.4 to 60 Gy in 33 or 30 fractions of 1.8 or 2.0 Gy. The target volume encompasses the resection cavity, contrast

1 enhancements on computed tomography (CT) or magnetic resonance imaging (MRI) T1-Sequences as  
 2 well as mass-like FLAIR hyperintensities defined as gross total volume (GTV). A 1.5-2.0cm margin is  
 3 added and adapted to account for anatomic barriers (CTV), the PTV encompasses additional 0.3-0.5  
 4 mm.

5 Different trials support the systemic treatment in anaplastic astrocytomas (AA) and anaplastic  
 6 oligodendrogliomas (AO). Based on the results of NOA-04, AO, as defined by a LOH 1p19q, seemed to  
 7 respond better to PCV as compared to TMZ [7]. As NOA-04 showed similar efficacies of mono RT and  
 8 mono ChT for AA and AO, thereby proofing the stand-alone efficacy of both modalities, the RTOG as  
 9 well as the EORTC started two trials to clarify the efficacy of a multimodal approach with RT with  
 10 neoadjuvant (RTOG 9402) or adjuvant (EORTC 26951) PCV [19, 20]. RTOG 9402 was formally  
 11 negative, with a similar OS of 4.6 and 4.7 years for patients treated with or without neoadjuvant PCV  
 12 before RT [20]. Noteworthy, patients with anaplastic oligo-astrocytoma (AOA) as well as with AO  
 13 could enter the trial. When the patients were stratified by LOH 1p19q, patients with a co-deletion  
 14 significantly benefited from a multimodal approach (OS 14.7 vs. 7.3 years) [20]. In opposite to RTOG  
 15 9402, EORTC 26951 was positive for the entire study cohort which also included patients with AO and  
 16 AOA as defined by histology (OS 42.3 months vs. 30.6 months for RT+PCV vs. RT) [19]. Also in EORTC  
 17 26951, the subgroup of patients with LOH 1p19q significantly benefited from the addition of PCV (OS  
 18 not reached vs. 112 months in RT+PCV vs. RT) [19]. When comparing the tolerability of both  
 19 regimens, EORTC 26951 had a better treatment compliance with 95% receiving the standard RT as  
 20 compared to 76% in RTOG 9402; ChT was tolerated equally with 12%, 9%, 22% and 54% receiving  
 21 1,2,3 or 4 cycles of PCV in RTOG 9402 vs. 11%, 22%, 17% and 49 receiving 1,2,3 or 4-6 cycles in EORTC  
 22 26951 [20, 21]. The reduced RT-compliance might be a reason for the primary negative result of  
 23 RTOG 9402. In conclusion, a multimodal approach with upfront RT followed by up to six cycles of PCV  
 24 is the current standard of care in AO. Whether there is a role of TMZ in AO is currently unknown, as  
 25 the final results from CODEL are still pending.

26 For AA the optimal management was less clear until the publication of the preliminary results of the  
 27 CATNON trial in 2017 [22]. The large phase 3 trial randomized patients to either receive RT, RT with  
 28 concomitant TMZ (RChT; 75 mg/m<sup>2</sup>/d), RT with adjuvant TMZ (12 cycles, 150-200 mg/m<sup>2</sup>/d1-5,q28)  
 29 or RChT with adjuvant TMZ. A planned interim analysis showed a significantly better five year-OS for  
 30 patients receiving an adjuvant ChT with TMZ as compared to patients not receiving an adjuvant ChT  
 31 (55.9 vs. 44.1%) [22]. The current EANO guidelines did already include these results [1]. As the data  
 32 for the concomitant ChT are not available yet, the current algorithm for AA is to treat patients with  
 33 an adjuvant RT with or without concomitant TMZ, followed by up to 12 cycles of adjuvant TMZ. The  
 34 treatment algorithm for WHO Grade III gliomas (AA, AO, AOA) are summarized in Figure 2.

35  
 36  
 37  
 38 INSERT FIGURE 2

### 39 Glioblastoma

40  
 41  
 42 The EORTC 26981/22981/NCIC trial by Stupp et al. defined in 2005 the current standard of care in  
 43 GBM patients younger than 70 years who are in a good performance status [23]. The trial tested an  
 44 adjuvant treatment with RT alone (60 Gy in 30 fractions) vs. a multimodal regimen with RT and  
 45 concomitant TMZ (75 mg/m<sup>2</sup>) followed by 6 adjuvant cycles of TMZ (150-200 mg/m<sup>2</sup>/d1-5, q28). The  
 46 PTV in this trial included the resection cavity as well as contrast enhancing lesions with a 1.5 – 2.0 cm  
 47 margin, adapted to anatomic barriers (CTV); for positioning uncertainties, a 0.5 cm margin was added  
 48 (an example can be found in Figure 3). The investigative arm gained a significantly improved survival  
 49 (12.1 vs. 14.6 months), with an additional advantage for patients with methylated MGMT promoter  
 50 [24]. Meanwhile, several trials investigated combination regimens with additional Bevacizumab  
 51 against this standard, but they of them failed to their primary objectives [25, 26]. Only recently, the  
 52 results of two trials were presented which suggested new treatment options for GBM patients. The  
 53 first one investigated the effect of the addition of TTFIELDS to the adjuvant treatment in GBM. The  
 54 trial randomized 695 patients – who did not progress after standard-of-care RT with concomitant  
 55

1 TMZ – to either receive up to six cycles of TMZ or the same number of TMZ cycles with the  
2 concomitant use of TTFIELDS. The trial was randomized but not blinded, which is one of the critics to  
3 the trial. However, after a median follow-up of 40 months, a significant median survival benefit of 4.9  
4 months was evident (20.9 vs. 16.0 months) [27]. The health-related quality of life was equal in both  
5 arms of the trial [28]. Secondly, the results from the German CeTeG-Trial have been reported at the  
6 SNO conference in 2017. The trial focussed on patients with a methylated MGMT promoter, and  
7 randomized these patients into two arms. Patients in the standard arm received a standard RT with  
8 60 Gy together with concomitant TMZ, followed by up to six cycles of adjuvant TMZ. Patients in the  
9 investigational arm received the same RT, but the systemic treatment consisted of up to six courses  
10 of CCNU (100 mg/m<sup>2</sup>, d1, q42) and TMZ (100 mg/m<sup>2</sup> d2-6, q42). The primary endpoint was the  
11 median OS in the modified intention to treat-cohort (mITT). The mITT included all patients who did  
12 receive the first dose of the study-drug. The survival was analysed after a pre-specified inverse  
13 probability weighting. This technique was used to up- or down-weight patients from centers with a  
14 distribution of RPA classes that was different from the whole population. The trial randomized 141  
15 patients and reported a significant improvement of the median OS in the mITT cohort from 31.4  
16 months in the standard arm to 48.1 months in the investigational arm. Noteworthy, there was  
17 neither a difference in median OS between the groups when all patients were compared in the  
18 unbalanced cohort nor there was a difference in the PFS. The toxicity was higher in the  
19 investigational arm, which was mostly driven by the hematologic toxicities. Taken together, the  
20 standard of care for all patients in a good performance status is a RT to 59.4-60 Gy with concomitant  
21 TMZ followed by up to six courses of adjuvant TMZ. Especially in patients with no progression after  
22 concomitant RT/ChT TTFIELDS can be considered. In thoroughly selected patients with methylated  
23 MGMT promoter and a very good performance status, the CeTeG regimen can be an option, too.  
24 Noteworthy, there are results from several promising trials pending, among these are trials with an  
25 intraoperative RT-boost (INTRAGO II, NCT02685605), early TTFIELDS (Bonn... ) and checkpoint-  
26 inhibition (BMS CA209-498 and BMS CA209-548).  
27 While the former algorithm is appropriate for patients in good performance status, a different  
28 algorithm should be used for elderly and/or frail patients. The best results are achieved when  
29 patients are treated with a short-course RT (40.67 Gy in 15 fractions) with concomitant TMZ (75  
30 mg/m<sup>2</sup>/d) followed by up to 12 cycles of TMZ (150-200 mg/m<sup>2</sup> d1-5, q28). This regimen is based on  
31 the results from NCIC CTG CE.6. The standard arm of this trial was based on the results from Roa et  
32 al., who compared a standard RT (60 Gy, 30 fractions) to a short-course RT (40.67 Gy in 15 fractions)  
33 without ChT. The mOS in this trial was not different (median OS 5.1 vs. 5.6 months) [29]. The  
34 investigational arm used the same RT but combined it with the ChT mentioned above. The  
35 investigational regimen resulted in a significant survival benefit (median OS 9.3 vs. 7.6 months), the  
36 subgroup of patients with a methylated MGMT promoter had a greater benefit (median OS 13.5 vs.  
37 7.7 months) [30]. When patients are not considered to be fit enough to undergo multimodal  
38 adjuvant treatment, mono-therapies should be considered. The efficacy of mono RT in elderly  
39 patients was proven by a French trial which randomized GBM patients older than 70 years to either  
40 receive a standard RT (60 Gy, 30 fractions) or best supportive care (BSC). The addition of RT resulted  
41 in an improved OS (29.1 vs. 16.9 weeks) [31]. Given the poor prognosis even after addition of RT,  
42 several trials investigated shorter course RT regimens with similar results with 40.67 Gy in 15  
43 fractions, 34 Gy in 10 fractions or 25 Gy in 5 fractions (median OS 5.6 vs. 5.1 months (34 Gy vs. 60  
44 Gy) [29], 7.5 vs. 6.0 (40.05 Gy vs. 60 Gy) months [32] and 7.9 vs. 6.4 months (25 Gy vs. 40.05 Gy)  
45 [33]). Despite this, treatment with ChT only can be considered especially in elderly and frail patients  
46 who are presenting with a methylated MGMT promoter [32, 34].  
47 According to the 2017 EANO guidelines, BSC is considered as a best choice, when patients are  
48 presenting with a very unfavorable prognosis, namely a Karnofsky performance score (KPS) of less  
49 than 50% (bed bound patients) as well as for patients who are not able to consent into the treatment  
50 [1]. These criteria should not be based on a single visit, as reversible conditions might, such as  
51 infections, epileptic convulsions or reversible brain edema might be present. Caregivers, therefore,  
52 should thoroughly search for this conditions, initiate an effective treatment and re-evaluate the  
53 performance status after an appropriate interval.  
54  
55

1 A summary of the presented algorithm can be found in figure 4.  
2 INSERT FIGURE 3 AND 4  
3

#### 4 Recurrent Glioma

5 Depending on the grade of the disease as well as of other risk factors, many patients are challenged  
6 by recurrences throughout their disease. Currently, it is not possible to define an ideal algorithm for  
7 the treatment of recurrences, yet a multimodal approach is increasingly considered as an option for  
8 these patients. While it is beyond the scope of this review to discuss all possible treatments of  
9 recurrent gliomas, it seems germane to briefly discuss the current evidence of second surgery with a  
10 focus on adjuvant treatments.  
11

12 A second surgery can be considered as a first step in the treatment whenever it is possible with a  
13 reasonable risk-profile, yet prospective data are missing for LGG and HGG. In GBM, the data from the  
14 DIRECTOR trial were secondarily analyzed for the value of a second surgery. While the trial did not  
15 show a benefit for all patients with second surgery, it did show an improved survival for a subgroup  
16 of patients with gross total resection (GTR) after second surgery (11.4 vs. 9.8 months). Patients with a  
17 subtotal resection after (STR) second surgery had a worse prognosis when compared to patients  
18 without surgery (6.5 vs. 9.8 months) [35]. The authors concluded that second surgery should only be  
19 performed when a safe GTR can be achieved. This conclusion is discussed very controversial, as  
20 pooled data from 20 centers came to different results, thereby supporting the efficacy also for STR  
21 [36].

22 While all patients in the DIRECTOR trial received an adjuvant ChT after the surgery, but no adjuvant  
23 Re-RT, the multi-institutional report from Ringel et al. reported on patients with surgery but with  
24 different adjuvant regimens. This enabled the authors to analyze their cohort for the value of  
25 adjuvant therapies after the second surgery. A significant benefit for patients who received an RT  
26 after second surgery was found (median OS after second surgery 8.5 vs. 13.4 months) [36]. This  
27 finding is in line with other reports [37–39]. Importantly, the indication for a second RT is re-thought  
28 currently. While Re-RT classically was prescribed only in cases with a macroscopic tumor, newer  
29 approaches now investigate the value of a second RT even after GTR of a recurrent glioma [38, 40,  
30 41]. This approach is supported by the efficacy of adjuvant RT in the primary situation, the efficacy of  
31 Re-RT in macroscopic disease as well as by the good safety profile of Re-RT with modern RT  
32 techniques. As by now, the evidence for a routine indication of adjuvant Re-RT after GTR of recurrent  
33 gliomas is still too weak for a general recommendation. Therefore, patients with recurrent gliomas  
34 should be discussed in an interdisciplinary context and included in clinical trials as possible.

35 Concerning systemic treatments after a second surgery, there is no standard defined yet and  
36 recommendations thus are based on the general efficacy of systemic protocols in recurrent GBM  
37 without previous surgery. Nitrosurea, a re-challenge with TMZ or Bevacizumab alone or a  
38 combination of these drugs are frequently used [1, 42]. Notably, Bevacizumab has no approval for  
39 this indication in the European Union, treatments therefore have to be requested at health insurance  
40 providers on an individual basis. Besides the traditional approaches of ChT and RT, alternating  
41 electric fields (TTFields) have proven equal efficacy as compared to best investigator choice ChT in  
42 recurrent HGG and GBM in a multicenter phase III trial [43].  
43  
44

#### 45 Conclusion

46 Adjuvant treatment of gliomas has evolved to follow complex and highly differentiated algorithms.  
47 All of those clearly state that adjuvant treatment in gliomas is essential, independently of primary  
48 histology. Radiation therapy has improved over time and is now highly precise and well targeted. So  
49 normal tissue can be spared, and treatment-related sequelae can be minimized. Chemotherapy can  
50 enhance the effect of radiotherapy, which has been confirmed to subgroups of low-grade gliomas  
51 comparable to high-grade gliomas.  
52  
53

54 In the future, molecular stratification will most likely increase and will further differentiate  
55 treatments. Until then, it is important that interdisciplinary decision making takes place and that any



1 adjuvant treatment is offered in a modern and state-of-the-art manor including advanced  
 2 radiotherapy techniques.  
 3  
 4

### 5 Disclosures

6 Christoph Straube: received a scholarship from Medac GmbH; received a travel grant from NovoCure  
 7 Ltd.;

8 Bernhard Meyer: work as consultants for BrainLab and Nexstim

9 Benedikt Wiestler: received speaker honoraria from Bayer AG

10 Friederike Schmidt-Graf: served as author for Medac GmbH

11 Claus Zimmer: has served on scientific advisory boards for Philips and Bayer Schering; serves as co-  
 12 editor on the Advisory Board of Clinical Neuroradiology; has received speaker honoraria from Bayer-  
 13 Schering and Philips and has received research support and investigator fees for clinical studies from  
 14 Biogen Idec, Quintiles, MSD Sharp & Dome, Boehringer Ingelheim, Inventive Health Clinical UK Ltd.,  
 15 Advance Cor, Brainsgate, Pfizer, Bayer-Schering, Novartis, Roche, Servier, Penumbra, WCT GmbH,  
 16 Syngis, SSS International Clinical Research, PPD Germany GmbH, Worldwide Clinical Trials Ltd.,  
 17 Phenox, Covidien, Actelion, Medivation, Medtronic, Harrison Clinical Research, Concentric,  
 18 Penumbra, Pharmtrace, Reverse Medical Corp., Premier Research Germany Ltd., Surpass Medical Ltd.  
 19 and GlaxoSmithKline.

20 Stephanie E. Combs: has served on Advisory Board of Bristol-Myers-Squibb (BMS); Advisory board  
 21 and Speaker's Bureau for BrainLab; Advisory Board of Roche, Daiichi Sankyo and Varian Medical  
 22 Systems. Has received Speakers Honoraria from BrainLab, Tomotherapy, Dr. Sennewald, BMS, Varian  
 23 Medical Systems, Elekta, Novocure and Medac GmbH.  
 24  
 25

### 26 References

- 27 1. Weller M, van den Bent M, Tonn J, Stupp R, Preusser M, Cohen-Jonathan-Moyal E, Henriksson  
 28 R, Le Rhun E, Balana C, Chinot O, Bendszus M, Reijneveld J, Dhermain F, French P, Marosi M,  
 29 Watts C, Oberg I, Pilkington G, Baumert B, Taphoorn M, Hegi M, Westphal M, Reifenberger R,  
 30 Soffietti S, Wick W (2017) For the European Association for Neuro-Oncology (EANO) Task  
 31 Force on Gliomas. EANO guideline on the diagnosis and treatment of adult astrocytic and  
 32 oligodendroglial gliomas. *Lancet Oncol*. doi: 10.1016/S1470-2045(17)30194-8
- 33 2. Weller M, van den Bent M, Hopkins K, Tonn JC, Stupp R, Falini A, Cohen-Jonathan-Moyal E,  
 34 Frappaz D, Henriksson R, Balana C, Chinot O, Ram Z, Reifenberger G, Soffietti R, Wick W  
 35 (2014) EANO guideline for the diagnosis and treatment of anaplastic gliomas and  
 36 glioblastoma. *Lancet Oncol* 15:e395–e403. doi: 10.1016/S1470-2045(14)70011-7
- 37 3. Brown TJ, Brennan MC, Li M, Church EW, Brandmeir NJ, Rakszawski KL, Patel AS, Rizk EB, Suki  
 38 D, Sawaya R, Glantz M (2016) Association of the Extent of Resection With Survival in  
 39 Glioblastoma: A Systematic Review and Meta-analysis. *JAMA Oncol* 2:1460–1469. doi:  
 40 10.1001/jamaoncol.2016.1373
- 41 4. Stummer W, Pichlmeier U, Meinel T, Wiestler OD, Zanella F, Reulen HJ (2006) Fluorescence-  
 42 guided surgery with 5-aminolevulinic acid for resection of malignant glioma: a randomised  
 43 controlled multicentre phase III trial. *Lancet Oncol* 7:392–401. doi: 10.1016/S1470-  
 44 2045(06)70665-9
- 45 5. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, Ohgaki  
 46 H, Wiestler OD, Kleihues P, Ellison DW (2016) The 2016 World Health Organization  
 47 Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol* 131:1–  
 48 18. doi: 10.1007/s00401-016-1545-1
- 49 6. Reuss DE, Kratz A, Sahm F, Capper D, Schrimpf D, Koelsche C, Hovestadt V, Bewerunge-Hudler  
 50 M, Jones DTW, Schittenhelm J, Mittelbronn M, Rushing E, Simon M, Westphal M, Unterberg  
 51 A, Platten M, Paulus W, Reifenberger G, Tonn JC, Aldape K, Pfister SM, Korshunov A, Weller  
 52 M, Herold-Mende C, Wick W, Brandner S, von Deimling A (2015) Adult IDH wild type  
 53 astrocytomas biologically and clinically resolve into other tumor entities. *Acta Neuropathol*  
 54  
 55

- 130:407–417. doi: 10.1007/s00401-015-1454-8
7. Wick W, Roth P, Hartmann C, Hau P, Nakamura M, Stockhammer F, Sabel MC, Wick A, Koeppe S, Ketter R, Vajkoczy P, Eyupoglu I, Kalff R, Pietsch T, Happold C, Galldiks N, Schmidt-Graf F, Bamberg M, Reifenberger G, Platten M, Von Deimling A, Meisner C, Wiestler B, Weller M (2016) Long-term analysis of the NOA-04 randomized phase III trial of sequential radiochemotherapy of anaplastic glioma with PCV or temozolomide. *Neuro Oncol* 18:1529–1537. doi: 10.1093/neuonc/nov133
  8. Straube C, Bette S, Pyka T, Einhellig H, Zimmer C, Schwaiger M, Combs SE (2017) Moderne Bildgebung in der Neuroonkologie. *Akt Neurol* 489–500. doi: 10.1055/s-0043-111588
  9. Bower M, Waxman J (2018) NCCN Clinical Practice Guidelines in Oncology: Central Nervous System Cancers. *Lect. Notes Oncol.*
  10. Niyazi M, Brada M, Chalmers AJ, Combs SE, Erridge SC, Fiorentino A, Grosu AL, Lagerwaard FJ, Minniti G, Mirimanoff R-O, Ricardi U, Short SC, Weber DC, Belka C (2016) ESTRO-ACROP guideline “target delineation of glioblastomas.” *Radiother Oncol* 118:35–42. doi: 10.1016/j.radonc.2015.12.003
  11. Straube C, Kessel KA, Schmidt-graf F, Krieg SM, Meyer B, Gempt J, Combs SE (2018) A trend towards a more intense adjuvant treatment of low-grade-gliomas in tertiary centers in Germany after RTOG 9802 – results from a multi-center survey. *BMC Cancer* 1–7. doi: 10.1186/s12885-018-4825-4
  12. Pignatti F, van den Bent M, Curran D, Debruyne C, Sylvester R, Therasse P, Afra D, Cornu P, Bolla M, Vecht C, Karim ABMF, European Organization for Research and Treatment of Cancer Brain Tumor Cooperative Group, European Organization for Research and Treatment of Cancer Radiotherapy Cooperative Group (2002) Prognostic factors for survival in adult patients with cerebral low-grade glioma. *J Clin Oncol* 20:2076–84.
  13. Buckner JC, Shaw EG, Pugh SL, Chakravarti A, Gilbert MR, Barger GR, Coons S, Ricci P, Bullard D, Brown PD, Stelzer K, Brachman D, Suh JH, Schultz CJ, Bahary J-P, Fisher BJ, Kim H, Murtha AD, Bell EH, Won M, Mehta MP, Curran WJJ (2016) Radiation plus Procarbazine, CCNU, and Vincristine in Low-Grade Glioma. *N Engl J Med* 374:1344–1355. doi: 10.1056/NEJMoa1500925
  14. Fisher BJ, Hu C, Macdonald DR, Lesser GJ, Coons SW, Brachman DG, Ryu S, Werner-Wasik M, Bahary JP, Liu J, Chakravarti A, Mehta M (2015) Phase 2 study of temozolomide-based chemoradiation therapy for high-risk low-grade gliomas: Preliminary results of radiation therapy oncology group 0424. *Int J Radiat Oncol Biol Phys* 91:497–504. doi: 10.1016/j.ijrobp.2014.11.012
  15. Bell EH, Zhang P, Fisher BJ, Macdonald DR, McElroy JP, Lesser GJ, Fleming J, Chakraborty AR, Liu Z, Becker AP, Fabian D, Aldape KD, Ashby LS, Werner-Wasik M, Walker EM, Bahary J-P, Kwok Y, Yu HM, Laack NN, Schultz CJ, Gray HJ, Robins HI, Mehta MP, Chakravarti A (2018) Association of MGMT Promoter Methylation Status With Survival Outcomes in Patients With High-Risk Glioma Treated With Radiotherapy and Temozolomide: An Analysis From the NRG Oncology/RTOG 0424 Trial. *JAMA Oncol.* doi: 10.1001/jamaoncol.2018.1977
  16. Baumert BG, Hegi ME, van den Bent MJ, von Deimling A, Gorlia T, Hoang-Xuan K, Brandes AA, Kantor G, Taphoorn MJB, Hassel M Ben, Hartmann C, Ryan G, Capper D, Kros JM, Kurscheid S, Wick W, Enting R, Reni M, Thiessen B, Dhermain F, Bromberg JE, Feuvret L, Reijneveld JC, Chinot O, Gijtenbeek JMM, Rossiter JP, Dif N, Balana C, Bravo-Marques J, Clement PM, Marosi C, Tzuk-Shina T, Nordal RA, Rees J, Lacombe D, Mason WP, Stupp R (2016) Temozolomide chemotherapy versus radiotherapy in high-risk low-grade glioma (EORTC 22033-26033): a randomised, open-label, phase 3 intergroup study. *Lancet Oncol.* doi: 10.1016/S1470-2045(16)30313-8
  17. Jaeckle K, Vogelbaum M, Ballman K, Anderson SK, Giannini C, Aldape K, Cerhan J, Wefel JS, Nordstrom D, Jenkins R, Klein M, Raizer J, Bent M Van Den, Wick W, Flynn P, Dhermain F, Cairncross J, Galanis E, Brown P (2016) CODEL (Alliance-N0577; EORTC-26081/22086; NRG-1071; NCIC-CEC-2): Phase III Randomized Study of RT vs. RT+TMZ vs. TMZ for Newly Diagnosed 1p/19q-Codeleted Anaplastic Oligodendroglial Tumors. Analysis of Patients Treated on the Original Protocol Design (PLO. Advanstar Communications

- 1 18. Shaw EG, Berkey B, Coons SW, Brachman D, Buckner JC, Stelzer KJ, Barger GR, Brown PD,  
2 Gilbert MR, Mehta M (2006) Initial report of Radiation Therapy Oncology Group (RTOG) 9802:  
3 Prospective studies in adult low-grade glioma (LGG). *J Clin Oncol* 24:1500. doi:  
4 10.1200/jco.2006.24.18\_suppl.1500
- 5 19. Van Den Bent MJ, Brandes AA, Taphoorn MJB, Kros JM, Kouwenhoven MCM, Delattre JY,  
6 Bernsen HJJA, Frenay M, Tijssen CC, Grisold W, Sipos L, Enting RH, French PJ, Dinjens WNM,  
7 Vecht CJ, Allgeier A, Lacombe D, Gorlia T, Hoang-Xuan K (2013) Adjuvant procarbazine,  
8 lomustine, and vincristine chemotherapy in newly diagnosed anaplastic oligodendroglioma:  
9 Long-term follow-up of EORTC brain tumor group study 26951. *J Clin Oncol* 31:344–350. doi:  
10 10.1200/JCO.2012.43.2229
- 11 20. Cairncross G, Wang M, Shaw E, Jenkins R, Brachman D, Buckner J, Fink K, Souhami L,  
12 Laperriere N, Curran W, Mehta M (2013) Phase III trial of chemoradiotherapy for anaplastic  
13 oligodendroglioma: Long-term results of RTOG 9402. *J Clin Oncol* 31:337–343. doi:  
14 10.1200/JCO.2012.43.2674
- 15 21. Van Den Bent MJ, Carpentier AF, Brandes AA, Sanson M, Taphoorn MJB, Bernsen HJJA, Frenay  
16 M, Tijssen CC, Grisold W, Sipos L, Haaxma-Reiche H, Kros JM, Van Kouwenhoven MCM, Vecht  
17 CJ, Allgeier A, Lacombe D, Gorlia T (2006) Adjuvant procarbazine, lomustine, and vincristine  
18 improves progression-free survival but not overall survival in newly diagnosed anaplastic  
19 oligodendrogliomas and oligoastrocytomas: A randomized European Organisation for  
20 Research and Treatment of Cancer p. *J Clin Oncol* 24:2715–2722. doi:  
21 10.1200/JCO.2005.04.6078
- 22 22. van den Bent MJ, Baumert B, Erridge SC, Vogelbaum MA, Nowak AK, Sanson M, Brandes AA,  
23 Clement PM, Baurain JF, Mason WP, Wheeler H, Chinot OL, Gill S, Griffin M, Brachman DG,  
24 Taal W, Rudà R, Weller M, McBain C, Reijneveld J, Enting RH, Weber DC, Lesimple T, Clenton S,  
25 Gijtenbeek A, Pascoe S, Herrlinger U, Hau P, Dhermain F, van Heuvel I, Stupp R, Aldape K,  
26 Jenkins RB, Dubbink HJ, Dinjens WNM, Wesseling P, Nuyens S, Golfopoulos V, Gorlia T, Wick  
27 W, Kros JM (2017) Interim results from the CATNON trial (EORTC study 26053-22054) of  
28 treatment with concurrent and adjuvant temozolomide for 1p/19q non-co-deleted anaplastic  
29 glioma: a phase 3, randomised, open-label intergroup study. *Lancet* 6736:1–9. doi:  
30 10.1016/S0140-6736(17)31442-3
- 31 23. Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJB, Belanger K, Brandes  
32 A a, Marosi C, Bogdahn U, Curschmann J, Janzer RC, Ludwin SK, Gorlia T, Allgeier A, Lacombe  
33 D, Cairncross JG, Eisenhauer E, Mirimanoff RO (2005) Radiotherapy plus concomitant and  
34 adjuvant temozolomide for glioblastoma. *N Engl J Med* 352:987–96. doi:  
35 10.1056/NEJMoa043330
- 36 24. Stupp R, Hegi ME, Mason WP, van den Bent MJ, Taphoorn MJ, Janzer RC, Ludwin SK, Allgeier  
37 A, Fisher B, Belanger K, Hau P, Brandes AA, Gijtenbeek J, Marosi C, Vecht CJ, Mokhtari K,  
38 Wesseling P, Villa S, Eisenhauer E, Gorlia T, Weller M, Lacombe D, Cairncross JG, Mirimanoff  
39 RO (2009) Effects of radiotherapy with concomitant and adjuvant temozolomide versus  
40 radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis  
41 of the EORTC-NCIC trial. *Lancet Oncol* 10:459–466. doi: 10.1016/S1470-2045(09)70025-7
- 42 25. Chinot OL, Wick W, Mason W, Henriksson R, Saran F, Nishikawa R, Carpentier AF, Hoang-Xuan  
43 K, Kavan P, Cernea D, Brandes A a, Hilton M, Abrey L, Cloughesy T (2014) Bevacizumab plus  
44 radiotherapy-temozolomide for newly diagnosed glioblastoma. *N Engl J Med* 370:709–22. doi:  
45 10.1056/NEJMoa1308345
- 46 26. Gilbert MR, Dignam JJ, Armstrong TS, Wefel JS, Blumenthal DT, Vogelbaum MA, Colman H,  
47 Chakravarti A, Pugh S, Won M, Jeraj R, Brown PD, Jaeckle KA, Schiff D, Stieber VW, Brachman  
48 DG, Werner-Wasik M, Tremont-Lukats IW, Sulman EP, Aldape KD, Curran WJ, Mehta MP  
49 (2014) A Randomized Trial of Bevacizumab for Newly Diagnosed Glioblastoma. *N Engl J Med*  
50 370:699–708. doi: 10.1056/NEJMoa1308573
- 51 27. Stupp R, Taillibert S, Kanner A, Read W, Steinberg DM, Lhermitte B, Toms S, Idbaih A,  
52 Ahluwalia MS, Fink K, Di Meo F, Lieberman F, Zhu J-J, Stragliotto G, Tran DD, Brem S,  
53 Hottinger AF, Kirson ED, Lavy-Shahaf G, Weinberg U, Kim C-Y, Paek S-H, Nicholas G, Burna J,  
54  
55

- 1 Hirte H, Weller M, Palti Y, Hegi ME, Ram Z (2017) Effect of Tumor-Treating Fields Plus  
2 Maintenance Temozolomide vs Maintenance Temozolomide Alone on Survival in Patients  
3 With Glioblastoma. *Jama* 318:2306. doi: 10.1001/jama.2017.18718
- 4 28. Taphoorn MJB, Dirven L, Kanner AA, Lavy-Shahaf G, Weinberg U, Taillibert S, Toms SA,  
5 Honnorat J, Chen TC, Sroubek J, David C, Idbaih A, Easaw JC, Kim C-Y, Bruna J, Hottinger AF,  
6 Kew Y, Roth P, Desai R, Villano JL, Kirson ED, Ram Z, Stupp R (2018) Influence of Treatment  
7 With Tumor-Treating Fields on Health-Related Quality of Life of Patients With Newly  
8 Diagnosed Glioblastoma: A Secondary Analysis of a Randomized Clinical Trial. *JAMA Oncol* 1–  
9 9. doi: 10.1001/jamaoncol.2017.5082
- 10 29. Roa W, Brasher PM, Bauman G, Anthes M, Bruera E, Chan A, Fisher B, Fulton D, Gulavita S,  
11 Hao C, Husain S, Murtha A, Petruk K, Stewart D, Tai P, Urtasun R, Cairncross JG, Forsyth P  
12 (2004) Abbreviated course of radiation therapy in older patients with glioblastoma  
13 multiforme: a prospective randomized clinical trial. *J Clin Oncol* 22:1583–1588. doi:  
14 10.1200/JCO.2004.06.082
- 15 30. Perry JR, Laperriere N, O'Callaghan CJ, Brandes AA, Menten J, Phillips C, Fay M, Nishikawa R,  
16 Cairncross JG, Roa W, Osoba D, Rossiter JP, Sahgal A, Hirte H, Laigle-Donadey F, Franceschi E,  
17 Chinot O, Golfinoopoulos V, Fariselli L, Wick A, Feuvret L, Back M, Tills M, Winch C, Baumert BG,  
18 Wick W, Ding K, Mason WP (2017) Short-Course Radiation plus Temozolomide in Elderly  
19 Patients with Glioblastoma. *N Engl J Med* 376:1027–1037. doi: 10.1056/NEJMoa1611977
- 20 31. Keime-Guibert F, Chinot O, Taillandier L, Cartalat-Carel S, Frenay M, Kantor G, Guillermo J-S,  
21 Jadaud E, Colin P, Bondiau P-Y, Meneï P, Loiseau H, Bernier V, Honnorat J, Barrié M, Mokhtari  
22 K, Mazon J-J, Bissery A, Delattre J-Y (2007) Radiotherapy for glioblastoma in the elderly. *N*  
23 *Engl J Med* 356:1527–1535. doi: 10.1056/NEJMoa065901
- 24 32. Malmström A, Grønberg BH, Marosi C, Stupp R, Frappaz D, Schultz H, Abacioglu U, Tavelin B,  
25 Lhermitte B, Hegi ME, Rosell J, Henriksson R (2012) Temozolomide versus standard 6-week  
26 radiotherapy versus hypofractionated radiotherapy in patients older than 60 years with  
27 glioblastoma: the Nordic randomised, phase 3 trial. *Lancet Oncol* 13:916–926. doi:  
28 10.1016/S1470-2045(12)70265-6
- 29 33. Socha J, Kepka L, Ghosh S, Roa W, Kumar N, Sinaika V, Matiello J, Lomidze D, de Castro DG,  
30 Hentati D, Fidarova E (2016) Outcome of treatment of recurrent glioblastoma multiforme in  
31 elderly and/or frail patients. *J Neurooncol* 126:493–498. doi: 10.1007/s11060-015-1987-7
- 32 34. Wick W, Platten M, Meisner C, Felsberg J, Tabatabai G, Simon M, Nikkhah G, Papsdorf K,  
33 Steinbach JP, Sabel M, Combs SE, Vesper J, Braun C, Meixensberger J, Ketter R, Mayer-  
34 Steinacker R, Reifenberger G, Weller M (2012) Temozolomide chemotherapy alone versus  
35 radiotherapy alone for malignant astrocytoma in the elderly: The NOA-08 randomised, phase  
36 3 trial. *Lancet Oncol* 13:707–715. doi: 10.1016/S1470-2045(12)70164-X
- 37 35. Suchorska B, Weller M, Tabatabai G, Senft C, Hau P, Sabel MC, Herrlinger U, Ketter R, Schlegel  
38 U, Marosi C, Reifenberger G, Wick W, Tonn JC, Wirsching H-G (2016) Complete resection of  
39 contrast-enhancing tumor volume is associated with improved survival in recurrent  
40 glioblastoma—results from the DIRECTOR trial. *Neuro Oncol* 18:nov326. doi:  
41 10.1093/neuonc/nov326
- 42 36. Ringel F, Pape H, Sabel M, Krex D, Bock HC, Misch M, Weyerbrock A, Westermaier T, Senft C,  
43 Schucht P, Meyer B, Simon M (2016) Clinical benefit from resection of recurrent  
44 glioblastomas: results of a multicenter study including 503 patients with recurrent  
45 glioblastomas undergoing surgical resection. *Neuro Oncol* 18:96–104. doi:  
46 10.1093/neuonc/nov145
- 47 37. Mandl ES, Dirven CMF, Buis DR, Postma TJ, Vandertop WP (2008) Repeated surgery for  
48 glioblastoma multiforme: only in combination with other salvage therapy. *Surg Neurol*  
49 69:506–509. doi: 10.1016/j.surneu.2007.03.043
- 50 38. Combs SE, Kessel KA, Hesse J, Straube C, Zimmer C, Schmidt-Graf F, Schlegel J, Gempt J, Meyer  
51 B (2018) Moving Second Courses of Radiotherapy Forward: Early Re-Irradiation After Surgical  
52 Resection for Recurrent Gliomas Improves Efficacy With Excellent Tolerability. *Neurosurgery*  
53 0:1–8. doi: 10.1093/neuros/nyx629  
54  
55

- 1 39. Chun SJ, Park SH, Park CK, Kim JW, Kim TM, Choi SH, Lee ST, Kim IH (2018) Survival gain with  
2 re-Op/RT for recurred high-grade gliomas depends upon risk groups. *Radiother Oncol*  
3 128:254–259. doi: 10.1016/j.radonc.2018.05.024
- 4 40. Straube C, Scherb H, Gempt J, Kirschke J, Zimmer C, Schmidt-Graf F, Meyer B, Combs SE  
5 (2018) Adjuvant stereotactic fractionated radiotherapy to the resection cavity in recurrent  
6 glioblastoma – the GlioCave study (NOA 17 – ARO 2016/3 – DTKK ROG trial). *BMC Cancer*  
7 18:15. doi: 10.1186/s12885-017-3928-7
- 8 41. Straube C, Elpula G, Gempt J, Gerhardt J, Bette S, Zimmer C, Schmidt-Graf F, Meyer B, Combs  
9 SE (2017) Re-irradiation after gross total resection of recurrent glioblastoma. *Strahlentherapie*  
10 *und Onkol.* doi: 10.1007/s00066-017-1161-6
- 11 42. Nava F, Tramacere I, Fittipaldo A, Bruzzone MG, Dimeco F, Fariselli L, Finocchiaro G, Pollo B,  
12 Salmaggi A, Silvani A, Farinotti M, Filippini G (2014) Survival effect of first- and second-line  
13 treatments for patients with primary glioblastoma: A cohort study from a prospective registry,  
14 1997-2010. *Neuro Oncol* 16:719–727. doi: 10.1093/neuonc/not316
- 15 43. Stupp R, Wong ET, Kanner AA, Steinberg D, Engelhard H, Heidecke V, Kirson ED, Taillibert S,  
16 Liebermann F, Dbalý V, Ram Z, Villano JL, Rainov N, Weinberg U, Schiff D, Kunschner L, Raizer  
17 J, Honnorat J, Sloan A, Malkin M, Landolfi JC, Payer F, Mehdorn M, Weil RJ, Pannullo SC,  
18 Westphal M, Smrcka M, Chin L, Kostron H, Hofer S, Bruce J, Cosgrove R, Paleologous N, Palti Y,  
19 Gutin PH (2012) NovoTTF-100A versus physician’s choice chemotherapy in recurrent  
20 glioblastoma: A randomised phase III trial of a novel treatment modality. *Eur J Cancer*  
21 48:2192–2202. doi: 10.1016/j.ejca.2012.04.011
- 22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55




Figure 1. Treatment algorithm in LGGs. \*Clinical high-risk features: subtotal resection and/or age  $\geq 40$  years. \*Molecular high-risk: IDHwt and/or MGMT not methylated.

PEER REVIEW COPY  
Journal of Neurosurgical Sciences

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55




Figure 2: Adjuvant treatment algorithm for high-grade gliomas. Solid lines represent 1<sup>st</sup> choice recommendation, dashed lines represent 2<sup>nd</sup> choice recommendations.

PEER REVIEW COPY  
Journal of Neurosurgical Sciences

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55




Figure 3: A case of a GBM patient treated in our hospital. A) contrast enhanced T1 sequence with overlying FET-PET image before resection. B) contrast enhanced T1 sequence after resection. There is a small remnant at the rostral edge of the resection cavity. C) Planning CT, PTV (red) resection cavity (yellow). D) Dose distribution of the RT-plan.






Figure 4. Adjuvant treatment in GBM. Solid lines represent the recommended strategies, whereas dashed lines are second and spotted lines are third choice options.

PEER REVIEWED  
Journal of Neurological Sciences

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55