

1
2 **ARTD1 in myeloid cells controls the IL-12/18-IFN- γ axis in a model of sterile**
3 **sepsis, chronic bacterial infection and cancer**

4
5
6 Friedrich A. Kunze^{1,2}, Michael Bauer^{3,4}, Juliana Komuczki^{2,5}, Margit Lanzinger^{2,5}, Kapila Gunasekera¹,
7 Ann-Katrin Hopp^{1,2}, Mareike Lehmann^{1,6}, Burkhard Becher⁵, Anne Müller³ and Michael O. Hottiger^{1,*}

8
9 ¹Department of Molecular Mechanisms of Disease, University of Zurich, Winterthurerstrasse 190, 8057
10 Zurich, Switzerland,

11 ²Molecular Life Science PhD Program of the Life Science Zurich Graduate School, Winterthurerstrasse
12 190, 8057 Zurich, Switzerland,

13 ³Institute of Molecular Cancer Research, University of Zurich, Winterthurerstrasse 190, 8057 Zurich,
14 Switzerland,

15 ⁴Cancer Biology PhD Program of the Life Science Zurich Graduate School, Winterthurerstrasse 190, 8057
16 Zurich, Switzerland,

17 ⁵Institute of Experimental Immunology, University of Zurich, Winterthurerstrasse 190, 8057 Zurich,
18 Switzerland,

19 ⁶Comprehensive Pneumology Center, Ludwig Maximilian University, University Hospital Grosshadern,
20 and Helmholtz Zentrum München, Max-Lebsche-Platz 31, 81377 Munich, Germany

21
22
23
24 Short title: Macrophage ARTD1 controls the IL-12/18-IFN γ axis

25 Key words: ADP-ribosylation, ARTD1/PARP1, LPS, IFN gamma, Helicobacter pylori, cancer, MC-38,

26
27
28 *Corresponding author: michael.hottiger@dmmmd.uzh.ch

29
30

31
32 **ABSTRACT**
33 **Mice deficient for ADP-ribosyltransferase diphteriatoxin-like 1 (ARTD1) are protected against**
34 **microbially induced inflammation. To address the contribution of ARTD1 to inflammation**
35 **specifically in myeloid cells, we generated an *Artd1* ^{Δ Myel} mouse strain with conditional ARTD1**
36 **deficiency in myeloid lineages and examined the strain in three disease models. We found that**
37 **ARTD1, but not its enzymatic activity, enhanced the transcriptional activation of distinct LPS-**
38 **induced genes that included IL-12, TNF- α and IL-6 in primary bone-marrow derived macrophages**
39 **and LPS-induced IL-12/IFN- γ signaling in *Artd1* ^{Δ Myel} mice. The loss of *Artd1* in myeloid cells also**
40 **reduced the T_H1 response to *Helicobacter pylori* and impaired immune control of the bacteria.**
41 **Furthermore, *Artd1* ^{Δ Myel} mice failed to control tumor growth in a subcutaneous MC-38 model of colon**
42 **cancer, which could be attributed to reduced T_H1 and CD8 responses. Together, these data provide**
43 **strong evidence for a cell-intrinsic role of ARTD1 in myeloid cells that is independent of its enzymatic**
44 **activity and promotes type I immunity by promoting IL-12/18 expression.**

45
46

47 INTRODUCTION

48 Inflammation is orchestrated by various cell types that integrate diverse stimuli for the generation of a
49 specific immune response to kill invading pathogens and/or regenerate damaged tissue (1). Pathogen
50 recognition receptors (PRRs) initiate innate immune responses and are expressed on macrophages and
51 dendritic cells, but also epithelial and endothelial cells and fibroblasts (2). PRRs recognize conserved
52 pathogen associated molecular patterns (PAMPs) such as bacterial and fungal cell wall components (i.e.
53 lipopolysaccharides, LPS) or danger associated molecular patterns (DAMPs) (1). The recognition of
54 PAMPs and DAMPs, often by the same receptors, activates intracellular signaling pathways, which result
55 in the expression of pro-inflammatory cytokines such as IL-12, IL-23, IL-6 and IL-1 β (3). IL-12 produced
56 by antigen-presenting cells (APCs) in response to microbial infection activates NK, CD4 and CD8 T cells
57 and promotes T_H1 differentiation and interferon - γ (IFN- γ) production (1).

58 Septic shock is the most common cause of death in intensive care units and is usually the result of
59 a systemic Gram-negative bacterial infection resulting in hypotension and failure of vital organs, in
60 particular the liver, kidney and heart (4). The bacterial membrane component lipopolysaccharide (LPS),
61 when injected into animals, causes a shock-like state that can lead to death. Mechanistically, LPS activates
62 the NF- κ B/Rel family of transcription factors, which trans-activate critical genes involved in the
63 pathogenesis of septic shock mediated by excessive T_H1/T_H17 immune responses (5). LPS binding to TLR4
64 induces the activation of several primary response genes such as *Ifnb1* and *Ccl5* (6). We and others have
65 reported that ADP-ribosyltransferase diphtheriatoxin-like 1 (ARTD1) promotes the NF- κ B-dependent
66 expression of pro-inflammatory cytokines such as IL-1 β , IL-6, TNF- α and CXCL-2, and of adhesion
67 molecules such as ICAM, VCAM or E-Selectin in LPS- or TNF- α -stimulated macrophages and fibroblasts
68 (7-9). ARTD1 belongs to the family of intracellular diphtheria toxin-like ADP-ribosyltransferases that
69 covalently attach ADP-ribose (ADPr) to amino acid residues of target proteins using nicotinamide adenine
70 dinucleotide (NAD⁺) as substrate. This process is called protein ADP-ribosylation and represents an ancient
71 posttranslational modification (PTM) found in a wide range of species (10). Protein ADP-ribosylation can
72 either affect the enzymatic activity of the modified protein or the interaction with nucleic acids and with
73 other proteins (11). ARTD1 is best known for its role during the DNA damage response (DDR), particularly
74 the base excision repair (BER) and double strand break (DSB) repair pathways (12). The concept that the
75 inhibition of the DNA-repair machinery could sensitize tumor cells to conventional DNA-damage-based
76 therapies such as chemo- or radiotherapy has led to the development of small-molecule ADP-ribosylation
77 inhibitors (i.e. PARP inhibitors; PARPi) (13). Today, FDA-approved PARPi's are clinically used to treat
78 breast and ovarian cancer patients. In addition, these compounds have also been reported to dampen various
79 types of inflammation, potentially contributing to their anti-tumor activity (13). Beneficial effects of
80 PARPi's have been reported in numerous inflammatory disease models confirming that ADP-ribosylation
81 broadly regulates inflammatory processes (14). ARTD1 function during transcriptional gene regulation in

82 various cells types is further supported by the fact that ARTD1 knock-out mice are resistant to a high dose
83 of LPS in an endotoxic shock model (15), suggesting a dominant role of ARTD1 during inflammatory
84 signaling. However, it is not clear which cell type(s) are contributing to this phenotype.

85 We have shown previously that the PARPi PJ34 prevents and cures *Helicobacter pylori* (*H. pylori*)-
86 associated, T cell-driven immunopathology that precedes gastric cancer development. *H. pylori* causes a
87 persistent mucosa-associated but non-invasive infection that is characterized by either regulatory T cell
88 (Treg)- or T_{H1}/T_{H17} immune responses (16). *H. pylori*-infected mice exhibit a striking effector T cell
89 infiltrate that limits the bacterial burden without clearing *H. pylori* completely (17, 18). Rather, the large
90 quantities of IFN- γ produced by *H. pylori*-specific T_{H1} cells are believed to directly cause pre-malignant
91 lesions, i.e. epithelial hyperplasia and intestinal metaplasia that precede the development of gastric cancer
92 (17). PJ34 exerts its anti-inflammatory effects by impairing T cell priming and T_{H1} polarization in the gut-
93 draining mesenteric lymph nodes (19). Our data indicated that PJ34 directly suppresses T cell effector
94 functions by blocking the IFN- γ production of mesenteric lymph node T cells. However, it was not clear
95 whether ARTD1 or another ARTD family member contributed to these observations and to which extent
96 ARTD1 regulates a specific cell type that contributed to the T_{H1} polarization. ARTD1's nuclear localization
97 in LPS-stimulated macrophages hinted at its possible role in mediating certain nuclear effects elicited by
98 LPS, a possibility also suggested by previous reports describing a transcriptional co-regulator activity of
99 ARTD1 (20-22).

100 Tumors are commonly infiltrated by heterogeneous populations of myeloid cells that are of
101 monocytic and granulocytic origin, and have considerable phenotypic plasticity with both positive and
102 negative effects on tumor growth and metastasis (23, 24). The balance between anti-tumor and pro-tumor
103 functions can depend on the polarization state, the interaction with the tumor microenvironment and/or the
104 tumor type (25). Subcutaneously growing, syngeneic MC-38 mouse colon adenocarcinoma cells have been
105 used extensively to study the role of macrophages in anti-tumor immunity (26). We have reported recently
106 that MC-38 cells express functional hypoxic (HIF-1 α) and inflammatory (p65/RelA) signaling pathways
107 (27). In contrast to myeloid cells, HIF-1 α levels remained unaffected in MC-38 cells treated with LPS, and
108 hypoxia failed to induce NF- κ B. The corresponding regulation of canonical HIF and NF- κ B target genes
109 confirmed these results (27). To which extent ARTD1 in the stroma contributes to the observed effects
110 remains to be uncovered in this model.

111 To address the role of ARTD1 in myeloid cells during inflammatory and anti-cancer immune
112 responses, we generated a mouse strain with a Cre-mediated deletion of *Artd1* specifically in myeloid cells
113 (*Artd1* ^{Δ Myel}). We observed that ARTD1 expression in murine macrophages regulates a specific set of genes
114 in a manner that is independent of its enzymatic activity, and that *Artd1* deficiency reduces LPS-induced
115 IL-12/IFN- γ signaling in an acute *in vivo* model of sterile sepsis. Moreover, in the above mentioned bacterial

116 infection model using *H. pylori* as infectious agent, *Artd1*^{ΔMyel} mice exhibited reduced IFN-γ expression in
 117 gastric CD4⁺ T cells and failed to control the bacteria. Finally, MC-38 tumor growth was strongly increased
 118 in *Artd1*^{ΔMyel} mice relative to their wild type littermates due to reduced activation of CD4⁺ and CD8⁺ T cells.
 119 Taken together, we provide evidence that ARTD1 in macrophages controls T_H1 responses through
 120 regulation of the IL-12/IFN-γ axis and this activity does not require the ADP-ribosylating function of the
 121 protein.

122

123 **Material and Methods**

124 **Animals and Animal Experiments**

125 The conditional targeting of the *Artd1* allele in ESCs as well as the generation of the *Artd1*^{lox/lox} mice was
 126 performed by Polygene as described (28). Briefly, the vector from EUCOMM (targeting project 45261) was
 127 electroporated in C57BL/6 ES cells and analyzed by PCR (*Artd1* localization 3' fwd:
 128 CACTGAACTGTCTCCTTAGCCAACCTCTGC, rev: GGAAGCTTCGGAATAGGAACTTCGGTTCC;
 129 *Artd1* localization 5' fwd: CTAGGATTCTGTGTCTTGACCATGCACTTG, rev:
 130 CGTATAGCATAACATTATACGAAGTTATGTCGAG). Correctly integrated ES cell clones were injected
 131 into blastocysts resulting in chimeric mice. Subsequent mating of the chimeric mice with the Flp-deleter
 132 mice resulted in the Flp-mediated deletion of the gene trap cassette and leaving only the loxP flanked exon
 133 4 of *Artd1* (Genotyping primer fwd: GCTTCTACTACCTCCCAAGAAAGAGCG, rev:
 134 GGCTTTAGTGTGGCAACTTATCCC). To generate whole-body *Artd1* knock-out mice (*Artd1*^{del/del}) the
 135 generated *Artd1*^{lox/lox} mouse strain was crossed to the CMV-Cre deleter strain (provided by R. Santoro)
 136 (Genotyping primer Deleted *Artd1* fwd: GCTTCTACTACCTCCCAAGAAAGAGCG, rev:
 137 CCTCTGCTGCGTGACTAAGGC). Myeloid specific *Artd1* knock-out mice (*Artd1*^{ΔMyel}) were generated
 138 by crossing *Artd1*^{lox/lox} mice to *LyzM*-Cre (The Jackson Laboratory, strain 004781) mice. For all
 139 experiments 6-12-week-old, age and sex matched mice were used. The sex ratio male:female among the
 140 animals used in the experiments was 1.01. LPS (*Escherichia coli* O111:B4, Sigma-Aldrich) was injected
 141 i.p. at a concentration of 4 mg per kg bodyweight.

142 *Helicobacter pylori* infections were performed as described (29). Briefly, mice were infected orally
 143 on 2 consecutive days with 10⁸ colony-forming units (CFUs) of *H. pylori* PMSS1 and analyzed at 1 months
 144 p.i.. The subcutaneous MC-38 tumor model was performed as described earlier (30). Briefly, colon
 145 adenocarcinoma cells (MC-38, 0.5x10⁶ cells in 100 μl phosphate buffered saline) were injected
 146 subcutaneously (s.c.) into both flanks and tumor progression was determined by caliper measurements every
 147 second day. After two weeks, mice were sacrificed, tumors were weighed and the volume was calculated
 148 using the formula (a² × b)/2, where a constitutes the shorter and b the longer dimension of the tumor.

149 All animals were housed under pathogen free conditions at the University of Zurich. All animal
 150 experiments were carried out in accordance with the Swiss and EU ethical guidelines and have been

151 approved by the local animal experimentation committee of the Canton Zurich under licenses 207/2015,
152 266/2014 and ZH140/2017.

153
154 **Cell culture and reagents**
155 Murine bone marrow derived macrophages (BMDM) were generated as previously described (31) and
156 maintained in RPMI1640 supplemented with 10 % fetal calf serum, 5 % penicillin/streptomycin and 20
157 ng/ml recombinant murine M-CSF (Preprotech). Thioglycolate (Sigma-Aldrich) elicited murine peritoneal
158 macrophages were isolated as described(32). Cell culture grade LPS (*Escherichia coli* O111:B4) was
159 purchased from Sigma-Aldrich and recombinant murine interferon gamma from Preprotech. The PARP
160 inhibitors PJ34 and Olaparib were purchased from Selleckchem. MC-38 cells were cultured as described
161 (30).

162
163 **Multiplex-bead array and ELISA**
164 Whole blood serum of mice and cell culture supernatant of BMDM was analyzed by ELISA (RnD-Systems)
165 and by ProcartaPlex Immunoassay (ThermoFisher) according to the protocol and measured on a Bio-Plex
166 instrument (Bio-Rad).

167
168 **Flow cytometry**
169 Flow cytometrical analyses of spleens, tumors and stomach tissue were performed as described earlier (29).
170 Briefly, tissues were cut into pieces and digested with 15 mM HEPES, 500 U/ml of type IV collagenase
171 (Sigma-Aldrich) and 0.05 mg ml⁻¹ DNase I in RPMI-1640 medium with 10% fetal bovine serum and 100
172 U ml⁻¹penicillin/streptomycin shaking at 37°C for 30-60 min. Subsequently, the samples were pushed
173 through a cell strainer (70µm) using a syringe plunger and red blood cell lysis was performed with a
174 hypotonic buffer (ACK buffer, Gibco). Intracellular cytokine staining was performed as described earlier
175 (29, 33). Briefly, cells were incubated in RPMI-1640 medium with 10% fetal bovine serum, 100 U
176 ml⁻¹penicillin/streptomycin and Brefeldin A (BD) shaking at 37°C for 4 hrs. In addition, gastric and tumor
177 suspensions were re-stimulated with PMA (50 ng/mL), ionomycin (500 ng/mL) or a MC-38-specific tumor
178 peptide. Single cell suspensions were stained with the respective antibodies and analyzed on LSR Fortessa
179 and FACSymphony (BD Bioscience). Cell sorting was performed analogously but analyzed by FACSARIA
180 III 5 L.

181
182 **RNA sequencing and bioinformatic analysis**
183 For the library preparation, the quality of the isolated RNA was determined with a Qubit® (1.0) Fluorometer
184 (Life Technologies) and a Bioanalyzer 2100 (Agilent). Only those samples with a 260 nm/280 nm ratio
185 between 1.8–2.1 and a 28S/18S ratio within 1.5–2 were further processed. The TruSeq RNA Sample Prep

186 Kit v2 (Illumina) was used in the subsequent steps. Briefly, total RNA samples (100-1000 ng) were poly-A
 187 enriched and then reverse-transcribed into double-stranded cDNA. The cDNA samples were fragmented,
 188 end-repaired and polyadenylated before ligation of TruSeq adapters containing the index for multiplexing.
 189 Fragments containing TruSeq adapters on both ends were selectively enriched with PCR. The quality and
 190 quantity of the enriched libraries were validated using Qubit® (1.0) Fluorometer and the Caliper GX
 191 LabChip® GX (Caliper Life Sciences). The product is a smear with an average fragment size of
 192 approximately 260 bp. Library DNA concentrations were normalized to 10 nM in Tris-Cl 10 mM, pH8.5
 193 with 0.1% Tween 20.

194 Cluster generation and sequencing were performed using the TruSeq PE Cluster Kit HS4000, or the
 195 TruSeq SR Cluster Kit HS4000 (Illumina) was used for cluster generation using 2 nM of pooled normalized
 196 libraries on the cBOT. Sequencing were performed on the Illumina HiSeq 4000 paired end at 2 x 150 bp or
 197 single end 125 bp using the TruSeq SBS Kit HS4000 (Illumina).

198 Reads were quality-checked with FastQC. Sequencing adapters were removed with Trimmomatic
 199 (34). Subsequently, reads of at least 20 base length, and with an overall average phred quality score greater
 200 than 10 were aligned to the reference genome and transcriptome of *Mus Musculus* (FASTA and GTF files,
 201 respectively, downloaded from GRCm38) with STAR v2.5.1(35) with default settings for single end reads.

202 Distribution of the reads across genomic isoform expression was quantified using the R package
 203 GenomicRanges (36) from Bioconductor Version 3.0. Differentially expressed genes were identified using
 204 the R package edgeR (37) from Bioconductor Version 3.0. A gene is marked as differentially expressed
 205 (DE) if it possesses the following characteristics: (1) at least 10 counts in at least half of the samples in one
 206 group; (2) $p \leq 0.05$; (3) fold change ≥ 1.5 .

207

208 **RNA extraction and quantitative real time PCR (qPCR) analysis.**

209 RNA isolation and qPCR were performed as described(38). Briefly, RNA extraction was performed with
 210 the NucleoSpin RNA II kit (Macherey-Nagel). RNA was quantified with a NanoDrop and reverse
 211 transcribed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems) according to
 212 the supplier's protocol. qPCRs were performed with the SYBR green KAPA SYBR® FAST (Sigma-
 213 Aldrich) and a Rotor-Gene Q 2plex HRM system (Qiagen). The following genes were quantified with the
 214 respective primer pairs: *Il12b* (fwd: GAAGTTCAACATCAAGAGCAGTAG, rev:
 215 AGGGAGAAGTAGGAATGGGG), *Cxcl3* (fwd: ACCCAGACAGAAGTCATAGCC, rev:
 216 ACACATCCAGACACCGTTGG), *Cd55* (fwd: CGGGCAAGGTCTCTTCTACC, rev:
 217 CAGTCTCCGCGTACAGTTGG), *Il23a* (fwd: ACCAGCGGGACATATGAATCT, rev:
 218 AGACCTTGCGGATCCTTTG), *Itgax* (fwd: GCAGACACTGAGTGATGCCA, rev:
 219 TCGGAGGTCACCTAGTTGGG), *Socs1* (fwd: CCGCCAGATGAGCCCAC, rev:
 220 GGTTGCGTGCTACCATCCTA), *Il18* (fwd: ATGCTTTCTGGACTCCTGCC, rev:

221 ATTGTTCTGGGCAAGAGG), *Il6* (fwd: CCAATTTCCAATGCTCTCCT, rev:
222 ACCACAGTGAGGAATGTCCA), *Tnfa* (fwd: GTCGTAGCAAACCACCAAGTGG, rev:
223 GAGATAGCAAATCGGCTGACGG), *Ifng* (fwd: CCTTCTTCAGCAACAGCAAGGCGA, rev:
224 TGGACCTGTGGGTTGTTGACCTCA). The relative amounts of each mRNA were normalized to the
225 *Rps12* (fwd: GAAGCTGCCAAAGCCTTAGA, rev: AACTGCAACCAACCACCTTC).

226

227 **Westernblot**

228 Western blotting was performed as described (39). For western blot analysis, proteins were separated by
229 SDS-PAGE, and bands were visualized using the Odyssey infrared imaging system (LI-COR). Antibodies
230 used for western blotting were anti-PARP-1 (1:1000, Santa Cruz sc-7150) and anti-Tubulin (1:10,000,
231 Sigma, #T6199).

232

233

234 **RESULTS**

235

236 **ARTD1 regulates IL-12/IL-18 gene expression in BMDMs in an enzymatically independent manner**

237 *Artd1*-deficient mice are protected against microbially induced inflammation. However, the relative
 238 contribution of ARTD1 deficiency in specific cell types to the overall phenotype is currently not known.
 239 Studies with whole-body *Artd1*-deficient mice in which ARTD1 is absent throughout development and in
 240 all cell types have often yielded conflicting results, as ARTD1 is expressed in all examined tissues and cell
 241 types (40). To avoid these complications while examining the specific role of ARTD1 in inflammation, and
 242 to address to which extent ARTD1 in myeloid cells regulates inflammation, we developed a mouse line with
 243 a conditional (“floxed”) exon 4 of *Artd1* (*Artd1^{flox/flox}*) (Fig. S1A). To verify the capacity for complete
 244 inactivation of the recombined floxed *Artd1* allele we first crossed *Artd1^{flox/flox}* mice with transgenic *CMV-*
 245 *Cre* deleter mice to generate a full body *Artd1* knockout (*Artd1^{flox/flox};CMV-cre*), hereafter called *Artd1^{del/del}*
 246 (Fig. S1A-C). The whole body ARTD1-deficient mouse developed normally and was fertile and thus
 247 reproduced observations made with the classical *Parp1^{-/-}* mouse (41). Although western blotting showed
 248 complete loss of ARTD1 in the tested organs (Fig. S1D), ARTD1 loss did not affect organ weight or organ
 249 structure (Fig. S1E and data not shown).

250 As we aimed to investigate the role of ARTD1 in macrophages and other myeloid cells, we first
 251 cultured bone marrow-derived macrophages (BMDMs) from *Artd1^{del/del}* mice and confirmed the genetic
 252 deletion of the *Artd1* gene by qPCR (Fig. S1F). BMDMs from *Artd1^{del/del}* mice and *Artd1^{flox/flox}* littermate
 253 controls were differentiated for 6 days, treated for 4 hrs with LPS/IFN- γ and subjected to RNA sequencing.
 254 LPS/IFN- γ stimulation induced a robust pro-inflammatory gene signature of approx. 2500 genes, (Fig. 1A),
 255 which included many NF- κ B target genes (e.g. *Ifnb1*, *Ccl5* and *Cxcl10*), whereas the expression of many
 256 basally expressed genes was reduced (approx. 3000 genes; Fig. S1G). Of the LPS-induced genes, 701 genes
 257 were found to be strictly dependent on ARTD1 (i.e. induction was reduced in BMDMs from *Artd1^{del/del}*
 258 *mice*), confirming that ARTD1 acts as a transcriptional co-activator of these genes (Fig. 1B, upper panel).
 259 Additionally, we observed that ARTD1 depletion also enhanced the expression of 449 genes (Fig. 1B, lower
 260 panel). Independent qPCR analyses of a defined set of genes belonging to either up- or down-regulated
 261 clusters after ARTD1 depletion confirmed the quantitative RNA-seq analysis (Fig. 1C). Furthermore, gene
 262 ontology (GO) analyses of the ARTD1 up-regulated genes revealed that ARTD1 activates genes regulating
 263 IL-12, IFN- γ and TNF- α production and adaptive immune responses (Fig 1D).

264 To investigate whether ARTD1’s transcriptional regulation of LPS/IFN- γ -induced target genes
 265 depends on its enzymatic activity, the stimulation of BMDMs was repeated in the presence of two ADP-
 266 ribosylation inhibitors (i.e. PARPi) (Fig. S1H). Although RNA-sequencing revealed that treatment of
 267 BMDMs with PARPis affected the expression of 267 LPS/IFN- γ -induced genes (Fig. S1H), the pro-

268 inflammatory genes observed above where not affected. GO analysis of PARPi-sensitive genes revealed no
 269 significantly enriched pathways but pH reduction and cell adhesion pathways only in the first cluster (C1,
 270 n = 95 genes) (Fig. S1I). These data suggest that the enzymatic activity of most likely ARTD1 in
 271 macrophages does not contribute to LPS/IFN- γ -induced innate immune responses under the examined
 272 conditions.

273 To confirm that the observed transcriptional changes translate into changes at the protein level, the
 274 expression levels of differentially regulated cytokines was confirmed by ELISA after treatment of BMDMs
 275 cultured from *Artd1*^{fllox/fllox} and *Artd1*^{del/del} mice for 18 hrs with LPS/IFN- γ (Fig. 1E). The levels of secreted
 276 IL-12p70, IL-18, IL-6 and TNF- α were all significantly reduced in BMDMs from ARTD1-deficient mice
 277 compared to wild-type mice. In addition, we analyzed whether the enzymatic activity of ARTD1 regulates
 278 pro-inflammatory cytokine expression at the protein level (Fig. S1J). ELISA quantification of BMDMs
 279 stimulated with LPS/IFN- γ in the presence or absence of the PARPi's PJ34 or Olaparib revealed no
 280 expression change for the abovementioned cytokines. These data implicate ARTD1 in the transcriptional
 281 control of innate cytokine genes, which is independent of its enzymatic activity, and likely to affect type I
 282 immune responses in inflammation and cancer models.

283
 284 **Myeloid cell-specific deletion of ARTD1 reduces LPS-induced IL-12/IFN- γ signaling in an *in vivo***
 285 **model of sterile sepsis**

286 To explore the myeloid cell-intrinsic role of ARTD1 *in vivo* in models driven by strong T_H1
 287 activation, we crossed the *Artd1*^{fllox/fllox} mice with transgenic mice harboring a *LyzM-Cre* cassette that resulted
 288 in a conditional ARTD1 knockout in lysozyme-expressing myeloid cells (*Artd1*^{fllox/fllox}; *LyzM-cre*), hereafter
 289 called *Artd1* ^{Δ Myel} (Fig. S2A). Mice lacking ARTD1 specifically in this compartment developed normally
 290 and were fertile as their ARTD1-proficient littermates (data not shown). The correct targeting was
 291 confirmed by western blotting for ARTD1 in *in vitro* cultured BMDMs and *ex vivo* in peritoneal
 292 macrophages (Fig. S2B).

293 ARTD1-deficient mice are known to be extremely resistant to LPS-induced endotoxic shock (15).
 294 We first set out to study the innate immune response to sublethal LPS exposure; to this end, 6–12 week old
 295 *Artd1*^{fllox/fllox} and *Artd1* ^{Δ Myel} littermates were intraperitoneally (i.p.) injected with 4 mg/kg body weight of LPS.
 296 Both *Artd1*^{fllox/fllox} and *Artd1* ^{Δ Myel} mice showed the expected clinical symptoms such as fever and lethargy, but
 297 none of the animals died during the 4 h time course of the experiments (data not shown). ELISA-based
 298 quantification of serum IFN- γ , IL-12p70, IL-18, TNF- α and IL-6 levels revealed the strong induction of
 299 these cytokines in *Artd1*^{fllox/fllox} mice upon LPS exposure relative to PBS treatment (Fig. 2A). Interestingly,
 300 IFN- γ levels in serum were significantly reduced in *Artd1* ^{Δ Myel} mice, while the other tested cytokines were
 301 only modestly reduced (Fig. 2A). Analyses of anti-inflammatory or Th2 cytokines revealed that the serum

302 levels of IL-10 and IL-13 were unchanged while the levels of IL-4 were reduced (Fig. S2C), suggesting that
 303 the observed differences in pro-inflammatory cytokine levels were not due to an enhanced anti-
 304 inflammatory or Th2-polarized response. Together these data confirmed the initiation of an innate immune
 305 response with the applied low LPS dose and support the *in vitro* data that ARTD1-deficient macrophages
 306 fail to express IFN- γ -inducing IL-12p70 and IL-18.

307
 308 **The reduced IFN- γ levels in *Artd1* ^{Δ Myel} mice are a consequence of the reduced production of IL-12 and**
 309 **IL-18 by macrophages**

310 Since IL-12 and IL-18 are known to strongly induce IFN- γ gene expression (42), we analyzed the transcript
 311 levels of several pro-inflammatory cytokines in spleen tissue. We found transcript levels of *Ifng*, *Il12b*, *Il18*,
 312 *Tnfa* and *Il6* to be strongly reduced in the spleen of *Artd1* ^{Δ Myel} mice (Fig. 2B), indicating that ARTD1 in
 313 macrophages regulates this inflammatory program *in vivo*. To exclude that the macrophage-specific deletion
 314 of ARTD1 affects the cellular composition of the spleen and thus alters cytokine production, the spleens
 315 from *Artd1*^{fllox/fllox} and *Artd1* ^{Δ Myel} were analyzed macroscopically and their splenocytes mice were analyzed
 316 by flow cytometry using cell type-specific markers. While the spleen size was not overtly altered (data not
 317 shown), the total number of splenocytes was increased in untreated *Artd1* ^{Δ Myel} compared to the *Artd1*^{fllox/fllox}
 318 control mice (Fig. S2D). This was fully attributable to an increase number of B and CD4⁺ T cells, whereas
 319 the frequencies of all other examined cell types, including CD8⁺ T cells, NK cells, and most importantly
 320 macrophages, monocytes and neutrophils, were not altered (Fig. S2E). These data indicate that the lack of
 321 ARTD1 in macrophages neither affected macrophage development nor drastically changed splenic immune
 322 cell composition.

323 IFN- γ is mainly expressed by NK and to a lower extent by T cells during acute inflammation and
 324 the onset of an innate immune response (4). To identify the cell type responsible for the reduced IFN- γ
 325 levels in the spleen (Fig. 2B), we performed intracellular cytokine staining (ICS) for IFN- γ in NK, CD4⁺
 326 and CD8⁺ T cells isolated from the spleens of LPS treated *Artd1*^{fllox/fllox} and *Artd1* ^{Δ Myel} mice (Fig. 2C,D).
 327 Although LPS treatment induced IFN- γ expression in all three analyzed cell types, the frequency of IFN- γ -
 328 expressing NK cells was considerably higher than the frequencies of CD4⁺ and CD8⁺ T cells. Thus, NK
 329 cells appear to be the main source of the differential IFN- γ expression in LPS-treated *Artd1*^{fllox/fllox} and
 330 *Artd1* ^{Δ Myel} mice (Fig. 2C,D). To exclude that NK cells from *Artd1*^{fllox/fllox} and *Artd1* ^{Δ Myel} mice were
 331 differentially responsive to IFN- γ -inducing IL-12/18 signaling, we isolated and stimulated primary NK cells
 332 from both genotypes with IL-12p70 (10 ng/ml) for 18 hrs to quantify their IFN- γ expression by ELISA (Fig.
 333 2E). Stimulation of NK cells from both mouse strains induced IFN- γ expression in an IL-12p70-dependent
 334 manner and to the same extent. The combined results indicate that the reduced IFN- γ levels produced by
 335 NK cells in *Artd1* ^{Δ Myel} mice exposed to LPS are a consequence of the reduced production of IL-12 and IL-

336 18 by macrophages and that ARTD1 is an important transcriptional activator of these genes.

337
 338 **ARTD1 expression in macrophages is required for T_H1-mediated immune control of bacterial**
 339 **infection**

340 Immunocompetent adult C57BL/6 mice infected with the human gastric pathogen *H. pylori* exhibit a
 341 striking T cell infiltration that is dominated by T_H1- and T_H17-polarized CD4⁺ T-cells and limits the bacterial
 342 burden (16). We therefore chose this model to assess the role of myeloid ARTD1 expression in anti-*H.*
 343 *pylori* T_H1 responses and infection control. To this end, we first cultured BMDMs from *Artd1*^{flox/flox} and
 344 *Artd1*^{del/del} littermates with the human *H. pylori* patient isolate PMSS1 for 6 h. Subsequent qPCR analysis
 345 revealed that *H. pylori* robustly induces *Il12b* and *Il1b* gene expression in cultured BMDMs in an -at least
 346 in part- ARTD1-dependent manner (Fig 3A). To investigate the role of ARTD1 expression in myeloid cells
 347 for the development of *H. pylori*-specific T_H1 and T_H17 responses *in vivo*, we infected *Artd1*^{flox/flox} and
 348 *Artd1*^{ΔMyel} mice with 10⁸ CFUs of *H. pylori* PMSS1. *Artd1*^{ΔMyel} mice were colonized at significantly higher
 349 levels than their *Artd1*^{flox/flox} counterparts at four weeks post infection (Fig. 3B), suggesting that ARTD1 in
 350 myeloid cells controls gastric colonization of *H. pylori in vivo*. *H. pylori* induced a strong infiltration of
 351 CD4⁺ T cells into the gastric mucosa that was similar in *Artd1*^{flox/flox} and *Artd1*^{ΔMyel} mice (Fig. 3C and S3).
 352 Interestingly, *H. pylori*-specific T_H1- but not T_H17 responses were dependent on ARTD1 expression in
 353 myeloid cells as assessed by intracellular staining for the signature cytokines IFN-γ and IL-17 (Fig. 3D,E).
 354 The combined results suggest that ARTD1 expression in myeloid cells is specifically required for IL-12-
 355 driven T_H1 polarization and infection control, but not for T_H17 responses or CD4⁺ T-cell recruitment.

356
 357 **Expression of ARTD1 in myeloid cells controls the T-cell-mediated immune control of**
 358 **MC-38 tumors**

359 Understanding the impact of myeloid cells on cancer development is essential in distinguishing and possibly
 360 manipulating positive and negative myeloid effectors. Polarization states of intratumoral myeloid
 361 populations contribute differentially to tumorigenesis (25). Tumor-associated macrophages (TAMs)
 362 infiltrate tumors and can serve as important initiators of T_H1-mediated anti-tumor immunity in some
 363 settings, but have also been described to promote tumor growth in others (43). To investigate a possible
 364 myeloid-specific ARTD1 contribution to anti-tumor immunity, we injected 0.5x10⁶ MC-38 colon
 365 adenocarcinoma cells subcutaneously into the flanks of *Artd1*^{flox/flox} and *Artd1*^{ΔMyel} mice. *Artd1*^{ΔMyel} mice
 366 sustained a significantly increased tumor burden from day 6 onwards relative to their *Artd1*^{flox/flox} littermates,
 367 which was evident until the study end point (Fig. 4A-C). Ly6C⁺ MHCII⁺ monocytes, identified according
 368 to the gating strategy shown in Fig. S3-4, infiltrated the tumors at a similar rate irrespective of ARTD1
 369 status, but expressed less *Il12b* and more *Arg1* as judged by qRT-PCR of FACS-sorted cells if lacking
 370 ARTD1 (Fig. 4D,E). F4/80⁺ macrophage frequencies in the tumor microenvironment were decreased in

371 *Artd1*^{ΔMyel} mice (Fig. 4F), and fewer macrophages expressed TNF- α (Fig. 4F) and *Iil2b* (Fig. 4G) as judged
372 by intracellular staining and qRT-PCR of sorted cells, respectively. As observed for monocytes, *Arg1* gene
373 expression was enhanced in macrophages as a consequence of ARTD1 deficiency (Fig. 4G). The data
374 suggests that ARTD1 potentially affects the differentiation of macrophages from inflammatory monocytes
375 as well as their M1 polarization. Interestingly, the loss of ARTD1 in myeloid cells further resulted in
376 strongly reduced intratumoral CD4⁺ and CD8⁺ T-cell activation, as judged by IFN- γ and TNF- α -specific
377 staining, although the recruitment of both populations to the tumor microenvironment was unaffected (Fig.
378 4H-K). The differences due to ARTD1 deficiency in CD8⁺ T-cell activation could be confirmed by re-
379 stimulation with MC-38 tumor-specific peptide (Fig. 4L,M)

380 The combined results implicate myeloid-intrinsic ARTD1 expression in the generation and/or
381 maintenance of type I immunity in models of LPS challenge, bacterial infection, and tumor growth.

382

383

384 **DISCUSSION**

385 Here, we investigated the role of ARTD1 in models of myeloid cell activation in response to innate immune
 386 stimulation, bacterial infection and MC-38 tumor growth. We show that ARTD1 expression in myeloid cells
 387 controls a transcriptional program that includes T_H1-polarizing cytokines and type I immunity; this function
 388 of ARTD1 appears to be independent of its ADP-ribosylating activity, as inhibitors targeting the enzymatic
 389 function of ARTD1 do not recapitulate the effects of loss of the protein.

390 Various reports have shown that ARTD1 acts as a transcriptional co-factor for NF- κ B and promotes
 391 gene expression by functional cooperation with the transcription machinery in response to pro-inflammatory
 392 stimuli (44). LPS- and TNF- α -induced NF- κ B-dependent gene expression in macrophages and fibroblasts
 393 was enhanced by ARTD1 independently of its enzymatic activity, through the initiation of a mediator
 394 complex that also contains p300 and NF- κ B. In the current study, treatment of BMDMs with ADP-
 395 ribosylation inhibitors (e.g. Olaparib or PJ34) did not affect the expression of LPS/IFN- γ -induced gene
 396 expression, although some non-inflammation-related genes involved in pH reduction and cell adhesion were
 397 susceptible to both inhibitors. In contrast, another report suggested that LPS treatment of macrophages
 398 induced ARTD1's enzymatic activity and nucleosome remodeling at promoters of pro-inflammatory genes,
 399 which directly destabilized histone-DNA interactions and facilitated NF- κ B binding and gene expression
 400 (45). The discrepancies could be explained by methodological differences such as the cell type (RAW267.4
 401 macrophages and primary BMDM or fibroblasts), the source of LPS (*S. enterica* and *E.coli*) or the serum
 402 starvation overnight prior to LPS stimulation.

403 The LPS-induced expression of IL-12/18 in primary macrophages was strongly dependent on
 404 ARTD1 *ex vivo* as well as *in vivo*. The *Il12b* gene is a well-studied example of a gene that requires chromatin
 405 remodeling during inflammation-induced gene expression (46). Studies of the *Il12b* promoter identified
 406 binding sites for various transcription factors including NF- κ B, C/EBP, AP-1 and NFATm (42). However
 407 their binding sites, -30 to -175 bp upstream of the transcription start site, are blocked by nucleosomes and
 408 require nucleosome remodeling prior to transcription (46). Indeed, the SWI-SNF complex that is responsible
 409 for nucleosome positioning is recruited to target genes in LPS-stimulated macrophages (47). Several models
 410 describe the targeting of chromatin remodeling machines to their site of action, i.e. acetylated histones that
 411 target and stabilize the SWI-SNF complex at target loci (48). Very recently, another report confirmed that
 412 small hairpin RNA-mediated knockdown of the endogenous ARTD1 expression resulted in reduced *Il12b*
 413 mRNA expression and *Il12b* promoter activity (49). BMDMs from ARTD1-deficient mice also exhibited
 414 decreased IL-12p40 expression at both mRNA and protein levels.

415 IL-12 connects innate and adaptive immune responses either indirectly via NK cell activation or
 416 directly by activating CD4⁺ and CD8⁺ T cells (5). In this study, we observed a reduced NK cell activation
 417 after LPS administration in *Artd1* ^{Δ Myel} compared to *Artd1*^{lox/lox} mice. Our analyses of the splenic cell
 418 composition of *Artd1* ^{Δ Myel} relative to *Artd1*^{lox/lox} mice revealed that, whereas B and CD4⁺ T cell counts were

419 increased, all other analyzed cell types including monocytes and macrophages were not altered, suggesting
420 that the deletion of ARTD1 in myeloid cells did not decrease the numbers of other cell types and their
421 functionality in this organ. ARTD1-deficient macrophages (or other myeloid cells) failed to activate NK
422 cells via IL-12 expression, ultimately leading to decreased IFN- γ serum levels and thus protecting against
423 the consequences of LPS administration. Several lines of evidence suggest that NK cells might be involved
424 in key functions during sepsis (4). Similar to the observations made in ARTD1-deficient mice, antibody-
425 mediated NK cell depletion *in vivo* protected against LPS-induced shock and significantly decreased IFN- γ
426 cytokine levels (15) During sepsis, NK cells promote and amplify the inflammatory response as a very early
427 and main source of IFN- γ , and thus represent a promising target for novel approaches in sepsis therapy (4).

428 Reduced NK cell activation significantly reduces the kinetics of pathogen clearance during sepsis
429 (4). Thus, by enhancing *Il12b* expression in myeloid cells, ARTD1 contributes to the generation of a potent
430 immune response to pathogens. This was very obvious in the second disease model we investigated.
431 Previous studies characterized the stomach under basal conditions to be a predominantly myeloid cell-
432 controlled organ with little or no lymphocyte immune surveillance (29). Myeloid cells centrally function as
433 initiators of immune responses against pathogens. Our study revealed that ARTD1-deficiency does not
434 affect gastric CD4⁺ T cell recruitment during *H. pylori* infection, but that the loss of ARTD1 results in
435 reduced T_H1 frequencies. Thus, ARTD1 drives pathogen control via IL-12 production and the initiation of
436 potent immune responses.

437 Macrophages play important yet bimodal roles in orchestrating tumor-associated immune responses
438 (43). On the one hand, they are involved in tumor killing and other effector functions, but they can also
439 promote tumor growth by skewing and suppressing T cell responses. We made several observations relating
440 to ARTD1 of macrophages in the MC38 tumor model. On the one hand, we find fewer macrophages in the
441 tumor microenvironment of *Artd1* ^{Δ Myel} mice, whereas monocyte frequencies are unchanged, which indicates
442 that macrophage differentiation is impaired in the absence of ARTD1 expression. On the other hand, those
443 macrophages that do infiltrate the tumors (or differentiate there from their monocyte precursors) express
444 less TNF- α and less IL-12, but more Arginase 1, indicating that they have adopted an M2-polarized state.
445 As a consequence of their lower intratumoral macrophage frequencies, combined with their dysregulated
446 macrophage activation and polarization, *Artd1* ^{Δ Myel} mice fail to generate appropriate T_H1 polarized CD4 and
447 cytotoxic CD8 T-cell responses and therefore cannot control the tumor burden as efficiently as wild type
448 littermates.

449 Taken together we identified ARTD1 expression in myeloid cell types as a critical regulator of pro-
450 inflammatory IL-12/18 cytokine expression. In particular, ARTD1 controls the initiation of potent immune
451 responses to LPS and the elimination of pathogens and tumors *in vivo*.

452

453

454 ACKNOWLEDGMENTS

455 We would like to thank Tobias Suter (University of Zurich) for the helpful discussions and for providing
456 editorial assistance. We thank Maria Domenica Moccia and Giancarlo Russo from the Functional Genomics
457 Center of the University of Zurich for the RNA Sequencing and data analysis. We also thank Raffaella
458 Santoro (University of Zurich) for providing the CMV-Cre mouse. Work in the lab of AM is funded by the
459 Swiss National Science Foundation (grant BSCGIO_157841/1). ADP-ribosylation research in the
460 laboratory of MOH is funded by the Kanton of Zurich and the Swiss National Science Foundation (grant
461 310030_157019).

462

463 AUTHOR CONTRIBUTIONS

464 F.A.K. generated the mice and performed the LPS *in vivo* models, RNA-Seq, ELISA, qPCR, Westernblot
465 analyses. J.K. and M.L performed ICS and immunophenotyping of the LPS models. M.B. performed the *H.*
466 *pylori* and MC-38 studies. K.G. analyzed the RNA sequencing. A.H. assisted with the LPS experiments.
467 M.L. helped in generating the floxed ARTD1 mouse. F.A.K. and M.O.H. prepared the manuscript. M.O.H.
468 and A.M. and B.B. directed and supervised all aspects of the study. All authors critically reviewed the
469 manuscript.

470

471 COMPETING FINANCIAL INTERESTS

472 The authors declare no competing financial interests.

473

474 **References**

- 475 1. Iwasaki, A., and R. Medzhitov. 2015. Control of adaptive immunity by the innate immune system. *Nat.*
476 *Immunol.* 16: 343–353.
- 477 2. Antoine, R., and C. Locht. 1990. Roles of the disulfide bond and the carboxy-terminal region of the S1
478 subunit in the assembly and biosynthesis of pertussis toxin. *Infect. Immun.* 58: 1518–1526.
- 479 3. Simon, N. C., K. Aktories, and J. T. Barbieri. 2014. Novel bacterial ADP-ribosylating toxins: structure
480 and function. *Nat. Rev. Microbiol.* 12: 599–611.
- 481 4. Chiche, L., J.-M. Forel, G. Thomas, C. Farnarier, F. Vely, M. Bléry, L. Papazian, and E. Vivier. 2011.
482 The role of natural killer cells in sepsis. *J. Biomed. Biotechnol.* 2011: 986491–8.
- 483 5. Chaudhry, H., J. Zhou, Y. Zhong, M. M. Ali, F. Mcguire, P. S. Nagarkatti, and M. Nagarkatti. 2013.
484 Role of Cytokines as a Double-edged Sword in Sepsis. *In Vivo* 27: 669–684.
- 485 6. Ramirez-Carrozzi, V. R., A. A. Nazarian, C. C. Li, S. L. Gore, R. Sridharan, A. N. Imbalzano, and S. T.
486 Smale. 2006. Selective and antagonistic functions of SWI/SNF and Mi-2beta nucleosome remodeling
487 complexes during an inflammatory response. *Genes Dev.* 20: 282–296.
- 488 7. Hassa, P. O., S. S. Haenni, M. Elser, and M. O. Hottiger. 2006. Nuclear ADP-Ribosylation Reactions in
489 Mammalian Cells: Where Are We Today and Where Are We Going? *Microbiology and Molecular*
490 *Biology Reviews* 70: 789–829.
- 491 8. Bai, P., and L. Virág. 2012. Role of poly(ADP-ribose) polymerases in the regulation of inflammatory
492 processes. *FEBS Letters* 586: 3771–3777.
- 493 9. Ba, X., and N. J. Garg. 2011. Signaling Mechanism of Poly(ADP-Ribose) Polymerase-1 (PARP-1) in
494 Inflammatory Diseases. *The American Journal of Pathology* 178: 946–955.
- 495 10. Hottiger, M. O., P. O. Hassa, B. Lüscher, H. Schüler, and F. Koch-Nolte. 2010. Toward a unified
496 nomenclature for mammalian ADP-ribosyltransferases. *Trends in Biochemical Sciences* 35: 208–219.
- 497 11. Gupte, R., Z. Liu, and W. L. Kraus. 2017. PARPs and ADP-ribosylation: recent advances linking
498 molecular functions to biological outcomes. *Genes Dev.* 31: 101–126.
- 499 12. Ray Chaudhuri, A., and A. Nussenzweig. 2017. The multifaceted roles of PARP1 in DNA repair and
500 chromatin remodelling. *Nat Rev Mol Cell Biol* 18: 610–621.
- 501 13. Lord, C. J., and A. Ashworth. 2017. PARP inhibitors: Synthetic lethality in the clinic. *Science* 355:
502 1152–1158.
- 503 14. Berger, N. A., V. C. Besson, A. H. Boulares, A. Bürkle, A. Chiarugi, R. S. Clark, N. J. Curtin, S.
504 Cuzzocrea, T. M. Dawson, V. L. Dawson, G. Haskó, L. Liaudet, F. Moroni, P. Pacher, P.
505 Radermacher, A. L. Salzman, S. H. Snyder, F. G. Soriano, R. P. Strosznajder, B. Sumegi, R. A.
506 Swanson, and C. Szabó. 2018. Opportunities for the repurposing of PARP inhibitors for the therapy of
507 non-oncological diseases. *Br. J. Pharmacol.* 175: 192–222.
- 508 15. Oliver, F. J. 1999. Resistance to endotoxic shock as a consequence of defective NF-kappa B activation
509 in poly (ADP-ribose) polymerase-1 deficient mice. *The EMBO Journal* 18: 4446–4454.
- 510 16. Koch, K. N., and A. Müller. 2015. Helicobacter pylori activates the TLR2/NLRP3/caspase-1/IL-18
511 axis to induce regulatory T-cells, establish persistent infection and promote tolerance to allergens. *Gut*
512 *Microbes* 6: 382–387.
- 513 17. Arnold, I. C., J. Y. Lee, M. R. Amieva, A. Roers, R. A. Flavell, T. Sparwasser, and A. Müller. 2011.
514 Tolerance rather than immunity protects from Helicobacter pylori-induced gastric preneoplasia.
515 *Gastroenterology* 140: 199–209.
- 516 18. Harris, P. R., S. W. Wright, C. Serrano, F. Riera, I. Duarte, J. Torres, A. Peña, A. Rollán, P. Viviani,
517 E. Guiraldes, J. M. Schmitz, R. G. Lorenz, L. Novak, L. E. Smythies, and P. D. Smith. 2008.
518 Helicobacter pylori gastritis in children is associated with a regulatory T-cell response.
519 *Gastroenterology* 134: 491–499.
- 520 19. Toller, I. M., M. Altmeyer, E. Kohler, M. O. Hottiger, and A. Müller. 2010. Inhibition of ADP
521 Ribosylation Prevents and Cures Helicobacter-Induced Gastric Preneoplasia. *Cancer Res* 70: 5912–
522 5922.
- 523 20. Hassa, P. O., and M. O. Hottiger. 1999. A role of poly (ADP-ribose) polymerase in NF-kappaB
524 transcriptional activation. *Biol. Chem.* 380: 953–959.
- 525 21. Hassa, P. O., M. Covic, S. Hasan, R. Imhof, and M. O. Hottiger. 2001. The enzymatic and DNA

- 526 binding activity of PARP-1 are not required for NF-kappa B coactivator function. *J. Biol. Chem.* 276:
527 45588–45597.
- 528 22. Hassa, P. O., S. S. Haenni, C. Buerki, N. I. Meier, W. S. Lane, H. Owen, M. Gersbach, R. Imhof, and
529 M. O. Hottiger. 2005. Acetylation of poly(ADP-ribose) polymerase-1 by p300/CREB-binding protein
530 regulates coactivation of NF-kappaB-dependent transcription. *J. Biol. Chem.* 280: 40450–40464.
- 531 23. Piccard, H., R. J. Muschel, and G. Opdenakker. 2012. On the dual roles and polarized phenotypes of
532 neutrophils in tumor development and progression. *Crit. Rev. Oncol. Hematol.* 82: 296–309.
- 533 24. Souto, J. C., L. Vila, and A. Brú. 2011. Polymorphonuclear neutrophils and cancer: intense and
534 sustained neutrophilia as a treatment against solid tumors. *Med Res Rev* 31: 311–363.
- 535 25. Gabrilovich, D. I., S. Ostrand-Rosenberg, and V. Bronte. 2012. Coordinated regulation of myeloid
536 cells by tumours. *Nat. Rev. Immunol.* 12: 253–268.
- 537 26. Roblek, M., M. Calin, M. Schlesinger, D. Stan, R. Zeisig, M. Simionescu, G. Bendas, and L. Borsig.
538 2015. Targeted delivery of CCR2 antagonist to activated pulmonary endothelium prevents metastasis. *J*
539 *Control Release* 220: 341–347.
- 540 27. Müller-Edenborn, K., K. Léger, J. F. Glaus Garzon, C. Oertli, A. Mirsaidi, P. J. Richards, H. Rehrauer,
541 P. Spielmann, D. Hoogewijs, L. Borsig, M. O. Hottiger, and R. H. Wenger. 2015. Hypoxia attenuates
542 the proinflammatory response in colon cancer cells by regulating IκB. *Oncotarget* 6: 20288–20301.
- 543 28. Skarnes, W. C., B. Rosen, A. P. West, M. Koutsourakis, W. Bushell, V. Iyer, A. O. Mujica, M.
544 Thomas, J. Harrow, T. Cox, D. Jackson, J. Severin, P. Biggs, J. Fu, M. Nefedov, P. J. de Jong, A. F.
545 Stewart, and A. Bradley. 2011. A conditional knockout resource for the genome-wide study of mouse
546 gene function. *Nature* 474: 337–342.
- 547 29. Arnold, I. C., X. Zhang, S. Urban, M. Artola-Borán, M. G. Manz, K. M. Ottemann, and A. Müller.
548 2017. NLRP3 Controls the Development of Gastrointestinal CD11b+Dendritic Cells in the Steady
549 State and during Chronic Bacterial Infection. *Cell Rep* 21: 3860–3872.
- 550 30. Lim, S. Y., A. E. Yuzhalin, A. N. Gordon-Weeks, and R. J. Muschel. 2016. Tumor-infiltrating
551 monocytes/macrophages promote tumor invasion and migration by upregulating S100A8 and S100A9
552 expression in cancer cells. *Oncogene* 35: 5735–5745.
- 553 31. Zhang, X., R. Goncalves, and D. M. Mosser. 2008. The isolation and characterization of murine
554 macrophages. *Curr Protoc Immunol* Chapter 14: Unit 14.1.
- 555 32. Fortier, A. H., and L. A. Falk. 2001. Isolation of murine macrophages. *Curr Protoc Immunol* Chapter
556 14: Unit 14.1–14.1.9.
- 557 33. Spath, S., J. Komuczki, M. Hermann, P. Pelczar, F. Mair, B. Schreiner, and B. Becher. 2017.
558 Dysregulation of the Cytokine GM-CSF Induces Spontaneous Phagocyte Invasion and
559 Immunopathology in the Central Nervous System. *Immunity* 46: 245–260.
- 560 34. Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina
561 sequence data. *Bioinformatics* 30: 2114–2120.
- 562 35. Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T.
563 R. Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15–21.
- 564 36. Lawrence, M., W. Huber, H. Pagès, P. Aboyoun, M. Carlson, R. Gentleman, M. T. Morgan, and V. J.
565 Carey. 2013. Software for computing and annotating genomic ranges. *PLoS Comput Biol* 9: e1003118.
- 566 37. Robinson, M. D., D. J. McCarthy, and G. K. Smyth. 2010. edgeR: a Bioconductor package for
567 differential expression analysis of digital gene expression data. *Bioinformatics* 26: 139–140.
- 568 38. Minotti, R., A. Andersson, and M. O. Hottiger. 2015. ARTD1 Suppresses Interleukin 6 Expression by
569 Repressing MLL1-Dependent Histone H3 Trimethylation. *Mol. Cell. Biol.* 35: 3189–3199.
- 570 39. Abplanalp, J., M. Leutert, E. Frugier, K. Nowak, R. Feurer, J. Kato, H. V. A. Kistemaker, D. V.
571 Filippov, J. Moss, A. Caflisch, and M. O. Hottiger. 2017. Proteomic analyses identify ARH3 as a
572 serine mono-ADP-ribosylhydrolase. *Nature Communications* 8: 2055.
- 573 40. Bürkle, A., and L. Virág. 2013. Poly (ADP-ribose): PARadigms and PARadoxes. *Molecular Aspects*
574 *of Medicine*.
- 575 41. Wang, Z. Q., B. Auer, L. Stingl, H. Berghammer, D. Haidacher, M. Schweiger, and E. F. Wagner.
576 1995. Mice lacking ADPRT and poly(ADP-ribosyl)ation develop normally but are susceptible to skin
577 disease. *Genes Dev.* 9: 509–520.

- 578 42. Ma, X., W. Yan, H. Zheng, Q. Du, L. Zhang, Y. Ban, N. Li, and F. Wei. 2015. Regulation of IL-10
579 and IL-12 production and function in macrophages and dendritic cells. *F1000Res* 4.
- 580 43. Mantovani, A., F. Marchesi, A. Malesci, L. Laghi, and P. Allavena. 2017. Tumour-associated
581 macrophages as treatment targets in oncology. *Nat Rev Clin Oncol* 14: 399–416.
- 582 44. Krishnakumar, R., and W. L. Kraus. 2010. The PARP Side of the Nucleus: Molecular Actions,
583 Physiological Outcomes, and Clinical Targets. *Molecular Cell* 39: 8–24.
- 584 45. Hassa, P. O., and M. O. Hottiger. 2002. The functional role of poly(ADP-ribose)polymerase 1 as novel
585 coactivator of NF-kappaB in inflammatory disorders. *Cell. Mol. Life Sci.* 59: 1534–1553.
- 586 46. Weinmann, A. S., D. M. Mitchell, S. Sanjabi, M. N. Bradley, A. Hoffmann, H. C. Liou, and S. T.
587 Smale. 2001. Nucleosome remodeling at the IL-12 p40 promoter is a TLR-dependent, Rel-independent
588 event. *Nat. Immunol.* 2: 51–57.
- 589 47. Lai, D., M. Wan, J. Wu, P. Preston-Hurlburt, R. Kushwaha, T. Grundström, A. N. Imbalzano, and T.
590 Chi. 2009. Induction of TLR4-target genes entails calcium/calmodulin-dependent regulation of
591 chromatin remodeling. *PNAS* 106: 1169–1174.
- 592 48. Becker, P. B., and W. Hörz. 2002. ATP-dependent nucleosome remodeling. *Annu. Rev. Biochem.* 71:
593 247–273.
- 594 49. Zhao, Q., Q. Du, F. Wei, J. Xie, and X. Ma. 2017. An Infectious Disease-Associated II12b
595 Polymorphism Regulates IL-12/23 p40 Transcription Involving Poly(ADP-Ribose) Polymerase 1. *J*
596 *Immunol* 198: 2935–2942.
- 597
- 598
- 599

600 **Figure legends:**601 **Figure 1: ARTD1 regulates IL-12 gene expression in BMDMs independent of its enzymatic activity.**

602 (A) RNA-sequencing of total RNA extracted from *Artd1^{flx/flx}* (floX) and *Artd1^{del/del}* (del) BMDMs either left
 603 untreated or treated with 10 ng/ml LPS and 2 ng/ml INF- γ for 4 h. The expression levels of all LPS/INF- γ -
 604 induced genes (approx. 2500 genes) (fold-change ≤ 2 , $P < 0.05$) were clustered. (B) Up-regulated genes
 605 identified in A) were clustered to identify ARTD1 co-activating (701 genes) and co-repressing (449 genes)
 606 function. (C) Quantitative Real-time PCR analysis of selective ARTD1 co-activated and co repressed genes
 607 in *Artd1^{flx/flx}* and *Artd1^{del/del}* BMDMs identified in B. Data presented as mean + SD of three biological
 608 replicates. *t*-Test: * $P < 0.05$, ** $P < 0.01$. (D) Gene enrichment analysis of ARTD1 co-activated and ARTD1
 609 co repressed genes (left and right panel, identified in B) in *Artd1^{flx/flx}* and *Artd1^{del/del}* BMDMs. (E) ELISA
 610 of cell culture supernatants for the quantification of the indicated cytokines produced by *Artd1^{flx/flx}* and
 611 *Artd1^{del/del}* BMDMs respectively. Cells were stimulated with 10 ng/ml LPS and 2 ng/ml INF- γ for 18 h. Data
 612 presented as mean concentration + SD of three independent experiments. *t*-Test: * $P < 0.05$, ** $P < 0.01$.

613
 614 **Figure 2: Myeloid-specific depletion of ARTD1 reduces the LPS-induced IL-12/INF- γ signaling *in***
 615 ***vivo***

616 (A) *Artd1^{flx/flx}* and *Artd1 ^{Δ Myel}* mice were intraperitoneally injected with PBS or 4 mg/kg LPS. After 4 h
 617 whole blood serum was collected and the selected cytokine levels were quantified by multiplex-bead array.
 618 Data shown as mean \pm SEM of three independent experiments. *t*-Test: * $P < 0.05$. (B) *Artd1^{flx/flx}* and
 619 *Artd1 ^{Δ Myel}* mice were intraperitoneally injected with PBS or 4 mg/kg LPS. After 4 h, RNA of total spleen
 620 tissue was isolated and gene expression levels quantified by qPCR. Data shown as mean \pm SEM of three
 621 independent experiments. *t*-Test: * $P < 0.05$, ** $P < 0.01$. (C, D) Intracellular cytokine staining of splenocytes
 622 from *Artd1^{flx/flx}* and *Artd1 ^{Δ Myel}* mice intraperitoneally injected with either PBS or 4 mg/kg LPS. After 4 h,
 623 intracellular INF- γ was detected by flow cytometry and the percentage of positively stained NK-, CD4⁺ T-
 624 and CD8⁺ T cells were quantified. Data shown as mean \pm SEM of two independent experiments. *t*-Test: * P
 625 < 0.05 . (E) ELISA for INF- γ , secreted from MACS-sorted splenic NK cells that were isolated from of
 626 *Artd1^{flx/flx}* and *Artd1 ^{Δ Myel}* mice. NK cells were stimulated with recombinant murine 10 ng/ml IL-12p70 for
 627 18 hrs and cell culture supernatant was analyzed.

628
 629 **Figure 3: ARTD1 in myeloid cells controls *Helicobacter* clearance and anti-*Helicobacter* Th1**
 630 **responses**

631 (A) qRT-PCR analysis of *Il12b* and *Il1b* expression in *Artd1^{flx/flx}* and *Artd1^{del/del}* BMDMs. Cells were either
 632 left untreated or incubated with *H. pylori* (MOI 50) for 6 h. Data are presented as median \pm SEM of three
 633 technical replicates; one representative experiment of two is shown. *t*-Test: * $P < 0.05$, ** $P < 0.01$. (B-E)
 634 Mice of the indicated genotypes were orally infected with *H. pylori*. Gastric colonization as assessed by

635 plating and colony counting is shown in *B*; gastric CD4⁺ T cell infiltration and IFN- γ and IL-17 expression
 636 by CD4⁺ T-cells, as determined by ICS, is shown in *C* and *D*. Horizontal lines in *B-D* indicate medians.
 637 Representative FACS plots are presented in *E*. Data in *B* are pooled from two independent studies; data in
 638 *C* and *D* are from one representative study of two independently conducted ones.

639

640 **Figure 4: ARTD1 controls macrophage infiltration and CD4⁺/CD8⁺ T cell activation in MC-38 tumors**

641 (*A-C*) Tumor volume and weight of MC-38 tumors, subcutaneously injected into the flanks of *Artd1*^{fl α /fl α}
 642 and *Artd1* ^{Δ Myel} mice, as determined by caliper measurements at the indicated time points and at the study end
 643 point. Each symbol represents one mouse. Data are pooled from two independent experiments and presented
 644 as mean \pm SEM (*A*). Horizontal lines in *B* and *C* indicate medians. *t*-Test: **P* < 0.05, ***P* < 0.01. (*D*)
 645 Monocyte infiltration of MC-38 tumors grown on the flanks of *Artd1*^{fl α /fl α} and *Artd1* ^{Δ Myel} mice, as
 646 quantified by flow cytometry. (*E*) *Il12b* and *Arg1* expression of FACS-sorted tumor infiltrating monocytes,
 647 as determined by qRT-PCR. Data in *E* are from one representative experiment of two; horizontal lines
 648 indicate medians. *t*-Test: **P* < 0.05, ***P* < 0.01. (*F*) Total and TNF- α ⁺ tumor-infiltrating macrophages as
 649 quantified by flow cytometry. (*G*) *Il12b* and *Arg1* expression of FACS-sorted tumor-infiltrating
 650 macrophages, as determined by qRT-PCR. Data are from one representative experiment of two, horizontal
 651 lines indicate medians. *t*-Test: **P* < 0.05, ***P* < 0.01. (*H, I*) Infiltration and IFN- γ and TNF- α expression
 652 of CD4⁺ T cells, as flow cytometrically determined after *ex vivo* re-stimulation with PMA/Ionomycin. Data
 653 are from one representative (left panel) or pooled from two independent experiments (right panels).
 654 Representative FACS plots are shown in *I*. *t*-Test: **P* < 0.05, ***P* < 0.01. (*J, K*) Infiltration and IFN- γ and
 655 TNF- α expression of CD8⁺ T cells, as flow cytometrically determined after *ex vivo* re-stimulation with
 656 PMA/ionomycin. Data in *J* are from one representative experiment of two. *t*-Test: **P* < 0.05, ***P* < 0.01.
 657 (*L, M*) Infiltration and IFN- γ and TNF- α expression of CD8⁺ T cells, as flow cytometrically determined
 658 after *ex vivo* re-stimulation with MC-38-specific peptide. Data are pooled from two experiments; horizontal
 659 lines indicate medians throughout. *t*-Test: **P* < 0.05, ***P* < 0.01.

660 **Supplementary Figure 1**

661 (A) Targeting strategy for the generation of the Cre-mediated whole body ARTD1-deficient mouse (adapted
 662 from Skarnes, Rosen et al. 2011(28)). Cre-mediated recombination deletes Exon 4 of the *Artd1* gene. (B)
 663 Successful modification of the *Artd1* locus. To verify the integration of the targeting vector specifically at
 664 the *Artd1* locus on chromosome 1, a long-range PCR amplifying the 3' and 5' homology arm of the *Artd1*
 665 gene was performed on genomic DNA from the initial breeding pair. Embryonic stem cell DNA used for
 666 the blastocyst injection served as control. Size of the bands are indicated as base pairs (bp). (C) Genotyping
 667 of mice. PCR amplification of mouse genomic DNA isolated from *Artd1^{flox/wt}*, *Artd1^{wt/wt}*, and *Artd1^{flox/flox}*
 668 mice. Size of the bands are indicated as base pairs (bp). (D) Western blot analysis of the indicated organs
 669 for ARTD1 and tubulin expression in 6 weeks old *Artd1^{wt/wt}*, *Artd1^{del/del}* and *Artd1^{wt/del}* mice. (E) Necropsy
 670 analysis of three *Artd1^{flox/flox}* and two *Artd1^{del/del}* male mice of 8 weeks. Body weights as well as indicated
 671 organ weights were determined. (F) Quantitative Real-time PCR analysis of *Artd1* expression in *Artd1^{flox/flox}*
 672 and *Artd1^{del/del}* BMDMs used for RNA Sequencing. (G) RNA-sequencing of total RNA extracted from
 673 *Artd1^{flox/flox}* and *Artd1^{del/del}* BMDMs either left untreated or treated with 10 ng/ml LPS and 2 ng/ml INF- γ for
 674 4 h. All LPS/INF- γ -repressed genes (approx. 3000 genes) (fold-change ≤ 2 , $P < 0.05$) were clustered. (H)
 675 RNA-sequencing of total RNA extracted from *Artd1^{flox/flox}* either left untreated or treated 5 μ M Olaparib for
 676 1 h prior the stimulation with 10 ng/ml LPS and 2 ng/ml INF- γ for 4 h. (fold-change ≤ 2 , $P < 0.05$) were
 677 clustered. (I) ELISA of cell culture supernatant for the quantification of the indicated cytokines from
 678 *Artd1^{flox/flox}* and *Artd1^{del/del}* BMDMs. Cells were treated either with DMSO or 5 μ M PARPi (PJ34/Olaparib)
 679 prior to stimulation with 10 ng/ml LPS and 2 ng/ml INF- γ for 18 h. Data presented as mean concentration
 680 + SD of three independent experiments. (J) Gene enrichment analysis of cluster 1 (n = 95, identified in H)
 681 in *Artd1^{flox/flox}* BMDMs.

682

683 **Supplementary Figure 2**

684 (A) Targeting strategy for the generation of myeloid specific ARTD1 knock-out mouse (adapted from
 685 Skarnes, Rosen et al. 2011(28)). Cre-mediated recombination deletes Exon 4 of the *Artd1* gene. (B) Western
 686 blot analysis of BMDMs and thioglycolate elicited peritoneal macrophages from *Artd1^{flox/flox}* and *Artd1^{AMyel}*
 687 mice. ARTD1 levels were quantified in whole cell lysates. Tubulin was used as loading control. (C)
 688 *Artd1^{flox/flox}* and *Artd1^{AMyel}* mice were intraperitoneally injected with PBS or 4 mg/kg of LPS. After 4 h whole
 689 blood serum was collected and the levels of selected cytokines were quantified by multiplex-bead array.
 690 Data shown as mean \pm SEM of three independent experiments. *t*-Test: * $P < 0.05$. (D,E)
 691 Immunophenotyping of the spleen from *Artd1^{flox/flox}* and *Artd1^{AMyel}* mice. Spleen single cell suspensions were
 692 stained with fluorescently labeled antibodies and cell counts were determined by flow cytometry. Data
 693 presented as mean \pm SEM of a representative experiment. *t*-Test: * $P < 0.05$.

694

695 **Supplementary Figure 3**

696 Gating strategy for intracellular cytokine staining of gastric leukocytes.

697

698 **Supplementary Figure 4**

699 Gating strategy for the quantification and sorting of myeloid cells from the tumor microenvironment.

700