






Immunometabolism and atherosclerosis: perspectives and clinical significance: a position paper from the Working Group on Atherosclerosis and Vascular Biology of the European Society of Cardiology

Daniel F.J. Ketelhuth ^{1,2†}, Esther Lutgens^{3,4,5†}, Magnus Bäck¹, Christoph J. Binder⁶, Jan Van den Bossche ⁷, Carolin Daniel⁸, Ingrid E. Dumitriu ⁹, Imo Hoefler¹⁰, Peter Libby¹¹, Luke O'Neill¹², Christian Weber ^{4,13}, and Paul C. Evans ^{14*}

¹Department of Medicine, Cardiovascular Medicine Unit, Center for Molecular Medicine, Karolinska Institute, Karolinska University Hospital, Stockholm, Sweden; ²Department of Cardiovascular and Renal Research, Institute of Molecular Medicine, University of Southern Denmark, Odense, Denmark; ³Amsterdam University Medical Centers, University of Amsterdam, Amsterdam Cardiovascular Sciences, Amsterdam, The Netherlands; ⁴Institute for Cardiovascular Prevention, Ludwig Maximilians University of Munich, Munich, Germany; ⁵German Centre for Cardiovascular Research (DZHK), Partner Site Munich Heart Alliance, Munich, Germany; ⁶Department of Laboratory Medicine, Medical University of Vienna, Vienna, Austria and CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences, Vienna, Austria; ⁷Department of Molecular Cell Biology and Immunology, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam Cardiovascular Sciences, Cancer Center Amsterdam, Amsterdam, The Netherlands; ⁸Division of Clinical Pharmacology, Department of Medicine IV, Ludwig Maximilians University of Munich, Munich, Germany; ⁹Molecular and Clinical Sciences Research Institute & Cardiology Clinical Academic Group, St. George's Hospital, University of London, Cranmer Terrace, London, UK; ¹⁰Laboratory of Clinical Chemistry and Hematology, University Medical Centre Utrecht, Utrecht, Netherlands; ¹¹Division of Cardiovascular Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA; ¹²School of Biochemistry and Immunology, Trinity Biomedical Sciences Institute, Trinity College Dublin, Dublin, Ireland; ¹³Department of Biochemistry, Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, Maastricht, The Netherlands; and ¹⁴Department of Infection, Immunity and Cardiovascular Disease, INSIGNEO Institute of In Silico Medicine and the Bateson Centre, University of Sheffield, Sheffield S10 2RX, UK

Received 17 December 2018; revised 19 May 2019; editorial decision 24 May 2019; accepted 18 June 2019; online publish-ahead-of-print 22 June 2019

Abstract

Inflammation is an important driver of atherosclerosis, and the favourable outcomes of the Canakinumab Anti-inflammatory Thrombosis Outcome Study (CANTOS) trial revealed the large potential of anti-inflammatory drugs for the treatment of cardiovascular disease, especially in patients with a pro-inflammatory constitution. However, the complex immune reactions driving inflammation in the vascular wall in response to an atherosclerotic microenvironment are still being unravelled. Novel insights into the cellular processes driving immunity and inflammation revealed that alterations in intracellular metabolic pathways are strong drivers of survival, growth, and function of immune cells. Therefore, this position paper presents a brief overview of the recent developments in the immunometabolism field, focusing on its role in atherosclerosis. We will also highlight the potential impact of immunometabolic markers and targets in clinical cardiovascular medicine.

Keywords

Atherosclerosis • Vascular • Inflammation • Immune system • Metabolism

This article is part of the **Spotlight Issue on Immunometabolism**.

1. Introduction

Cardiovascular disease (CVD), including myocardial infarction and stroke, is a major cause of morbidity and mortality in the western world. Despite the management of risk factors, including cessation of smoking, treatment of hypertension, and lipid-lowering regimens using HMG-CoA reductase inhibitors or the recently developed proprotein

convertase subtilisin/kexin type 9 (PCSK9) inhibitors, a substantial proportion of the population still suffers from CVD.¹ Over the past years, it became clear that atherosclerosis, the underlying cause of the majority of CVDs, is not only driven by lipids, but also by inflammation.^{2,3} The research aimed at understanding the complex immune reactions driving inflammation in the vascular wall has shown that the infiltration, retention, and accumulation of lipoproteins in the arterial intima elicit maladaptive

* Corresponding author. Tel: +44(0)1142159525; E-mail: paul.evans@sheffield.ac.uk

† These authors contributed equally to this work.

immune responses that influence the development, progression, and stability of atherosclerotic lesions.^{4–6}

Atherosclerotic plaques recruit many different immune cells, including macrophages, T cells, B cells, dendritic cells, neutrophils, and mast cells that change in composition during atherogenesis.^{7,8} Together, they determine atherosclerotic plaque progression through the secretion of cytokines, chemokines, proteases, pro-thrombotic factors, and other bioactive substances. The balance between pro-inflammatory and anti-inflammatory responses in the plaque will dictate the rate of disease development as well as the size and complexity of lesions. Large atherosclerotic lesions presenting unresolved inflammation, extensive matrix remodelling, large necrotic cores, and thin fibrous caps are at risk of rupture leading to acute thrombosis and subsequent vascular occlusion.⁴

Compelling evidence that the immune system plays a pivotal role in atherosclerotic CVD in humans was reported in 2017 when the results from the Canakinumab Anti-inflammatory Thrombosis Outcome Study (CANTOS) trial were released.⁹ Here, immunotherapy with an interleukin-1 β (IL-1 β) antibody effectively reduced CVD risk and mortality, without affecting low-density lipoprotein (LDL) cholesterol concentrations.⁹ The initiation and favourable outcomes of the CANTOS trial revealed the potential of anti-inflammatory drugs to combat CVD. Although Canakinumab was not approved by the FDA for treatment of atherosclerotic CVD, CANTOS is an important proof of principle study, revealing the potential of anti-inflammatory therapies in atherosclerosis, especially in CVD patients with residual inflammation.¹⁰ An important lesson was learned from the CIRT trial where treatment of CVD patients with methotrexate, a broad-spectrum anti-inflammatory agent, failed to reduce CVD or mortality.¹¹ Collectively, the data from CANTOS and CIRT have taught us that we need to identify drug targets that block atherosclerosis-specific inflammatory pathways.

Recently, it was found that alterations in intracellular metabolic pathways in immune cells occur during activation and that these pathways are key regulators of the immune responses. These data led to a novel research field in immunology, termed 'immunometabolism'.¹² Recent data show a pivotal role for immunometabolism in disease progression, especially in cancer, but also in obesity and Type 2 diabetes, diseases driving CVD atherosclerosis.^{13,14} Therefore, this position paper presents a brief overview of the recent developments in the field and the impact of 'immunometabolic' changes on atherosclerosis.

2. Immune cell metabolic pathways—brief overview

Immune cells are highly dynamic and face different metabolic demands in an inflammatory environment. Upon activation, immune cells switch between different metabolic traits (metabolic reprogramming) and can adapt to variations in environmental cues (e.g. oxygen, nutrients, growth factors), as well as to energy and biosynthetic requirements.

At least seven major cellular metabolic pathways have been described in immune cells. The interconnecting pathways of glycolysis, the pentose phosphate pathway (PPP), the tricarboxylic acid (TCA) cycle, oxidative phosphorylation (OXPHOS), mitochondrial fatty acid β -oxidation (FAO), fatty acid synthesis, and the metabolism of amino acids regulate the survival, growth, and activation of immune cells.¹²

The interconnectivity of these pathways was reviewed by O'Neill et al.¹² In brief, immune cells take-up glucose via the GLUT1 receptor, which is then converted into glucose-6-phosphate and metabolized via glycolysis, thereby converted into pyruvate and two molecules of

adenosine triphosphate (ATP). Pyruvate is converted to lactate by lactate dehydrogenase or enters the TCA cycle.

The PPP runs parallel to the glycolytic pathway and is responsible for nucleotide and NADPH production. NADPH is used by NADPH oxidase to generate reactive oxygen species (ROS), which is counterbalanced by the generation of glutathione and other antioxidants. The latter are major mediators of antimicrobial immunity and prevent tissue damage induced by activated macrophages and neutrophils. NADPH generated by the PPP is also required for *de novo* fatty acid synthesis for expansion of the endoplasmic reticulum (ER) and the golgi to facilitate enhanced cytokine secretion.^{12,14}

After glycolysis, pyruvate enters the TCA cycle where it can be converted, together with fatty acids, into acetyl-coA. The TCA cycle results in the generation of NADH and flavin adenine dinucleotide (FADH₂), that transfer electrons to the electron-transport chain, where OXPHOS takes place to yield ATPs. FAO is the most effective way to produce large amounts of ATP. Short-chain fatty acids diffuse into mitochondria where they become oxidized, whereas long-chain fatty acids conjugate to palmitoyltransferase 1 for transport. FAO yields acetyl-coA, NADH, and FADH₂ that enter the TCA cycle/OXPHOS to generate ATP.^{12,14}

Via *fatty acid synthesis*, metabolic intermediates are converted into triacylglycerols and phospholipids, needed to (re)build cellular structures. *Amino acid metabolism* yields important mediators that exert effector and regulatory functions on immune cells. Examples include glutamine, arginine, and tryptophan.^{12,14}

3. Immunometabolic signalling and regulation: macrophages and T cells

Atherosclerotic plaques are characterized by the presence of a plethora of macrophage and T cell subtypes. The entire spectrum between classically activated M1, and alternatively activated M2 macrophages is present.^{7,8,15} Likewise, the majority of T cell subsets including Th1, Th17, and regulatory T cells have been detected within the atherosclerotic plaque.⁵ Activated macrophages and T cells display a stronger metabolic bias towards aerobic glycolysis than towards mitochondrial metabolism, while immune regulatory cells including M2 macrophages and Tregs exhibit a mixed metabolism involving glycolysis, fatty acid oxidation, and OXPHOS.^{16,17} Here, we provide a brief overview of the metabolic alterations that go hand in hand with the different macrophage and T cell activation states (summarized in *Figure 1*).

3.1 Macrophages

Macrophages form an important plaque constituent. Plaque macrophages take-up modified LDL and most of them become lipid-laden foam cells but can also be activated by lipoprotein-derived antigens, e.g. phospholipids, cholesterol crystals, and apolipoprotein B peptides. In addition, part of the macrophage population in the plaque also has anti-inflammatory properties.^{18,19} To fulfil this broad range of functions, macrophages are highly plastic cells and able to acquire a wide array of activation states. For instance, during inflammation, their high-plasticity enables macrophages to initiate inflammatory responses and to switch off inflammatory responses when no longer needed. Recent studies have elucidated how some of the metabolic processes in macrophages are wired and how metabolism shapes macrophage inflammatory responses.¹⁴

The importance of metabolism in macrophage activation is illustrated by the fact that metabolism of the amino acid arginine formed the basis




Figure 1 The role of immunometabolism in atherosclerosis. Pro-inflammatory and anti-inflammatory immune cells present distinct metabolic phenotypes. In general, M1 type macrophages, Th1, and Th17 cells are characterized by more catabolic metabolism, while M2 type macrophages and Tregs present bias towards more anabolic metabolism. Alterations in metabolism of immune cells carry the potential to influence plaque progression and stabilization. FA, fatty acid; FAO, fatty acid oxidation; IFN γ , interferon gamma; NLRP3, NACHT, LRR, and PYD domains-containing protein 3; OXPPOS, oxidative phosphorylation; PPP, pentose phosphate pathway; ROS, reactive oxygen species; TGF β , growth factor beta; TNF, tumour necrosis factor.

of the dichotomous M1/M2 classification. Whereas anti-inflammatory M2 macrophages use arginase to convert arginine to urea and ornithine, inflammatory M1 macrophages metabolize arginine using inducible NO synthase to convert arginine into the pro-inflammatory NO and citrulline.²⁰ Once M1 macrophages have formed, NO damages the mitochondrial electron-transport chain and M1 macrophages cannot repolarize towards M2 macrophages, whereas M2 macrophages can easily switch to an M1 phenotype.²¹ The arginine pathway is only one of many metabolic pathways that drive macrophage activation states.

Inflammatory macrophages require a rapid supply of energy and biosynthetic products as they need to release their inflammatory contents fast, whereas anti-inflammatory macrophages need a more sustained source of energy for long-lasting repair responses.¹⁴ Therefore,

inflammatory macrophages have an increased glucose uptake via the glucose transporter GLUT1 and exhibit enhanced aerobic glycolysis, whereas OXPPOS via the TCA cycle is impaired.²² During this process, pyruvate, produced by the glycolytic pathway, and generated through dimerization of pyruvate kinase iso-enzyme 2 (PKM2),²³ which catalyses the final step of glycolysis, is converted to lactate, resulting in two ATP molecules and induction of ROS.¹² At the same time, the PPP is enhanced through increased flux of glucose intermediates, resulting in increased NADPH synthesis, which is key for cholesterol and fatty acid synthesis, needed for phagocytosis as well as expansion of the ER and golgi, which results in enhanced production of inflammatory cytokines.²⁴

Other metabolic intermediaries can regulate macrophage function, including succinate, an intermediate of the TCA cycle, which accumulates

in M1 macrophages and drives inflammation via succinylation of intracellular proteins, including the hypoxia-inducible factor 1- α that augments IL-1 β production.²⁵ Succinate can also propagate inflammatory responses extracellularly through direct binding to the succinate receptor GPR91.²⁶ The endogenous metabolite itaconate has been recently shown to oppose the deleterious metabolic rewiring, allowing nuclear factor (erythroid-derived)-like 2 (NRF2) to induce downstream anti-inflammatory and anti-oxidant genes.²⁷ Other TCA cycle intermediates, including fumarate and citrate, can contribute to histone acetylation and methylation, thereby affecting epigenetic marks that drive innate immune memory.²⁸ Increased fatty acid synthesis is also associated with macrophage activation, as fatty acid synthesis activates the NLRP3 inflammasome, thereby promoting the release of IL-1 β .^{6,28}

Anti-inflammatory macrophages have a less well-understood phenotype but are characterized by increased rates of OXPHOS and FAO. In these macrophages, both pyruvate and fatty acids enter the intact TCA cycle as acetyl-coA, resulting in sustained ATP production via OXPHOS, which leads to up-regulation of genes associated with tissue repair.²² Interestingly, when OXPHOS prevails, the enzyme glyceraldehyde 3-phosphate dehydrogenase (GAPDH), a crucial glycolytic enzyme, moonlights and represses tumour necrosis factor and IFN γ through the binding to their mRNA.²⁹ FAO generates more/a larger number of ATP molecules than glycolysis and is an effective way for anti-inflammatory macrophages to provide a long-lasting energy source. However, the importance of FAO in anti-inflammatory macrophage function has recently been questioned, as etomoxir mediated inhibition of FAO or deficiency in CPT2, an enzyme required for fatty acid import does not alter the M2 phenotype.^{30,31}

The availability and metabolism of various amino acids have also been shown to regulate innate immune cell responses. In macrophages, glutamine can regulate IL-1 β secretion, the production of NO, as well as M2 polarization.^{22,32–34} Arginine metabolism, via the citrulline pathway and iNOS induction, leading to nitric oxide formation is associated with the M1 phenotype.³⁵ Through a different mechanism, acetyl-CoA and S-adenosylmethionine can regulate epigenetic enzymes to enable histone acetylation and methylation, thereby translating metabolic rewiring into regulation of gene expression and macrophage function.³⁶

3.2 T cells

T cells migrate to tissues followed by tissue-specific adaptations and carry the ability to respond to environmental and metabolic signals. In this context, high-calorie consumption, obesity, or T2D have been characterized by infiltration of T cells in metabolically relevant organs in both mice and humans.

The adipose tissue, which stores and senses the availability of nutrients, is an important site of immunometabolic crosstalk. While lean visceral adipose tissue (VAT) is usually enriched in Th2 cells and Tregs, VAT from obese individuals has been associated with reduced Treg and increased effector T-cell numbers.^{37,38} Brown adipose tissue, which has a high-energy expenditure, as well as browning/beiging of VAT have been associated with increased Treg numbers.^{37,38} Interestingly, nutrient metabolism, as well as adipokines such as leptin, have been linked with different T-cell subset differentiation.³⁹ Leptin has been implicated to directly induce glycolysis in T cells, promoting effector responses and increasing inflammation.⁴⁰ Leptin-deficiency results in an increase in Tregs that are also more suppressive than Treg from wild type mice. Transfer of leptin-deficient Tregs into an experimental model of atherosclerosis caused a significant reduction of plaque size and a marked reduction of IFN γ production, compared to transfer of wild type Tregs.⁴¹

Similar to macrophages, glycolysis plays an important role in T-cell responses. Inhibition of glycolysis with 2-deoxyglucose (2DG) shifts the polarization of naïve T cells from Th17 towards Tregs.⁴² Hence, increased OXPHOS metabolism has been linked to the induction Tregs.⁴³ Interestingly, the glycolytic enzyme enolase has been shown to promote Foxp3 splicing and the generation of Treg cells through a non-anticipated 'moonlight' function,⁴³ and 3-phosphate dehydrogenase (GAPDH) has been shown to modulate Th1 responses through repression of interferon- γ (IFN γ) mRNA in a low glycolytic activity status.⁴⁴ Hence, only upon increased glycolytic activity Th1 cells can mount full pro-inflammatory response and secrete IFN γ .⁴⁵

The balance between effector and Treg is also influenced by FAO and fatty acid synthesis. Effector T cells have been shown to have reduced FAO upon activation,⁴⁶ while Tregs have increased expression of FAO enzymes, including carnitine palmitoyltransferase 1A (CPT1A).⁴⁷ Inhibition of acetyl-CoA carboxylase 1 (ACC1), the rate-limiting enzyme in fatty acid synthesis, restrains Th17 polarization and promotes the development of Tregs.^{48,49}

Recent studies reveal that specific amino acids and amino acid transporters regulate homeostasis and activation of the adaptive immune system.^{48,49} Down-regulation of large neutral amino acid transporter (LAT1) can impair Th1 and Th17 differentiation *in vivo*.⁵⁰ Moreover, indoleamine 2,3-dioxygenase-1 (IDO1), the rate-limiting enzyme catalysing tryptophan (Trp) degradation can modulate T cell effector responses, the expansion of Tregs, and the degree of vascular inflammation and atherosclerosis.^{51–53} Upon activation, T and B cells increase glutamine usage.⁵⁴ Glutamine metabolism has been implicated in the balance between effector T cells, Th1 and Th17, and Tregs.⁵⁵ Moreover, arginine, as well as glutamine and Trp availability are known regulators of immune function via the mTOR pathway.⁵⁶

Several studies demonstrate an essential role for cholesterol and fatty acids in activation, differentiation, and function of T cells.^{49,57,58} Hypercholesterolaemia can influence T cell receptor (TCR) signalling and Treg numbers, while Tregs can tightly regulate lipoprotein metabolism and influence hepatic inflammation.^{59–61} Recent data suggest that apolipoprotein A1 (ApoA1), the main protein component of HDL, modulates the conversion of Tregs into T follicular helper cells influencing atherosclerosis.⁶² The crosstalk between the immune system and lipid metabolism is an area of increasing interest due to increasing prevalence of the metabolic syndrome, together with chronic inflammatory liver diseases such as non-alcoholic fatty liver disease (NAFLD)/non-alcoholic steatohepatitis (NASH)—all representing additive risk to CVD.

4. Immunometabolism in atherosclerosis

4.1 Plaque metabolism

The metabolic blueprints for macrophages and T cells described above are obtained from experimental model systems. Although they are of utmost importance to understanding metabolic rewiring of immune cell types in activated and modulatory states, these data may not fully reflect the changes that occur in tissues during pathogenesis. Immunometabolic data in atherosclerotic CVD are still sparse, but recent reports have revealed insights into some of the immune-metabolic patterns that mediate atherosclerosis.

The first proof that alterations in immunometabolism are an important feature of atherogenesis was provided by the nuclear imaging

community. ^{18}F -fluorodeoxyglucose (FdG) positron-emission tomography imaging, which reflects glucose transport into cells, is widely used clinically to detect diseased tissue conditions exhibiting increased glucose metabolisms, such as tumours or infections. As a radionucleotide analogue of glucose, FdG is taken-up by cells via glucose transporters (GLUTs) and phosphorylated into FdG-6-phosphate that cannot be further metabolized and accumulates inside of the cell. Increased glucose metabolism has been also considered as the basis for imaging the burden of atherosclerosis. High FdG uptake has been suggested to reflect the degree of vascular inflammation and plaque vulnerability, but may also reflect hypoxia.^{63,64}

A recent study has revealed that besides an increase in glucose uptake, atherosclerotic disease is characterized by changes in multiple intracellular metabolic pathways in the arterial wall. Metabolomics analysis of 159 plaques of symptomatic (TIA and stroke) and asymptomatic patients obtained during carotid endarterectomy revealed a distinct metabolite profile in inflammatory plaques of patients with symptomatic carotid artery disease. These symptomatic plaques revealed a cluster of metabolites and enzymes that are associated with increased glycolysis, elevated amino acid utilization, and decreased FAO. Moreover, this cluster was highly associated with plaque inflammation.⁶⁵

4.2 Systemic changes in metabolism affecting atherosclerosis

Work in experimental atherosclerosis models has confirmed the functional importance of immunometabolic pathways in CVD. For example, it has been shown that atherosclerotic mice deficient in hematopoietic GLUT1 have a decreased glycolytic flux in their bone marrow and atherosclerotic plaques, resulting in a decrease in atherosclerosis.⁶⁶ Likewise, deficiency of glucose-6-phosphate dehydrogenase, a key enzyme in the PPP reduced vascular superoxide levels and also decreased atherosclerosis.⁶⁷

Amino acid metabolism also proved of importance in atherosclerosis. Ablation of IDO-dependent Trp metabolism leads to a substantial increase in vascular inflammation and acceleration of atherosclerosis in *Apoe*^{-/-} mice.^{51,53} In line with this data, IDO induction has been linked to atheroprotection and increased plaque stability.^{52,68,69} Nevertheless, the role of IDO in health and disease seems to be sensitive to alterations in the gut microbiome that may impair its anti-inflammatory and anti-atherosclerotic effects.^{70–72}

The microbiome also plays an important role in mediating immunometabolism in atherosclerosis. Changes in composition of the microbiome also affect the release of immunopotent metabolites that have been associated with CVD. One example is Trimethylamine N-oxide (TMAO), a plasma metabolite that is formed through the conversion of microbiome generated trimethylamine (TMA) into TMAO via the host's hepatic flavin monooxygenase.⁷³ The generation of TMAO involves nutrient precursors that are highly abundant in a western diet, and TMAO was shown to induce platelet activation and vascular inflammation in experimental models and patients.^{73–75} TMAO levels are associated with atherosclerosis burden⁷⁶ and predict both near- and long-term risk of major adverse cardiovascular events.⁷⁷

4.3 Monocytes–macrophages

Atherosclerosis is associated with several changes in monocytes and macrophages. Analysis of monocytes isolated from healthy individuals or individuals suffering from atherosclerotic CVD revealed that monocytes obtained from patients had a higher oxygen consumption rate, a higher

glycolytic acidification rate, and glycolytic flux. These monocytes had an enhanced glucose uptake, produced more mitochondrial ROS, and had enhanced inflammatory signalling. It was found that monocytes from atherosclerotic CVD patients switched to the glucose-ROS-PKM2-STAT3 pathway through which glucose utilization led to unbalanced ROS generation from the mitochondrial chain, that induced translocation of the enzyme PKM2 and induction of STAT3 signalling, resulting in inflammation.²³

The mechanism for altered monocyte–macrophage metabolic activity may relate to alterations in LDL levels which are characteristic of atherosclerosis.^{78,79} *In vitro* studies have shown that oxLDL results in an increase in glycolysis, inflammation, and oxidative damage in macrophages.^{80–82} *In vivo* studies confirm these findings. High cholesterol-diet-fed LDLR^{-/-} mice indeed show epigenetic and metabolic reprogramming of myeloid (progenitor) cells, with profound up-regulation of the inflammasome.⁸³ Exposure to high levels of modified LDL triggers epigenetic and metabolic reprogramming of macrophages and exacerbate inflammatory responses.⁸³ Altered macrophage metabolism can provoke prolonged responses, a phenomenon called 'innate immune memory' or 'trained immunity'.

⁸⁴ Glycolysis, glutaminolysis, and cholesterol synthesis can influence the activity of methyltransferases and demethylases, acetyltransferases, and deacetylases, which by targeting DNA and histones promote increased inflammatory gene transcription. Interestingly, detailed analysis of distinct plaque macrophage subsets uncovered that non-foamy, rather than lipid-loaded foamy macrophages are pro-inflammatory and are likely the cells that drive lesion inflammation.⁸⁵ Mechanistically, increased activation of the liver-X-receptor, trying to mediate cholesterol efflux, in parallel to suppressed activity of the PPP in foam cells, may explain the reduced inflammatory responses in those lipid-loaded macrophages.^{86,87}

Although much work needs to be done to understand the intertwined and dynamic metabolic changes and pathways that occur in atherosclerosis, it is clear that the atherosclerotic microenvironment causes immunometabolic changes that can drive progression or regression and stabilization of atherosclerotic disease (illustrated in Figure 1). Therefore, targeting of immunometabolic pathways is a promising approach to combat atherosclerotic CVD.

Unfortunately, a big gap still exists between the experimental work on immunometabolism and implementation of these experimental results in the clinic. Compelling evidence substantiates that targeting inflammation in CVD improves outcome,⁹ and it is increasingly known that existing therapies, including HMG-CoA reductase inhibitors⁸⁸ and anti-diabetic drugs, can also reduce arterial inflammation, most likely by affecting immunometabolic pathways. Therefore, we are approaching an exciting future for scientific advances in immunometabolism which may lead to novel therapies to address residual cardiovascular risk in the clinic.

5. Conclusions and perspectives

This position paper illustrates the potential of immunometabolism to identify new targets for prevention and treatment of CVDs. Although experimental data are promising, implementation of experimental therapies in the clinic presents challenges. More early-stage investments should help to develop this field further. Ultimately, large scale randomized clinical trials will be necessary to evaluate therapies that target immunometabolism and ascertain their effectiveness and possible unwanted actions.

6. Consensus statements

- Metabolites are not just 'fuels' in their pathways, they are also effectors and signalling molecules that regulate the immune system.
- Cellular metabolic pathways are tightly regulated by several pro-atherogenic factors including lipids, glucose, amino acids, and pro- and anti-inflammatory cytokines.
- Systemic and microenvironment-induced changes in basic metabolic pathways can skew the balance between pro- and anti-inflammatory responses in atherosclerosis.
- The identification of the key immunometabolic reactions governing plaque development and stability will give a new understanding of disease processes, and likely lead to novel therapeutic approaches to prevent and treat atherosclerotic CVDs.
- We now recognize a new frontier, immunometabolism, which presents further opportunities for the CVD field to expand fundamental understanding and furnish new therapeutic avenues for our patients.

Author contributions

All authors contributed substantially to the conception, design and organization of the paper. All authors provided important intellectual content and helped write (draft, critique, revise) the manuscript.

Conflict of interest: CJB has received honoraria for consultancy and lectures from Amgen and AOP Pharma. CJB is board member of Technoclone GmbH. PL's laboratory has received research support from Novartis. Other authors declared none.

Funding

D.F.J.K. is supported by the Swedish Heart-Lung Foundation and the Novo Nordisk Foundation (NNF15CC0018346). E.L. is funded by the NWO (VICI), the European Research Council (ERC-con) and the DFG (SFB1123), and the CVON consortium GENIUS-II. M.B. is funded by Swedish Research Council (2014-2312); and the Swedish Heart and Lung Foundation (20150600 and 20150683). J.V.d.B. received a VENI grant from NWO [91615052], a junior postdoctoral grant [2013T003] and senior fellowship [2017T048] from the Netherlands Heart Foundation, a CCA (Cancer Center Amsterdam) PhD project and proof of concept grant, and a joint VUmc/AMC PhD project from ACS (Amsterdam Cardiovascular Sciences). I.E.D. is funded by the British Heart Foundation (grant number: PG/10/50/28434; PG/13/24/30115; PG/14/18/30724; PG/17/15/32845) and St George's Hospital Charity (grant number 16.03.05). CB received SFB-54 "InThro" of the Austrian Science Fund (FWF). P.L. has received funding from the National Heart, Lung, and Blood Institute (R01HL080472), the American Heart Association (18CSA34080399), and the RRM Charitable Fund. C.W. is funded by the DFG (SFB1123) and by ERC (AdG °692511). P.C.E. is funded by the British Heart Foundation.

References

1. Cholesterol Treatment Trialists' (CTT) Collaborators, Mihaylova B, Emberson J, Blackwell L, Keech A, Simes J, Barnes EH, Voysey M, Gray A, Collins R, Baigent C. The effects of lowering LDL cholesterol with statin therapy in people at low risk of vascular disease: meta-analysis of individual data from 27 randomised trials. *Lancet* 2012;**380**:581–590.
2. Weber C, Noels H. Atherosclerosis: current pathogenesis and therapeutic options. *Nat Med* 2011;**17**:1410–1422.
3. Legein B, Janssen EM, Theelen TL, Gijbels MJ, Walraven J, Klarquist JS, Hennies CM, Wouters K, Seijkens TTP, Wijnands E, Sluimer JC, Lutgens E, Zenke M, Hildner K, Biessen EAL, Temmerman L. Ablation of CD8 α (+) dendritic cell mediated cross-presentation does not impact atherosclerosis in hyperlipidemic mice. *Sci Rep* 2015;**5**:15414.
4. Libby P, Lichtman AH, Hansson GK. Immune effector mechanisms implicated in atherosclerosis: from mice to humans. *Immunity* 2013;**38**:1092–1104.

5. Ketelhuth DJF, Hansson GK. Adaptive response of T and B cells in atherosclerosis. *Circ Res* 2016;**118**:668–678.
6. Tabas I, Bornfeldt KE. Macrophage phenotype and function in different stages of atherosclerosis. *Circ Res* 2016;**118**:653–667.
7. Cole JE, Park I, Ahern D, Kassiteridi C, Danso Abeam D, Goddard M, Green P, Maffia P, Monaco C. Immune cell census in murine atherosclerosis: cytometry by time of flight illuminates vascular myeloid cell diversity. *Cardiovasc Res* 2018;**390**:1151.
8. Winkels H, Ehinger E, Vassallo M, Buscher K, Dinh H, Kobiyama K, Hamers A, Cochain C, Vafadarnejad E, Saliba A-E, Zernecke A, Pramod A, Ghosh A, Anto Michel N, Hoppe N, Hilgendorf I, Zirikli A, Hedrick C, Ley K, Wolf D. Atlas of the immune cell repertoire in mouse atherosclerosis defined by single-cell RNA-sequencing and mass cytometry. *Circ Res* 2018;**122**:1675. [10.1161/CIRCRESAHA.117.312513]
9. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, Fonseca F, Nicolau J, Koenig W, Anker SD, Kastelein JJP, Cornel JH, Pais P, Pella D, Genest J, Cifkova R, Lorenzatti A, Forster T, Kobalava Z, Vida-Simiti L, Flather M, Shimokawa H, Ogawa H, Dellborg M, Rossi PRF, Troquay RPT, Libby P, Glynn RJ; CANTOS Trial Group. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *N Engl J Med* 2017;**377**:1119–1131.
10. Ridker PM, MacFadyen JG, Everett BM, Libby P, Thuren T, Glynn RJ, Ridker PM, MacFadyen JG, Everett BM, Libby P, Thuren T, Glynn RJ, Kastelein J, Koenig W, Genest J, Lorenzatti A, Varigos J, Siostrzonek P, Sinnaeve P, Fonseca F, Nicolau J, Gotcheva N, Yong H, Urina-Triana M, Milicic D, Cifkova R, Vetter R, Anker SD, Manolis AJ, Wyss F, Forster T, Sigurdsson A, Pais P, Fucili A, Ogawa H, Shimokawa H, Veze I, Petrauskiene B, Salvador L, Cornel JH, Klemsdal TO, Medina F, Budaj A, Vida-Simiti L, Kobalava Z, Otasevic P, Pella D, Lainscak M, Seung K-B, Commerford P, Dellborg M, Donath M, Hwang J-J, Kultursay H, Flather M, Ballantyne C, Bilazarian S, Chang W, East C, Forghash L, Harris B, Ligueros M; CANTOS Trial Group. Relationship of C-reactive protein reduction to cardiovascular event reduction following treatment with canakinumab: a secondary analysis from the CANTOS randomised controlled trial. *Lancet* 2018;**391**:319–328.
11. Ridker PM, Everett BM, Pradhan A, MacFadyen JG, Solomon DH, Zaharris E, Mam V, Hasan A, Rosenberg Y, Iturrriaga E, Gupta M, Tsigoulis M, Verma S, Clearfield M, Libby P, Goldhaber SZ, Seagle R, Ofori C, Saklayen M, Butman S, Singh N, Le May M, Bertrand O, Johnston J, Paynter NP, Glynn RJ; CIRT Investigators. Low-dose methotrexate for the prevention of atherosclerotic events. *N Engl J Med* 2019;**380**:752–762.
12. O'Neill LAJ, Kishton RJ, Rathmell J. A guide to immunometabolism for immunologists. *Nat Rev Immunol* 2016;**16**:553–565.
13. Font-Burgada J, Sun B, Karin M. Obesity and cancer: the oil that feeds the flame. *Cell Metab* 2016;**23**:48–62.
14. Koelwyn GJ, Corr EM, Erbay E, Moore KJ. Regulation of macrophage immunometabolism in atherosclerosis. *Nat Immunol* 2018;**19**:526–537.
15. Cochain C, Vafadarnejad E, Arampatzis P, Jaroslav P, Winkels H, Ley K, Wolf D, Saliba A-E, Zernecke A. Single-Cell RNA-Seq reveals the transcriptional landscape and heterogeneity of aortic macrophages in murine atherosclerosis. *Circ Res* 2018;**122**:1661. CIRCRESAHA.117.312509.
16. Mills EL, Kelly B, O'Neill L. Mitochondria are the powerhouses of immunity. *Nat Immunol* 2017;**18**:488–498.
17. Raud B, Roy DG, Divakaruni AS, Tarasenko TN, Franke R, Ma EH, Samborska B, Hsieh WY, Wong AH, Stüve P, Arnold-Schrauf C, Guderian M, Lochner M, Rampertaap S, Romito K, Monsale J, Brønstrup M, Bensinger SJ, Murphy AN, McGuire PJ, Jones RG, Sparwasser T, Berod L. Etomoxir actions on regulatory and memory T cells are independent of Cpt1 α -mediated fatty acid oxidation. *Cell Metab* 2018;**28**:504–515.e507.
18. Moore KJ, Tabas I. Macrophages in the pathogenesis of atherosclerosis. *Cell* 2011;**145**:341–355.
19. Subramanian M, Tabas I. Dendritic cells in atherosclerosis. *Semin Immunopathol* 2014;**36**:93–102.
20. Ley K. M1 means kill; M2 means heal. *J Immunol* 2017;**199**:2191–2193.
21. Van den Bossche J, Baardman J, Otto NA, van der Velden S, Neele AE, van den Berg SM, Luque-Martin R, Chen H-J, Boshuizen MCS, Ahmed M, Hoeksma MA, de Vos AF, de Winther M. Mitochondrial dysfunction prevents repolarization of inflammatory macrophages. *Cell Rep* 2016;**17**:684–696.
22. Jha AK, Huang S-C, Sergushichev A, Lampropoulou V, Ivanova Y, Lognischeva E, Chmielewski K, Stewart KM, Ashall J, Everts B, Pearce EJ, Driggers EM, Artyomov MN. Network integration of parallel metabolic and transcriptional data reveals metabolic modules that regulate macrophage polarization. *Immunity* 2015;**42**:419–430.
23. Shirai T, Nazarewicz RR, Wallis BB, Yanes RE, Watanabe R, Hilhorst M, Tian L, Harrison DG, Giacomini JC, Assimes TL, Goronzy JJ, Weyand CM. The glycolytic enzyme PKM2 bridges metabolic and inflammatory dysfunction in coronary artery disease. *J Exp Med* 2016;**213**:337–354.
24. Van den Bossche J, O'Neill LA, Menon D. Macrophage immunometabolism: where are we (going)? *Trends Immunol* 2017;**38**:395–406.
25. Tannahill GM, Curtis AM, Adamik J, Palsson-McDermott EM, McGettrick AF, Goel G, Frezza C, Bernard NJ, Kelly B, Foley NH, Zheng L, Gardet A, Tong Z, Jany SS, Corr SC, Haneklaus M, Caffrey BE, Pierce K, Walmsley S, Beasley FC, Cummins E, Nizet V, Whyte M, Taylor CT, Lin H, Masters SL, Gottlieb E, Kelly VP, Clish C,

- Auron PE, Xavier RJ, O'Neill LAJ. Succinate is an inflammatory signal that induces IL-1 β through HIF-1 α . *Nature* 2013;**496**:238–242.
26. Littlewood-Evans A, Sarret S, Apfel V, Loesle P, Dawson J, Zhang J, Muller A, Tigani B, Kneuer R, Patel S, Valeaux S, Gommermann N, Rubic-Schneider T, Junt T, Carballido JM. GPR91 senses extracellular succinate released from inflammatory macrophages and exacerbates rheumatoid arthritis. *J Exp Med* 2016;**213**:1655–1662.
 27. Mills EL, Ryan DG, Prag HA, Dikovskaya D, Menon D, Zaslon Z, Jedrychowski MP, Costa ASH, Higgins M, Hams E, Szpyt J, Runtsch MC, King MS, McGouran JF, Fischer R, Kessler BM, McGettrick AF, Hughes MM, Carroll RG, Booty LM, Knatko EV, Meakin PJ, Ashford MLJ, Modis LK, Brunori G, Sévin DC, Fallon PG, Caldwell ST, Kunji ERS, Chouchani ET, Frezza C, Dinkova-Kostova AT, Hartley RC, Murphy MP, O'Neill LA. Itaconate is an anti-inflammatory metabolite that activates Nrf2 via alkylation of KEAP1. *Nature* 2018;**556**:113–117.
 28. Groh L, Keating ST, Joosten LAB, Netea MG, Riksen NP. Monocyte and macrophage immunometabolism in atherosclerosis. *Semin Immunopathol* 2018;**40**:203–214.
 29. Millet P, Vachharajani V, McPhail L, Yoza B, McCall CE. GAPDH binding to TNF- α mRNA contributes to posttranscriptional repression in monocytes: a novel mechanism of communication between inflammation and metabolism. *J Immunol* 2016;**196**:2541–2551.
 30. Tan Z, Xie N, Cui H, Moellering DR, Abraham E, Thannickal VJ, Liu G. Pyruvate dehydrogenase kinase 1 participates in macrophage polarization via regulating glucose metabolism. *J Immunol* 2015;**194**:6082–6089.
 31. Nomura M, Liu J, Rovira II, Gonzalez-Hurtado E, Lee J, Wolfgang MJ, Finkel T. Fatty acid oxidation in macrophage polarization. *Nat Immunol* 2016;**17**:216–217.
 32. Wallace C, Keast D. Glutamine and macrophage function. *Metab Clin Exp* 1992;**41**:1016–1020.
 33. Murphy C, Newsholme P. Importance of glutamine metabolism in murine macrophages and human monocytes to L-arginine biosynthesis and rates of nitrite or urea production. *Clin Sci* 1998;**95**:397–407.
 34. Bellows CF, Jaffe BM. Glutamine is essential for nitric oxide synthesis by murine macrophages. *J Surg Res* 1999;**86**:213–219.
 35. Rath M, Müller I, Kropf P, Closs EI, Munder M. Metabolism via arginase or nitric oxide synthase: two competing arginine pathways in macrophages. *Front Immunol* 2014;**5**:532.
 36. Baardman J, Licht I, de Winther MPJ, Van den Bossche J. Metabolic-epigenetic crosstalk in macrophage activation. *Epigenomics* 2015;**7**:1155–1164.
 37. Becker M, Levings MK, Daniel C. Adipose-tissue regulatory T cells: critical players in adipose-immune crosstalk. *Eur J Immunol* 2017;**47**:1867–1874.
 38. Kálin S, Becker M, Ott VB, Serr I, Hosp F, Mollah MMH, Keipert S, Lamp D, Rohner-Jeanraud F, Flynn VK, Scherm MG, Nascimento LFR, Gerlach K, Popp V, Dietzen S, Bopp T, Krishnamurthy P, Kaplan MH, Serrano M, Woods SC, Tripal P, Palmisano R, Jastroch M, Blüher M, Wolfrum C, Weigmann B, Ziegler A-G, Mann M, Tschöp MH, Daniel C. A Stat6/Pten axis links regulatory T cells with adipose tissue function. *Cell Metab* 2017;**26**:475–492.e477.
 39. Lord GM, Matarese G, Howard JK, Baker RJ, Bloom SR, Lechler RI. Leptin modulates the T-cell immune response and reverses starvation-induced immunosuppression. *Nature* 1998;**394**:897–901.
 40. Gerriets VA, Danzaki K, Kishton RJ, Eisner W, Nichols AG, Saucillo DC, Shinohara ML, MacIver NJ. Leptin directly promotes T-cell glycolytic metabolism to drive effector T-cell differentiation in a mouse model of autoimmunity. *Eur J Immunol* 2016;**46**:1970–1983.
 41. Taleb S, Herbin O, Ait-Oufella H, Verreth W, Gourdy P, Barateau V, Merval R, Esposito B, Clément K, Holvoet P, Tedgui A, Mallat Z. Defective leptin/leptin receptor signaling improves regulatory T cell immune response and protects mice from atherosclerosis. *Arterioscler Thromb Vasc Biol* 2007;**27**:2691–2698.
 42. Shi LZ, Wang R, Huang G, Vogel P, Neale G, Green DR, Chi H. HIF1 α -dependent glycolytic pathway orchestrates a metabolic checkpoint for the differentiation of TH17 and Treg cells. *J Exp Med* 2011;**208**:1367–1376.
 43. De Rosa V, Galgani M, Porcellini A, Colamatteo A, Santopaulo M, Zuchegna C, Romano A, De Simone S, Procaccini C, La Rocca C, Carrieri PB, Maniscalco GT, Salvetti M, Buscarinu MC, Franzese A, Mozzillo E, La Cava A, Matarese G. Glycolysis controls the induction of human regulatory T cells by modulating the expression of FOXP3 exon 2 splicing variants. *Nat Immunol* 2015;**16**:1174–1184.
 44. Chang C-H, Curtis JD, Maggi LB, Faubert B, Villarino AV, O'Sullivan D, Huang SC-C, van der Windt GJW, Blagih J, Qiu J, Weber JD, Pearce EJ, Jones RG, Pearce EL. Posttranscriptional control of T cell effector function by aerobic glycolysis. *Cell* 2013;**153**:1239–1251.
 45. Mukhopadhyay R, Jia J, Arif A, Ray PS, Fox PL. The GAIT system: a gatekeeper of inflammatory gene expression. *Trends Biochem Sci* 2009;**34**:324–331.
 46. Wang R, Dillon CP, Shi LZ, Milasta S, Carter R, Finkelstein D, McCormick LL, Fitzgerald P, Chi H, Munger J, Green DR. The transcription factor Myc controls metabolic reprogramming upon T lymphocyte activation. *Immunity* 2011;**35**:871–882.
 47. Gerriets VA, Kishton RJ, Nichols AG, Macintyre AN, Inoue M, Ilkayeva O, Winter PS, Liu X, Priyadarshini B, Slawinska ME, Haeblerl L, Huck C, Turka LA, Wood KC, Hale LP, Smith PA, Schneider MA, MacIver NJ, Loscalce JW, Newgard CB, Shinohara ML, Rathmell JC. Metabolic programming and PDHK1 control CD4+ T cell subsets and inflammation. *J Clin Invest* 2015;**125**:194–207.
 48. Michalek RD, Gerriets VA, Jacobs SR, Macintyre AN, MacIver NJ, Mason EF, Sullivan SA, Nichols AG, Rathmell JC. Cutting edge: distinct glycolytic and lipid oxidative metabolic programs are essential for effector and regulatory CD4+ T cell subsets. *J Immunol* 2011;**186**:3299–3303.
 49. Berod L, Friedrich C, Nandan A, Freitag J, Hagemann S, Harmrolfs K, Sandouk A, Hesse C, Castro CN, Bähre H, Tschirner SK, Gorinski N, Gohmert M, Mayer CT, Huehn J, Ponimaskin E, Abraham W-R, Müller R, Lochner M, Sparwasser T. *De novo* fatty acid synthesis controls the fate between regulatory T and T helper 17 cells. *Nat Med* 2014;**20**:1327–1333.
 50. Buck MD, O'Sullivan D, Pearce EL. T cell metabolism drives immunity. *J Exp Med* 2015;**212**:1345–1360.
 51. Polyzos KA, Ovchinnikova O, Berg M, Baumgartner R, Agardh H, Pirault J, Gisterà A, Assinger A, Laguna-Fernández A, Bäck M, Hansson GK, Ketelhuth D. Inhibition of indoleamine 2,3-dioxygenase promotes vascular inflammation and increases atherosclerosis in ApoE^{-/-} mice. *Cardiovasc Res* 2015;**106**:295–302.
 52. Forteza MJ, Polyzos KA, Baumgartner R, Suur BE, Mussbacher M, Johansson DK, Hermansson A, Hansson GK, Ketelhuth DJ. Activation of the regulatory T-cell/indoleamine 2,3-dioxygenase axis reduces vascular inflammation and atherosclerosis in hyperlipidemic mice. *Front Immunol* 2018;**9**:950.
 53. Cole JE, Astola N, Cribbs AP, Goddard ME, Park I, Green P, Davies AH, Williams RO, Feldmann M, Monaco C. Indoleamine 2,3-dioxygenase-1 is protective in atherosclerosis and its metabolites provide new opportunities for drug development. *Proc Natl Acad Sci USA* 2015;**112**:13033–13038.
 54. Carr EL, Kelman A, Wu GS, Gopaul R, Senkevitch E, Aghvanyan A, Turay AM, Frauwrith KA. Glutamine uptake and metabolism are coordinately regulated by ERK1/2 MAPK during T lymphocyte activation. *J Immunol* 2010;**185**:1037–1044.
 55. Nakaya M, Xiao Y, Zhou X, Chang J-H, Chang M, Cheng X, Blonska M, Lin X, Sun S-C. Inflammatory T cell responses rely on amino acid transporter ASCT2 facilitation of glutamine uptake and mTORC1 kinase activation. *Immunity* 2014;**40**:692–705.
 56. Bar-Peled L, Sabatini DM. Regulation of mTORC1 by amino acids. *Trends Cell Biol* 2014;**24**:400–406.
 57. Wang F, Beck-García K, Zorzín C, Schamel WWA, Davis MM. Inhibition of T cell receptor signaling by cholesterol sulfate, a naturally occurring derivative of membrane cholesterol. *Nat Immunol* 2016;**17**:844–850.
 58. Molnár E, Swamy M, Holzer M, Beck-García K, Worch R, Thiele C, Guigas G, Boye K, Luescher IF, Schwille P, Schubert R, Schamel W. Cholesterol and sphingomyelin drive ligand-independent T-cell antigen receptor nanoclustering. *J Biol Chem* 2012;**287**:42664–42674.
 59. Mailer RKW, Gisterà A, Polyzos KA, Ketelhuth DJF, Hansson GK. Hypercholesterolemia Enhances T Cell Receptor Signaling and Increases the Regulatory T Cell Population. *Sci Rep* 2017;**7**:15655.
 60. Maganto-García E, Tarrío ML, Grabie N, Bu D-X, Lichtman AH. Dynamic changes in regulatory T cells are linked to levels of diet-induced hypercholesterolemia. *Circulation* 2011;**124**:185–195.
 61. Butcher MJ, Filipowicz AR, Waseem TC, McGary CM, Crow KJ, Maglínck N, Boldin M, Lundberg PS, Galkina EV. Atherosclerosis-driven Treg plasticity results in formation of a dysfunctional subset of plastic IFN γ + Th1/Tregs. *Circ Res* 2016;**119**:1190–1203.
 62. Gaddis DE, Padgett LE, Wu R, McSkimming C, Romines V, Taylor AM, McNamara CA, Kronenberg M, Crotty S, Thomas MJ, Sorci-Thomas MG, Hedrick CC. Apolipoprotein AI prevents regulatory to follicular helper T cell switching during atherosclerosis. *Nat Commun* 2018;**9**:1095.
 63. Folco EJ, Sheikine Y, Rocha VZ, Christen T, Shvartz E, Sukhova GK, Di Carli MF, Libby P. Hypoxia but not inflammation augments glucose uptake in human macrophages: implications for imaging atherosclerosis with 18fluorine-labeled 2-deoxy-D-glucose positron emission tomography. *J Am Coll Cardiol* 2011;**58**:603–614.
 64. Joseph P, Tawakol A. Imaging atherosclerosis with positron emission tomography. *Eur Heart J* 2016;**37**:2974–2980.
 65. Tomas L, Edsfeldt A, Mollet IG, Perisic Matic L, Prehn C, Adamski J, Paulsson-Berne G, Hedin U, Nilsson J, Bengtsson E, Gonçalves I, Björkbacka H. Altered metabolism distinguishes high-risk from stable carotid atherosclerotic plaques. *Eur Heart J* 2018;**39**:2301–2310.
 66. Sarrazy V, Viaud M, Westertep M, Ivanov S, Giorgetti-Peraldi S, Guinamard R, Gautier EL, Thorp EB, De Vivo DC, Yvan-Charvet L. Disruption of Glut1 in hematopoietic stem cells prevents myelopoiesis and enhanced glucose flux in atheromatous plaques of ApoE(-/-) mice. *Circ Res* 2016;**118**:1062–1077.
 67. Matsui R, Xu S, Maitland KA, Mastroianni R, Leopold JA, Handy DE, Loscalzo J, Cohen RA. Glucose-6-phosphate dehydrogenase deficiency decreases vascular superoxide and atherosclerotic lesions in apolipoprotein E(-/-) mice. *Arterioscler Thromb Vasc Biol* 2006;**26**:910–916.
 68. Daissormont I, Christ A, Temmerman L, Sampedro Millares S, Seijkens T, Manca M, Rousch M, Poggi M, Boon L, van der Loos C, Daemen M, Lutgens E, Halvorsen B, Aukrust P, Janssen E, Biessen E. Plasmacytoid dendritic cells protect against atherosclerosis by tuning T-cell proliferation and activity. *Circ Res* 2011;**109**:1387–1395.
 69. Yun TJ, Lee JS, Machmach K, Shim D, Choi J, Wi YJ, Jang HS, Jung I-H, Kim K, Yoon WK, Miah MA, Li B, Chang J, Bego MG, Pham TNQ, Loschko J, Fritz JH, Krug AB, Lee S-P, Keler T, Guimond JV, Haddad E, Cohen EA, Sirois MG, El-Hamamsy I, Colonna M, Oh GT, Choi J-H, Cheong C. Indoleamine 2,3-dioxygenase-expressing aortic plasmacytoid dendritic cells protect against atherosclerosis by induction of regulatory T cells. *Cell Metab* 2016;**23**:852–866.

70. Metghalchi S, Ponnuswamy P, Simon T, Haddad Y, Laurans L, Clement M, Dalloz M, Romain M, Esposito B, Koropoulos V, Lamas B, Paul J-L, Cottin Y, Kotti S, Bruneval P, Callebert J, Ruijter den H, Launay J-M, Danchin N, Sokol H, Tedgui A, Taleb S, Mallat Z. Indoleamine 2,3-dioxygenase fine-tunes immune homeostasis in atherosclerosis and colitis through repression of interleukin-10 production. *Cell Metab* 2015;**22**: 460–471.
71. Laurans L, Venticlef N, Haddad Y, Chajadine M, Alzaid F, Metghalchi S, Sovran B, Denis RGP, Dairou J, Cardellini M, Moreno-Navarrete J-M, Straub M, Jegou S, McQuitty C, Viel T, Esposito B, Tavitian B, Callebert J, Luquet SH, Federici M, Fernandez-Real JM, Burcelin R, Launay J-M, Tedgui A, Mallat Z, Sokol H, Taleb S. Genetic deficiency of indoleamine 2,3-dioxygenase promotes gut microbiota-mediated metabolic health. *Nat Med* 2018;**24**:1113–1120.
72. Fatkhullina AR, Peshkova IO, Dzutsev A, Aghayev T, McCulloch JA, Thovarai V, Badger JH, Vats R, Sundt P, Tang H-Y, Kossenkov AV, Hazen SL, Trinchieri G, Grivennikov SI, Koltsova EK. An interleukin-23-interleukin-22 axis regulates intestinal microbial homeostasis to protect from diet-induced atherosclerosis. *Immunity* 2018;**49**:943–957.e949.
73. Zhu W, Gregory JC, Org E, Buffa JA, Gupta N, Wang Z, Li L, Fu X, Wu Y, Mehrabian M, Sartor RB, McIntyre TM, Silverstein RL, Tang WHW, DiDonato JA, Brown JM, Lusa AJ, Hazen SL. Gut microbial metabolite TMAO enhances platelet hyperreactivity and thrombosis risk. *Cell* 2016;**165**:111–124.
74. Seldin MM, Meng Y, Qi H, Zhu W, Wang Z, Hazen SL, Lusa AJ, Shih DM. Trimethylamine N-oxide promotes vascular inflammation through signaling of mitogen-activated protein kinase and nuclear factor- κ B. *J Am Heart Assoc* 2016;**5**: e002767.
75. Haghikia A, Li XS, Liman TG, Bledau N, Schmidt D, Zimmermann F, Kränkel N, Widera C, Sonnenschein K, Haghikia A, Weissenborn K, Fraccarollo D, Heimesaat MM, Bauersachs J, Wang Z, Zhu W, Baveniek U, Hazen SL, Endres M, Landmesser U. Gut microbiota-dependent trimethylamine N-oxide predicts risk of cardiovascular events in patients with stroke and is related to proinflammatory monocytes. *Arterioscler Thromb Vasc Biol* 2018;**38**:2225–2235.
76. Senthong V, Li XS, Hudec T, Coughlin J, Wu Y, Levison B, Wang Z, Hazen SL, Tang WHW. Plasma trimethylamine N-oxide, a gut microbe-generated phosphatidylcholine metabolite, is associated with atherosclerotic burden. *J Am Coll Cardiol* 2016;**67**: 2620–2628.
77. Li XS, Obeid S, Klingenberg R, Gencer B, Mach F, Räber L, Windecker S, Rodondi N, Nanchen D, Muller O, Miranda MX, Matter CM, Wu Y, Li L, Wang Z, Alamri HS, Gogonea V, Chung Y-M, Tang WHW, Hazen SL, Lüscher TF. Gut microbiota-dependent trimethylamine N-oxide in acute coronary syndromes: a prognostic marker for incident cardiovascular events beyond traditional risk factors. *Eur Heart J* 2017;**38**:814–824.
78. Tuñón J, Bäck M, Badimón L, Bochaton-Piallat M-L, Cariou B, Daemen MJ, Egido J, Evans PC, Francis SE, Ketelhuth DF, Lutgens E, Matter CM, Monaco C, Steffens S, Stroes E, Vindis C, Weber C, Hofer IE; ESC Working Group on Atherosclerosis and Vascular Biology. Interplay between hypercholesterolaemia and inflammation in atherosclerosis: translating experimental targets into clinical practice. *Eur J Prev Cardiol* 2018;**25**:948–955.
79. Lacy M, Atzler D, Liu R, de Winther M, Weber C, Lutgens E. Interactions between dyslipidemia and the immune system and their relevance as putative therapeutic targets in atherosclerosis. *Pharmacol Ther* 2019;**193**:50–62.
80. Lee SJ, Thien Quach CH, Jung K-H, Paik J-Y, Lee JH, Park JW, Lee K. H. Oxidized low-density lipoprotein stimulates macrophage 18 F-FDG uptake via hypoxia-inducible factor-1 α activation through Nox2-dependent reactive oxygen species generation. *J Nucl Med* 2014;**55**:1699–1705.
81. Bekkering S, Quintin J, Joosten LAB, van der Meer JWM, Netea MG, Riksen NP. Oxidized low-density lipoprotein induces long-term proinflammatory cytokine production and foam cell formation via epigenetic reprogramming of monocytes. *Arterioscler Thromb Vasc Biol* 2014;**34**:1731–1738.
82. Wang Y, Wang GZ, Rabinovitch PS, Tabas I. Macrophage mitochondrial oxidative stress promotes atherosclerosis and nuclear factor- κ B-mediated inflammation in macrophages. *Circ Res* 2014;**114**:421–433.
83. Christ A, Günther P, Lauterbach MAR, Duewell P, Biswas D, Pelka K, Scholz CJ, Oosting M, Haendler K, Baßler K, Klee K, Schulte-Schrepping J, Ulas T, Moorlag S, Kumar V, Park MH, Joosten LAB, Groh LA, Riksen NP, Espevik T, Schlitzer A, Li Y, Fitzgerald ML, Netea MG, Schultze JL, Latz E. Western diet triggers NLRP3-dependent innate immune reprogramming. *Cell* 2018;**172**:162–175.e14.
84. Bekkering S, Arts RJW, Novakovic B, Kourtzelis I, van der Heijden C, Li Y, Popa CD, Horst Ter R, van Tuijl J, Netea-Maier RT, van de Veerdonk FL, Chavakis T, Joosten LAB, van der Meer JWM, Stunnenberg H, Riksen NP, Netea MG. Metabolic induction of trained immunity through the mevalonate pathway. *Cell* 2018;**172**:135–146.e139.
85. Kim K, Shim D, Lee JS, Zaitsev K, Williams JW, Kim K-W, Jang M-Y, Seok Jang H, Yun TJ, Lee SH, Yoon WK, Prat A, Seidah NG, Choi J, Lee S-P, Yoon S-H, Nam JW, Seong JK, Oh GT, Randolph GJ, Artyomov MN, Cheong C, Choi J-H. Transcriptome analysis reveals nonfoamy rather than foamy plaque macrophages are proinflammatory in atherosclerotic murine models. *Circ Res* 2018;**123**:1127–1142.
86. Spann NJ, Garmire LX, McDonald JG, Myers DS, Milne SB, Shibata N, Reichart D, Fox JN, Shaked I, Heudobler D, Raetz CRH, Wang EW, Kelly SL, Sullards MC, Murphy RC, Merrill AH, Brown HA, Dennis EA, Li AC, Ley K, Tsimikas S, Fahy E, Subramaniam S, Quehenberger O, Russell DW, Glass CK. Regulated accumulation of desmosterol integrates macrophage lipid metabolism and inflammatory responses. *Cell* 2012;**151**:138–152.
87. Baardman J, Verberk SGS, Prange KHM, van Weeghel M, van der Velden S, Ryan DG, Wüst RCI, Neele AE, Speijer D, Denis SW, Witte ME, Houtkooper RH, O'Neill LA, Knatko EV, Dinkova-Kostova AT, Lutgens E, de Winther MPJ, Van den Bossche J. A defective pentose phosphate pathway reduces inflammatory macrophage responses during hypercholesterolemia. *Cell Rep* 2018;**25**:2044–2052.e2045.
88. Tuñón J, Badimón L, Bochaton-Piallat M-L, Cariou B, Daemen MJ, Egido J, Evans PC, Hofer IE, Ketelhuth DFJ, Lutgens E, Matter CM, Monaco C, Steffens S, Stroes E, Vindis C, Weber C, Bäck M. Identifying the anti-inflammatory response to lipid lowering therapy: a position paper from the working group on atherosclerosis and vascular biology of the European Society of Cardiology. *Cardiovasc Res* 2019;**115**:10–19.