



Full length article

Mother-child transfer rates of organohalogen compounds up to four years of age



Mercè Garí^{a,b,*}, Joan O. Grimalt^a, Esther Vizcaino^c, Adonina Tardón^{d,e,f},
Ana Fernández-Somoano^{e,d,f}

^a Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Barcelona, Catalonia, Spain

^b Institute of Computational Biology, Helmholtz Zentrum München for Environmental Health, Neuherberg, Germany

^c Agència de Qualitat i Avaluació Sanitàries de Catalunya (AQUAS), Generalitat de Catalunya, Barcelona, Catalonia, Spain

^d IUOPA Medicine Department, University of Oviedo, Asturias, Spain

^e Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Spain

^f Institute of Health Research of the Principality of Asturias-Foundation for Biosanitary Research of Asturias (ISPA-FINBA), Oviedo, Asturias, Spain

ARTICLE INFO

Handling Editor: Heather Stapleton

Keywords:

Organochlorine compounds

Polybrominated diphenyl ethers

Breastfeeding

Maternal and children body burdens

Concentration gains of organochlorine compounds at four years

Physical-chemical constants influencing in maternal-child transfer

ABSTRACT

Background: Breastfed children absorb persistent and toxic chemicals such as organohalogen compounds (OHCs) during the entire lactation period. Nursing is a main contributor to the burden of these pollutants in the first years of life, hence further assessments on the OHC load processes are needed.

Objectives: To identify the determinants of OHC increase in children at four years of age, considering concentration gains, maternal venous concentrations and breastfeeding time.

Methods: Concentrations of 19 organochlorine compounds (OCs) and 14 polybrominated diphenyl ethers (PBDEs) were analyzed in maternal venous (n = 466), cord blood (n = 326) and children venous serum at four years of age (n = 272) in the Asturias INMA cohort representing the Spanish general population. Data were evaluated considering the socio-demographic and individual information collected at recruitment and follow up surveys, as well as the OHC physical-chemical constants.

Results: The four years-old children concentration gains of the most abundant OHCs showed strong correlations ($R^2 = 0.65-0.93$) with the maternal concentrations during pregnancy and lactation period. The child gain/maternal transfer rates of most correlated pollutants were similar.

Discussion: Between 65 and 93% of the variance of OCs in four years-old children was explained by the maternal concentrations during pregnancy and the lactation period. The compounds with $\log(Kow) > 3.7$ (hydrophobic) showed analogous child gain/maternal transfer rates indicating similar processes of membrane lipid dissolution and passive diffusion from the epithelial cells into the milk. Molecular weight of these pollutants did not influence on these rates. Compounds with low $\log(Koa)$ such as hexachlorobenzene are more volatile and less retained, involving lower child gain/maternal transfer rates. These results may be useful to anticipate the increase of the concentrations of OCs in children using the maternal concentration of these compounds during pregnancy and the planned lactation period and to implement prophylactic measures in mothers with high venous pollutant concentrations.

1. Introduction

The burden of organohalogen compounds (OHCs) in children during the first years of age is conveyed from the mothers by transplacental delivery (Barr et al., 2005; Vizcaino et al., 2014; Wolff et al., 2007) and breastfeeding (Karmaus et al., 2001; Carrizo et al., 2006). These pollutants include organochlorines (OCs) and polybrominated diphenyl ethers (PBDEs), among others.

OCs were used intensively in agriculture and industry for several decades. They include polychlorinated biphenyls (PCBs), DDT and its metabolites, hexachlorocyclohexanes (HCHs), hexachlorobenzene (HCB) and others. Their lipophilic nature and high chemical stability lead to bioaccumulation in food chains and human tissues (Junqué et al., 2017, 2018; Bravo et al., 2019). Due to their adverse effects in humans and the environment, they were progressively banned in many countries since the 1970s and, with a few exceptions, finally banned

* Corresponding author at: Ingolstädter Landstr. 1. 85764 Neuherberg in Munich, Germany.

E-mail address: merce.gari@helmholtz-muenchen.de (M. Garí).

<https://doi.org/10.1016/j.envint.2019.105241>

Received 1 August 2019; Received in revised form 2 October 2019; Accepted 2 October 2019

0160-4120/© 2019 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

worldwide by the Stockholm Convention in 2001 (Stockholm Convention on Persistent Organic Pollutants, 2017).

PBDEs have been used as flame retardants in a wide range of products. They are semi-volatile, environmentally persistent, hydrophobic and biomagnify through the food web (Johnson-Restrepo et al., 2005). They have been found in environmental samples and human fluids and tissues (Antignac et al., 2009; Covaci and Voorspoels, 2005; Hites, 2004; Jin et al., 2009; Lunder et al., 2010; Uemura et al., 2010; Vizcaino et al., 2011; Zhu et al., 2009), including newborns (Costa et al., 2016; Vizcaino et al., 2011). There is public health concern for the potential human exposure to these pollutants (Sjödin et al., 2003; Birnbaum and Staskal, 2004), e.g. effects on neurodevelopment (Gascón et al., 2011). Accordingly, production and use of penta- and octa-BDE formulations were banned in the European Union in 2004. Only deca-BDE is still permitted, but the use in electronic applications was banned in Europe in 2008.

Concentrations of these pollutants have been reported in placenta, breastmilk, maternal, cord and newborn blood serum (Ando et al., 1986; Bravo et al., 2017; Carrizo et al., 2006; Grimalt et al., 2010; Llop et al., 2010; Ribas-Fitó et al., 2003; Sala et al., 2001; Vizcaino et al., 2014). Breastfed children incorporate them during the entire lactation period as they tend to accumulate in fat (Hotham and Hotham, 2015). OHC blood concentrations are higher in breastfed than formula fed children even some years after discontinuation of breastfeeding, e.g. 3.5 years (Lanting et al., 1998), 4 years (Carrizo et al., 2006) and 7 years afterward (Karmaus et al., 2001). Children are more vulnerable to environmental pollutants than adults due to their greater exposure by high consumption of water, food and air in relation to their body weight, the immaturity and weakness of their metabolic system and longer lifetime to develop chronic diseases (Landrigan, 2016; Landrigan and Goldman, 2011). In addition, there are sensitive windows of exposure and development, in specific life stages, such as pregnancy and early childhood, in which the organism is more susceptible or vulnerable to the adverse health effects caused by exposure to environmental pollutants (Markris et al., 2008). These remarks outline the need for improving our knowledge on the main processes determining the intake of OHCs through maternal feeding within the first years of child development.

The present study is devoted to increase our understanding on the processes that determine the accumulation of OHCs in children up to four years of age. Specifically, the study is aimed to identify the influence of breastfeeding once the transplacental transfer contribution is assessed. Accordingly, the concentrations of OHCs in maternal venous serum during pregnancy (12th week), cord blood and venous blood serum at four years of age have been determined in mothers and their children ($n = 272$). Comparison of the maternal and children body burdens allowed us to identify the determinants of increase of these pollutants in children at four years of age, including the dependence of the concentration gain from maternal venous concentrations and breastfeeding time.

2. Methods

2.1. Study population and sampling

In 2004, the INMA Asturias cohort was established by the University of Oviedo in San Agustín Hospital (Avilés, Asturias, North-West Spain). Between 2004 and 2007, 494 pregnant women were recruited and their children were followed-up until 8 years of age (Fernández-Somoano et al., 2011; Fernández-Somoano and Tardon, 2014). Maternal serum during the 12th week of gestation ($n = 466$) and cord blood samples ($n = 326$) were collected and analysed for OHCs (Vizcaino et al., 2014b, 2014a). The health status of these children was followed until four years of age ($n = 453$) and serum samples were collected ($n = 272$). These compounds have now been analysed in venous blood serum of the four years-old children. Written informed consent was

obtained from the parents of each child before the study, which was approved by the Asturias Regional Ethics Committee.

2.2. Analysis of persistent organic pollutants

The analyses of maternal venous, cord blood and children serum were performed with methods described elsewhere (Grimalt et al., 2010; Vizcaino et al., 2009). Pentachlorobenzene (PeCB), hexachlorobenzene (HCB), four hexachlorocyclohexane isomers (α -, β -, δ - and γ -HCH), 4,4'-DDT, 2,4'-DDT, 2,4'-DDD, 4,4'-DDD, 2,4'-DDE, 4,4'-DDE, and seven PCB congeners (28, 52, 101, 118, 138, 153 and 180) were analyzed by gas chromatography with electron capture detection (GC-ECD). Possible coelutions and structural confirmation was performed by GC coupled to mass spectrometry (GC-MS). Fourteen PBDE congeners, including deca-BDE (BDE-209), were analyzed by GC-MS in negative chemical ionization mode (GC-MS-NICI). Limits of detection and quantification, LOD and LOQ, respectively, were calculated as described in Garí and Grimalt (2010). Method uncertainties calculated from repeatability were 0.3–4.5% and 0.9–9.1% of the OC and PBDE measurements, respectively (Vizcaino et al., 2009; Grimalt et al., 2010). The methods performed satisfactorily in repeated international inter-calibration exercises within the Arctic Monitoring and Assessment Program (AMAP Ring Test, 2014).

2.3. Lipid adjustment and body burden calculations

OHC concentrations were expressed in ng/ml (crude concentrations) and in ng/g lipid (lipid adjusted) using the equation described in Phillips et al. (1989). Total cholesterol and triglycerides in maternal, cord and 4-year old children samples were determined using colorimetric enzymatic methods in the General Biochemistry Laboratory of San Agustín Hospital. Total serum burdens (ng) of OHCs were estimated based on total blood volumes (ml) per weight in mothers (~ 65 ml/kg) and children at different times (~ 85 ml/kg at birth and ~ 75 ml/kg at 4 years) and the individual weights. The samples were processed using a Roche Diagnostics COBAS C711.

2.4. Covariates

Information on socio-demographic characteristics, parity, maternal age and pre-pregnancy weight was collected at recruitment. Time of gestation was recorded at birth and type and duration of breastfeeding was obtained in follow-up surveys. Information on children's height and weight was gathered at 4 years post-partum visit questionnaires.

Children BMI categories were based on child growth standards set by the World Health Organization (WHO, 2009), involving recommended BMI: < 16.7 kg/m² (boys) and < 16.8 kg/m² (girls), overweight: 16.7 – 18.2 kg/m² (boys) and 16.8 – 18.5 kg/m² (girls) and obesity: > 18.2 kg/m² (boys) and > 18.5 kg/m² (girls).

Parity at child's birth was categorized as no siblings, one sibling and ≥ 2 siblings. Duration of breastfeeding was divided in 3 categories: < 2 weeks (mainly formula-fed children), short-term (2–16 weeks) and long-term (> 16 weeks).

2.5. Data analysis

Data analysis and graphics were performed using the statistical software R (R Development Core Team, 2018). Medians and geometric means (GMs) with 95% confidence intervals (CIs) were used for descriptive analysis. Statistical differences between covariates were tested for significance using the Chi-square test. Spearman's correlation coefficients (ρ) were used to assess the correlations.

Multivariate linear regression analyses were used to assess the association of socio-demographic covariates with OHC concentrations. All variables were standardized (centred at zero and scaled to two standard deviations) for inclusion in the model (Gelman, 2008). The

Table 1
Socio-demographic characteristics of the study population.

	Children included ^a (n = 272)	Children not included (n = 213)	p-value ^b
	n (%)	n (%)	
<i>Children's characteristics</i>			
Sex			0.71
Female	127 (47)	104 (49)	
Male	145 (53)	109 (51)	
BMI			0.40
Recommended weight	173 (64)	99 (71)	
Overweight	64 (24)	26 (18)	
Obesity	32 (12)	15 (11)	
<i>Maternal characteristics</i>			
Age at delivery			0.070
< 30 years	63 (23)	65 (30)	
30–34 years	97 (36)	80 (38)	
> 35 years	112 (41)	68 (32)	
Parity at child's birth			1
Primiparous	166 (61)	130 (61)	
Multiparous	106 (39)	83 (39)	
Breastfeeding			0.023
No (< 2 weeks)	73 (29)	79 (39)	
Short (2–16 weeks)	60 (23)	49 (24)	
Long (> 16 weeks)	123 (48)	73 (36)	
Maternal educational level			0.30
Up to primary school	43 (16)	46 (21)	
Secondary school	126 (46)	91 (41)	
University degree	103 (38)	85 (38)	
Socioeconomic status			0.70
I-II (highest)	64 (24)	45 (21)	
III	54 (20)	48 (23)	
IV-V (lowest)	153 (56)	120 (56)	

^a OHC concentrations available.

^b p-value from Chi-square *t*-test between included and not included children in the OHC determination at age 4 years.

concentrations and breastfeeding duration were transformed into natural logarithms. The model included the following covariates: children's sex, body mass index and breastfeeding duration, maternal age, parity, socio-economic status and maternal educational level. This model was tested for interactions between maternal OHCs * breastfeeding duration. The final model was selected by both AIC (Akaike Information Criteria) and BIC (Bayesian Information Criteria).

3. Results

3.1. Socio-demographic characteristics of the studied population

The characteristics of the population are described in Table 1. Forty-seven percent of the children were girls. About 64% of the children had recommended weight, 24% were overweight and 12% obese. Only 23% of the mothers were younger than 30 years, and two thirds were primiparous (61%). Concerning maternal breastfeeding, one third of the children did not received maternal breastfeeding (< 2 weeks), twenty-three percent had short breastfeeding (2–16 weeks) and almost half (48%) were breastfed for more than four months. A low percentage of mothers (16%) only attended primary school, 46% had a secondary school degree and approximately one third had a university degree. The socio-economic status encompassed a large spectrum of cases including the least affluent social class (IV-V, 56%), the middle affluent one (III; 20%) and in the most affluent levels (I-II, 24%).

3.2. Organohalogen concentrations in four years-old children

The concentrations of the most abundant OHCs in both crude and lipid-adjusted values are shown in Tables S1 and S2. 4,4'-DDE (median 64.1 ng/g lipid) was the most abundant organochlorine compound in the venous serum of the four years-old children and was found above LOQ in all samples analysed. PCB-153, PCB-138 and HCB (medians 23.6 ng/g lipid, 20.4 ng/g lipid and 18.7 ng/g lipid, respectively), were found in more than 96% of the samples analysed. β -HCH, 4,4'-DDT, PCB-118 and PCB-180 were encountered in 70–79% of the samples and their median concentration ranged between 4.3 ng/g lipid and 14.5 ng/g lipid (Table S2). These OC concentrations were much lower than those found in four-year old children from other INMA cohorts and European populations performed some years before the current study (Gascón et al., 2015; Karlsen et al., 2017), which is consistent with a decrease in human OC levels (Jakszyn et al., 2009; Schuhmacher et al., 2009; Thomas et al., 2017).

The most abundant PBDE was BDE-209, with a median concentration of 3.1 μ g/g lipid (detected above LOQ in 32% of the samples). This compound was followed by BDE-28 and BDE-99 (medians 1.7 μ g/g lipid and 1.5 μ g/g lipid, respectively), and found in 75–82% of the samples analysed. BDE-47 (median 0.55 μ g/g lipid) was found above LOQ in 24% of the samples and BDE-153 (median 0.44 μ g/g lipid) was found in a higher percentage, 42% (Table S2). These concentrations were similar to those found in previous European studies (Carrizo et al., 2007; Caspersen et al., 2016), but still lower than those found in children from the US, China or Australia recruited between 2004 and 2013 (Vuong et al., 2017; Eskenazi et al., 2013; Erkin-Cakmak et al., 2015; Xu et al., 2014; Toms et al., 2018). Higher PBDE concentrations in children than adults have been observed in some studies (e.g., Catalonia, recruited in 2002; Garí and Grimalt, 2013; Australia, recruited in 2006–2007; Toms et al., 2009) but in other sites such as Australia (recruited in 2010–2013) and Northern Quebec (recruited in 2006–2010) the trend is the opposite (Toms et al., 2018; Turgeon O'Brien et al., 2019). The current study also found similar or even higher PBDE concentrations in 4-year old children than in their mothers (Table S1).

4. Discussion

4.1. Maternal and children body burdens of the organohalogen compounds

Body burdens of the OHC in mothers, newborns and four years-old children have been calculated from the individual concentrations of these compounds and the weight of each cohort participant (see Methods section for details of body burden calculations). Box plots of the resulting distributions are shown in Fig. 1. As expected, the highest values correspond to the mothers and the lowest to the newborns. The occurrence of OHC in the newborns involves a transplacental transport from mother to foetus as already stated (Vizcaino et al., 2014b). The higher body burdens in four years-old children than in newborns involves an additional intake of these pollutants in this first life period.

In some cases, the body burden increase is small, e.g. DDT, which reflects a low compound incorporation and is consistent with the low presence of this insecticide in the environment and human tissues because of the ban of the Stockholm Convention. The relatively rapid transformation of 4,4'-DDT into 4,4'-DDE and other metabolites in many environmental and biological processes also leads to a depletion of this compound in the absence of recent use.

The Stockholm convention also banned PCBs, HCHs and HCB. In Spain, the use of these OHCs was discontinued in the 80 s. These restrictions likely decreased the exposure of the population to these compounds. However, human intake from environmental and diet sources still occur as observed in recent studies (Martí-Cid et al., 2010; Bosch et al., 2015; Rodríguez-Hernández et al., 2016; Junqué et al., 2017).




Fig. 1. Boxplot of OHC body burden (ng) in pregnant women ($n = 467$), at birth (cord blood, $n = 326$) and children at four years of age ($n = 272$) in the INMA Asturias birth cohort.

4.2. Determinants of organohalogen compounds in children

Univariate analyses between OHC concentrations in four years-old children and several socio-demographic and life-style determinants are shown in Fig. 2. Children sex, place of residence and parity at child's birth did not involve statistically significant concentration differences for any of the compounds analysed. In general, OHC concentrations among obese children were lower than in children with recommended weight or overweight and these differences were statistically significant for total PCBs (Fig. 2).

Breastfeeding duration was positively associated with the venous concentrations of OCs in four years-old children, while no statistically significant differences were found for PBDEs (Fig. 2). Multivariate regression models confirmed that duration of breastfeeding was the main driver of OHC in these children (Fig. 3). Maternal age was also found to be an important determinant of OHC concentrations in children, although with a minor influence (standardized beta-coefficients ranging between -0.25 and 0.25 ; Fig. 3). Maternal educational level and socio-economic status were found to be statistically significantly associated with total PCBs, with higher levels among children whose mothers had a university degree and belonged to the most affluent social class (Fig. 2). This observed higher exposure of the most affluent and higher educated women may respond to distinct dietary habits (Tarasuk et al., 2010) since these groups tend to have a higher proportion of fish and seafood in the diet and fish has been shown to be a preferred source of incorporation of these pollutants (Junqué et al., 2018). However, there are also studies showing lower OC concentrations in the highest educated population (Cerrillo et al., 2006; Cao et al., 2011) or lack of correlation between social class and OC concentrations (Porta et al., 2010). Studies on the influence of socio-economic status and OC exposure in pregnant women have attributed only 1–5% of the concentration variability to this factor (Vrijheid et al., 2012). Overall, these determinants show a maternal influence on the OHC concentrations and burdens even at four years which was likely mediated by breastfeeding.

4.3. Influence of breastfeeding on infant exposure to organohalogen compounds

The present cohort study is one of the few in which the OHC concentrations are available in maternal venous serum during pregnancy, cord blood serum and four years-old venous serum over a high number of cases (Table S1). These comprehensive data allow to evaluate the dependence of the OHC concentration gain at four years from breastfeeding transfer. This gain can be defined with Eq. (1).

$$BG_{ij} = (C_{4ij} * W_{4i} - C_{0ij} * W_{0i}) / W_{4i} \quad (1)$$

in which BG_{ij} is the concentration gain (ng/ml) of pollutant j in children i ; C_{4ij} and C_{0ij} are the concentrations of pollutant j (ng/ml) in children at four years and at birth, respectively; W_{4i} and W_{0i} are the weights (kg) of children i at four years and at birth, respectively.

The breastfeeding transfer of OHC can be related to the maternal concentrations of these compounds and the lactation time. The influence of these two parameters can be described with Eq. (2).

$$BE_{ij} = CM_{ij} * BT_i \quad (2)$$

in which BE_{ij} is the estimation of breastfeeding transfer (ng-month/ml) of pollutant j in children i ; CM_{ij} is the concentration of pollutant j in the mother of children i ; BT_i is the breastfeeding time of children i .

Representation of BG_{ij} vs BE_{ij} for the mother-child pairs with available OHC concentrations ($n = 272$) shows strong correlations for the compounds present in highest abundance in maternal serum, 0.14 – 1.4 ng/ml or 26 – 270 μ g/g lipid (Fig. 4). The correlation coefficients (R^2) of the linear curve fittings range between 0.65 and 0.93 and are statistically significant ($p < 0.0001$). These high coefficients indicate that most of the variability of the concentration gain is related to the breastfeeding transfer. That is, the measured maternal concentration of these pollutants in the 12th week of pregnancy and breastfeeding time.

One interesting feature emerging from these results is the strong similarity of the slopes of these OCs, between 0.056 and 0.062 $\text{ng}_{\text{children}} / (\text{ng}_{\text{mother}} \text{ month})$ (Table 2). This strong uniformity is




Fig. 2. Univariate distributions of geometric means and 95% confidence intervals (ng/g lipid) of the OHC concentrations in four years-old children for several socio-demographic characteristics. Socioeconomic status defined as I-II, and IV-V are the most and least affluent, respectively.

consistent with common transfer mechanisms and delivery rates between mother and infant during breastfeeding. Only HCB shows a significantly different lower slope, $0.028 \text{ ng}_{\text{children}}/(\text{ng}_{\text{mother}} \text{ month})$. Examination of the physical-chemical properties of these compounds, namely the octanol-water and octanol-air coefficients, K_{ow} and K_{oa} , respectively, (Table 2) shows that HCB has a $\log(K_{oa})$ value that is significantly lower, 6.9 at 36°C , than those of the other compounds, 8.3–9.7 (36°C).

In contrast, the $\log(K_{ow})$ values outline β -HCH, 3.7 at 36°C (Table 2), from the more common range 5.4–7.1 at 36°C of the other compounds. Hydrophobic compounds are transferred from maternal

plasma to breast milk by passive diffusion and binding to lipids (Anderson and Sauberman, 2016; Hotham and Hotham, 2015; Quezada and Vafai, 2014). K_{ow} is the main property determining water solubility and therefore the compound distribution between aqueous and organic phases. The similar slopes of β -HCH, 4,4'-DDE and PCBs indicate that compounds with $\log(K_{ow}) \geq 3.7$ (36°C) are hydrophobic enough for efficient dissolution into the lipid membranes and diffusion across the cells interior as to transfer from epithelial cells into the milk.

Small molecular weight also enhances compound excretion into human milk (Sachs et al., 2019). HCB has the smallest molecular weight of the compounds showing strong correlation between organochlorine




Fig. 3. Standardized beta-coefficients from multivariate regression models for several socio-demographic characteristics on the levels of OHCs in children at 4 years of age. Models are adjusted by sex, body mass index, breastfeeding duration, parity, maternal age, socio-economic status and maternal educational level.




Fig. 4. Correlations of the concentration gain of organochlorine compounds in four years-old children (ng/ml) with the maternal transfer (concentration of these compounds during pregnancy and lactation time; ng-month/ml).

Table 2

Constants of the correlations from linear curve fitting of the estimated gains of organochlorine compounds at four years of age vs maternal transfer.

Compound	Kow ^a 25 °C	Kow ^a 36 °C	Koa ^a 25 °C	Koa ^a 36 °C	Slope	Intercept	R ²	Molecular weight
HCB	5.5	5.4	7.4	6.9	0.028	0.027	0.77	284
β-HCH	3.8	3.7	8.9	8.3	0.062	0.025	0.65	291
4,4'-DDE	6.0	5.9	9.3	8.7	0.061	-0.008	0.93	318
PCB138	6.7	6.6	9.7	9.4	0.056	0.06	0.67	361
PCB153	6.8	6.6	9.6	9.1	0.059	0.06	0.79	361
PCB180	7.2	7.1	10.2	9.7	0.062	0.013	0.82	395

^a Calculated from Schecter et al., 1989 and Beyer et al., 2002.

compound gain at four years of age and maternal transfer. The lower slope of this compound in comparison with the others included in Table 2 is therefore not influenced by this property. The distinct behaviour of HCB can be related to the log(Koa). Higher volatility involves lower retention in human fluids, including blood, and the amounts transferred to breast milk are smaller.

The distinct behaviour of HCB is consistent with the observations of a previous study from the INMA cohort in Menorca (Carrizo et al., 2006) in which lower increases of HCB and PeCB were observed in breastfeeders when comparing to the other OCs and the difference was also related to the lower log(Koa) of these two compounds. In this Menorca study, only OC data on serum from cord blood and venous blood at four years-old children was available. The additional data on OC concentrations in maternal venous blood from the INMA Asturias cohort shows that the lower log(Koa) of HCB is specifically relevant for the mother-infant transfer of this pollutant upon breastfeeding, involving lower maternal retention of the more volatile compounds and therefore lower transference to infants.

These results highlight the influence of breastfeeding in the first years of life. Through breastfeeding, women transfer certain pollutants into their newborns, and its duration is directly related to the children's pollutant concentration gain. In this regard, women with higher serum concentrations than expected (e.g. based on reference/exposure limit values, or on biomonitoring equivalents; Steckling et al., 2018) should modify their diet during gestation and lactation, in order to reduce infant's exposure to chemicals (Mead, 2008). Nonetheless, breastfeeding is highly beneficial for both infants and mothers, and even if maternal's pollutant concentrations are high, breastfeeding should be encouraged during the newborn's first 6 months of life (WHO, 2007).

5. Conclusions

The most abundant OHCs in maternal serum (> 0.18 ng/ml) show strong correlations between the concentration gains of these compounds in four years-old children and maternal transfer. Between sixty-five and ninety-three percent of the variance can be explained from the maternal concentrations of these compounds during pregnancy (12th week in this case) and lactation period. The compounds with log (Kow) > 3.7 (hydrophobic) show the same child gain/maternal transfer rates indicating similar processes of dissolution into the lipid membranes and diffusion across the cells interior to pass from the epithelial cells into the milk. Compounds with low log(Koa) are more volatile and less retained, involving lower child gain/maternal transfer rates. These results are useful to anticipate the increase of the concentrations of OHCs in children at the age of four years using the maternal concentration of these compounds during pregnancy and the planned lactation period. This information may be useful for implementation of prophylactic measures (e.g. maternal diet or breastfeeding duration) in cases of mothers with high venous concentrations of these pollutants.

Declaration of Competing Interest

The authors declare they have no actual or potential competing financial interests.

Acknowledgments

The authors would particularly like to thank all the participants for their generous collaboration and the staff from Hospital San Agustín in Aviles for their effort. This study was funded by grants from CIBERESP, FIS-FEDER (PI04/2018, PI09/02311, PI13/02429, PI18/00909), Obra Social Cajastur/Fundación Liberbank and Oviedo University. This paper was also supported by funding from the European Union projects: HEALS (FP7-ENV-2013- 603946), Neurosome (H2020-MSCA-ITN-2017_SEP-210411486) and EDC-MET (H2020-HEALTH/0490-825762).

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2019.105241>.

References

- AMAP, 2014. Ring test. URL www.inspq.qc.ca/en/ctq/eqas/amap/description.
- Anderson, P.O., Sauberman, J.B., 2016. Modeling drug passage into human milk. *Clin. Pharmacol. Ther.* 100 (1), 42–52.
- Ando, M., Saito, H., Wakisaka, I., 1986. Gas chromatographic and mass spectrometric analysis of polychlorinated biphenyls in human placenta and cord blood. *Environ. Res.* 41, 14–22.
- Antignac, J.P., Cariou, R., Zalko, D., Berrebi, A., Cravedi, J.P., Maume, D., Marchand, P., Monteau, F., Riu, A., Andre, F., Le Bizet, B., 2009. Exposure assessment of French women and their newborn to brominated flame retardants: determination of tri- to deca- polybromodiphenylethers (PBDE) in maternal adipose tissue, serum, breast milk and cord serum. *Environ. Pollut.* 157 (1), 163–173.
- Barr, D.B., Wang, R.Y., Needham, L.L., 2005. Biologic monitoring of exposure to environmental chemicals throughout the life stages: Requirements and issues for consideration for the National Children's Study. *Environ. Health Perspect.* 113, 1083–1091.
- Beyer, A., Wania, F., Gouin, T., Mackay, D., Matthies, M., 2002. Selecting internally consistent physicochemical properties of organic compounds. *Environ. Toxicol. Chem.* 21, 941–953.
- Birnbaum, L.S., Staskal, D.F., 2004. Brominated flame retardants: cause for concern? *Environ. Health Perspect.* 112, 9–17.
- Bosch, C., Grimalt, J.O., Fernandez, P., 2015. Enantiomeric fraction and isomeric composition to assess sources of DDT residues in soils. *Chemosphere* 138, 40–46.
- Bravo, N., Hansen, S., Okland, I., Garí, M., Alvarez, M.V., Matiocevič, S., Odland, J.O., Grimalt, J.O., 2017. Influence of maternal and sociodemographic characteristics on the accumulation of organohalogen compounds in Argentinian women. *The EMASAR study. Environ. Res.* 158, 759–767.
- Bravo, N., Grimalt, J.O., Chashchin, M., Chashchin, V.P., Odland, J.O., 2019. Drivers of maternal accumulation of organohalogen pollutants in Arctic areas (Chukotka, Russia) and 4,4'-DDT effects on the newborns. *Environ. Int.* 124, 541–552.
- Cao, L.L., Yan, C.H., Yu, X.D., Tian, Y., Zhao, L., Liu, J.X., Shen, X.M., 2011. Relationship between serum concentrations of polychlorinated biphenyls and organochlorine pesticides and dietary habits of pregnant women in Shanghai. *Sci. Total Environ.* 409, 2997–3002.
- Carrizo, D., Grimalt, J.O., Ribas-Fito, N., Sunyer, J., Torrent, M., 2006. Physical-chemical and maternal determinants of the accumulation of organochlorine compounds in four-year-old children. *Environ. Sci. Technol.* 40, 1420–1426.
- Carrizo, D., Grimalt, J.O., Ribas-Fito, N., Sunyer, J., Torrent, M., 2007. Influence of breastfeeding in the accumulation of polybromodiphenyl ethers during the first years of child growth. *Environ. Sci. Technol.* 41, 4907–4912.
- Caspersen, I.H., Kvale, H.E., Haugen, M., Brantsaeter, A.L., Meltzer, H.M., Alexander, J., Thomsen, C., Froshaug, M., Bremnes, N.M., Broadwell, S.L., Granum, B., Kogevinas, M., Knutsen, H.K., 2016. Determinants of plasma PCB, brominated flame retardants, and organochlorine pesticides in pregnant women and 3 years old children in The Norwegian Mother and Child Cohort Study. *Environ. Res.* 146, 136–144.
- Cerrillo, I., Olea-Serrano, M.F., Ibarluzea, J., Exposito, J., Torne, P., Laguna, J., Pedraza, V., Olea, N., 2006. Environmental and lifestyle factors for organochlorine exposure among women living in Southern Spain. *Chemosphere* 62, 1917–1924.
- Costa, O., López-Espinosa, M.J., Vizcaino, E., Murcia, M., Iñiguez, C., Navarrete-Muñoz, E.M., Grimalt, J.O., Tardón, A., Ballester, F., Fernández-Somoano, A., 2016. Dietary and Household Sources of Prenatal Exposure to Polybrominated Diphenyl Ethers (PBDEs) in the INMA Birth Cohort (Spain). *Environ. Sci. Technol.* 50 (11), 5935–5944.
- Covaci, A., Voorspoels, S., 2005. Optimization of the determination of polybrominated diphenyl ethers in human serum using solid-phase extraction and gas chromatography-electron capture negative ionization mass spectrometry. *J. Chromatogr. B* 827, 216–223.

- Erkin-Cakmak, A., Harley, K.G., Chevrier, J., Bradman, A., Kogut, K., Eskenazi, B., 2015. In utero and childhood polybrominated diphenyl ether exposures and body mass at age 7 years: the CHAMACOS study. *Environ. Health Perspect.* 123 (6), 636–642.
- Eskenazi, B., Chevrier, J., Rauch, S.A., Kogut, K., Harley, K.G., Johnson, C., Trujillo, C., Sjödin, A., Bradman, A., 2013. In utero and childhood polybrominated diphenyl ether (PBDE) exposures and neurodevelopment in the CHAMACOS study. *Environ. Health Perspect.* 121 (2), 257–262.
- Fernández-Somoano, A., Estarlich, M., Ballester, F., Fernández-Patier, R., Aguirre-Alfaro, A., Herce-Garraleta, M.D., Tardón, A., 2011. Outdoor NO₂ and benzene exposure in the INMA (Environment and Childhood) Asturias cohort (Spain). *Atmos. Environ.* 45 (29), 5240–5246.
- Fernández-Somoano, A., Tardón, A., 2014. Socioeconomic status and exposure to outdoor NO₂ and benzene in the Asturias INMA birth cohort, Spain. *J. Epidemiol. Community Health* 68 (1), 29–36.
- Garí, M., Grimalt, J.O., 2010. Use of proficiency testing materials for the calculation of detection and quantification limits in the analysis of organochlorine compounds in human serum. *Anal. Bioanal. Chem.* 397, 1383–1387.
- Garí, M., Grimalt, J.O., 2013. Inverse age-dependent accumulation of decabromodiphenyl ether and other PBDEs in serum from a general adult population. *Environ. Int.* 54, 119–127.
- Gascón, M., Vrijheid, M., Martínez, D., Forns, J., Grimalt, J.O., et al., 2011. Effects of pre and postnatal exposure to low levels of polybromodiphenyl ethers on neurodevelopment and thyroid hormone levels at 4 years of age. *Environ. Int.* 37, 605–611.
- Gascón, M., Vrijheid, M., Garí, M., Fort, M., Grimalt, J.O., Martínez, D., Torrent, M., Guxens, M., Sunyer, J., 2015. Temporal trends in concentrations and total serum burdens of organochlorine compounds from birth until adolescence and the role of breastfeeding. *Environ. Int.* 74, 144–151.
- Gelman, A., 2008. Scaling regression inputs by dividing by two standard deviations. *Stat. Med.* 27 (15), 2865–2873.
- Grimalt, J.O., Howsam, M., Carrizo, D., Otero, R., De Marchi, M.R.R., Vizcaino, E., 2010. Integrated analysis of halogenated organic pollutants in sub-millilitre volumes of venous and umbilical cord blood sera. *Anal. Bioanal. Chem.* 396, 2265–2272.
- Hites, R.A., 2004. Polybrominated diphenyl ethers in the environment and in people: a metaanalysis of concentrations. *Environ. Sci. Technol.* 38, 945–956.
- Hotham, N., Hotham, E., 2015. Drugs in breastfeeding. *Aust. Prescr.* 38 (5), 156–159.
- Jakszyn, P., Goñi, F., Etzeandia, A., Vives, A., Millán, E., López, R., Amiano, P., Ardanaz, E., Barricarte, A., Chirlaque, M.D., Dorronsoro, M., Larrañaga, N., Martínez, C., Navarro, C., Rodríguez, L., Sánchez, M.J., Tormo, M.J., González, C.A., Agudo, A., 2009. Serum levels of organochlorine pesticides in healthy adults from five regions of Spain. *Chemosphere* 76 (11), 1518–1524.
- Jin, J., Wang, Y., Yang, C., Hu, J., Liu, W., et al., 2009. Polybrominated diphenyl ethers in the serum and breast milk of the resident population from production area, China. *Environ. Int.* 35, 1048–1052.
- Johnson-Restrepo, B., Kannan, K., Rapaport, D.P., Rodan, B.D., 2005. Polybrominated diphenyl ethers and polychlorinated biphenyls in human adipose tissue from New York. *Environ. Sci. Technol.* 39, 5177–5182.
- Junqué, E., Garí, M., Arce, A., Torrent, M., Sunyer, J., Grimalt, J.O., 2017. Integrated assessment of infant exposure to persistent organic pollutants and mercury via dietary intake in a central western Mediterranean site (Menorca Island). *Environ. Res.* 156, 714–724.
- Junqué, E., Garí, M., Lull, R.M., Grimalt, J.O., 2018. Drivers of the accumulation of mercury and organochlorine pollutants in Mediterranean lean fish and dietary significance. *Sci. Total Environ.* 634, 170–180.
- Karlsen, M., Grandjean, P., Weihe, P., Steuerwald, U., Oulhote, Y., Valvi, D., 2017. Early-life exposures to persistent organic pollutants in relation to overweight in preschool children. *Reprod. Toxicol.* 68, 145–153.
- Karmaus, W., DeKoning, E.P., Kruse, H., Witten, J., Osius, N., 2001. Early childhood determinants of organochlorine concentrations in school-aged children. *Pediatr. Res.* 50 (3), 331–336.
- Landrigan, P.J., 2016. Children's environmental health: a brief history. *Acad. Pediatr.* 16, 1–9.
- Landrigan, P.J., Goldman, L.R., 2011. Children's vulnerability to toxic chemicals: a challenge and opportunity to strengthen health and environmental policy. *Health Aff. (Millwood)* 30, 842–850.
- Lanting, C.I., Patandin, S., Fidler, V., Weisglas-Kuperus, N., Sauer, P.J.J., et al., 1998. Neurological condition in 42-month-old children in relation to pre- and postnatal exposure to polychlorinated biphenyls and dioxins. *Early Hum. Dev.* 50, 283–292.
- Llop, S., Ballester, F., Vizcaino, E., Murcia, M., Lopez-Espinosa, M.J., Rebagliato, M., Vioque, J., Marco, A., Grimalt, J.O., 2010. Concentrations and determinants of organochlorine levels among pregnant women in Eastern Spain. *Sci. Total Environ.* 408, 5758–5767.
- Lunder, S., Hovander, L., Athanassiadis, I., Bergman, A., 2010. Significantly higher polybrominated diphenyl ether levels in young U.S. children than in their mothers. *Environ. Sci. Technol.* 44, 5256–5262.
- Markris, S.L., Thompson, C.M., Euling, S.Y., Selevan, S.G., Sonawane, B., 2008. A life-stage-specific approach to hazard and dose-response characterization for children's health risk assessment. *Birth Defects Res. B Dev. Reprod. Toxicol.* 83, 530–546.
- Martí-Cid, R., Huertas, D., Nadal, M., Linares, V., Schuhmacher, M., Grimalt, J.O., Domingo, J.L., 2010. Dietary Exposure to Organochlorine Compounds in Tarragona Province (Catalonia, Spain): Health Risks. *Human Ecol. Risk Assess* 16, 588–602.
- Mead, M.N., 2008. Contaminants in Human Milk: Weighing the Risks against the Benefits of Breastfeeding. *Environ. Health Perspect.* 116 (10), A426–A434.
- Phillips, D.L., Pirkle, J.L., Burse, V.W., Bernert Jr, J.T., Henderson, L.O., Needham, L.L., 1989. Chlorinated hydrocarbon levels in human serum: effects of fasting and feeding. *Archiv. Environ. Contam. Toxicol.* 18 (4), 495–500.
- Porta, M., Gasull, M., Puigdomènech, E., Garí, M., Bosch de Basea, M., Guillén, M., López, T., Bigas, E., Pumarega, J., Llebaria, X., Grimalt, J.O., Tresserras, R., 2010. Distribution of blood concentrations of persistent organic pollutants in a representative sample of the population of Catalonia. *Environ. Int.* 36, 655–664.
- Quezada, A., Vafai, K., 2014. Modeling and analysis of transport in the mammary glands. *Phys. Biol.* 11 (4), 045004.
- R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Ribas-Fitó, N., Cardo, E., Sala, M., Eulàlia de Muga, M., Mazón, C., Verdú, A., Kogevinas, M., Grimalt, J.O., Sunyer, J., 2003. Breastfeeding, exposure to organochlorine compounds, and neurodevelopment in infants. *Pediatrics* 111 (5 Pt 1), e580–e585.
- Rodríguez-Hernández, A., Camacho, M., Henríquez-Hernández, L.A., Boada, L.D., Ruiz-Suárez, N., Valerón, P.F., Almeida, M.G., Zaccaroni, A., Zumbado, M., Luzardo, O.P., 2016. Assessment of human health hazards associated with the dietary exposure to organic and inorganic contaminants through the consumption of fishery products in Spain. *Sci. Total Environ.* 557–558, 808–818.
- Sachs HC and Committee on Drugs, 2013. The transfer of drugs and therapeutics into human breast milk: An update on selected topics. *Am. Acad. Pediatr.* pp. e796–e809.
- Sala, M., Ribas-Fitó, N., Cardo, E., de Muga, M.E., Marco, E., Mazón, C., Verdú, A., Grimalt, J.O., Sunyer, J., 2001. Levels of hexachlorobenzene and other organochlorine compounds in cord blood: exposure across placenta. *Chemosphere* 43 (4–7), 895–901.
- Schecter, A., Furst, P., Kruger, C., Meemker, H.A., Groebel, W., Constable, J.D., 1989. Levels of polychlorinated dibenzofurans, dibenzodioxins, PCBs, DDT and DDE, hexachlorobenzene (HCB), dieldrin, hexachlorocyclohexane and oxychlordane in human breast milk from the United States, Thailand, Vietnam, and Germany. *Chemosphere* 118, 445–454.
- Schuhmacher, M., Kiviranta, H., Ruokojärvi, P., Nadal, M., Domingo, J.L., 2009. Concentrations of PCDD/Fs, PCBs and PBDEs in breast milk of women from Catalonia, Spain: a follow-up study. *Environ. Int.* 35 (3), 607–613.
- Sjödin, A., Patterson, D.G., Bergman, A., 2003. A review on human exposure to brominated flame retardants—particularly polybrominated diphenyl ethers. *Environ. Int.* 29, 829–839.
- Steckling, N., Gotti, A., Bose-O'Reilly, S., Chapizanis, D., Costopoulou, D., De Vocht, F., Garí, M., Grimalt, J.O., Heath, E., Hiscock, R., Jagodic, M., Karakitsios, S.P., Kedikoglou, K., Kosjek, T., Leondiadis, L., Maggos, T., Mazej, D., Polańska, K., Povey, A., Rovira, J., Schoierer, J., Schuhmacher, M., Spiric, Z., Staknjo, A., Stierum, R., Tratnik, J.S., Vassiliadou, I., Annesi-Maesano, I., Horvat, M., Sarigiannis, D.A., 2018. Biomarkers of exposure in environment-wide association studies – Opportunities to decode the exposome using human biomonitoring data. *Environ. Res.* 164, 597–624.
- Stockholm Convention on Persistent Organic Pollutants (POPs), 2017. URL <https://chm.pops.int/TheConvention/ThePOPs/AllPOPs/tabid/2509/Default.aspx>.
- Tarasuk, V., Fitzpatrick, S., Ward, H., 2010. Nutrition inequities in Canada. *Appl. Physiol. Nutr. Metab.* 35, 172–179.
- Thomas, A., Toms, L.M., Harden, F.A., Hobson, P., White, N.M., Mengersen, K.L., Mueller, J.F., 2017. Concentrations of organochlorine pesticides in pooled human serum by age and gender. *Environ. Res.* 154, 10–18.
- Toms, L.M.L., Sjödin, A., Harden, F., Hobson, P., Jones, R., Edenfield, E., Mueller, J.F., 2009. Serum Polybrominated Diphenyl Ether (PBDE) Levels are Higher in Children (2–5 Years of Age) than in Infants and Adults. *Environ. Health Perspect.* 117 (9), 1461–1465.
- Toms, L.M.L., Sjödin, A., Hobson, P., Harden, F.A., Aylward, L.L., Mueller, J.F., 2018. Temporal trends in serum polybrominated diphenyl ether concentrations in the Australian population, 2002–2013. *Environ. Int.* 121, 357–364.
- Turgeon O'Brien, H., Gagné, D., Lauzière, J., Blanchet, R., Vézina, C., Ayotte, P., 2019. Temporal trends of legacy and emerging persistent organic pollutants in Inuit preschoolers from Northern Quebec (Canada). *Int. J. Environ. Health Res.* 1–14.
- Uemura, H., Arisawa, K., Hiyoshi, M., Dakeshita, S., Kitayama, A., et al., 2010. Congener-specific body burden levels and possible determinants of polybrominated diphenyl ethers in the general Japanese population. *Chemosphere* 79, 706–712.
- Vizcaino, E., Arellano, L., Fernandez, P., Grimalt, J.O., 2009. Analysis of whole congener mixtures of polybromodiphenyl ethers by gas chromatography-mass spectrometry in both environmental and biological samples at femtogram levels. *J. Chromatogr. A* 1216 (25), 5045–5051.
- Vizcaino, E., Grimalt, J.O., Fernández-Somoano, A., Tardón, A., 2014b. Transport of persistent organic pollutants across the human placenta. *Environ. Int.* 65, 107–115.
- Vizcaino, E., Grimalt, J.O., Glomstad, B., Fernández-Somoano, A., Tardón, A., 2014a. Gestational weight gain and exposure of newborns to persistent organic pollutants. *Environ. Health Perspect.* 122 (8), 873–879.
- Vizcaino, E., Grimalt, J.O., Lopez-Espinosa, M.-J., Llop, S., Rebagliato, M., Ballester, F., 2011. Polybromodiphenyl ethers in mothers and their newborns from a non-occupationally exposed population (Valencia, Spain). *Environ. Int.* 37, 152–157.
- Vrijheid, M., Martínez, D., Aguilera, I., Ballester, F., Basterrechea, M., 2012. Socioeconomic status and exposure to multiple environmental pollutants during pregnancy: evidence for environmental inequity? *J. Epidemiol. Community Health* 66, 106–113.
- Vuong, A.M., Yolton, K., Xie, C., Webster, G.M., Sjödin, A., Braun, J.M., Dietrich, K.N., Lanphear, B.P., Chen, A., 2017. Childhood polybrominated diphenyl ether (PBDE) exposure and neurobehavior in children at 8 years. *Environ. Res.* 158, 677–684.
- Wolff, M.S., Engel, S., Berkowitz, G., Teitelbaum, S., Siskind, J., Barr, D.B., Wetmur, J., 2007. Prenatal pesticide and PCB exposures and birth outcomes. *Pediatr. Res.* 61, 243–250.
- World Health Organization, 2007. Fourth WHO-Coordinated Survey of Human Milk for Persistent Organic Pollutants in Cooperation with UNEP. Guidelines for Developing a National Protocol. URL <https://www.who.int/foodsafety/chem/POPprotocol.pdf> (Last accessed September 2019).

World Health Organization, United Nations Children's Fund. 2009. WHO child growth standards and the identification of severe acute malnutrition in infants and children. A Joint Statement. URL <https://www.who.int/nutrition/publications/severemalnutrition/9789241598163/en/> (Last accessed November 2018).

Xu, X., Liu, J., Zeng, X., Lu, F., Chen, A., Huo, X., 2014. Elevated serum polybrominated

diphenyl ethers and alteration of thyroid hormones in children from Guiyu, China. *PLoS One* 9 (11), e113699.

Zhu, L., Ma, B., Hites, R.A., 2009. Brominated flame retardants in serum from the general population in northern China. *Environ. Sci. Technol.* 43 (18), 6963–6968.