

Integrative Metabolomic and metallomic analysis in a case control cohort with Parkinson´s disease

Authors: Marianna Lucio^{1*¶}, Desiree Willkommen^{1¶}, Michael Schroeter², Ali Sigaroudi^{3,4}, Philippe Schmitt-Kopplin^{1,5}, Bernhard Michalke¹

¹ Analytische BioGeoChemie, Helmholtz Zentrum München, Neuherberg, Germany

² Uniklinik Köln, Klinik und Poliklinik für Neurologie, Kerpener Str. 62; 50937 Köln

³ Universitätsspital Zürich, Klinik für Klinische Pharmakologie und Toxikologie, Rämistr. 100, 8091 Zürich

⁴ Uniklinik Köln, Institut I für Pharmakologie, Zentrum für Pharmakologie, Gleueler Str. 24, 50931 Köln

⁵ TU Munich, Chair of analytical food chemistry, Science center Weihenstephan, Freising, Germany

¶Equally contributed

* Corresponding author:

Marianna Lucio

marianna.lucio@helmholtz-muenchen.de

Keywords: Data integration, Metabolomics, Metallomics, Block-sPLS-DA, Parkinson´s disease.

Supporting Information

Table S1: Assigned and significantly differentiating compounds of metabolomics investigation.

Table S2: List of metabolites obtained after the feature selection algorithm, published in (Willkommen *et al.*, 2018a)

Figure S1: Illustration of correlation between metallomic and metabolomic data out of the same sample set with respective correlation value.

