

Published in final edited form as:

Cochrane Database Syst Rev. ; (6): CD005653. doi:10.1002/14651858.CD005653.pub2.

Pharmacological interventions for borderline personality disorder

Jutta Stoffers², Birgit A Völm³, Gerta Rücker⁴, Antje Timmer⁵, Nick Huband³, and Klaus Lieb¹

¹Department of Psychiatry and Psychotherapy, University Medical Center Mainz, Mainz, Germany

²Department of Psychiatry and Psychotherapy, Freiburg, & Department of Psychiatry and Psychotherapy, Mainz, Germany

³Section of Forensic Mental Health, Institute of Mental Health, Nottingham, UK

⁴German Cochrane Centre, Department of Medical Biometry and Statistics, Freiburg, Germany

⁵Institute of Epidemiology, Helmholtz Zentrum München Research Center for Health and Environment, München, Germany

Abstract

Background—Drugs are widely used in borderline personality disorder (BPD) treatment, chosen because of properties known from other psychiatric disorders (“off-label use”), mostly targeting affective or impulsive symptom clusters.

Objectives—To assess the effects of drug treatment in BPD patients.

Search methods—We searched bibliographic databases according to the Cochrane Developmental, Psychosocial and Learning Problems Group strategy up to September 2009, reference lists of articles, and contacted researchers in the field.

Copyright © 2010 The Cochrane Collaboration. John Wiley & Sons, Ltd.

Contact address: Klaus Lieb, Department of Psychiatry and Psychotherapy, University Medical Center Mainz, Untere Zahlbacherstr 8, Mainz, D-55131, Germany. klaus.lieb@ukmainz.de.

Editorial group: Cochrane Developmental, Psychosocial and Learning Problems Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 6, 2010.

Review content assessed as up-to-date: 24 January 2010.

CONTRIBUTIONS OF AUTHORS

Jutta Stoffers: wrote protocol and final report, selected studies, obtained papers, extracted data, entered data.

Birgit Völm: helped write the protocol and final report, selected studies, obtained papers, extracted data, entered data.

Gerta Rücker: helped write the protocol and final report, gave statistical support and helped extract data.

Antje Timmer: helped write the protocol, corrected final report.

Nick Huband: helped examine literature search retrievals.

Klaus Lieb: sought funds, helped write the protocol, revised final report.

DECLARATIONS OF INTEREST

We are not aware of any personal, political, academic or financial conflicts. KL was involved in the Schulz 2007 trial by recruiting and assessing participants as one of the 39 study sites. KL and JS have been involved in the process of preparation of the WFSBP guidelines for the treatment of personality disorders (Herpertz 2007). KL has not received any payments from industry since the beginning of 2007 to avoid any kind of conflicts of interest. We are not aware of any secondary or personal interests.

Selection criteria—Randomised studies comparing drug versus placebo, or drug versus drug(s) in BPD patients. Outcomes included total BPD severity, distinct BPD symptom facets according to DSM-IV criteria, associated psychopathology not specific to BPD, attrition and adverse effects.

Data collection and analysis—Two authors selected trials, assessed quality and extracted data, independently.

Main results—Twenty-eight trials involving a total of 1742 trial participants were included. First-generation antipsychotics (flupenthixol decanoate, haloperidol, thiothixene); second-generation antipsychotics (aripirazole, olanzapine, ziprasidone), mood stabilisers (carbamazepine, valproate semisodium, lamotrigine, topiramate), antidepressants (amitriptyline, fluoxetine, fluvoxamine, phenelzine sulfate, mianserin), and dietary supplementation (omega-3 fatty acid) were tested. First-generation antipsychotics were subject to older trials, whereas recent studies focussed on second-generation antipsychotics and mood stabilisers. Data were sparse for individual comparisons, indicating marginal effects for first-generation antipsychotics and antidepressants.

The findings were suggestive in supporting the use of second-generation antipsychotics, mood stabilisers, and omega-3 fatty acids, but require replication, since most effect estimates were based on single studies. The long-term use of these drugs has not been assessed.

Adverse event data were scarce, except for olanzapine. There was a possible increase in self-harming behaviour, significant weight gain, sedation and changes in haemogram parameters with olanzapine. A significant decrease in body weight was observed with topiramate treatment. All drugs were well tolerated in terms of attrition.

Direct drug comparisons comprised two first-generation antipsychotics (loxapine versus chlorpromazine), first-generation antipsychotic against antidepressant (haloperidol versus amitriptyline; haloperidol versus phenelzine sulfate), and second-generation antipsychotic against antidepressant (olanzapine versus fluoxetine). Data indicated better outcomes for phenelzine sulfate but no significant differences in the other comparisons, except olanzapine which showed more weight gain and sedation than fluoxetine. The only trial testing single versus combined drug treatment (olanzapine versus olanzapine plus fluoxetine; fluoxetine versus fluoxetine plus olanzapine) yielded no significant differences in outcomes.

Authors' conclusions—The available evidence indicates some beneficial effects with second-generation antipsychotics, mood stabilisers, and dietary supplementation by omega-3 fatty acids. However, these are mostly based on single study effect estimates. Antidepressants are not widely supported for BPD treatment, but may be helpful in the presence of comorbid conditions. Total BPD severity was not significantly influenced by any drug. No promising results are available for the core BPD symptoms of chronic feelings of emptiness, identity disturbance and abandonment. Conclusions have to be drawn carefully in the light of several limitations of the RCT evidence that constrain applicability to everyday clinical settings (among others, patients' characteristics and duration of interventions and observation periods).

Medical Subject Headings (MeSH)

Antidepressive Agents [therapeutic use]; Antipsychotic Agents [adverse effects; *therapeutic use]; Borderline Personality Disorder [* drug therapy; psychology]; Fatty Acids, Omega-3 [therapeutic use]; Randomized Controlled Trials as Topic

MeSH check words

Humans

BACKGROUND

The disorder is a condition first recognised in the 19th century. The term 'Borderline Personality Disorder' (BPD) was coined by A. Stern describing a condition in the "borderland" between psychosis and neurosis (Stern 1938). Subsequent psychoanalytic contributions (especially that of Kernberg 1975) have reaffirmed this distinction emphasising that the capacity to test reality remains grossly intact but is subject to subtle distortions, especially under stress.

According to current diagnostic criteria, BPD is characterised by a pervasive pattern of instability in affect regulation, impulse control, interpersonal relationships, and self-image. Clinical hallmarks include emotional dysregulation, impulsive aggression, repeated self-injury, and chronic suicidal tendencies (Lieb 2004). Whereas some authors have suggested that it is a variant of affective disorders (Akiskal 2004), others claim only partially overlapping etiologies (Paris 2007). Despite the difficulties in defining the condition, borderline personality disorder is the focus of great interest. Its importance stems from the huge suffering of the persons concerned, functional impairment (Skodol 2002), and from the significant impact it has on mental health services (Zanarini 2004a).

The definition of BPD in the Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV, also DMS-IV-TR; APA 1994; APA 2000a) comprises nine criteria that cover the above features, with a definite diagnosis requiring that five criteria are met, and probable diagnosis requiring four. The competing International Classification of Diseases in its 10th edition (ICD-10) refers to the condition of Emotionally Unstable Personality Disorder (F60.3) of which there is an impulsive type (F60.30) and a borderline type (F60.31) (WHO 1993). The latter essentially overlaps with the DSM-IV definition. A significant problem with this type of polythetic definition is that it is possible for two people to satisfy the criteria and yet have very different personalities. This heterogeneity is a major problem in assessing the impact of an intervention. In addition to the specific BPD criteria, DSM-IV and ICD-10 provide general diagnostic criteria for personality disorders that also must be met.

The prevalence of BPD is estimated to be about 1.5% in the general population (most recent data: Lenzenweger 2007; for a survey of epidemiologic studies see Torgersen 2005), but higher (up to 20%) among psychiatry inpatients, and predominantly diagnosed in women (75%; APA 2000a). There are particular problems in its diagnosis in adolescents and young adults where existential dilemmas may be mistakenly classified as BPD (DSM-IV). BPD commonly co-occurs with mood disorders, substance misuse, eating disorders, post-traumatic stress disorder (PTSD) and is also associated with other personality disorders (McGlashan 2000). Suicidal behaviour is reported to occur in up to 84% of patients with BPD (Soloff 2002), comorbid mood disorders or substance use being the most relevant risk factors for completion (Black 2004).

Although the short to medium-term outcome of BPD is poor - similar to that of schizophrenia - there is some evidence that long term follow-up shows a more favourable course, with remission rates of about 88% within ten years (Zanarini 2007). However, remission here only means that diagnostic criteria are not fulfilled and doesn't indicate the absence of any symptoms. Indeed, whereas acute symptoms such as self-mutilation, help-seeking suicide threats or attempts and impulsivity in most cases decrease with time, affective symptoms reflecting areas of chronic dysphoria, such as chronic feelings of emptiness, intense anger or profound abandonment largely remain (Zanarini 2007). Therefore, the majority of people with BPD still have significant levels of symptoms. Risk factors for a poorer long term outcome are comorbid substance use disorders, PTSD, and anxious cluster disorders (Zanarini 2005; Zanarini 2007), and also a family history of psychiatric disorder (especially mood disorders and substance use disorders), demographic issues, such as older age, longer treatment history, pathological childhood experiences, temperament issues, and adult psychosocial functioning (Zanarini 2007). It is estimated that about 60% to 78% of BPD patients make suicide attempts (Links 2009), but the rate of completed suicides is far less. Zanarini and colleagues found suicide rates of 4% during follow-up of ten years (Zanarini 2007), whereas Stone 1993 reported a suicide rate of 8.5% after 16.5 years. Study estimates of the lifetime risk of suicide among patients with BPD range from 3% to 10% (Links 2009).

The direct costs of BPD are considerable in that many people so affected make major demands on health professionals. The problem of deliberate self-harm is a particular issue in this group (Linehan 1997). In medical settings, people with BPD often present after self-harming behaviour or in suicidal crisis and are treated in emergency settings, often involving repeated psychiatric hospitalizations. Additionally, more than 80% of BPD patients are in individual psychotherapy for at least half of a six year period, and the same number is taking standing medication (Zanarini 2004a). Treatment settings and provisions for BPD patients may vary across different countries. Nevertheless, pharmacological interventions are increasingly being used to treat different facets of the BPD pathology spectrum, such as affective instability, impulsivity, dissociative states, or cognitive-perceptual symptoms. Associated pathology, such as depression, can likewise be the target of psychopharmacological interventions. Therefore, different classes of agents are used in the treatment of BPD patients, such as mood stabilisers, antipsychotics, or antidepressants (Lieb 2004).

In summary, BPD is a condition that has been extensively studied. It has a major impact on health facilities as those affected often present in crisis. Its long-term course leads to improvement but people continue to have considerable problems. The polythetic nature of the diagnosis is likely to lead to heterogeneity making it difficult to assess treatment efficacy.

OBJECTIVES

To evaluate the effects of pharmacological interventions in BPD.

METHODS

Criteria for considering studies for this review

Types of studies—All relevant randomised comparisons testing pharmacological interventions in BPD were included. Likewise, data from randomised cross-over studies up to the point of first cross-over (first period only) were eligible. We excluded outcomes of following periods since carry-over effects of the preceding treatments were likely. Furthermore, since BPD characteristically has no stable course but comprises rapid mood shifts, it seemed inappropriate for subjects to serve as their own controls (i.e. within-subject comparisons). Thus, we decided to use first period data only (Elbourne 2002).

At least 70% of study participants had to have a formal diagnosis of BPD. Studies including BPD patients as a subsample were included as well, if separate data on these patients were available. Studies were eligible if they stated both provider and recipient blinding. The adequacy of relevant arrangements was judged subsequently.

Types of participants—Adult patients with a formal diagnosis of BPD according to DSM criteria (see table below). Since its introduction in 1980, the criteria have only changed marginally.

DSM-III (APA 1980) 301.83 Borderline Personality Disorder	DSM-IV-TR (APA 2000a) 301.83 Borderline Personality Disorder
<i>Diagnostic criterion A (5 of the following are required)</i>	<i>Diagnostic criterion A: A pervasive pattern of instability of interpersonal relationships, self-image, and affects, and marked impulsivity beginning by early adulthood and present in a variety of contexts, as indicated by five (or more) of the following:</i>
(6) intolerance of being alone, e.g., frantic efforts to avoid being alone, depressed when alone	(1) frantic efforts to avoid real or imagined abandonment - note: do not include suicidal or self-mutilating behavior covered in criterion 5
(2) a pattern of unstable and intense interpersonal relationships, e.g., marked shifts of attitude, idealization, devaluation, manipulation (consistently using others for one's own ends)	(2) a pattern of unstable and intense interpersonal relationships characterized by alternating between extremes of idealization and devaluation
(4) identity disturbance manifested by uncertainty about several issues relating to identity, such as self-image, gender identity, long-term goals or career choice, friendship patterns, values, and loyalties, e.g., 'Who am I', 'I feel like I am my sister when I am good'	(3) identity disturbance: markedly and persistently unstable self-image or sense of self
(1) impulsivity or unpredictability in at least two areas that are potentially self-damaging, e.g., spending, sex, substance use, shoplifting, overeating, physically self-damaging acts	(4) impulsivity in at least two areas that are potentially self-damaging (e.g., spending, sex, substance abuse, reckless driving, binge eating) - note: do not include suicidal or self-mutilating behavior covered in criterion 5
(7) physically self-damaging acts, e.g., suicidal gestures, self-mutilation, recurrent accidents or physical fights	(5) recurrent suicidal behavior, gestures, or threats, or self-mutilating behavior
(5) affective instability: marked shifts from normal mood to depression, irritability, or anxiety, usually lasting a few hours and only rarely more than a few days, with a return to normal mood	(6) affective instability due to a marked reactivity of mood (e.g., intense episodic dysphoria, instability, or anxiety usually lasting a few hours and only rarely more than a few days)
(8) chronic feelings of emptiness or boredom	(7) chronic feelings of emptiness
(3) inappropriate, intense anger or lack of control of anger, e.g., frequent displays of temper, constant anger	(8) inappropriate, intense anger or difficulty controlling anger (e.g., frequent displays of temper, constant anger, recurrent physical fights)

DSM-III (APA 1980) 301.83 Borderline Personality Disorder	DSM-IV-TR (APA 2000a) 301.83 Borderline Personality Disorder
	(9) transient, stress-related paranoid ideation or severe dissociative symptoms
<i>Diagnostic criterion B: If under 18, does not meet the criteria for Identity Disorder</i>	

Types of interventions—Any drug or a defined combination of drugs administered on a long-term basis (i.e. not only in case of crisis only) with the intention to treat BPD pathology.

Comparison treatments were classified in four categories:

- placebo;
- active comparator drug;
- combination of drugs;
- combined treatment, i.e. drug plus concomitant psychotherapeutic treatment or counselling.

Types of outcome measures—Outcomes could either be self-rated by patients or interviewer-assessed. Only adequately validated measures were included. Studies were only included if they provided data that could be used for effect size calculation for at least one of the primary or secondary outcomes defined below.

If a trial provided more than one measure for the same outcome construct (e.g. several questionnaires for the assessment of depression) the one most often used in the whole pool of included studies was used for effect size calculation, in order to minimise heterogeneity of outcomes in form and content. If a study reported the data of two assessment instruments that were equally frequently used, two reviewers (JS, BV) discussed the issue and chose the one which was in its content most appropriate for assessing BPD patients. Self-rated measures were preferred.

Primary outcomes

- Overall BPD severity.
- Severity of single BPD criteria according to DSM (avoidance of abandonment, dysfunctional interpersonal patterns, identity disturbance, impulsivity, suicidal ideation, suicidal behaviour, self-mutilating behaviour, affective instability, feelings of emptiness, anger, psychotic paranoid symptoms, dissociative symptoms).

Secondary outcomes

- Depression.
- Anxiety.

- General psychiatric pathology: comprehensive measures.
- Mental health status.
- Attrition.
- Adverse effects.

Search methods for identification of studies

Electronic searches—A qualified librarian searched the following electronic databases:

- CENTRAL (*The Cochrane Library*, 2009, issue 3);
- MEDLINE (January 1966 to 11 September 2009);
- CINAHL (1982 to September 2009);
- EMBASE (1980 to 37th week 2009);
- BIOSIS (1985 to 16 September 2009);
- PsycINFO (1872 to 2nd week September 2009);
- Sociological Abstracts (1963 to September 2009);
- ASSIA (1987 to June 2008);
- WEB OF SCIENCE (1981 to 12 September 2009);
- SIGLE (1980 to April 2006);
- COPAC (September 2009);
- Dissertation Abstracts (September 2009);
- ASSIA (1987 to September 2009).

For detailed search strategies and periods searched, see Appendix 1 to Appendix 13.

The following trial registers were searched via the WHO International Clinical Trials Registry Platform (ICTRP), using “borderline personality disorder” as search term:

- ISRCTN (International Standard Randomised Controlled Trial Number);
- ClinicalTrials.gov;
- ACTR (Australian Clinical Trials Registry).

Searching other resources—Relevant journals such as the *Journal of Personality Disorders*, the *American Journal of Psychiatry*, the *Archives of General Psychiatry*, the *British Journal of Psychiatry* and the *Journal of Clinical Psychiatry* were surveyed on a regular basis. Additionally, researchers in the field were contacted by e-mail and asked for unpublished data. Cross-references from relevant literature were also traced.

Data collection and analysis

Selection of studies—On the basis of publication abstracts, a first estimation of study eligibility was made. After that, the studies were critically appraised by two reviewers (JS, BV), independently, in order to decide about inclusion or exclusion of studies according to the above mentioned criteria. The RefMan bibliography management software was used in order to keep track of appraised trials and decisions. If the reviewers' judgements did not match, a third person (KL) was called upon to finally discuss inclusion or exclusion. To ensure transparency of study selection, flow charts were provided according to the QUOROM statement, showing how many hits had been excluded for a certain reason (Moher 1999).

Data extraction and management—Data were independently extracted by two reviewers (JS, BV). For this purpose, standardized data extraction forms were used, and data were double entered into the Review Manager software. If discrepancies arose that were not due to oversights, they were again resolved by discussion and adjudication by a third person (KL). In case of incomplete data reporting in publications, or where relevant subsample data were lacking, we contacted the study authors for more information.

Assessment of risk of bias in included studies—Again, two reviewers (JS, BV) independently rated the included trials in terms of their risk of bias. A standardized rating form was used in order to judge the probability of different risks of bias. Using The Cochrane Collaboration's tool for assessing risk of bias, the following questions were judged: Was the allocation sequence adequately generated? Was allocation adequately concealed? Was knowledge of the allocated intervention adequately prevented during the study (this question was judged separately for observer- and self-rated outcomes)? Were incomplete outcome data adequately addressed? Are reports of the study free of suggestion of selective outcome reporting? Was the study apparently free of other problems that could put it at a high risk of bias? Relevant text passages were quoted and, if necessary, commented upon. After that, the overall risk of bias was rated either as low (question answered 'Yes') or high (question answered 'No'). If insufficient detail was reported, or sufficient detail was known but the actual risk of bias was unknown, the judgement was 'Unclear'. Both reviewers (JS, BV) tried to reach a concerted estimation taking into account the information available. In case of disagreement, a third person (KL) was called in again.

Measures of treatment effect—Standardized mean differences (SMDs) were calculated on the basis of post-treatment results and follow-up data, respectively. Follow-up data were to be subsumed in 6 month steps. In case the direction of a scale was opposite to most of the other scales, the corresponding mean values were multiplied by -1 to ensure adjusted values.

For some trials, effect sizes could not be calculated as intended, i.e. as SMDs as described above, because relevant information was lacking. However, we decided to include these data by calculating alternative estimates, and discussed the peculiar risk of over- or underestimating the effects.

The following effect sizes were used alternatively:

- **Pre-standardized mean differences (MDs):** The effects were calculated by using the post-treatment means as intended, but the standard deviations (SDs) of pre-treatment means. This may have led to an overestimation of effect sizes, as the pre-SDs are commonly smaller than post-SDs. This kind of effect size had to be used for the Goldberg 1986 outcome data.
- **Standardised mean changes:** The effects were calculated by using the pre-post mean change scores and their SDs. This is also a common method for preparing standardized effect sizes, but these data cannot be pooled with the common SMDs due to statistical assumptions (Higgins 2008). Standardized mean changes were calculated for Bogenschutz 2004; Schulz 2007 (partly) and Zonarini 2007 (partly).
- **Mean change differences:** For some outcomes of Schulz 2007 and Zonarini 2007, data allowed only for the calculation of the differences in mean baseline changes experienced by the two groups. Its standard errors (SE) were derived from the pair-wise P-values of the ANCOVA, as provided in the study reports. This is, therefore, a non-standardized measure reflecting the mere difference in reduction of assessment instrument scores. Both studies used the same assessment instrument.

Effect sizes were preferably calculated on the basis of intention-to-treat (ITT) data. If means and standard deviations from intention-to-treat analysis with missing values replaced were available, we used these data. In other cases we used analysis based on available data. Regarding dichotomous outcomes, the risk ratio (RR) was computed on an intention-to-treat basis. We acted on the conservative assumption that all participants who were lost to post-treatment assessment had an unfavourable outcome, e.g. they had left because the treatment had not been acceptable for them. We specified in the Characteristics of included studies risk of bias tables if continuous data of a certain study referred to the intention-to-treat or per-protocol sample.

All calculations were done using the latest release of the Review Manager software (RevMan 2008).

Unit of analysis issues

Cross-over trials: We planned to include data from randomised cross-over studies up to the point of first cross-over (first period only). We decided not to consider outcomes of following periods due to the likelihood of carry-over effects of the preceding treatment(s).

Repeated observations: We did not plan to combine repeated observations on participants in one meta-analysis. Data from different points of measurement (i.e. post-treatment, catamnestic data of 6-months-steps) were subject to separate analyses. Interim observations were not used.

Studies with multiple treatment groups: If a trial compared more than two intervention groups, all pairwise comparisons were included as long as they were not subject to the same meta-analysis. If, for example, two different doses of a certain drug were tested against placebo, only the one comparison of placebo to the group with the dosage most similar to either recommended dosage standards or (if available) other trials testing this comparison

was included. Thus, we avoided including the same group of participants twice in the same meta-analysis. If the experimental groups received different treatments with regard to contents, such as different drugs or combinations of drugs, and were not subject to the same meta-analysis, we included all comparisons.

Dealing with missing data: Where there was incomplete reporting of outcomes stated as having been assessed, we contacted the study authors. If data were not reported in an immediately usable way but required processing before being analysed, a statistician (GR) was consulted. Results derived from processed data were reported in sensitivity analyses.

Assessment of heterogeneity: Both visual inspection of the graphs and the I^2 statistics (Higgins 2003) were used to investigate statistical heterogeneity within a certain comparison. Besides the I^2 statistic, the number of studies and study characteristics such as duration, dose, and participants were taken into account to judge if heterogeneity was more probable due to clinical, i.e. explainable factors, or to unknown factors. In case of substantial heterogeneity, we made up subgroups, depending on study characteristics such as study size, duration, dose, or participants, and discussed the most apparent sources of heterogeneity.

Assessment of reporting biases: Funnel plots were to be provided for comparisons with sufficient primary studies. However, the poor numbers of study effects per comparison did not allow for constructing interpretable figures.

Data synthesis—If data pooling seemed feasible, the primary studies effects were pooled and their 95% confidence interval (CI) was calculated. A random-effects model was used, as some degree of clinical heterogeneity was present in most cases, though confined by study inclusion criteria and not regarded as preventing from pooling in principle.

As a basic rule, I^2 scores of up to 75% were regarded as indicating possibly substantial, but accountable degrees of heterogeneity permitting statistical pooling. In case of I^2 scores exceeding 75%, we discussed if diversity of specific study characteristics (dose, duration, participants, outcome assessment, size) was likely to cause heterogeneity and tried to investigate this by setting up subgroups, the number of effect estimates permitting. If heterogeneity could not be explained, the estimates were not pooled.

Sensitivity analysis—Sensitivity analyses for the primary outcomes were planned to be performed as follows:

- trials requiring patients to have a certain psychiatric comorbidity in addition to BPD were to be excluded;
- only ITT data based outcomes were to be included.

Given the small numbers of effect estimates per comparison and outcome, we did not conduct sensitivity analyses, as this would only have led to omitting results. Instead, we strived to make all potential shortcomings of methodological quality explicit (see Characteristics of included studies tables and the “Risk of bias in included studies” section of the Description of studies) and to critically discuss all findings.

RESULTS

Description of studies

See: Characteristics of included studies; Characteristics of excluded studies; Characteristics of ongoing studies.

Results of the search

The study searches were re-run several times for updates. Due to overlaps in time periods covered and the use of sensitive search strategies (see Appendix 1 to Appendix 13), a large number of references was retrieved during preparation of this review. Study searches generated 13,972 references, 3723 of which were identified as duplicates. After screening of titles and abstracts of the remaining 10,249 hits, 489 citations merited closer inspection, and the full texts were ordered and scrutinized by two reviewers (JS, BV). Of these, 425 citations were excluded because they did not meet the inclusion criteria. Seven references referred to currently ongoing trials (see Characteristics of ongoing studies). A total of 57 different citations were included, relating to 28 RCTs.

Included studies

Setting of studies/study sample—Included studies were published between the years 1979 and 2009, with 20 of the 28 included trials dating from 2000 or later. The studies were conducted in either the USA (14 studies; Bogenschutz 2004; Frankenburg 2002; Goldberg 1986; Hollander 2001; Leone 1982; Linehan 2008; Reich 2009; Salzman 1995; Simpson 2004; Soloff 1989; Soloff 1993; Zanzarini 2001; Zanzarini 2003; Zanzarini 2004) or in Western European countries (12 studies; 5 in Germany and/or Austria (Loew 2006; Nickel 2004; Nickel 2005; Nickel 2006; Tritt 2005), two each in the UK (Montgomery 1979/82; Montgomery 81/82/83) and Spain (Pascual 2008; Soler 2005), and one each in Belgium (De la Fuente 1994), Ireland (Hallahan 2007) and the Netherlands (Rinne 2002)). There were two international multicentre trials: The Schulz 2007 trial was carried out in 39 study centres located in the USA and Western European countries. The RCT by Zanzarini 2007 took place in 13 study centres in the USA, South America, and Eastern Europe.

Study samples ranged from N = 16 (Hollander 2001) to N = 314 (Schulz 2007) in size. In the Zanzarini 2007 trial, even more patients had been involved altogether but there were three treatment groups, only two of which could be included in this meta-analysis, leaving 301 patients (see Characteristics of included studies). In total, the included studies provided data from 1742 patients.

Characteristics of participants

Demographic data—Most studies were not restricted to any gender, but nine studies included female patients only (Frankenburg 2002; Linehan 2008; Loew 2006; Rinne 2002; Simpson 2004; Tritt 2005; Zanzarini 2001; Zanzarini 2003; Zanzarini 2004). The study of Nickel and colleagues reported the study data of a female (Nickel 2004) and a male sample (Nickel 2005) in separate publications. Patients were at least 18 years of age with the exception of two studies (Hallahan 2007; Nickel 2006) where participants had to be at least

16 years old. The mean participants' age ranged from 21.7 (Nickel 2006) to 38.6 (Hollander 2001) years, with 14 of the 28 studies having a mean age below 30 years.

Treatment settings—Study participants were mostly outpatients. The participants of only one trial were inpatients (De la Fuente 1994), while in two others participants were initially treated as inpatients for a minimum of two and three weeks, respectively (Soloff 1989; Soloff 1993), but could continue as outpatients afterwards.

Five trials dating from before 1990 diagnosed the participants according to DSM-III (Goldberg 1986; Leone 1982; Montgomery 1979/82; Montgomery 81/82/83; Leone 1982; Goldberg 1986; Soloff 1989), three studies used DSM-III-R criteria (De la Fuente 1994; Soloff 1993; Salzman 1995; Soloff 1993). Diagnoses of all 20 remaining studies were based on DSM-IV or DSM-IV-TR.

Psychiatric comorbidity—Most study samples were clearly defined as BPD patients with a formal diagnosis of BPD as the main inclusion criterion (Bogenschutz 2004; De la Fuente 1994; Hollander 2001; Leone 1982; Linehan 2008; Loew 2006; Nickel 2004; Nickel 2005; Nickel 2006; Pascual 2008; Reich 2009; Rinne 2002; Salzman 1995; Schulz 2007; Simpson 2004; Soler 2005; Soloff 1989; Soloff 1993; Tritt 2005; Zanarini 2001; Zanarini 2003; Zanarini 2004; Zanarini 2007). However, there were a few exceptions: The Goldberg 1986 study required patients to have a diagnosis of BPD and/or schizotypal personality disorder (PD). Of the 50 patients included, 17 were diagnosed as having BPD, 13 as having schizotypal PD, and 20 as satisfying both sets of criteria. Hence, 74% of the study sample were BPD patients. The Montgomery 1979/82 and Montgomery 81/82/83 studies included patients admitted to a general hospital after a suicidal act, who had a history of two or more previous documented suicidal acts. BPD patients constituted 76.6% and 78.9%, respectively, of all included participants of the two studies. Similarly, all patients of the Hallahan 2007 trial were recruited from the accident and emergency department, where they had presented acutely with self-harm. Additionally, all had to have a lifetime history of at least one other episode. Of all participants, 71% were diagnosed as having BPD. Only one trial required patients to satisfy another diagnosis besides BPD: All patients of the Frankenburg 2002 study additionally had a bipolar II disorder.

Exclusion criteria—Exclusion criteria varied between studies. Commonly, patients particularly prone to pharmacotoxic effects (i.e. pregnant or breastfeeding women, persons with known allergic reactions or intolerances) were excluded, as were patients with severe somatic illnesses, or neurological disorders (especially seizure disorders). Organic brain syndrome or mental retardation were also listed as exclusions by most studies.

The most common exclusion criteria relating to psychiatric conditions were schizophrenia, bipolar disorders, major depressive disorder and substance related disorders. Patients with any comorbid Axis-I disorder were excluded in two studies (De la Fuente 1994; Salzman 1995), as were patients with any unstable Axis-I disorder in another trial (Soler 2005).

In the remaining 25 trials, patients suffering from schizophrenia were excluded in 15 trials (Bogenschutz 2004; Goldberg 1986; Hallahan 2007; Hollander 2001; Linehan 2008; Loew

2006; Montgomery 1979/82; Montgomery 81/82/83; Nickel 2004; Nickel 2005; Nickel 2006; Pascual 2008; Reich 2009; Schulz 2007; Tritt 2005). Another nine trials specified that even the lifetime diagnosis of schizophrenia was an exclusion criterion (Frankenburg 2002; Reich 2009; Simpson 2004; Soloff 1989; Soloff 1993; Zanarini 2001; Zanarini 2003; Zanarini 2004; Zanarini 2007). Eight studies also excluded patients with current (Bogenschutz 2004; Linehan 2008) or lifetime schizoaffective disorder (Frankenburg 2002; Soloff 1993; Zanarini 2001; Zanarini 2003; Zanarini 2004; Zanarini 2007).

Patients with the diagnosis of any bipolar disorder were excluded in 20 of all 28 included trials. However, Hallahan 2007; Leone 1982; Loew 2006; Montgomery 1979/82; Montgomery 81/82/83; Nickel 2006 and Rinne 2002 did not exclude bipolar patients. Additionally, all patients of the Frankenburg 2002 study sample had a bipolar II disorder as an inclusion criterion (here, patients with bipolar I disorder were excluded). In the Soler 2005 trial, bipolar patients could be included if they were in a stable condition.

Patients with current major depressive disorder were not allowed in the majority of trials (besides the De la Fuente 1994 and Salzman 1995 trials that excluded any Axis-I disorder, and Soler 2005 that excluded any unstable Axis-I disorder): Bogenschutz 2004; Frankenburg 2002; Hollander 2001; Montgomery 1979/82; Montgomery 81/82/83; Nickel 2004; Nickel 2005; Pascual 2008; Schulz 2007; Soloff 1989; Soloff 1993; Tritt 2005; Zanarini 2003; Zanarini 2004; Zanarini 2007). The Goldberg 1986 trial excluded patients with melancholia, and Linehan 2008 excluded patients currently suffering from major depressive disorder with psychotic features.

Another frequent exclusion criterion was substance related disorder. Besides the two aforementioned trials that did not include patients with any Axis-I disorder (De la Fuente 1994; Salzman 1995), and the one trial that excluded patients with any unstable Axis-I condition (Soler 2005), there were ten trials (Bogenschutz 2004; Goldberg 1986; Hallahan 2007; Linehan 2008; Pascual 2008; Reich 2009; Schulz 2007; Simpson 2004; Soloff 1993; Zanarini 2007) that did not include patients who currently satisfied criteria for alcohol or drug dependence. Another eight trials did not even include patients abusing alcohol or drugs at the time of recruitment (Frankenburg 2002; Hollander 2001; Loew 2006; Nickel 2004; Nickel 2005; Tritt 2005; Zanarini 2001; Zanarini 2007). Therefore, only seven of the 28 trials did not state any substance related disorder as hindering patients from entering the trial (Leone 1982; Montgomery 1979/82; Montgomery 81/82/83; Nickel 2006; Rinne 2002; Soloff 1989; Zanarini 2003). Current suicidality was an explicit exclusion criterion in 13 trials (Bogenschutz 2004; Frankenburg 2002; Hollander 2001; Linehan 2008; Loew 2006; Nickel 2004; Nickel 2005; Nickel 2006; Salzman 1995; Tritt 2005; Zanarini 2001; Zanarini 2004; Zanarini 2007). Eleven trials did not explicitly specify suicidality as an exclusion criterion (De la Fuente 1994; Goldberg 1986; Leone 1982; Pascual 2008; Reich 2009; Rinne 2002; Schulz 2007; Simpson 2004; Soloff 1989; Soloff 1993; Zanarini 2003). However, all patients of the Montgomery 1979/82 and Montgomery 81/82/83 trials were recruited following admittance to hospital due to a suicidal act, so these patients can be assumed to have been acutely suicidal when entering the trial.

Severity of illness at baseline—The study participants' baseline severity of illness varied between studies. Seven studies used the Global Assessment scale (GAS; Endicott 1976) to assess individuals' level of functioning, and seven used the Global Assessment of Functioning scale (GAF; APA 1994). Both are 100-point single item rating scales used to rate functioning; on a hypothetical continuum from intact mental health to mental illness. The scale values range from 1, which represents the hypothetically most impaired individual, to 100, the hypothetically healthiest individual (APA 2000b). The GAS and GAF scores ranged from 42.2 to 72.4 and were, therefore, typical for psychiatric outpatients (APA 2000b). The average functioning in one study (Salzman 1995) was located at the lower end of the interval range or 71 to 80 ("slight impairment in functioning"), while the average level of functioning of the Goldberg 1986 study participants was rated between 61 and 70 ("some mild symptoms"). The participants of most studies (De la Fuente 1994; Hollander 2001; Frankenburg 2002; Reich 2009; Schulz 2007; Zanarini 2003; Zanarini 2004; Zanarini 2007) were located in the interval range from 51 to 60, defined as "having moderate symptoms or generally functioning with some difficulty". The samples of four other studies (Linehan 2008; Simpson 2004; Soloff 1989; Soloff 1993) had a lower level of functioning and were rated between 41 and 50, i.e. as having "any serious symptomatology or impairment in functioning that most clinicians would think obviously requires treatment or attention".

The Clinical Global Impressions Severity of Illness Scale (CGI-S; Guy 1976) was used in two trials to estimate participants' severity of illness. This scale covers seven items from 1 "not ill at all" to 7 "among the most extremely ill". Here, the average ratings ranged from 4.2 to 5.14. The average CGI-S ratings of Bogenschutz 2004 (4.3) was closest to item 4, "moderately ill", while the participants of the Soler 2005 trial were rated with an average of 5.14, which fits best with item 5, "markedly ill". A similar estimation was found by Pascual 2008, who used the Clinical Global Impressions Border-line Personality Disorder (CGI-BPD) scale specifically referring to the rating of BPD severity (Perez 2007). These patients had an average CGI-BPD severity of illness of 4.8.

Rinne 2002 provided data specifically concerning BPD severity of illness. On average, the participants met 6.95 (SD = 1.3) DSM BPD criteria. Additionally, the BPDSI (Borderline Personality Disorder Severity Index; Arntz 2003) was used to assess BPD severity. The BPDSI is a fully structured interview measuring the frequency of occurrence of all DSM-IV BPD criteria during the last three months. Each of the nine DSM criteria is operationalized as a subscale, and the sum of all subscales constitutes the BPDSI-total, with a possible range of 0 (no occurrence) to 90 (most severe). A BPDSI total score above 15 signifies BPD pathology (Arntz 2003). For inclusion, a BPDSI total score of 20 was required, and the average baseline mean of all participants was 32.9 (SD = 7.7), indicating moderate severity.

For the samples of the Loew 2006 and Nickel 2006 trials, the t-transformed baseline SCL-90-R global severity index scores (SCL-90-R-GSI) were reported. The SCL-90-R scale is a measure of the status of psychopathology along nine symptom constructs: somatization, obsessive-compulsive symptoms, interpersonal sensitivity, depression, anxiety, hostility, phobic-anxiety, paranoid ideation, and psychoticism. The GSI is essentially a mean of all scores. With t-transformed baseline GSI scores above 70 (Loew 2006: mean GSI at baseline

72.25; Nickel 2006: mean GSI at baseline: 74.7), the participants of these trials can be considered as having “high to very high mental stress” (Franke 2002). For the participants of the Hallahan 2007 trial it is reported that the ‘mean scores for all psychometric instruments [i.e. concerning depression, impulsivity, perceived stress] were well in excess of published normative data’.

Concerning the trials of Nickel 2004; Nickel 2005; Tritt 2005; and Zanarini 2001, there were no psychometric data available relating to the overall severity of illness, psychopathologic burden or impairment. All samples were described as “moderately ill”, and treatment histories were given (Nickel 2004: 10.3% had previously been hospitalized for psychiatric reasons, 58.6% had a history of psychotherapeutic treatment, and 69.0% had received pharmacotherapeutic treatment previously; Nickel 2005: 7.1% had previously been hospitalized for psychiatric reasons, 23.8% had been in psychotherapeutic treatment, and 57.1% had received pharmacotherapeutic treatment; Tritt 2005: 18.5% had previously been hospitalized for psychiatric reasons, 44.4% had been in psychotherapeutic treatment, and 71.1% had received pharmacotherapeutic treatment; Zanarini 2001: 14.3% had previously been hospitalized for psychiatric reasons, 82.1% had been in psychotherapeutic treatment, and 64.3% had received pharmacotherapeutic treatment). Treatment use may depend on availability and health care system specifics, though. However, all three trials excluded patients with bipolar disorders, substance-related disorders, schizophrenia, and especially acutely suicidal patients.

No psychometric data concerning the patients’ severity of illness were available concerning the Montgomery 1979/82 and Montgomery 81/82/83 studies. However, all participants were included after admission to hospital following a suicidal act, and had a history of at least two more documented suicidal acts. Therefore, the severity of illness can be considered very serious.

Concerning the Leone 1982 trial sample, the only information available was that participants had to meet four or more of the diagnostic BPD characteristics described by Gunderson and Kolb, and two of these had to be rated as severe, two as at least moderate. Therefore, patients with rather lower levels of pathology may have been included.

Interventions

Comparisons—Older studies focused mainly on first-generational antipsychotics and antidepressants. Since the mid 1990s, second-generation antipsychotics, mood stabilisers, and selective serotonin reuptake inhibitor (SSRI) antidepressants have gained more attention.

The majority of studies involved two comparison groups. However, there were some studies with three comparison groups: Soloff 1989 tested two active groups, i.e. haloperidol and amitriptyline, against placebo, and Soloff 1993 tested haloperidol and phenelzine sulfate against placebo. Zanarini 2004 compared two active drugs in three different combinations, i.e. fluoxetine alone versus olanzapine alone versus fluoxetine plus olanzapine. Therefore, each of the three comparison groups was involved twice within this review. The different testings belonged to different comparison categories, and were therefore not pooled.

Additionally, Zanarini 2007 also compared three conditions, i.e. olanzapine in two different dosages, to placebo. Since the comparison of each of the two olanzapine groups to placebo would have belonged to the same comparison category, and would have led to pooling dependent data, we decided to include only one of the two olanzapine groups. Therefore, we chose the one olanzapine group with the dosage most closely matching the remaining olanzapine versus placebo comparisons.

In total, included studies comprised the following comparisons.

Active drug versus placebo

1. First-generation antipsychotics: thiothixene (Goldberg 1986), flupenthixol decanoate (Montgomery 1979/82), haloperidol (Soloff 1989; Soloff 1993).
2. Second-generation antipsychotics: aripiprazole (Nickel 2006), olanzapine (Bogenschutz 2004; Linehan 2008; Schulz 2007; Soler 2005; Zanarini 2001; Zanarini 2007), ziprasidone (Pascual 2008).
3. Mood stabilisers: carbamazepine (De la Fuente 1994), valproate semisodium (Frankenburg 2002; Hollander 2001), lamotrigine (Reich 2009; Tritt 2005), topiramate (Loew 2006; Nickel 2004; Nickel 2005).
4. Antidepressants: amitriptyline (Soloff 1989), fluoxetine (Salzman 1995; Simpson 2004), fluvoxamine (Rinne 2002), phenelzine sulfate (Soloff 1993), mianserin (Montgomery 81/82/83).
5. Miscellaneous: omega-3 fatty acids (Hallahan 2007; Zanarini 2003).

Active drug versus active comparator drug

1. First-generation antipsychotic versus first-generation antipsychotic: loxapine versus chlorpromazine (Leone 1982).
2. First-generation antipsychotic versus antidepressant: haloperidol versus amitriptyline (Soloff 1989), haloperidol versus phenelzine sulfate (Soloff 1993).
3. Second-generation antipsychotic versus antidepressant: olanzapine versus fluoxetine (Zanarini 2004).

Active drug versus combination of drugs

1. Second-generation antipsychotic versus second-generation antipsychotic plus antidepressant: olanzapine versus olanzapine plus fluoxetine (Zanarini 2004).
2. Antidepressant versus antidepressant plus second generation antipsychotic: fluoxetine versus fluoxetine plus olanzapine (Zanarini 2004).

Study duration

The intervention times ranged from 32 days to 24 weeks, with a mean duration of 84.0 days (SD = 43.6), i.e. approximately 12 weeks.

Concomitant medication

In 13 of the 28 studies, patients were not taking any concomitant psychotropic medication (Bogenschutz 2004; Frankenburg 2002; Goldberg 1986; Loew 2006; Nickel 2004; Nickel 2005; Nickel 2006; Rinne 2002; Soloff 1993; Tritt 2005; Zanarini 2001; Zanarini 2003; Zanarini 2004). Four studies did not specify whether psychotropic medication was allowed or not (Hollander 2001; Linehan 2008; Montgomery 1979/82; Montgomery 81/82/83). Two trials gave no details on permissible drug treatment, but there was a washout period for tricyclic antidepressants (TCAs) and monoamine oxidase inhibitors (MAOIs) in one (De la Fuente 1994) and a one week placebo run-in, probably without any other psychotropic treatment, in the other (Salzman 1995). Some studies specified permissible drugs that could be taken in order to manage adverse effects, or to address certain symptoms that were not addressed by the study drugs. Mostly, these were drugs with sedative or anxiolytic effects.

In the Leone 1982 study, nighttime sedatives (flurazepam and chloral hydrate) could be taken. In the case of insomnia, participants of the Simpson 2004 study were allowed to take 50 to 100 mg/day of Trazodone. In cases of extrapyramidal reactions, participants of the Soloff 1989 study could take 2 mg/day of biperiden hydrochloride. Patients of both the Pascual 2008 and Soler 2005 studies could continue treatment with benzodiazepines, antidepressants or mood stabilisers, if initiated prior to study inclusion, but doses could not be modified. Participants of the Reich 2009 study were allowed to take one antidepressant but had to have been on a stable dose for at least one month. Patients of the Schulz 2007 and the Zanarini 2007 trials were allowed to take benzodiazepines and hypnotics.

Concomitant psychotherapeutic treatment

Concerning the permission of concomitant psychotherapy, six studies gave no details at all on this (Goldberg 1986; Hollander 2001; Leone 1982; Schulz 2007; Zanarini 2001; Zanarini 2003). In eight trials, psychotherapeutic treatment was an exclusion criterion (Hallahan 2007; Loew 2006; Nickel 2004; Nickel 2005; Nickel 2006; Reich 2009; Rinne 2002; Tritt 2005). Although not cited as a reason for exclusion in two trials, no participants of the Frankenburg 2002 and Zanarini 2004 studies received psychotherapy at the time. In two further studies, very few patients received concomitant psychotherapy (Salzman 1995: two out of 22; Zanarini 2007: 10 out of 415). The Bogenschutz 2004 trial allowed patients to continue psychotherapeutic treatment if initiated more than three months prior to randomization, but there was no specification as to how many subjects actually did. Four trials provided supportive non-specific psychotherapeutic treatment to all their participants (De la Fuente 1994: "supportive atheoretical psychotherapy"; Montgomery 1979/82 and Montgomery 81/82/83: follow-up with support from social workers, community nurses and a crisis intervention team after admission due to a suicidal act; Pascual 2008: weekly two hour, non-specific group psychotherapy). There were three trials (Linehan 2008; Simpson 2004; Soler 2005) in which all participants received specific psychotherapeutic treatment, i.e. Dialectic Behavioural Therapy. All patients of the Soloff 1989 and Soloff 1993 studies started as inpatients for three and two weeks, respectively, but it was not specified whether they received psychotherapeutic treatment during this time and thereafter.

Outcome measures

As a rule, higher scores indicate more severe pathology. There are only two exceptions: GAF and GAS scores (mental health status assessments) are oppositely directed, i.e. higher scores indicate higher or better levels of functioning.

Some trials reported several measures relating to the same outcome, as defined for this review (e.g. for depression there were both Beck Depression Inventory (BDI) and Hamilton Depression Scale (HAM-D) scores available). To avoid an unnecessary inflation of type-I error, only one relevant result out of each study was used for effect size calculation. BPD-specific assessment instruments were first choice for primary outcome assessment. If none was available, the measure most often used in the whole pool of included studies was chosen for effect size calculation, in order to minimise the heterogeneity of outcomes in form and content. If there was no difference in the frequency of use, we chose the measure that we thought was in its contents most adequately reflecting the particular outcome in BPD patients. Self-rated measures were also preferred.

Concerning adverse events, objective data were preferred (i.e. weight increase in kg was used instead of the ratio of patients with perceived weight gain). The ratios of patients experiencing a certain adverse event in each group were only statistically compared if the event occurred more than once in at least one of the two groups. Table 1 (FGAs versus placebo), Table 2; Table 3 and Table 4 (SGAs versus placebo), Table 5; Table 6; and Table 7 (mood stabilisers versus placebos), Table 8 and Table 9 (antidepressants versus placebo), Table 10 (miscellaneous active agents versus placebo); Table 11 (FGAs versus FGAs); Table 12 (FGAs versus antidepressants); Table 13 (SGAs versus antidepressants); Table 14 (SGAs versus SGA+antidepressant) and Table 15 (antidepressants versus antidepressant+SGA) specify the measures the effect sizes were calculated from for each comparison category. If there were several measures available for the same outcome, the reasons for choosing a particular one were indicated.

In the following, a survey of the assessment instruments finally used in the review is given. Measures used in the included studies to assess outcomes that were not relevant to this review are not considered, as are data that were of relevance but could not be used for effect size calculation due to the format of reporting.

Primary outcomes

(1) BPD severity

- (a) Borderline Syndrome Index (BSI): Soloff 1993.
- (b) Clinical Global Impression (CGI) scale for use in borderline personality disorder patients (CGI-BPD), global: Pascual 2008.
- (c) Schedule of Interviewing Schizotypal Personalities (SIB), subscale "borderline score": Goldberg 1986.
- (d) Zanarini Rating Scale for borderline personality disorder (Zan-BPD) total score: Schulz 2007; Reich 2009; Zanarini 2007.

(2) Avoidance of abandonment

- (a) CGI-BPD, subscale “abandonment”: Bogenschutz 2004; Pascual 2008.
- (b) ZAN-BPD, subscale “frantic efforts to avoid abandonment”: Schulz 2007; Zanarini 2007.

(3) Interpersonal problems

- (a) Atypical Depression Inventory, subscale “rejection sensitivity”: Soloff 1993.
- (b) CGI-BPD, subscale “unstable relationships”: Bogenschutz 2004; Pascual 2008.
- (c) Hopkins Symptom Checklist (HSCL), Symptom Checklist-90 (SCL-90) or Symptom Checklist-90-Revised (SCL-90-R), subscale “interpersonal sensitivity”: De la Fuente 1994; Frankenburg 2002; Goldberg 1986; Loew 2006; Nickel 2006; Soloff 1989; Zanarini 2001.
- (d) ZAN-BPD, subscale “unstable interpersonal relationships”: Schulz 2007; Zanarini 2007.

(4) Identity disturbance

- (a) CGI-BPD, subscale “identity disturbance”: Bogenschutz 2004; Pascual 2008.
- (b) ZAN-BPD, subscale “identity disturbance”: Schulz 2007; Zanarini 2007.

(5) Impulsivity

- (a) Acting out-Scale, ratio of patients with status quo or worsened after treatment: De la Fuente 1994.
- (b) Barrett Impulsiveness Scale (BIS): Soloff 1989; Soloff 1993.
- (c) Behavioural reports of numbers of episodes of impulsivity/aggressive behaviour: Soler 2005.
- (d) Borderline Personality Disorder Severity Index (BPDSI), subscale “impulsivity”: Rinne 2002.
- (e) CGI-BPD, subscale “impulsivity”: Bogenschutz 2004; Pascual 2008.
- (f) Modified Overt Aggression Scale (MOAS), total score: Frankenburg 2002; Zanarini 2003.
- (g) Overt Aggression Scale-Modified (OAS-M), subscale “aggression”: Hollander 2001; Simpson 2004.
- (h) State-Trait Anger Expression Inventory (STAXI), subscale “anger out”: Nickel 2004 and Nickel 2005; Nickel 2006; Tritt 2005.
- (i) ZAN-BPD, subscale “impulsivity that are self-damaging”: Reich 2009; Schulz 2007; Zanarini 2007.

(6) Suicidal behaviour/suicidal ideation

- (a) Behavioural reports of numbers of episodes of self-injuring behaviour/suicide attempts: Soler 2005.
- (b) CGI-BPD, subscale “recurrent suicidal ideation”: Bogenschutz 2004; Pascual 2008.
- (c) OAS-M, subscale “suicidality”: Hallahan 2007; Hollander 2001; Simpson 2004.
- (d) OAS-M, subscale “suicidality”: number of patients with high suicidality, i.e. frequent suicide ideation and/or planning or behaviour: Linehan 2008.
- (e) Ratio of patients with suicidal act during treatment: Montgomery 1979/82; Montgomery 81/82/83.
- (f) ZAN-BPD, subscale “suicidal or self-mutilating behaviour”: Schulz 2007; Zhanarini 2007.

(7) Self-injurious behaviour

- (a) Ratio of patients with self-injury during treatment period: Hallahan 2007; Linehan 2008; Nickel 2006.
- (b) OAS-M, subscale “auto aggression”: Simpson 2004.

(8) Affective instability

- (a) BPDSI, subscale “rapid mood shifts”: Rinne 2002.
- (b) CGI-BPD, subscale “affective instability”: Bogenschutz 2004; Pascual 2008.
- (c) ZAN-BPD, subscale “affective instability”: Reich 2009; Schulz 2007; Zhanarini 2007.

(9) Chronic feelings of emptiness

- (a) CGI-BPD, subscale “chronic feelings of emptiness”: Bogenschutz 2004; Pascual 2008.
- (b) ZAN-BPD, subscale “chronic feelings of emptiness”: Schulz 2007; Zhanarini 2007.

(10) Inappropriate anger

- (a) BPDSI, subscale “anger”: Rinne 2002.
- (b) CGI-BPD, subscale “inappropriate anger”: Bogenschutz 2004; Pascual 2008.
- (c) HSCL, SCL-90 or SCL-90-R, subscale “hostility”: De la Fuente 1994; Frankenburg 2002; Goldberg 1986; Loew 2006; Nickel 2006; Soloff 1989; Soloff 1993; Zhanarini 2001.
- (d) OAS-M, subscale “irritability”: Hollander 2001.
- (e) Profile of Mood States (POMS), subscale “anger”: Salzman 1995.

- (f) STAXI, subscale “trait anger”: Nickel 2004 and Nickel 2005; Tritt 2005.
- (g) ZAN-BPD, subscale “intense anger”: Schulz 2007; Zanarini 2007.

(11) Psychotic/paranoid symptoms or dissociation

- (a) CGI-BPD, subscale “transient paranoia or dissociation”: Bogenschutz 2004; Pascual 2008.
- (b) Dissociative Experiences Scale (DES): Simpson 2004.
- (c) HSCL, SCL-90 or SCL-90-R, subscale “paranoid ideation”: De la Fuente 1994; Loew 2006; Nickel 2006; Soloff 1989; Soloff 1993.
- (d) SIB, subscale “suspicious/paranoid”: Goldberg 1986.
- (e) ZAN-BPD, subscale “paranoid ideation of dissociation”: Schulz 2007; Zanarini 2007.

Secondary outcomes

(1) Depression

- (a) BDI: Hallahan 2007; Hollander 2001; Simpson 2004; Soloff 1989; Soloff 1993.
- (b) HAM-D: Linehan 2008; Nickel 2006; Pascual 2008; Salzman 1995; Soler 2005.
- (c) HSCL, SCL-90 or SCL-90-R, subscale “depression”: De la Fuente 1994; Frankenburg 2002; Goldberg 1986; Loew 2006.
- (d) Montgomery-Asberg Depression Rating Scale (MADRS): Schulz 2007; Zanarini 2003; Zanarini 2004; Zanarini 2007.

(2) Anxiety

- (a) Hamilton Anxiety Rating Scale (HARS): Nickel 2006; Pascual 2008; Soler 2005.
- (b) HSCL, SCL-90 or SCL-90-R, subscale “anxiety”: De la Fuente 1994; Loew 2006; Soloff 1989; Soloff 1993; Zanarini 2007.
- (c) State-Trait Anxiety Inventory (STAI), trait score: Simpson 2004.

(3) General psychiatric pathology

- (a) HSCL, SCL-90 or SCL-90-R, Global Severity Index (GSI): De la Fuente 1994; Nickel 2006; Loew 2006; Pascual 2008; Schulz 2007; Soloff 1989; Soloff 1993; Zanarini 2001; Zanarini 2007.

(4) Mental health status

- (a) Clinical Global Impressions Scale (CGI), subscale “severity of illness”: Soler 2005.
- (b) Global Assessment Scale (GAS): De la Fuente 1994; Goldberg 1986; Salzman 1995; Soloff 1989; Soloff 1993.

- (c) Global Assessment of Functioning (GAF): Schulz 2007;Simpson 2004; Zanarini 2007.
- (d) Ratio of patients with Clinical Global Impressions Scale - improvement (CGI-I) score of 3 or more (i.e. minimally improved to very much worse): Hollander 2001.

(5) Attrition

- (a) Ratio of patients lost after randomisation in each group: De la Fuente 1994; Frankenburg 2002; Goldberg 1986; Hallahan 2007;Hollander 2001; Leone 1982; Linehan 2008; Loew 2006; Nickel 2004; Nickel 2005; Pascual 2008; Reich 2009; Rinne 2002; Schulz 2007; Simpson 2004; Soler 2005; Soloff 1989; Soloff 1993; Tritt 2005; Zanarini 2001; Zanarini 2003; Zanarini 2004; Zanarini 2007.

(6) Adverse events - body weight change

- (a) Total weight at endpoint (kg): Loew 2006; Nickel 2004; Nickel 2005; Tritt 2005.
- (b) Baseline to endpoint weight change (kg): Bogenschutz 2004;Frankenburg 2002; Linehan 2008; Schulz 2007; Soler 2005;Zanarini 2001; Zanarini 2004; Zanarini 2007.
- (c) Atypical Depression Inventory (ADS), subscale "weight gain":Soloff 1993.

(7) Patient-reported adverse events (AE)

- (a) Any AE: Leone 1982; Pascual 2008; Schulz 2007; Zanarini 2007.
- (b) Akathisia: Zanarini 2004.
- (c) Anxiety: Schulz 2007.
- (d) Constipation: Zanarini 2001.
- (e) Disturbed attention: Zanarini 2007.
- (f) Dizziness: Loew 2006.
- (g) Dry mouth: Schulz 2007; Zanarini 2007.
- (h) Fainting spells: Leone 1982.
- (i) Fatigue: Loew 2006; Schulz 2007; Zanarini 2007.
- (j) Headache: Loew 2006; Schulz 2007; Zanarini 2007.
- (k) Increased appetite: Schulz 2007; Zanarini 2007.
- (l) Insomnia: Schulz 2007; Zanarini 2007.
- (m) Memory problems: Loew 2006.
- (n) Menstrual pain: Loew 2006.
- (o) Muscle spasms: Leone 1982.

- (p) Nausea: Schulz 2007; Zanmarini 2007.
- (q) Paraesthesia: Loew 2006.
- (r) Restlessness: Leone 1982.
- (s) Sedation: Schulz 2007; Zanmarini 2001; Zanmarini 2004.
- (t) Sleepiness/drowsiness: Leone 1982.
- (u) Somnolence: Schulz 2007; Zanmarini 2007.
- (v) Trouble in concentrating: Loew 2006.

(8) Laboratory values

- (a) Lipids: High-density lipoprotein (HDL) cholesterol baseline to endpoint mean change (mmol/L): Zanmarini 2007.
- (b) Lipids: Low-density lipoprotein (LDL) cholesterol baseline to endpoint mean change (mmol/L): Schulz 2007.
- (c) Lipids: total cholesterol baseline to endpoint mean change (mmol/L): Schulz 2007; Zanmarini 2007.
- (d) Lipids: triglycerides, fasting, baseline to endpoint mean change (mmol/L): Zanmarini 2007.
- (e) Liver function: gamma-glutamyl transferase (GGT) baseline to endpoint mean change Units per litre (U/L): Zanmarini 2007.
- (f) Liver function: Alanine transaminase (ALT)/serum glutamic pyruvic transaminase (SGPT) baseline to endpoint mean change (U/L): Schulz 2007; Zanmarini 2007.
- (g) Liver function: Aspartate transaminase (AST)/serum glutamic oxaloacetic transaminase (SGOT) baseline to endpoint mean change (U/L): Schulz 2007; Zanmarini 2007.
- (h) Liver function: total bilirubin baseline to endpoint mean change ($\mu\text{mol/L}$): Schulz 2007.
- (i) Liver function: direct bilirubin baseline to endpoint mean change ($\mu\text{mol/L}$): Schulz 2007.
- (j) Prolactin: baseline to endpoint mean change ($\mu\text{g/L}$): Schulz 2007; Zanmarini 2007.
- (k) Blood values: leukocyte count baseline to endpoint mean change (GI/L): Zanmarini 2007.
- (l) Blood values: monocytes baseline to endpoint mean change (GI/L): Zanmarini 2007.
- (m) Blood values: neutrophils, segmented, baseline to endpoint mean change (GI/L): Zanmarini 2007.

- (n) Blood values: platelet count baseline to endpoint mean change (GI/L): Zanarini 2007.

Risk of bias in included studies

The assessment of the risk of bias caused several problems, mainly because about one third of trials dated from before publication of the CONSORT statement, and may, therefore, have paid less attention to reporting all relevant issues. However, we tried to be consistent in judging methodological quality throughout all included trials, old or not, which may have resulted in a more 'liberal' judgment.

The judgments for each single study can be found in the Characteristics of included studies tables, and are summarised in Figure 1 and Figure 2.

Allocation

All included trials stated treatment allocation as "randomised". Some trials (Frankenburg 2002; Hallahan 2007; Linehan 2008; Reich 2009; Zanarini 2001) reported the use of a randomised number sequence. Participants of the Simpson 2004 trial were randomised "blocked on the presence of a diagnosis of major depressive disorder or post-traumatic stress disorder (PTSD)", which seems justifiable in the light of an overall small sample size. In the Pascual 2008 trial, allocation was carried out "in blocks of four generated using the SPSS software". The Schulz 2007 and Zanarini 2007 trials were both carried out in parallel multicentre studies by sponsorship of Eli Lilly and Company. The publications only make mention of the use of a randomisation code. However, as one of the reviewers (KL) had been involved at one of the study centres, we know that randomisation was carried out centrally, and investigators were strictly kept blinded to the patients' allocation. These trials were rated 'Yes' with regard to adequacy of sequence generation.

Loew 2006; Nickel 2004; Nickel 2005; Nickel 2006 and Tritt 2005 specified that randomisation had been performed confidentially by the clinic administration, but there were no further details of how this was actually done. Leone 1982 stated that "subjects [...] were selected randomly", but in the light of the identical numbers of men and women in the two groups, the use of some matching procedure seems probable. All remaining trials were only described as having used a randomisation procedure, without giving further details. Thus, it remains unclear if sequence generation was adequate or not.

The actual concealment of allocation was judged adequate for twelve trials where relevant details were given, such as involvement of a third, independent person to disperse medication or to adjust dosages (especially in case of agents with very peculiar adverse effects that could disclose treatment allocation to the clinician) or the use of numbered tablet boxes (Frankenburg 2002; Hallahan 2007; Linehan 2008; Loew 2006; Nickel 2004; Nickel 2005; Nickel 2006; Schulz 2007; Soloff 1989; Tritt 2005; Zanarini 2001; Zanarini 2007). In the Zanarini 2004 trial, the actual numbers of participants in each group were not concordant with the intended group sizes (45 participants should be allocated "in equal numbers" to three groups, but the group sizes differed in an irreproducible way). Hollander 2001 stated that "although the planned patient assignment ratio was 2:1 [...], the ratio was actually 3:1". Here, allocation seems not to have been conducted adequately. For the remaining 14 trials,

no information was given how adequate allocation concealment was ensured, but there were also no indications for inadequate concealment (Bogenschutz 2004; De la Fuente 1994; Goldberg 1986; Leone 1982; Montgomery 1979/82; Montgomery 81/82/83; Pascual 2008; Reich 2009; Rinne 2002; Salzman 1995; Simpson 2004; Soler 2005; Soloff 1993; Zanarini 2003).

Blinding

Self-rated outcomes—All trials were stated as “double-blind” by their authors. In cases where details were given to assure that patients were kept blind, e.g. by using opaque capsules, blinding was judged as adequate. The majority of trials either did so, or there was no risk of bias since there were no self-rated outcomes assessed (Bogenschutz 2004; Frankenburg 2002; Goldberg 1986; Hallahan 2007; Leone 1982; Linehan 2008; Loew 2006; Montgomery 1979/82; Nickel 2004; Nickel 2005; Nickel 2006; Reich 2009; Salzman 1995; Schulz 2007; Soloff 1993; Tritt 2005; Zanarini 2001; Zanarini 2004; Zanarini 2007). The remaining nine trials gave no details, but there were also no indications for non-blindness of participants (e.g. by possibly experiencing very peculiar adverse effects, or by joining the same therapy groups as other participants; Bogenschutz 2004; De la Fuente 1994; Hollander 2001; Montgomery 81/82/83; Pascual 2008; Rinne 2002; Simpson 2004; Soler 2005; Soloff 1989; Zanarini 2003).

Blinding of outcome assessors—The majority of trials reported that outcome observers were blinded or did not use observer-rated outcomes, so the risk of bias was rated as improbable in this regard (De la Fuente 1994; Goldberg 1986; Hallahan 2007; Hollander 2001; Linehan 2008; Loew 2006; Nickel 2004; Nickel 2005; Reich 2009; Salzman 1995; Schulz 2007; Simpson 2004; Soloff 1989; Soloff 1993; Tritt 2005; Zanarini 2001; Zanarini 2004; Zanarini 2007). For the remaining trials, it was not apparent if the person who actually assessed outcomes was blinded, and the risk of bias was judged unclear (Bogenschutz 2004; Frankenburg 2002; Leone 1982; Montgomery 1979/82; Montgomery 81/82/83; Nickel 2006; Pascual 2008; Rinne 2002; Soler 2005; Zanarini 2003).

Incomplete outcome data—Incomplete outcome data were rated as adequately handled for the trials of Goldberg 1986; Leone 1982; Montgomery 1979/82; Montgomery 81/82/83; Nickel 2006; Rinne 2002; Soloff 1989; Soloff 1993. In these cases, only data referring to the intention-to-treat (ITT) sample were used. Mostly, a last-observation-carried-forward (LOCF) approach was used in primary studies. This item was also judged ‘Yes’ if the primary study reported on completers only but drop-outs could be imputed ex post as having the negative outcome for the purpose of this review.

The risk of bias due to inadequate handling of incomplete outcome data was judged ‘unclear’ for studies that used a LOCF approach but had a total drop-out of more than 20% of the initial sample (Bogenschutz 2004; Frankenburg 2002; Hallahan 2007; Linehan 2008; Soler 2005; Zanarini 2001; Zanarini 2007). Two trials used a LOCF approach, but it was not clear how the trial participants were chosen out of eligible subjects (Loew 2006; Tritt 2005). For another three trials it was not clear if continuous data referred to the ITT or completer samples (De la Fuente 1994; Reich 2009; Schulz 2007). The risk of bias due to incomplete

outcome data was therefore judged 'unclear'. The item was also judged 'unclear' for studies that reported on completers only, but the overall dropout rate did not exceed 10%, and reasons for dropping out were specified, not related to treatments and balanced across groups. This was the case for Nickel 2004; Nickel 2005; Salzman 1995; Zamarini 2003; Zamarini 2004.

One trial that both had high drop-out rates (i.e. more than 10%) and excluded non-completers from analyses were judged 'No', i.e. as having a high risk of bias due to incomplete outcome data (Simpson 2004). Two trials with very high attrition rates (i.e. more than 50%) plus unclear selection of study participants out of eligible patients were judged 'No' as well (Hollander 2001; Pascual 2008).

Selective reporting—For the majority of cases no study protocols were available, so there was not enough information to judge if selective reporting was present or not. These trials were rated as 'Unclear' in terms of being biased due to selective reporting (Bogenschutz 2004; De la Fuente 1994; Frankenburg 2002; Goldberg 1986; Hallahan 2007; Hollander 2001; Leone 1982; Linehan 2008; Loew 2006; Montgomery 1979/82; Montgomery 81/82/83; Nickel 2004; Nickel 2005; Nickel 2006; Rinne 2002; Salzman 1995; Simpson 2004; Soloff 1989; Soloff 1993; Zamarini 2003; Zamarini 2004). Four studies that protocols were available for with no major differences of final reporting to the pre-specified way were judged 'Yes', i.e. as having a low risk of bias with this regard (Pascual 2008; Reich 2009; Soler 2005; Zamarini 2007). In one case reported outcomes and the study protocol differed (Schulz 2007), for another study the authors said they only reported significant findings (Zamarini 2001), and a third one provided data from one assessment instrument only, but it seems implausible that in such a complex trial only one assessment instrument had been used (Tritt 2005). These three trials were rated 'No'.

Other potential sources of bias

Carry-over effects from previous pharmacological treatment: To avoid carry-over effects from additional psychotropic medication, concomitant psychotropic treatment was not allowed during the experimental treatment, and, in the main part, a washout-phase or placebo run-in preceded the experimental phase. Internal validity was judged as not threatened by concomitant medication for the trials of Bogenschutz 2004; De la Fuente 1994; Frankenburg 2002; Goldberg 1986; Loew 2006; Nickel 2004; Nickel 2005; Nickel 2006; Rinne 2002; Salzman 1995; Simpson 2004; Soloff 1989; Soloff 1993; Tritt 2005; Zamarini 2001; Zamarini 2003; Zamarini 2004.

The risk of bias due to co-medication was judged unclear for 10 studies because of the following reasons: no details were given whether concomitant psychotropic drug use was allowed or not, or if there was a drug washout (Hollander 2001; Linehan 2008; Montgomery 1979/82; Montgomery 81/82/83; Zamarini 2007); participants were allowed to continue previous psychotropic treatment if initiated prior to study participation (Pascual 2008; Reich 2009; Soler 2005); participants were allowed to take sedatives/hypnotics concomitantly (Leone 1982; Schulz 2007). For the case of Hallahan 2007, bias seemed to be probable, as

concomitant medication was allowed without restrictions, and changes could also be made anytime.

Bias due to sponsoring: Two studies (Soloff 1989; Soloff 1993) declared financial support solely from national non-profit organisations. Another study (Tritt 2005) claimed that there was no funding at all. Hallahan 2007 explicitly declared that the active preparation and placebo were provided by a certain company, but that it was not otherwise involved in the study. These four trials were rated as having a low risk of bias due to sponsoring.

For six studies (Bogenschutz 2004; Frankenburg 2002; Leone 1982; Linehan 2008; Reich 2009; Zanarini 2004) the authors declared support by pharmaceutical companies, seven studies were supported in part by pharmaceutical companies (Hollander 2001; Pascual 2008; Rinne 2002; Simpson 2004; Soler 2005; Zanarini 2001; Zanarini 2003). Another two studies were sponsored by a pharmaceutical company, and the company's trial reports were used in this review (Schulz 2007; Zanarini 2007). These 15 studies were rated 'No' in terms of bias to sponsoring being unlikely.

No sufficient information about funding and sponsoring was available for the remaining nine studies (De la Fuente 1994; Goldberg 1986; Loew 2006; Montgomery 1979/82; Montgomery 81/82/83; Nickel 2004; Nickel 2005; Nickel 2006; Salzman 1995). These were rated 'unclear'.

In summary, 14 out of 28 included trials were at least partly supported by pharmaceutical companies, with no further specification of the companies' roles in conducting and evaluating. For these, bias due to sponsoring cannot be ruled out.

Effects of interventions

Generally, SMDs with a negative value indicate a greater reduction of pathology by the first treatment in line (mostly: verum treatment) in contrast to the alternate treatment (mostly: placebo). Should the opposite be the case, i.e. positive values favour the first mentioned treatment, this will be indicated.

Risk ratios (RRs) with a value lower than one indicate that the risk of a certain event in the first treatment (mostly: active agent) group is lower than that in the comparison treatment (mostly: placebo) group.

For a survey of all outcomes and assessment instruments, see the Description of studies/ Outcome measures section. In addition, tables are provided showing per comparison which assessment instruments were used for assessment of the results that the final effect estimates are based upon, and in case several measures were available for a certain outcome, why the definite one was chosen (Table 1 to Table 10).

1. Drug versus placebo comparisons—For corresponding analyses of drug versus placebo comparisons, refer to Analysis 15.1 to Analysis 65.1.

Primary outcomes

1.1 BPD severity: There were two single study estimates for first-generation antipsychotics, one comparing haloperidol to placebo, the other one thiothixene. Both indicated less favourable results for the groups receiving first-generation antipsychotics (haloperidol: $N = 58$, 1 RCT, SMD 0.30, 95% confidence interval (CI) -0.22 to 0.82 ; thiothixene: $N = 50$, 1 RCT, SMD calculated on basis of post-means and pre-SD 0.28, 95% CI -0.28 to 0.83).

Two large RCTs assessed the impact of olanzapine treatment on BPD severity. The pooled SMDs, based on change scores, indicated olanzapine treated patients to be slightly better-off, but not significantly ($N = 596$, 2 RCTs, SMD calculated on basis of changes scores -0.15 , 95% CI -0.41 to 0.10 , $I^2 = 60\%$). For ziprasidone, data also indicated better results for verum treated patients, but the effect was not significant ($N = 60$, 1 RCT, SMD -0.47 , 95% CI -0.98 to 0.05).

Data for mood stabiliser treatment were provided by one RCT ($N = 27$) that tested lamotrigine. There was a non-significant effect estimate of moderate size (SMD calculated on basis of change scores -0.43 , 95% CI -1.20 to 0.34).

For treatment with antidepressants, only one RCT provided data for BPD severity. Here, the group with phenelzine sulfate treatment had slightly better results, but the difference was, again, not significant ($N = 62$, 1 RCT, SMD -0.15 , 95% CI -0.65 to 0.35).

In summary, none of the investigated agents (i.e. first- and second-generation antipsychotics, one MAOI antidepressant) yielded a significant effect on overall BPD severity.

1.2 Avoidance of abandonment: Data were available for second-generation antipsychotics only. There was almost no difference between ziprasidone and placebo treated patients ($N = 60$, 1 RCT, SMD -0.08 , 95% CI -0.58 to 0.43) and neither did data indicate a substantial impact for olanzapine treatment ($N = 631$, 3 RCTs, SMD calculated on basis of change scores -0.01 , 95% CI -0.22 to 0.21 , $I^2 = 35\%$).

In summary, the data did not suggest substantial effects of second-generation antipsychotics for this outcome. The outcome was not assessed for any other agent.

1.3 Interpersonal problems: First- and second-generation antipsychotics, mood stabilisers and antidepressants were investigated with regard to possible amelioration of interpersonal problems (see Figure 3 for SMDs, and Analysis 3.2 to Analysis 3.4 for additional effect sizes).

As can be seen, most estimates favoured drug treatment, with exception of phenelzine sulfate, for which less favourable results were reported. Significant effects were found for the second-generation antipsychotic aripiprazole (SMD -0.77 , $N = 52$, 1 RCT, 95% CI -1.33 to -0.20) and the mood stabilisers valproate semisodium (SMD -1.04 , $N = 30$, 1 RCT, 95% CI -1.85 to -0.23) and topiramate (SMD -0.91 , $N = 56$, 1 RCT, 95% CI -1.46 to -0.35). All significant effects were derived from one single study each.

In summary, there were significant effects of medium to large size for aripiprazole, valproate semisodium, and topiramate, but all were based on single studies only.

1.4 Identity disturbance: The pooled mean change SD for olanzapine was -0.06 ($N = 631$, 3 RCTs, 95% CI -0.21 to 0.10 , $I^2 = 0\%$). The single study estimate (SMD) for ziprasidone was -0.38 ($N = 60$, 1 RCT, 95% CI -0.90 to 0.13).

In summary, data for this outcome were only available for the second-generation antipsychotics olanzapine and ziprasidone, with no significant results.

1.5 Impulsivity: Impulsivity had been assessed in trials investigating first- and second-generation antipsychotics, mood stabilisers, antidepressants, and omega-3 fatty acids. SMDs are displayed in Figure 4, additional effect sizes were calculated for olanzapine and carbamazepine (see Analysis 5.2 to Analysis 5.4).

Again, most findings were based on single study estimates. Large, significant effects were found for the second-generation antipsychotic aripiprazole ($N = 52$, 1 RCT, SMD -1.84 , 95% CI -2.49 to -1.18), and the mood stabilisers lamotrigine (two RCTs the estimates of which could not be pooled: $N = 27$, 1 RCT, SMD -1.62 , 95% CI -2.54 to -0.69 ; $N = 27$, 1 RCT, SMD on basis of baseline to post mean changes -1.41 , 95% CI -2.27 to -0.55) and topiramate ($N = 71$, 2 RCTs, SMD -3.36 , 95% CI -4.44 to -2.27 , $I^2 = 51\%$). Available data indicated no beneficial effects for the first-generation antipsychotic haloperidol, the second-generation antipsychotics olanzapine and ziprasidone, the mood stabiliser valproate semisodium, the antidepressants amitriptyline, fluoxetine, fluvoxamine, and phenelzine sulfate, or omega-3 fatty acids.

In summary, data consistently indicated significant beneficial effects for mood stabilisers, and the second-generation antipsychotic aripiprazole. The direction of study estimates indicates no beneficial effects for first-generation antipsychotics and antidepressants.

1.6 Suicidal ideation: Usable data concerning the effect of the second-generation antipsychotic olanzapine on suicidal ideation were provided by four RCTs (Bogenschutz 2004; Linehan 2008; Schulz 2007; Zanarini 2007). Due to different formats of data reporting, only two of these could be pooled, and several kinds of effect sizes had to be calculated. There was a small effect for one RCT in terms of mean change difference ($N = 291$, 1 RCT, MCD -0.10 , 95% CI -0.20 to -0.00). However, the remaining three RCTs indicated more suicidal ideation in olanzapine treated patients, resulting in one nonsignificant effect (RR of having high suicidality scores 1.20 , $N = 24$, 1 RCT, 95% CI 0.50 to 2.88), and even another significant one: the pooled mean change SD of the remaining two trials was 0.29 ($N = 340$, 2 RCTs, 95% CI 0.07 to 0.50 , $I^2 = 0\%$). For ziprasidone, another second-generation antipsychotic, there was a single study estimate of SMD -0.27 ($N = 60$, 1 RCT, 95% CI -0.78 to 0.23), indicating a tendency of better outcomes in verum than placebo treated patients.

For mood stabilisers and antidepressants, there were two single estimates of small studies available. Both tended to suggest a worse outcome following drug treatment, but neither was

significant (valproate semisodium: SMD 0.52, N = 16, 1 RCT, 95% CI -0.63 to 1.67; fluoxetine: SMD 0.44, N = 20, 1 RCT, 95% CI -0.46 to 1.33).

The impact of omega-3 fatty acids on suicidal ideation was assessed by one RCT. There, significantly less patients who had received omega-3 fatty acids reported at least slight or more severe suicidal tendencies (RR 0.52, N = 49, 1 RCT, 95% CI 0.28 to 0.95).

In summary, the findings indicate that drug treatment may not only have no substantial effect of decreasing suicidal ideation but may even result in worsening of suicidal ideation, or at least in less favourable outcomes, as compared to placebo treatment. However, this estimation is only based on single study findings for valproate semisodium and fluoxetine. For olanzapine, several estimates are available, with one small significant effect in favour of olanzapine and a medium significant effect against it, and yet another study supporting this tendency. There was a significant beneficial effect for omega-3 fatty acids as reported by one study.

1.7 Suicidal behaviour: There was a significant single study estimate for the reduction of suicidal behaviour by the first-generation antipsychotic flupenthixol decanoate (RR of suicidal behaviour 0.49, N = 37, 1 RCT, 95% CI 0.26 to 0.92).

Another RCT of olanzapine assessed the frequency of suicidal episodes. Again, olanzapine treated patients had unfavourable results as compared to placebo, resulting in a non-significant SMD of 0.15 (N = 60, 1 RCT, 95% CI -0.36 to 0.65). No significant effect was found for the antidepressant mianserin sulfate (RR of suicidal behaviour 1.00, N = 58, 1 RCT, 95% CI 0.71 to 1.41).

In summary, there was a significant reduction in suicidal behaviour during flupenthixol decanoate treatment, a first-generation antipsychotic given as a long acting depot. Additionally, there was another study effect supporting by its direction the possible unfavourable effects of olanzapine for self-damaging tendencies in general, as previously seen for suicidality (cf. to 1.6 Suicidal ideation, above). The prevalence of suicidal behaviour was reported to be lower in mianserin treated patients, but not significantly.

1.8 Self-mutilating behaviour: There were two single study estimates for the second-generation antipsychotics aripiprazole and olanzapine, respectively. Both were non-significant but had opposite directions. Data indicated that patients treated with aripiprazole were less likely to engage in self-mutilating behaviour (RR 0.29, N = 52, 1 RCT, 95% CI 0.07 to 1.25), whereas olanzapine treated patients were not (RR 1.20, N = 24, 1 RCT, 95% CI 0.50 to 2.88). A comparable effect size was found for omega-3 fatty acids by one study (RR 1.23, N = 49, 1 RCT, 95% CI 0.51 to 2.97).

For the SSRI antidepressant fluoxetine, a SMD of 0.03 was found (N = 20, 1 RCT, 95% CI -0.85 to 0.92), indicating almost no difference between experimental and control group.

In summary, none of the available single study estimates yielded a significant effect. However, the possibility of more self-damaging behaviour in general under olanzapine treatment was, again, fortified (cf. to 1.6 Suicidal ideation and 1.7 Suicidal behaviour). Self-

mutilating behaviour also occurred more often under omega-3 fatty acid supplementation as compared to placebo.

1.9 Affective instability: There was a significant decrease in affective instability by olanzapine treatment (mean change SD -0.16 , $N = 631$, 3 RCTs, 95% CI -0.32 to -0.01 , $I^2 = 0\%$), but the effect was small in size. Another small but non-significant effect was found by one RCT for the second-generation antipsychotic ziprasidone (SMD -0.10 , $N = 60$, 1 RCT, 95% CI -0.61 to 0.41).

One trial indicated a medium to large effect of the mood stabiliser lamotrigine (mean change SD -0.61 , $N = 27$, 95% CI -1.39 to 0.17) and another trial showed a moderate to large effect of fluvoxamine, (SMD -0.64 , $N = 38$, 1 RCT, 95% CI -1.30 to 0.01). In summary, data indicated a significant (but small) effect of olanzapine in ameliorating affective instability, but no substantial effect for ziprasidone. Available data suggest that both lamotrigine and fluvoxamine may also be effective, but there are only single study effect estimates in each case with possibly too low power to detect statistical significance.

1.10 Chronic feelings of emptiness: This outcome was only assessed in RCTs of second-generation antipsychotics, i.e. olanzapine and ziprasidone. For olanzapine, there was a minimal non-significant difference between olanzapine and placebo in terms of mean change SDs (-0.03 , $N = 631$, 3 RCTs, 95% CI -0.22 to 0.16 , $I^2 = 23\%$). Ziprasidone treated patients felt slightly worse compared to patients who had been given placebo (SMD 0.18 , $N = 60$, 1 RCT, 95% CI -0.32 to 0.69).

In summary, the evidence available for this outcome suggests no substantial effect of second-generation antipsychotics for this outcome.

1.11 Anger: SMDs are provided in Figure 5. For additional effect sizes, see Analysis 11.2 and Analysis 11.3.

There was a significant effect for haloperidol treatment (SMD -0.46 , $N = 114$, 2 RCTs, 95% CI -0.84 to -0.09 , $I^2 = 0\%$). Another first-generation antipsychotic, thiothixene, yielded no significant effect, as investigated by one RCT (SMD on basis of post-means and pre-SDs -0.07 , $N = 50$, 1 RCT, 95% CI -0.63 to 0.48).

Usable data were also available for the second-generation antipsychotics aripiprazole, olanzapine, and ziprasidone. There was a large, significant effect for aripiprazole (SMD -1.14 , $N = 52$, 1 RCT, 95% CI -1.73 to -0.55), and another significant effect for olanzapine (mean change SD -0.27 , $N = 631$, 3 RCTs, 95% CI -0.43 to -0.12 , $I^2 = 0\%$). For ziprasidone, data suggested no beneficial effect (SMD 0.08 , $N = 60$, 1 RCT, 95% CI -0.43 to 0.58). For mood stabilisers, data indicated significant beneficial effects for any agent investigated here, with the exception of carbamazepine, where there was a positive but non-significant effect. Two RCTs tested valproate semisodium, but we did not pool the study estimates due to considerable heterogeneity ($I^2 = 78\%$). Both RCTs indicated better results for their experimental groups as compared to placebo, but the difference was only significant in one case (Hollander 2001: SMD -1.83 , $N = 16$, 1 RCT, 95% CI -3.17 to

–0.48). There was another large effect for lamotrigine treatment (SMD –1.69, $N = 27$, 1 RCT, 95% CI –2.62 to –0.75). For topiramate, there were three RCTs available, two including women only, and one men. Each of the three study estimates favoured topiramate treatment significantly, but the size of effects varied. Therefore, all three estimates were considerably heterogeneous ($I^2 = 93\%$), and we decided not to pool them. Instead, the two female samples were pooled, yielding a large overall effect estimate of SMD –3.00 ($N = 85$, 2 RCTs, 95% CI –3.64 to –2.36, $I^2 = 0\%$). The effect of topiramate in the remaining male sample was smaller, but also significant (SMD –0.65, 1 RCT, 95% CI –1.27 to –0.03).

There were no significant effects for antidepressant treatment, i.e. the TCA amitriptyline, the SSRIs fluoxetine and fluvoxamine, and the MAOI phenelzine sulfate. Each estimate was based on one single study, though. Effect sizes were small to moderate in size (SMD –0.26 for amitriptyline to –0.65 for fluoxetine, see Figure 5).

In summary, data were available for first- and second-generation antipsychotics, mood stabilisers, and antidepressants. Significant effects were found for mood stabilisers (topiramate, valproate semisodium, lamotrigine) and second-generation antipsychotics (aripiprazole, olanzapine).

1.12 Psychotic symptoms: Findings indicated no significant beneficial effects for first-generation antipsychotics, mood stabilisers or antidepressants. With exception of haloperidol, all estimates were derived from single studies. However, all suggest better results for the experimental groups (see Figure 6 and Analysis 12.3), except of one trial of thiothixene (see Analysis 12.2).

There were significant effects for the second-generation antipsychotics aripiprazole (SMD –1.05, $N = 52$, 1 RCT, 95% CI –1.64 to –0.47) and olanzapine (mean change SD –0.18, $N = 631$, 3 RCTs, 95% CI –0.34 to –0.03, $I^2 = 0\%$), but not for ziprasidone (see Figure 6 and Analysis 12.3).

In summary, data indicated significant benefits for second-generation antipsychotics only, i.e. for aripiprazole and olanzapine.

1.13 Dissociation: This outcome was only assessed by one RCT investigating the SSRI antidepressant fluoxetine. The study estimate indicated unfavourable results for fluoxetine treated patients, but the effect was not significant (SMD 0.42, $N = 20$, 1 RCT, 95% CI –0.47 to 1.32).

In summary, data for treatment of dissociative symptoms are scarce. Available data suggest that the antidepressant fluoxetine may not be beneficial in this regard.

Secondary outcomes

1.14 Depression: SMDs are provided in Figure 7. For additional effect sizes concerning thiothixene and olanzapine, see Analysis 14.2 to Analysis 14.4.

No significant effects were found for the first-generation antipsychotics haloperidol and thiothixene.

There was a large significant effect for the second-generation antipsychotic aripiprazole (SMD -1.25 , $N = 52$, 1 RCT, 95% CI -1.85 to -0.65). No significant effects were found for olanzapine or ziprasidone.

Another significant effect was found for the mood stabiliser valproate semisodium (SMD -0.66 , $N = 46$, 2 RCTs, 95% CI -1.31 to -0.01 , $I^2 = 0\%$). Single study estimates indicated better results for carbamazepine and topiramate as compared to placebo, but none yielded a significant effect.

Among antidepressant agents, a significant effect was only found for the TCA amitriptyline (SMD -0.59 , $N = 57$, 1 RCT, 95% CI -1.12 to -0.06). For phenelzine sulfate, a MAOI, the direction of effect pointed to better outcomes for the experimental group as well, but not to a significant effect. The pooled estimate for the SSRI fluoxetine, however, indicated worse results for fluoxetine treated patients as compared to placebo (SMD 0.12 , $N = 42$, 2 RCTs, 95% CI -1.13 to 1.36 , $I^2 = 74\%$).

Omega-3 fatty acids were found to have beneficial effects by two trials. A non-significant yet favourable difference between the active treatment and placebo of SMD -0.34 ($N = 30$, 1 RCT, 95% CI -1.11 to 0.42) was found by one RCT. This finding was supported by another RCT reporting a significantly lower risk of nonresponding in terms of a 50% reduction of depressive pathology if having received omega-3 fatty acids (RR 0.48 , $N = 49$, 1 RCT, 95% CI 0.28 to 0.81).

In summary, agents of different classes of drugs were found to be effective in the treatment of depression (second-generation antipsychotic aripiprazole, mood stabiliser valproate semisodium, TCA amitriptyline, omega-3 fatty acids).

1.15 Anxiety: SMDs are given in Figure 8, for additional effect size calculations see Analysis 15.2.

No significant beneficial effects were found for the first-generation antipsychotic haloperidol or the antidepressant agents amitriptyline, fluoxetine, or phenelzine sulfate. Of the second-generation antipsychotics, aripiprazole was found to be significantly beneficial (SMD -0.73 , $N = 52$, 1 RCT, 95% CI -1.29 to -0.17) as was olanzapine (mean change difference -0.22 , $N = 274$, 1 RCT, 95% CI -0.41 to -0.03).

Additionally, a large significant effect was found for topiramate (SMD -1.40 , $N = 56$, 1 RCT, 95% CI -1.99 to -0.81). Data indicated favourable results for carbamazepine treatment as well, with an effect of medium size, but this was not significant (SMD -0.51 , $N = 19$, 1 RCT, 95% CI -1.43 to 0.41).

In summary, significant effects were found for second-generation antipsychotics (aripiprazole and olanzapine), and the mood stabiliser topiramate. These estimates are single study findings only.

1.16 General psychiatric pathology: SMDs are given in Figure 9, for additional effect size calculations see Analysis 16.2.

No significant beneficial effects were found for the first-generation antipsychotic haloperidol or the antidepressant agents amitriptyline and phenelzine sulfate. Of second-generation antipsychotics, aripiprazole was found to be significantly beneficial with a large effect of SMD -1.27 ($N = 52$, 1 RCT, 95% CI -1.87 to -0.67), but not so for olanzapine and ziprasidone. For these, the direction of effect did favour drug treatment though.

Additionally, another large significant effect was found for topiramate (SMD -1.19 , $N = 56$, 1 RCT, 95% CI -1.76 to -0.61). Data indicated favourable results for carbamazepine treatment as well, with an effect of medium size, but this was not significant (SMD -0.57 , $N = 19$, 1 RCT, 95% CI -1.49 to 0.36).

Small and non-significant effects were found for the antidepressant agents amitriptyline and phenelzine sulfate.

In summary, significant effects were found for the second-generation antipsychotic aripiprazole and the mood stabiliser topiramate. These estimates were derived from single studies each.

1.17 Mental health status: SMDs are given in Figure 10, for additional effect size calculations see Analysis 17.2 to Analysis 17.4. For this outcome, positive values of effect sizes indicate an amelioration, i.e. an increase of functioning by drug treatment.

Data were available for first-generation antipsychotics haloperidol and thiothixene, the second-generation antipsychotic olanzapine, the mood stabilisers carbamazepine and valproate semisodium, and the antidepressant agents amitriptyline, fluoxetine, and phenelzine sulfate. The effect sizes were small to medium in size but none was significant.

In summary, available data do not suggest a significant increase of overall functioning by any of the investigated drugs.

1.18 Attrition: Overall tolerability was assessed in terms of the risk of not completing the study per protocol. Risk ratios of leaving the study early are given in Figure 11.

Attrition did not differ significantly between experimental and control groups for any other drug versus placebo comparison. Lower drop-out rates for active drug treatment as compared to placebo were found for olanzapine, valproate semisodium, lamotrigine, topiramate, amitriptyline, fluvoxamine, phenelzine sulfate, and omega-3 fatty acids. No usable data of attrition were available for the comparison of aripiprazole versus placebo.

In summary, available data indicated that none of the active drugs was less well tolerated than placebo.

1.19 Adverse effects: Adverse effects outcomes will be reported separately by drug classes.

1.19.1 First-generation antipsychotics: There was a non-significant effect of haloperidol reducing body weight (SMD -0.18 , $N = 58$, 1 RCT, 95% CI -0.70 to 0.34). For thiothixene or flupenthixol decanoate treatment, no adverse effects data were reported.

1.19.2 Second-generation antipsychotics: No usable data were available for aripiprazole treatment. Detailed data were available for olanzapine. There was a significant effect of weight gain (SMD 1.05, N = 752, 6 RCTs, 95% CI 0.90 to 1.20, $I^2 = 0\%$). The ratio of study participants reporting any adverse event was not significantly increased among olanzapine treated patients as compared to placebo (RR 1.13, N = 615, 2 RCTs, 95% CI 1.00 to 1.28, $I^2 = 0\%$), but single events were: increased appetite was significantly more often reported in olanzapine groups (RR 2.78, N = 615, 2 RCTs, 95% CI 1.75 to 4.34, $I^2 = 0\%$) as was somnolence (RR 2.97, N = 215, 2 RCTs, 95% CI 1.75 to 5.03, $I^2 = 0\%$) and mouth-dryness (RR 2.24, N = 615, 2 RCTs, 95% CI 1.08 to 4.67, $I^2 = 0\%$). Sedation had been assessed by two trials (Schulz 2007; Zanarini 2001), but we decided not to pool the two estimates because of considerable statistical heterogeneity ($I^2 = 82\%$), that may have been due to the different observation periods (12 weeks versus 24 weeks) and sample sizes (N = 314 versus N = 28). Schulz 2007 reported sedation as significantly more often experienced by olanzapine treated patients (RR 9.23, N = 314, 1 RCT, 95% CI 2.18 to 39.12), while Zanarini 2001's findings supported the direction of effect (RR 1.26, N = 27, 1 RCT, 95% CI 0.44 to 3.66). The following adverse events were also reported but not found to occur significantly more often under olanzapine treatment: headache (RR 0.91, N = 615, 2 RCTs, 95% CI 0.43 to 1.92, $I^2 = 67\%$), disturbed attention (RR 11.37, N = 301, 1 RCT, 95% CI 0.63 to 203.81), fatigue (RR 2.04, N = 615, 2 RCTs, 95% CI 0.79 to 5.23, $I^2 = 54\%$), insomnia (RR 0.68, N = 615, 2 RCTs, 95% CI 0.33 to 1.37, $I^2 = 15\%$), anxiety (RR 0.90, N = 314, 1 RCT, 95% CI 0.33 to 2.42), nausea (RR 0.83, N = 615, 2 RCTs, 95% CI 0.43 to 1.59, $I^2 = 1\%$), constipation (RR 6.50, N = 28, 1 RCT, 95% CI 0.41 to 104.20), and nasopharyngitis (RR 0.62, N = 301, 1 RCT, 95% CI 0.23 to 1.66).

Detailed data were also available for laboratory value and vital sign changes. Therefore, all following effect estimates are based on baseline to endpoint change data. Significant changes were found for liver function tests, blood lipids, the haemogram, and calcium. For liver function parameters, the following effect estimates were found (all significant): AST/SGOT change: SMD 0.35, N = 526, 2 RCTs, 95% CI 0.18 to 0.52, $I^2 = 0\%$; ALT/SGPT change: SMD 0.46, N = 530, 2 RCTs, 95% CI 0.29 to 0.63, $I^2 = 0\%$; GGT (GGPT/SGGT/YGGT) change: SMD 0.26, N = 268, 1 RCT, 95% CI 0.02 to 0.50; total bilirubin change: SMD -0.29, N = 264, 1 RCT, 95% CI -0.53 to -0.05; direct bilirubin change: SMD -0.35, N = 158, 1 RCT, 95% CI -0.60 to -0.11). The following blood lipid changes were reported (all significant): total cholesterol change: SMD 0.42, N = 327, 2 RCTs, 95% CI 0.20 to 0.64, $I^2 = 0\%$; LDL cholesterol change: SMD 0.35, N = 259, 1 RCT, 95% CI 0.10 to 0.59; HDL cholesterol change: SMD -0.28, N = 269, 1 RCT, 95% CI -0.52 to -0.04; triglycerides (fasting) change: SMD 0.37, N = 203, 1 RCT, 95% CI 0.09 to 0.64. There was also a significant effect for prolactin change (SMD 0.41, N = 528, 2 RCTs, 95% CI 0.23 to 0.59, $I^2 = 10\%$). There were some significant differences in changes of haemogram parameters: leukocyte count change: SMD -0.40, N = 262, 1 RCT, 95% CI -0.65 to -0.16; neutrophils (segmented) change: SMD -0.39, N = 262, 1 RCT, 95% CI -0.63 to -0.14; basophils change: SMD -0.28, N = 262, 1 RCT, 95% CI -0.53 to -0.04; monocytes change: SMD -0.28, N = 262, 1 RCT, 95% CI -0.53 to -0.04. No significant differences in baseline to endpoint change were found for the following: erythrocyte count change: SMD -0.18, N = 262, 95% CI -0.42 to 0.06; haemoglobin change: SMD -0.21, N = 262, 1 RCT, 95% CI

−0.45 to 0.03 and mean cell haemoglobin concentration change: SMD 0.03, N = 260, 1 RCT, 95% CI −0.22 to 0.27). For the platelet count change, conflicting results were reported, and effect estimates were not pooled because of considerable heterogeneity ($I^2 = 90\%$). Despite having the same treatment durations, participants characteristics, and treatment doses, Schulz 2007 found a significant increase in the platelet count (SMD 0.32, N = 257, 1 RCT, 95% CI 0.07 to 0.56), whereas Zanarini 2007 reported a significant decrease (SMD −0.26, N = 260, 1 RCT, 95% CI −0.50 to −0.01). There was a significant effect for calcium change (SMD −0.33, N = 268, 1 RCT, 95% CI −0.57 to −0.09), but not for albumin change (SMD −0.21, N = 269, 1 RCT, 95% CI −0.45 to 0.03). There were no significant effects of olanzapine concerning kidney function parameters (creatinine phosphokinase change: SMD −0.21, N = 268, 1 RCT, 95% CI −0.45 to 0.03; urea nitrogen change: SMD −0.14, N = 269, 1 RCT, 95% CI −0.38 to 0.10) or vital signs changes (pulse, standing: SMD 0.08, N = 290, 1 RCT, 95% CI −0.15 to 0.31; pulse, supine: SMD 0.02, N = 290, 1 RCT, 95% CI −0.21 to 0.25; diastolic blood pressure, standing: SMD −0.03, N = 290, 1 RCT, 95% CI −0.26 to 0.20; diastolic blood pressure, supine: SMD −0.01, N = 290, 1 RCT, 95% CI −0.24 to 0.22; systolic blood pressure, standing: SMD 0.03, N = 290, 1 RCT, 95% CI −0.20 to 0.26; systolic blood pressure, supine: SMD −0.04, N = 290, 1 RCT, 95% CI −0.27 to 0.19).

For ziprasidone, data on single adverse events as reported by patients were available. There was no increased risk of experiencing any adverse event under ziprasidone treatment (RR 2.75, N = 60, 1 RCT, 95% CI 0.99 to 7.98). In detail, the following symptoms were reported more frequently by participants who had received ziprasidone, with no significant differences of frequency: dizziness (RR 9.00, N = 60, 1 RCT, 95% CI 0.51 to 160.17), sedation (RR 6.00, N = 60, 1 RCT, 95% CI 0.77 to 46.87), “uneasy feeling” (RR 7.00, N = 60, 1 RCT, 95% CI 0.38 to 129.93).

1.19.3 Mood stabilisers: For carbamazepine, no detailed data were available concerning adverse effects. For valproate semisodium and lamotrigine, body weight changes were given. No significant changes of body weight were observed for valproate semisodium (SMD 0.68, N = 30, 1 RCT, 95% CI −0.10 to 1.47) or lamotrigine (SMD −0.13, N = 27, 1 RCT, 95% CI −0.93 to 0.67).

There was a significant effect of body weight change by topiramate treatment, indicating significant weight loss (SMD −0.55, N = 127, 3 RCTs, 95% CI −0.91 to −0.19, $I^2 = 0\%$). The following single adverse events were reported, with no significantly increased risk: memory problems (RR 2.00, N = 56, 1 RCT, 95% CI 0.55 to 7.22), trouble in concentrating (RR 2.00, N = 56, 1 RCT, 95% CI 0.55 to 7.22), headache (RR 1.00, N = 56, 1 RCT, 95% CI 0.15 to 6.61), fatigue (RR 2.00, N = 56, 1 RCT, 95% CI 0.40 to 10.05), dizziness (RR 1.50, N = 56, 1 RCT, 95% CI 0.27 to 8.30), menstrual pain (RR 1.67, N = 56, 1 RCT, 95% CI 0.44 to 6.31), and paraesthesia (RR 3.00, N = 56, 1 RCT, 95% CI 0.33 to 27.12).

1.19.4 Antidepressants: No detailed data of adverse effects were available for amitriptyline, fluoxetine, fluvoxamine, and mianserin. For phenelzine sulfate, a non-significant effect of weight gain was reported (SMD 0.11, N = 62, 1 RCT, 95% CI −0.39 to 0.61).

1.19.5 Miscellaneous active agents: No detailed data were available for the remaining active agent that had been included in this review, i.e. omega-3 fatty acids.

2. Drug versus drug comparisons: For corresponding analyses to drug versus drug comparisons, see Analysis 66.1 to Analysis 82.1.

In the following, results will be reported by comparison category.

2.1 First-generation antipsychotic versus first-generation antipsychotic: One RCT compared two first-generational antipsychotics, i.e. loxapine versus chlorpromazine (Leone 1982).

The participants of this trial were administered either loxapine (N = 40; mean daily dose 14.5 mg) or chlorpromazine (N = 40; mean daily dose 110 mg) for six weeks. Both male and female outpatients were included. Their mean age was 30.8 years. Severity of illness was rather low, as participants had to fulfil only four of the diagnostic BPD characteristics of Gunderson et al. (Gunderson 1981). Two of them had to be rated as severe and two as at least moderate.

Only tolerability and adverse events data were usable for effect size calculation. The ratio of patients who did not complete at least three weeks of treatment or were removed due to side effects did not differ significantly between either of the groups (RR of loxapine treated patients as compared to chlorpromazine treated patients 1.14, N = 80, 95% CI 0.46 to 2.85). Neither did the frequency of any adverse events differ significantly between the two groups (RR 1.14, N = 80, 95% CI 0.46 to 2.85), nor did any of the most frequent adverse events in particular (sleepiness/drowsiness: RR 0.80, N = 80, 95% CI 0.23 to 2.76; restlessness: RR 1.50, N = 80, 95% CI 0.26 to 8.50; muscle spasms: RR 3.00, N = 80, 95% CI 0.33 to 27.63; fainting spells: RR 0.14, N = 80, 95% CI 0.01 to 2.68).

2.2 First-generation antipsychotic versus antidepressant: Two RCTs compared haloperidol, a first-generational antipsychotic agent, with antidepressant medication. In the Soloff 1989 trial, the comparison treatment was the TCA amitriptyline, in the Soloff 1993 trial it was the MAOI phenelzine sulfate..

Patients in the Soloff 1989 trial received 4 to 16 mg/day of haloperidol (mean daily dose 4.8 mg/day, average plasma level 8.66 ng/ml, SD 3.7 ng/mL) or 100 to 175 mg/day of amitriptyline (mean daily dose 149.1 mg/day, average plasma level of 240.4 ng/mL amitriptyline + nortriptyline, SD 99.4). Patients (N = 61, both male and female, mean age 25.1 years) started as inpatients and were allowed to leave the hospital after two weeks. Nevertheless, almost two thirds (62%) completed as inpatients. Study duration was five weeks. With average GAS scores of 42.2 at baseline, the severity of illness was serious.

Data were available for interpersonal problems, impulsivity, anger, and psychotic paranoid symptoms. There were no significant effects for any primary outcome. Results indicated that patients tended to profit more from haloperidol treatment concerning interpersonal problems (SMD -0.14, N = 57, 95% CI -0.66 to 0.38), anger (SMD -0.36, N = 57, 95% CI -0.89 to 0.16), and psychotic paranoid symptoms (SMD -0.35, N = 57, 95% CI -0.87 to 0.18); and

more from amitriptyline concerning impulsivity (SMD 0.20, N = 57, 95% CI -0.32 to 0.72). Neither drug proved to be significantly superior to the other one for any other pathology related outcome. Favourable results were found for haloperidol concerning anxiety (SMD -0.18, N = 57, 95% CI -0.70 to 0.34) and general psychiatric pathology (SMD -0.07, N = 57, 95% CI -0.59 to 0.45) as well as for amelioration of mental health status (SMD 0.29, N = 57, 95% CI -0.23 to 0.81). Depressive pathology responded slightly better in amitriptyline treated patients (SMD 0.08, N = 57, 95% CI -0.44 to 0.59). The risk of dropping out was higher in the haloperidol treated group (RR 2.90, N = 61, 95% CI 0.32 to 26.38), but again this was not statistically significant. Soloff 1993 tested up to 4 mg of haloperidol (average dose 3.93 mg/day, SD 0.65, mean plasma level 5.29 ng/mL, SD 5.05) against up to 90 mg/day of the MAOI phenelzine sulfate (average dose 60.45 mg/day, SD 9.55, 77.54% mean platelet MAO inhibition after three weeks) in male and female BPD patients (N = 64, mean age: 26.7 years, SD = 7.2) for a duration of five weeks. All patients started as inpatients and remained in hospital for at least two weeks. With average GAS scores of 43.9 at baseline, the severity of illness was serious.

There were no significant differences between the two drugs concerning primary outcomes. The results indicated a tendency for haloperidol treated patients to suffer less from interpersonal problems as compared to phenelzine sulfate treated patients (SMD -0.46, N = 64, 95% CI -0.96 to 0.04). In all other cases, i.e. concerning BPD severity (SMD 0.46, N = 64, 95% CI -0.03 to 0.96), impulsivity (SMD 0.09, N = 64, 95% CI -0.04 to 0.58), anger (SMD 0.08, N = 64, 95% CI -0.41 to 0.57), and psychotic/paranoid symptoms (SMD 0.15, N = 64, 95% CI -0.34 to 0.64), phenelzine sulfate treated patients were better off.

Significant effects were found in favour of phenelzine sulfate treatment for depression (SMD 0.68, N = 64, 95% CI 0.17 to 1.19), anxiety (SMD 0.66, N = 64, 95% CI 0.15 to 1.16), general psychiatric pathology (SMD 0.53, N = 64, 95% CI 0.03 to 1.03) and mental health status (SMD -0.51, N = 64, 95% CI -1.01 to -0.01), with medium effect sizes. The risk of dropping out was higher in the haloperidol group (RR 1.58, N = 74, 95% CI 0.49 to 5.15), but the effect was not significant. Haloperidol treated patients experienced less weight gain than amitriptyline treated patients did (SMD -0.29, N = 64, 95% CI -0.78 to 0.21), but the effect was not significant.

2.3 Second-generation antipsychotic versus antidepressant: One RCT (Zanarini 2004) compared the second-generation antipsychotic olanzapine to the SSRI antidepressant fluoxetine. All participants were female outpatients (N = 30, mean age: 23.0 years, SD = 5.7), receiving either 2.5 mg/day of olanzapine or 10.0 mg/day of fluoxetine for a duration of eight weeks. With average GAF scores of 52.5 (SD = 6.9), the patients were moderately ill. All SMD effect sizes were calculated on the basis of mean baseline change scores. Dichotomous data were calculated on basis of the ITT sample as intended.

There were no significant differences between the two drugs for any pathology-related outcome. Usable data were provided for impulsivity, with a small, non-significant effect favouring olanzapine (mean change SMD -0.20, N = 29, 95% CI -0.93 to 0.53). Olanzapine treated patients also experienced a greater decrease in depressive pathology (SMD -0.73, N = 29, 95% CI -1.49 to 0.03), but the effect was, again, not significant. Attrition did not

differ significantly between the groups (RR 0.29, N = 30, 95% CI 0.20 to 1.76). The only significant findings referred to adverse effects of treatment, with higher weight gain (SMD 0.98, N = 29, 95% CI 0.20 to 1.76) and more cases of mild sedation (RR 3.50, N = 30, 95% CI 1.23 to 9.92) in olanzapine treated patients. Akathisia was also more often reported by olanzapine treated patients, but the effect estimate was not significant (RR 0.70, N = 30, 95% CI 0.23 to 2.11).

3. Drug versus combination of drugs: The Zanarini 2004 RCT comprised three experimental groups. One group received the second-generation antipsychotic olanzapine (2.5 mg/day; N = 16), another group received the SSRI antidepressant fluoxetine (10 mg/day; N = 14), and the third group received both drugs (2.5 mg/day of olanzapine plus 10.0 mg/day; N = 15) for eight weeks. All participants were female outpatients (mean age: 23.0 years, SD = 5.7). With average GAF scores of 52.5 (SD = 6.9), they were moderately ill.

All SMD effect sizes were calculated on basis of mean baseline change scores. Dichotomous data were calculated on basis of the ITT sample as intended.

Only small and non-significant differences were found between olanzapine treatment alone and combined treatment with fluoxetine (impulsivity: mean change SMD 0.02, N = 29, 95% CI -0.71 to 0.76, “favouring” combined treatment; depression: mean change SMD -0.26, N = 29, 95% CI -1.00 to 0.47, favouring olanzapine alone). Neither attrition (RR 0.19, N = 31, 95% CI 0.01 to 3.63), nor weight gain (SMD 0.70, N = 29, 95% CI -0.05 to 1.46) differed significantly between the two groups. There were no significant differences in the ratio of participants reporting sedation (RR 1.61, N = 31, 95% CI 0.87 to 2.96) or akathisia (RR 0.75, N = 31, 95% CI 0.25 to 2.28) between the groups.

For the comparison of fluoxetine versus combined treatment with olanzapine, again there were no significant differences. However, both effect estimates of pathology-related outcomes indicated better results for combined treatment than fluoxetine alone (impulsivity: mean change SMD 0.25, N = 26, 95% CI -0.53 to 1.02; depression: mean change SMD 0.54, N = 26, 95% CI -0.24 to 1.33). For tolerability, body weight change and sedation, data indicated better results for the group that had received fluoxetine alone (attrition: RR 0.54, N = 29, 95% CI 0.05 to 5.28; body weight change: SMD -0.54, N = 29, 95% CI -1.32 to 0.25; sedation: RR 0.46, N = 29, 95% CI 0.15 to 1.44). Akathisia was more often experienced by the participants with single treatment (RR 1.07, N = 29, 95% CI 0.39 to 2.92).

DISCUSSION

Summary of main results

1. Drug versus placebo—The following placebo comparisons were investigated in the identified RCTs.

- (1) First-generation antipsychotics:
 - (a) thiothixene (Goldberg 1986, N = 50);
 - (b) flupenthixol (Montgomery 1979/82, N = 30);

- (c) haloperidol (Soloff 1989, N=60; Soloff 1993, N = 58).
- (2) Second-generation antipsychotics:
 - (a) aripiprazole (Nickel 2006, N = 52);
 - (b) olanzapine (Bogenschutz 2004, N = 40; Linehan 2008, N = 24; Schulz 2007, N = 314; Soler 2005, N = 60; Zanarini 2001, N = 28; Zanarini 2007, N = 301);
 - (c) ziprasidone (Pascual 2008, N = 60).
- (3) Mood stabilisers:
 - (a) carbamazepine (De la Fuente 1994, N = 20);
 - (b) valproate semisodium (Frankenburg 2002, N = 30; Hollander 2001, N = 16);
 - (c) lamotrigine (Reich 2009; Tritt 2005, N = 27);
 - (d) topiramate (Loew 2006, N = 56; Nickel 2004 and Nickel 2005, N = 31 + N = 44).
- (4) Antidepressants:
 - (a) amitriptyline (Soloff 1989, N = 59);
 - (b) fluoxetine (Salzman 1995, N = 22, Simpson 2004, N = 25);
 - (c) fluvoxamine (Rinne 2002, N = 38);
 - (d) phenelzine sulfate (Soloff 1993, N = 72);
 - (e) mianserin (Montgomery 81/82/83, N = 38).
- (5) Miscellaneous:
 - (a) omega-3 fatty acid (Hallahan 2007, N = 49; Zanarini 2003, N = 30).

1.1 Pathology related outcomes: Of the first-generation antipsychotics under investigation, haloperidol had a significant effect concerning the reduction of anger, and flupenthixol treated patients were significantly less likely to get engaged in suicidal acts. No proof of efficacy was found for thiothixene.

Of the second-generation antipsychotics, aripiprazole had significant effects in the reduction of interpersonal problems, impulsivity, anger, psychotic paranoid symptoms, depression, anxiety, and general psychiatric pathology. For olanzapine, no significant effects were found for any pathology related outcome in primary analyses. Secondary analyses indicated significant decreases in affective instability, anger, psychotic paranoid symptoms, and anxiety. A significantly greater decrease in anxiety by olanzapine was found by one trial. Concerning suicidal ideation and self-mutilating behaviour, only two of the five relevant study results could be pooled due to different formats of reporting. The pooled effect of these two estimates suggests that the olanzapine-treated group experienced a significantly lower degree of amelioration of recurrent suicidal ideation as compared to the placebo

group. Of the remaining three trials reporting on self-harming behaviour, two also found non-significant tendencies of unfavourable outcomes for olanzapine. No significant effects were found for ziprasidone treatment.

There were also significant effects for the mood stabilisers valproate semisodium, lamotrigine, and topiramate. Valproate semisodium had significant effects concerning the reduction of interpersonal problems and depression. A significant effect in the reduction of anger was found by one study, and the positive direction of effect was supported by the findings of another study. Lamotrigine was significantly superior to placebo concerning impulsivity and anger. Topiramate had significant effects concerning interpersonal problems, impulsivity, anger (as assessed by three single, significant study effects, only two of which could be pooled), anxiety, and general psychiatric pathology. No significant effects were found for carbamazepine treatment.

For antidepressants, there was only a significant effect for the TCA amitriptyline concerning the reduction of depression. No significant effects were found for mianserin, the SSRI agents fluoxetine and fluvoxamine, nor for the MAOI agent phenelzine sulfate.

Omega-3 fatty acid was found to have a significant effect on suicidality. For depression, a significant effect was found by one study, with the second study (that could not be pooled with the first one due to different formats of data reporting) supporting the direction of effect.

1.2 Adverse effects: Tolerability did not differ for any drug placebo comparison, i.e. drug treatment was not associated with a higher rate of non-completers than was placebo treatment.

Most trials did not provide numerical data on specific adverse effects, with the exception of body weight changes. Haloperidol and phenelzine treatment had no significant effects on body weight, nor did valproate semisodium or lamotrigine. However, olanzapine treatment was associated with significant weight gain, and topiramate treatment with significant weight loss. The ratio of olanzapine, ziprasidone and lamotrigine treated patients reporting any adverse event did not differ significantly as compared to the placebo groups.

Numerous data on additional specific adverse effects were only available for a few trials. For the placebo comparisons of ziprasidone and topiramate, single adverse events were reported, with no significant differences in occurrence between the groups. Detailed data were available for olanzapine, including even changes in laboratory values. Here, the ratio of participants reporting any adverse event in each group did not differ significantly between olanzapine and placebo treatment. However, olanzapine treated patients reported significantly more often increased appetite, somnolence, and mouth-dryness. One trial reported significantly more sedation in olanzapine treated patients, and another one (that could not be pooled with the first one due to substantial heterogeneity) supported this direction of effect. Additionally, significant effects on liver values, blood lipids, prolactin levels, and full blood counts were found, but there were no significant effects on kidney function values or cardiovascular system parameters.

However, little is known about adverse events increasing the risk of patients not completing treatment or experiencing body weight changes, except for olanzapine treatment. Therefore, the above cited significant effects should be regarded with caution. Known adverse effects of the remaining drugs have certainly also to be considered when choosing a treatment option for a certain patient, though the data were too sparse to calculate effect sizes in this review for most drugs.

2. Drug versus drug—The following drug versus drug comparisons were investigated (numeration as in results section for ease of comparison).

- (6) First-generation antipsychotic versus first-generation antipsychotic:
 - (a) loxapine versus chlorpromazine (Leone 1982, N = 40).
- (7) First-generation antipsychotic versus antidepressant:
 - (a) haloperidol versus amitriptyline (Soloff 1989, N = 61);
 - (b) haloperidol versus phenelzine sulfate (Soloff 1993, N = 74).
- (8) Second-generation antipsychotic versus antidepressant:
 - (a) olanzapine versus fluoxetine (Zanarini 2004, N = 30).

2.1 Pathology related outcomes: Concerning the comparison of the two first-generation antipsychotics loxapine versus chlorpromazine, there were no usable data available regarding any pathology related outcome.

The first-generation antipsychotic haloperidol and the antidepressant amitriptyline did not differ significantly concerning any primary or secondary outcome. The antidepressant phenelzine sulfate proved to be superior to haloperidol in reducing depression, anxiety, general psychiatric pathology, and improving the overall mental health status.

No significant differences were found for the comparison of the second-generation antipsychotic olanzapine with the antidepressant fluoxetine for any pathology related outcome.

2.2 Adverse effects: Tolerability, i.e. attrition, did not differ significantly for any of the investigated drug versus drug comparisons.

The comparison of the frequencies of adverse events (i.e. any adverse event, sleepiness, restlessness, muscle spasms, fainting spells) in loxapine and chlorpromazine treated patients yielded no significant differences.

No data of adverse effects were available for the comparison of haloperidol versus amitriptyline. For the haloperidol versus phenelzine sulfate comparison, weight change was reported, with no significant difference between the two treatments.

However, olanzapine and fluoxetine treatment differed significantly concerning weight gain, with more weight gain in the olanzapine treated group. Additionally, a higher ratio of olanzapine treated patients reported mild sedation, as compared to the fluoxetine group.

3. Active drug versus combination of drugs—The following comparisons were investigated (numeration following on from 2. Drug versus drug, above).

- (9) Second-generation antipsychotic versus second-generation antipsychotic plus antidepressant:
 - (a) olanzapine versus olanzapine plus fluoxetine (Zanarini 2004, N = 31).
- (10) Antidepressant versus antidepressant plus second-generation antipsychotic:
 - (a) fluoxetine versus fluoxetine plus olanzapine (Zanarini 2004, N = 29).

3.1 Pathology related outcomes: For both the comparisons olanzapine versus olanzapine plus fluoxetine as well as fluoxetine versus fluoxetine plus olanzapine, data on impulsivity and depressive pathology were available. There were no significant differences indicating any benefits from combined treatment versus treatment with olanzapine or fluoxetine alone.

3.2 Adverse effects: There were no significant differences for both comparisons in terms of tolerability, body weight change, and the frequency of restlessness or mild sedation.

Overall completeness and applicability of evidence

Participants—As described earlier (see Characteristics of included studies table), most study participants exhibited mild to moderate levels of illness. However, there was a broad range from seriously ill (e.g. Soloff 1989: mean GAS value pre-treatment 42.2, reflecting “serious symptomatology or impairment in functioning”) to very mildly impaired patients (e.g. Zanarini 2001: “patients [...] leading active social and vocational lives”). Acutely suicidal patients were not included in most trials, with the exception of Montgomery 1979/82 and Montgomery 81/82/83, including patients immediately following a severe suicidal act that had led to hospital admission.

Of concern regarding applicability to clinical settings might be the psychiatric exclusion criteria of most studies. Besides acutely suicidal patients, people with comorbid schizophrenia or schizoaffective disorders, bipolar disorders, alcohol or drug dependence and sometimes even alcohol or drug abuse were often not eligible for study participation. What is more, a current major depressive episode or severe depression was also a criterion of exclusion in the majority of trials. As comorbid axis-I disorders are highly prevalent in BPD patients, especially mood disorders (96.9%) and substance use disorders (62.1%; Zanarini 2004b), the exclusion of those participants renders applicability difficult. However, eating disorders, which are highly prevalent in BPD patients as well (53%, Zanarini 2004b), were no reason for exclusion in any study (with the exception of De la Fuente 1994; Salzman 1995, who excluded patients with any comorbid axis-I disorder). Anxiety disorders, which are prevalent in 89.0% of BPD patients (Zanarini 2004b), were only excluded in two trials (i.e. current PTSD, panic disorder, or obsessive-compulsive disorder; Schulz 2007; Zanarini 2007).

One third of trials was restricted to female patients (nine out of 27), one trial was restricted to men (Nickel 2005), and within the remaining samples, women were always predominant. Thus, women constituted the majority of all participants involved within this review. This

reflects the overall higher prevalence of BPD diagnosis in women as compared to men (although the “real” prevalence is supposed to be even; Torgersen 2005), but may make the applicability to male patients difficult.

Interventions—Study duration ranged from 32 days to 24 weeks, with a mean duration of approximately 12 weeks. These observation periods may be sufficient to judge treatment efficacy in the single patient. However, drug treatment often lasts longer in clinical settings. Therefore, especially adverse effects must be monitored cautiously. Since most BPD patients are taking psychotropic medication continuously, future RCTs on this topic should cover appropriate observation periods of longer duration (e.g. six months as a rough navigation) to allow for better applicability to clinical settings.

Catamnestic data are only available for Loew 2006; Nickel 2004; Nickel 2005 (all: topiramate versus placebo), Tritt 2005 (lamotrigine versus placebo in female patients), and Nickel 2006 (aripiprazole versus placebo in both male and female patients) studies. As blinding was broken after the initial 8- and 10-week treatment phases in each of these trials, we decided not to include the catamnestic data here, as the break of blinding is likely to introduce bias on efficacy findings, especially on self-rated or self-reported data. For aripiprazole and lamotrigine, all significant findings of the post-treatment comparisons were still present after an additional 18 months of open treatment. Significant changes throughout the whole observation period were reported for all topiramate trials. All findings at post-treatment were corroborated by the catamnestic data, with exception of two outcome variables: There was no significant change for general psychiatric pathology concerning the overall observation period, and a significant change for depressive pathology emerged at the end of the 18-month follow-up.

Another difference to clinical settings may be that patients often receive several psychotropic drugs at a time. Polypharmacy is the rule rather than the exception (Zanarini 2004a). With the exception of the comparison of combined olanzapine and fluoxetine treatment to olanzapine and fluoxetine treatment alone (Zanarini 2004; no beneficial effects for combined treatment), there are no data from RCTs available supporting or even investigating polypharmacological treatment. As combined drug treatment cannot be considered as having the additional effect to that of each single drug treatment, it should always be considered that the administration of several drugs is not empirically supported by any RCT, and, to our knowledge, not by any trial of lower evidence level either.

With the exception of the Linehan 2008; Simpson 2004 and Soler 2005 trials, all patients of whom were in DBT treatment, the study participants did not receive specific concomitant psychotherapy, either because it was not allowed by the study protocol, or patients were allowed to receive psychotherapy but did not. However, mental health service treatment options vary internationally, i.e. receiving psychotherapy may be more characteristic for some countries and out of character for others.

There are some more substances that are currently discussed for use in BPD patients (especially second-generation antipsychotics), that could not be included in this review as RCTs are currently not available. However, there are some ongoing trials (see

Characteristics of ongoing studies), the results of which will hopefully be included in subsequent versions of this review. Additionally, there are findings from lower-evidence studies on further second-generation antipsychotics (clozapine, quetiapine, risperidone), mood stabilisers (divalproex-extended release, lithium, oxcarbazepine), antidepressants (MAOI tranylcypromine, NRI reboxetine, SNRI venlafaxine, the SSRIs sertraline and paroxetine, the TCAs desipramine and imipramine), anxiolytics (alprazolam), and miscellaneous drugs (clonidine, opioid antagonists naloxone and naltrexone, riluzole, and trifluoperazine).

Outcomes—Unfortunately, there was little consensus between primary studies on which outcome variables are crucial to rate therapy efficacy upon, even for those testing the same drug comparisons. Additionally, the use of different assessment instruments for one outcome variable renders comparability more difficult than necessary, and increases heterogeneity.

Mostly, outcome assessment was restricted to target variables that were not assessed with BPD-specific assessment instruments. For example, psychotic pathology was a common outcome, and common unspecific assessment instruments were used (i.e. SCL-90-subscale “psychoticism”), but BPD-specific psychotic pathology, i.e. stress-related paranoid ideation and dissociation, was not assessed. Hence, some domains of BPD core pathology were almost completely neglected, e.g. affective instability, dissociation, or chronic feelings of emptiness. Fortunately, relevant assessment instruments have been developed lately, reflecting each of the BPD core criteria (e.g. the BPDSI scale by Arntz 2003, the CGI-BPD scale by Perez 2007, or the ZAN-BPD scale by Zanarini 2003a). Additionally, there were very few numerical data provided concerning adverse events. Some studies reported no details at all beyond attrition and body weight changes, whereas others only stated that adverse events were few and comparably frequent in all groups. We therefore appreciate the detailed reporting of adverse effects allowing for the calculation of treatment effects, and warn against regarding the other drugs, where relevant data are lacking, as safe, especially with regard to long-term therapy. We strongly recommend considering known adverse effects of all drugs when choosing a certain treatment option, although we were not able to report them here because of incomplete assessment or reporting of the primary studies investigated here.

Specificity of treatment effects—We cannot exclude that unspecific sedating effects, e.g. of valproate semisodium or olanzapine, may have contributed to study results. However, effects were also seen with non-sedating substances such as aripiprazole or lamotrigine questioning the broad conclusion that sedating effects are central for treatment effects.

Quality of the evidence

Altogether, 28 RCTs have been included, covering 22 different comparisons in ten comparison categories (see “Description of studies”). In the presence of the multitude of different comparisons and outcome variables, most results are based on single study findings only. The study sample sizes were rather small, and ranged, with exception of two large trials (Schulz 2007; N = 314; Zanarini 2007; N of patient data used here: 301), between 16 (Hollander 2001) and 108 (Soloff 1993; divided into three groups). Depending on the

randomisation algorithm, i.e. if study groups were equal in size or one group twice as large as the other, and the overall number of treatment groups, the minimum group size was $N = 4$ (Hollander 2001) and the maximum group size was $N = 43$ (Nickel 2004; Nickel 2005). Therefore, the power to detect significant effects was quite low.

In addition, the overall robustness of findings must be considered low for the majority of comparisons. Because the current evidence embraces only one single RCT effect, further findings would be likely to affect the actual results, especially if including larger study samples. However, the influence of further trials cannot definitely be predicted: On the one hand, further primary studies would enhance power and therefore make the detection of significant effects more likely. On the other hand, the actual data are based on very few to single observations, so it is impossible to judge about publication biases (e.g. by depicting funnel plots as intended), even if the concealment of negative, non-significant findings is much more likely. In Figure 12, a funnel plot was drawn for the comparison category with most single study effects (comparison 11.1, SMDs of active drug versus placebo comparisons for the outcome “anger”). It is most difficult to draw definite conclusions from that figure, as it embraces a heterogeneous sample of effect sizes for diverse drug-placebo comparisons. On the one hand, there seems to be an overall tendency of lacking non-significant findings (no effect estimates at bottom right corner). On the other hand, the publication of additional RCTs matching the comparisons already investigated here is rather unlikely, as we are only aware of ongoing trials testing different drugs (see Characteristics of ongoing studies).

Summary of findings tables including ratings of the evidence quality are provided for all drug versus placebo comparisons (see Appendix 14 to Appendix 29).

Potential biases in the review process

We strived to identify all relevant published and unpublished evidence (see Search methods for identification of studies). The search was not restricted to any language. In spite of the great efforts to avoid publication bias, we were not able to include any unpublished data.

As relating to our inclusion criteria, we tried to retain a most homogeneous pool of primary studies. However, there were some inconsistencies between studies particularly pertaining to psychiatric comorbidity of study participants. For example, bipolar disorders were a common exclusion criterion, whereas one study (Frankenburg 2002) required its participants to have a diagnosis of bipolar II disorder. Also, acute suicidal patients were not eligible for most studies, but the participants of the Montgomery 1979/82 and Montgomery 81/82/83 trials were recruited immediately following a suicidal act that had led to hospital admission. In nine studies, only women were included (Frankenburg 2002; Loew 2006; Nickel 2004; Rinne 2002; Simpson 2004; Tritt 2005; Zanarini 2001; Zanarini 2003; Zanarini 2004), whereas the remaining study samples consisted of both male and female patients.

The severity of illness also varied between studies, mostly from mild to moderate. However, we tried to exactly specify and describe all studies with regard to their crucial characteristics (see Description of studies, Characteristics of included studies), in order to let the reader decide about applicability of relevant study characteristics to his or her decisive situation.

Another point of concern is reporting bias. Most studies provide only a fragmentary outcome pattern, making the concealment of non-significant findings likely. We tried to deal with this by first defining all patient-relevant outcome variables that are directly (primary outcomes) or indirectly (secondary outcomes) associated with BPD treatment, i.e. all outcome variables that a consumer and his or her therapist are likely to be interested in. We have tried not only to stress reported findings but also outcome gaps, such as outcome variables for which the effects of a certain treatment cannot be judged due to a lack of data.

Agreements and disagreements with other studies or reviews

Other reviews—This is an update and new citation version of the preceding Cochrane Collaboration review ‘Pharmacological interventions for BPD’ by Binks 2006. Its literature searches covered the period up to October 2002, and the latest included study dates from 2001. Since then, there have been further research activities, and new substances have been investigated in BPD. The preceding review included ten RCTs, whereas we were aware of 28 includable studies at the point of last literature search updates (September 2009).

As concerns other systematic reviews and meta-analysis on the topic of pharmacotherapy for BPD, we did not review this type of evidence systematically. However, there are three recent works, each with a similar focus, that should be referred to at this point (Duggan 2008; Ingenhoven 2010; Nosè 2006). Nosè 2006 Duggan 2008; Ingenhoven 2010

Both Nosè 2006 Nosè 2006 and Ingenhoven 2010 Ingenhoven 2010 included placebo-controlled RCTs. Mixed study samples with primarily BPD patients were includable in the Nosè 2006 Nosè 2006 review, participants with both BPD and/or schizotypal PD were includable in the Ingenhoven 2010 Ingenhoven 2010 review, and people with any PD were included in the Duggan 2008 Duggan 2008 review. The most recent literature searches were done in June 2006, December 2007 and December 2006, respectively. Due to different inclusion criteria and different search periods, the study pools differ from ours. Mainly, these reviews had less RCTs of antipsychotic drugs available, but included more RCTs of antidepressants since these drugs have been tested in mixed samples that were not includable in this review (if less than 70% of participants had a diagnosis of BPD, see Types of studies). Outcomes were, by and large, comparable to those of our review. All three reviews conducted meta-analyses across classes of drugs, i.e. effect estimates referring to a certain class of drugs (any antipsychotic, any antidepressant, or any mood stabiliser) were pooled. In this review, study effects were only pooled if referring to the same substance.

Both reviews report several findings of effectiveness for antidepressants. This differs from our findings that are only based on RCTs conducted in study samples of more than 70% BPD patients, and were not derived from accumulation of findings from different (antidepressant) substances.

Guidelines—This systematic review is not a guideline, which provides treatment recommendations. It is meant to help providers, practitioners and patients make informed decisions. However, we will now comment on the main guidelines that give recommendations for pharmacotherapy treatment of BPD in light of the results of this systematic review.

The American Psychiatric Association Practice Guidelines (APA 2001; updated in 2005, APA 2005) are commonly cited when recommending pharmacological treatment strategies for BPD. However, these are based on literature searches covering the literature up to 1998. Since then, 20 RCTs have been published, investigating mood stabilisers (valproate semisodium, lamotrigine, topiramate), antidepressants (fluvoxamine, fluoxetine), second-generation antipsychotics (aripiprazole, olanzapine, ziprasidone), and omega-3 fatty acids.

Taking the findings of this review into account with regard to the APA guidelines, some differences are apparent: The up-to-date RCT evidence presented here does not support the recommendation of SSRIs as first-line treatment for affective dysregulation and impulsive-behavioural dyscontrol symptoms. Instead, beneficial effects were found for mood stabilisers (topiramate, valproate semisodium, lamotrigine) and second-generation antipsychotics (aripiprazole, olanzapine) for affective dysregulative symptoms. Beneficial effects indicating a reduction of impulsive-behavioural dyscontrol symptoms are available for mood stabilisers (topiramate, lamotrigine) or second-generation antipsychotics (aripiprazole). The APA guidelines recommend low-dose antipsychotics in general for the treatment of cognitive-perceptual symptoms, whereas our findings support the use of SGA (aripiprazole, olanzapine) in particular. This development, a shift towards second-generation antipsychotics, has been foreshadowed by John M. Oldham in his guideline watch of 2005 (APA 2005 APA 2005), but to our knowledge, the original guideline recommendations have not been modified since.

The World Federation of Societies of Biological Psychiatry (WFSBP) Guidelines for Biological Treatment of Personality Disorders (Herpertz 2007), that are based upon RCTs, open trials and individual clinical experiences, refer to all evidence that was identified in MEDLINE searches up to June 2007. Although their aim was to grade the evidence for the use of drugs in BPD treatment, and to give recommendations based not only on RCT but also lower-level evidence, they conclude that there is no evidence at either level of evidence that any drug improves BPD psychopathology in general. In addition, they did not find any beneficial evidence for the use of a the combination of several drugs. In contrast to this review, the WFSBP guidelines conclude that SSRIs “are best shown to influence emotional dysregulation such as depressive mood, anxiety and mood swings and [...] appear to extend the improvement of comorbid conditions of mood and anxiety disorders.” (Herpertz 2007, p. 214). This recommendation is not corroborated by the RCT evidence, as investigated in this review. Additionally, some trials were included in the WFSBP guidelines as randomised controlled trials, that were not included in this review due to stricter inclusion criteria (see Criteria for considering studies for this review and Excluded studies).

The NICE guideline on treatment and management of BPD (NICE 2009) is based on a similar pool of RCTs, even if the NICE literature searches were last updated in May 2008 (NICE 2009, p. 56). The guideline developers recommend that “Drug treatment should not be used specifically for borderline personality disorder or for the individual symptoms or behaviour associated with the disorder (for example, repeated self-harm, marked emotional instability, risk-taking behaviour and transient psychotic symptoms).” (NICE 2009, p. 297). This seems somehow contradictory to the findings of this review. However, the scope of NICE and this Cochrane Collaboration review differ in asking different questions and using

different means and methods to answer them. In relation to the question or topic, this Cochrane review aims at reviewing all of the available valid RCT evidence concerning pharmacotherapy of BPD treatment, whereas NICE aims at providing specific recommendations for a defined clinical setting, with pharmacotherapy being only one component within a comprehensive framework of possible health care provisions. Regarding methods, NICE considers somewhat different sources of evidence and applies additional criteria to weigh the costs and benefits of treatments (“NICE has always been focused on providing guidance on the most effective way to use NHS resources”, NICE 2010) and it is consensus-based. In contrast, the aim of this review is, according to Cochrane Collaboration standards, “to present information, rather than to offer advice” (Higgins 2008, p. 67).

AUTHORS' CONCLUSIONS

Implications for practice

The current RCT evidence supporting the use of pharmacotherapy for BPD is very sparse when compared to its widespread usage. Despite the remarkable growth in RCT evidence (this review includes 18 more RCTs than its previous version of 2006, with 9 different substances under test), the conclusion that pharmacotherapy in BPD “is not based on good evidence from trials” (Binks 2006, p. 19) still stands. There are only a few study results per comparison, with small numbers of included participants. However, it is important to remember that no evidence of an effect is not evidence of no effect. Current findings from RCTs presented in this review are not robust and can easily be changed by future research.

The findings suggest there is support for the use of second-generation antipsychotics, mood stabilisers, and omega-3 fatty acids, but these require replication since most effect estimates were based on single study effects. The small amount of available information for individual comparisons indicated marginal effects for first-generation antipsychotics and antidepressants.

Notably, avoidance of abandonment, chronic feelings of emptiness, identity disturbance, and dissociation were not found to be affected significantly by any drug. This finding may be the result of the use of non-BPD specific assessment instruments that do not reproduce these very specific outcomes, but it also reflects that these symptoms are broadly not regarded treatable by pharmacotherapeutic interventions and remain subject to non-pharmaceutical treatments such as psychotherapy.

It is important to consider the adverse effects for these interventions. Most trials did not provide detailed data of adverse effects, but these can be assumed to be similar to those experienced by patients with other conditions. However, the data available suggest an increase in self-harming behaviour when using olanzapine. In addition, toxic effects in case of overdosing (e.g. TCA antidepressants) and the potential for misuse or substance dependence (e.g. hypnotics and sedatives) need special attention in BPD treatment. In the presence of a comorbid eating disorder, possible effects on body weight changes (especially weight gain by olanzapine treatment and weight loss by topiramate treatment) should be taken into account and discussed between the treating physician and the patient (“shared decision making”).

Currently, there is no evidence from RCTs that any drug reduces overall BPD severity, but there are distinct pathology facets. Therefore, pharmacotherapeutic treatment of BPD should be targeted at defined symptoms. Drug treatment should last a sufficient period of time (according to pharmacokinetic and dynamic properties of a certain substance) to judge if there are any benefits, and should be stopped or changed if there are none. Polypharmacy is not supported by the up-to-date evidence and should be avoided wherever possible.

As discussed above, the evidence is not robust. The studies may not adequately reflect several characteristics of clinical settings (among others, patients' characteristics and duration of interventions and observation periods). Further research is urgently needed to provide reliable recommendations. Furthermore, there are some difficulties stemming from the polythetic nature of BPD. Different patients with BPD are likely to experience different facets of the disorder, and clinicians working with these patients are acquainted with different subtypes. The question "What works for whom?" remains broadly unanswered. Consequently, there is little consensus among researchers about a common battery of outcome variables and measures, even for primary studies testing the same drugs with putatively the same treatment targets and effects. Thus, we have a fragmentary view on drug effects, and it remains uncertain how the modulation of one symptom affects another.

People with BPD and their carers should lobby for and facilitate good research.

Implications for research

In recent years, a shift of attention in BPD treatment research has been observed towards SGAs and mood stabilisers, which may be a consequence of study sponsoring by pharmaceutical companies.

Some other classes of drugs have been paid much less attention to than their actual importance in clinical settings suggests. For example, antidepressants, especially SSRIs, play a major role in everyday practice but currently only three placebo-controlled RCTs exist that tested SSRIs in BPD. These drugs urgently need further attention in future placebo-controlled RCTs of BPD treatment. However, replicative studies for all comparisons would be desirable in order to enhance the robustness of findings. On the other hand, placebo-controlled RCTs testing different mood stabilisers (such as oxcarbazepine) and SGAs (e.g. clozapine, quetiapine, risperidone) that have lately been investigated in several non-randomised studies with promising results would be of interest. Currently, there is no RCT evidence-based "gold standard", so any future research endeavour should comprise a placebo condition. Longer observation periods would be sensible, this would enhance external validity and the applicability of findings to primary care settings. Additionally, patients with comorbid axis-I disorders should not be excluded, as psychiatric comorbidities are common in BPD patients. Another point for future research may be the update of popular algorithms to follow in this patient group, e.g. the "Soloff-algorithm" (Soloff 1998 Soloff 1998) or the APA guidelines algorithm (Oldham 2004 Oldham 2004).

There is a huge heterogeneity of outcome variables and assessment instruments. A consensus on a minimum set of therapy outcome variables that are most likely to be of interest for any BPD patient would be desirable. Outcome assessment should be more

specific and sensitive to BPD relevant pathology. For example, psychotic pathology should be assessed in terms of BPD relevant symptoms, i.e. stress-related paranoid ideation. Fortunately, several assessment instruments have been developed lately to reflect BPD core pathology as described precisely by the DSM-IV criteria (e.g. the BPDSI scale by Arntz 2003, the CGI-BPD scale by Perez 2007, or the ZAN-BPD scale by Zanarini 2003a). However, some DSM-IV BPD criteria embrace several symptoms, e.g. the criterion of “stress-related paranoid ideation OR dissociation”. The possibility of more differentiated outcome assessment may stimulate further research on drugs that may affect BPD core symptoms but have been neglected in the existing RCTs. In particular, drugs targeting affective instability, an important hallmark of BPD pathology, would be of interest. Outcome assessment should also embrace a thorough, standardized assessment of adverse events. Spontaneous reporting of patients may not be as valid and comprehensive as would be desirable.

Acknowledgments

We would like to thank the previous authors of this review, Claire Binks, Mark Fenton, Lucy McCarthy, Tracy Lee, Clive Adams, and Conor Duggan. Additionally, we thank the editorial team of the Cochrane Developmental, Psychosocial and Learning Problems Group (CDPLPG), Bristol, especially Chris Champion, Jane Dennis, Jo Abbott, and Geraldine Macdonald, and the German Cochrane Centre for supporting this work. We are grateful to Gerd Antes, director of the German Cochrane Centre, who made contact with CDPLPG and helped in gaining grants for financing this work. We are also grateful to Martin Schumacher, director of the Institute of Biostatistics and Medical Informatics at the University Medical Center Freiburg, who gave support in application submission. Thanks also to Nicolas Rüsch, who helped translating studies during the study selection phase.

We are grateful to the German Ministry of Education and Research (BMBF; grant no. 01KG0609), and the research committee of the University Medical Center Freiburg for supporting this work.

SOURCES OF SUPPORT

Internal sources

- Research Committee of the University Hospital Freiburg, Germany.

External sources

- German Federal Ministry of Education and Research, grant no. 01KG0609, Germany.

CHARACTERISTICS OF STUDIES

Characteristics of included studies [*ordered by study ID*]

Bogenschutz 2004

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double, no further details Duration: 12 weeks (patients had to be free of mood stabilisers, antipsychotics, benzo-diazepines, and antidepressants for at least 2 weeks prior to participation) Setting: Outpatient
Participants	Diagnosis: BPD (DSM-IV, SCID-II) Age: Mean 32.6 years (SD = 10.3) Sex: 25 F, 15 M Exclusions: Schizophrenia, schizoaffective disorder, bipolar affective disorder, current major depressive episode, psychotic disorder due to substance or general medical condition, substance dependence not in full or partial remission, suicide

attempts in past 6 months, having current suicidal intent or definite plan, pregnancy, neurologic impairment		
Interventions	1 Olanzapine: flexible dose (2.5-20 mg/day), mean dose at endpoint: 6.9 mg/day (SD = 3.2) N = 20* 2 Placebo: no further details, mean pseudo-dose at endpoint: 10.2 mg/day (SD = 5.3) N = 20* Concomitant psychotherapy: Allowed to continue if initiated more than 3 months prior to randomization Concomitant pharmacotherapy: Medication for stable, chronic medical conditions such as hypertension	
Outcomes	Avoidance of abandonment: CGI-BPD-abandonment Interpersonal problems: CGI-BPD-unstable interpersonal relationships Identity disturbance: CGI-BPD-identity disturbance Impulsivity: CGI-BPD-impulsivity, OAS-M-aggression Suicidal ideation: CGI-BPD-recurrent suicidal ideation Affective instability: CGI-BPD-affective instability Feelings of emptiness: CGI-BPD-chronic feelings of emptiness Anger: CGI-BPD-inappropriate anger, AIAQ Dissociative symptoms: CGI-BPD-transient paranoia or dissociation Anxiety: HARS General psychiatric pathology: SCL-90 Mental health status: CGI Attrition Adverse effects: weight	
Notes	*as randomised	
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Patients were randomly assigned in equal numbers" (Bogenschutz 2004, p. 105)
Allocation concealment?	Unclear	Comment: No information provided
Blinding? self-rated outcomes	Unclear	Quote: "double-blind trial", "pseudo-dose [...] for patients receiving placebo" (Bogenschutz 2004, p. 105)
Blinding? observer-rated outcomes	Unclear	Quote: "double-blind trial", "pseudo-dose [...] for patients receiving placebo" (Bogenschutz 2004, p. 105) Comment: No information provided on who actually adjusted the dose and if this person was blind to the patients' allocation
Incomplete outcome data addressed? All outcomes	Unclear	Comment: "evaluable patients" refers to all patients remaining at least 2 weeks in the study, i.e. who attended both baseline and preliminary assessment after two weeks; reasons for early termination specified (Bogenschutz 2004, p.106). However, 2 patients dropped out of the olanzapine group due to "violation of protocol" (Bogenschutz 2004, p. 106) Of the 40 patients enrolled, 23 completed the full 12 weeks of the trial (10 in olanzapine group, 13 in placebo group) Reasons for early termination: Lost to follow-up: 2 in the olanzapine group, 5 in the placebo group Lack of efficacy: 2/2 Weight gain: 2/0 Sedation: 2/0 Violation of protocol: 2/0 Continuous data based on LOCF of patients remaining at least 2 weeks in the study dichotomous data based on ITT sample
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "No other psychotropic medications could be taken during the study or in the 2 weeks prior to the study." (Bogenschutz 2004, p. 105)
Bias due to sponsoring improbable?	No	Quote: "Supported by a grant from Eli Lilly and Co, Indianapolis, Ind." (Bogenschutz 2004, p. 104)

De la Fuente 1994

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double. (One clinician was blind to drug treatment and performed all clinical and psychometric assessments, the other one was not blind to drug treatment and correctly adjusted the plasma levels of carbamazepine and asked for adverse side effects. Patients were instructed not to communicate side effects to the blind clinician) Duration: 32 days (after at least 10 days psychotropic drug washout; 15 days for TCAs and MAOIs, no patient had taken neuroleptics in the 2-month period before the study) Setting: Inpatients	
Participants	Diagnosis: BPD (DSM-III-R; DIB) Age: Mean 32.7 years (range 22-45) Sex: 14 F, 6M Exclusions: DSM-III-R axis I disturbances (not excluded: patients who were depressed for less than 2 weeks), inability to stop alcohol or psychoactive substances, suspected poor treatment compliance, standard physical or neurological examinations abnormal, perturbed standard biological blood tests, perturbed electrocardiogram, positive history for epilepsy or standard traits for epilepsy, antecedents of encephalitis or cranial trauma	
Interventions	1	Carbamazepine: therapeutic range of blood drug levels required for the management of epileptic and affectively ill patients; averages were continuously between 6.44-7.07 micrograms/mL for carbamazepine and 1.07-1.24 micrograms/mL for 10.11-epoxycarbamazepine) N = 10*
	2	Placebo: no further details N = 10* Concomitant psychotherapy: Supportive atheoretical psychotherapy was administered to all patients Concomitant pharmacotherapy: No further details
Outcomes	Interpersonal problems: SCL-90-INT Impulsivity: Acting-out scale Anger: SCL-90-H05 Psychotic symptoms: BPRS, SCL-90-PSY, SCL-90-PAR Depression: HDRS-24 Anxiety: SCL-90-ANX General psychiatric pathology: GAS, SCL-90-PST Attrition Adverse effects: asked for by the non-blind clinician	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Of the 20 patients enrolled in the trial, we randomized 10 into the CBZ [carbamazepine] group and 10 received PLC [placebo]" (De la Fuente 1994, p. 481)
Allocation concealment?	Unclear	Quote: "The study was carried out by two clinicians. One of them [...] was blind and performed all the clinical and psychometric assessments. The other one [...] who was not blind to the drug treatment, correctly adjusted the plasma levels of CBZ and asked for adverse side effects." (De la Fuente 1994 p. 480) Comment: Unclear, who exactly enrolled patients.
Blinding? self-rated outcomes	Unclear	Quote (Eur Neuropsychopharmacol, 1994, 4): "[...] active or placebo treatment. We administered it in a single daily dose at 10 p.m. [...]] The study was carried out by two clinicians. One of them [...] was blind [...]. The

		other one who was not blind to the drug treatment, correctly adjusted the plasma levels of CBZ and asked for adverse side effects.” (p. 480) Comment: Unclear, if opaque capsules were used, and if a pseudo-adjustment of the placebo dose was done
Blinding? observer-rated outcomes	Yes	Quote: “The study was carried out by two clinicians. One of them [...] was blind to the drug treatment and performed all the clinical and psychometric assessments. [...] We instructed the patients to not communicate side effects to the blind clinician.” (De la Fuente 1994, p. 480)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: “Of the 20 patients enrolled in the trial, we randomized 10 into the CBZ group and 10 received PLC. [...] Two patients receiving CBZ dropped out. [...] No patient on PLC dropped out.” (De la Fuente 1994, p. 481). Reasons for early termination: Dramatic increase in frequency and intensity of aggression (against self and others) : 2 in carbamazepine group, 0 in placebo group Comment: Reasons for early termination specified (De la Fuente 1994, p.481). However, it remains unclear if the reported continuous outcomes confer to LOCF analyses. We decided to use the same numbers of patients for which dichotomous outcomes were reported. For dichotomous outcomes, lacking numbers of patients were imputed as having the unfavourable results
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of ‘Yes’ or ‘No’
Free of other bias?	Yes	Quote: “Patients underwent a psychotropic washout period of at least 10 days before the beginning of the treatment period (15 days for tricyclic antidepressants and monoamine oxidase inhibitor agents) . No patient had taken neuroleptics in the 2-month period before the study.” (De la Fuente 1994, p. 480)
Bias due to sponsoring improbable?	Unclear	No details about funding or sponsoring provided.

Frankenburg 2002

Methods	Design: RCT Allocation: Randomised in a 2:1 manner; tablets were supplied in numbered bottles containing drug or placebo as determined by a prearranged random number sequence Blinding: Double-blind; one investigator was given either real drug blood levels or sham levels (in case of placebo) and adjusted the dose according to perceived response, reported or sham level, and side effects Duration: 6 months Setting: Outpatient
Participants	Diagnosis: BPD (DSM-IV; DIPD-IV borderline module) + bipolar II disorder (DSM-IV) Age: Valproate semisodium group: mean age 27.3 (SD 7.4), placebo group: mean age 26.4 (SD 7.3) Sex: 30 F Exclusions: Acutely suicidal patients (i.e. having clear-cut and pressing intent to commit suicide in near future); actively abusing alcohol or drugs; current criteria for major depressive episode or hypomanic episode met; current or lifetime schizophrenia, schizoaffective disorder, psychotic disorder not otherwise specified, bipolar I disorder; patients formerly been treated with valproate semisodium;

subjects who were pregnant, breastfeeding or not using reliable forms of contraception; medically ill, seizure disorder		
Interventions	1 Valproate semisodium: Adjusted to achieve a serum valproate semisodium level of between 50 and 100 mg/L; actual average dose: 850 mg/day (SD 249) or 3.4 tablets/day (SD .9) N = 20* 2 Placebo: tablets; actual average dose: 2.6 tablets/day (SD .5) N = 10* Concomitant pharmacotherapy: No other psychotropic medication allowed Concomitant psychotherapy: No patient was in psychotherapy	
Outcomes	Interpersonal problems: SCL-90-INT Impulsivity: MOAS Anger: SCL-90-HOS Depression: SCL-90-DEP Attrition Adverse effects: weight, menstrual changes, tremors, diarrhea, hair loss, increase in hepatic transaminases, thrombocytopenia	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Quote: "Prearranged random number sequence" (p. 443)
Allocation concealment?	Yes	Quote: "Tablets were supplied in numbered bottles containing drug or placebo as determined by a prearranged random number sequence" (p.443)
Blinding? self-rated outcomes	Yes	Quote: "Tablets were supplied in numbered bottles [...] Each tablet contained either 250 mg of valproate semisodium or matching inert placebo. [...] One of the investigators [...] was given either the real or a sham level (if the subject was receiving placebo). This same investigator met with the subjects for [...] medication checks and adjusted the dose according to perceived response, reported or sham level, and side effects
Blinding? observer-rated outcomes	Unclear	Comment: No information given on who exactly assessed outcomes
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "Endpoint values [...] are based on last observation carried forward." (p. 444) Comment: reasons for early termination specified (p.444). For dichotomous outcomes, lacking numbers of patients were imputed as having the unfavourable result Of the 30 patients enrolled, 11 completed the full 24 weeks of the trial (7 in valproate semisodium group, 4 in placebo group) Reasons for early termination: Lost to follow-up: 9 in the valproate semisodium group, 3 in the placebo group Moved out of the area: 1/0 Inability to use reliable forms of contraception: 1/0 Withdrawal of consent: 1/0 Diarrhea and tremors: 1/0 Development of a major depressive episode: 0/2 Hair loss: 0/1
Free of selective reporting?	Unclear	Insufficient information to permit judgment of 'Yes' or 'No'

Free of other bias?	Yes	Quote: "No other psychotropic medication was allowed during this study." (p. 443)
Bias due to sponsoring improbable?	No	Quote: "Supported by a grant from Abbott Laboratories, Chicago, Ill." (p. 442)

Goldberg 1986

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double Duration: 12 weeks (after 1 week placebo washout) Setting: Outpatient
Participants	Diagnosis: BPD and/or schizotypal personality disorder (DSM-III; SIB) and having at least one psychotic symptom Age: Mean 32 years (no SD given) Sex: 29 F, 21 M Exclusions: Current alcoholism or drug addiction, schizophrenia, mania, melancholia, severe hepatic, renal, or cardiovascular disease, organic brain syndrome, mental retardation, history of epilepsy or seizures, glaucoma, severe hypertensive or hypotensive cardiovascular disease, severe metabolic disorders
Interventions	1 Thiothixene: mean final dose 8.67 mg/day (range 2 mg/day - 35 mg/day) N = 24* 2 Placebo: no further details, mean pseudo-dose at endpoint 26.38 mg/day (not further specified) N = 26* Concomitant psychotherapy: Not specified Concomitant pharmacotherapy: Patients had to pass one week placebo washout; no further details
Outcomes	BPD severity: SIB-borderline score Interpersonal problems: HSCL-INT Anger: HSCL-HOS Psychotic symptoms: SIB-psychotic Depression: HSCL-DEP Mental health status: GAS Attrition Adverse effects

Notes

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Randomly allocated to thiothixene or placebo" (p. 681)
Allocation concealment?	Unclear	Comment: No information given
Blinding? self-rated outcomes	Yes	Quote: "Both agents were provided in identical-appearing capsules containing 5 mg of thiothixene hydrochloride or an equivalent amount of lactose for placebo. The initial dose for all patients was one capsule [...] and on each succeeding visit the dose was increased by one capsule unless side-effects or marked improvement intervened. A maximum dose of 40 mg, or eight capsules, was to be allowed [...]. (p. 682)
Blinding? observer-rated outcomes	Yes	Quote: "Both agents were provided in identical-appearing capsules containing 5 mg of thiothixene hydrochloride or an equivalent amount of lactose for placebo. The initial dose for all patients was one capsule [...] and on each succeeding visit the dose was increased

		by one capsule unless side-effects or marked improvement intervened. A maximum dose of 40 mg, or eight capsules, was to be allowed [...]. (p. 682) Comment: Trial described as “double-blind” (p. 681)
Incomplete outcome data addressed? All outcomes	Yes	Quote: “Patients who terminated their participation early were assessed at that point and those assessments were taken as their endpoints.” (p. 682) Of the 50 patients enrolled, 40 completed treatment (17 in thiothixene group, 23 in placebo group) Reasons for early termination: Adverse effects: 7 in thiothixene group, 0 in placebo group Lack of efficacy: 0/3 Continuous data based on LOCF
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of ‘Yes’ or ‘No’
Free of other bias?	Yes	Quote: “One-week placebo washout” (p. 681)
Bias due to sponsoring improbable?	Unclear	No details about funding or sponsoring provided

Hallahan 2007

Methods	Design: RCT Allocation: Randomised, according to computer-generated list Blinding: Double-blind; an independent colleague dispensed identically looking capsules according to computer-generated list, code was only revealed after completion of data collection Duration: 12 weeks Setting: Outpatient	
Participants	Diagnosis: Patients with recurrent self-harm, recruited at an accident and emergency department where they had presented acutely with self-harm; additionally, participants had to have a lifetime history of at least one other self-harm episode. Actually, 71% of all participants satisfied DSM-IV BPD criteria as assessed by SCID-II Age: Mean 30.6 years Sex: 17 M, 32 F Exclusions: Current addiction, substance misuse, psychosis, eating disorder, dyslipidaemia, treatment, diet or illness known to interfere with study drug, weight loss > 10% during previous 3 months, taking supplements containing omega-3 fatty acids of consuming fish more than once per week, changes to, or introduction of psychotropic medication during previous 6 weeks pregnancy	
Interventions	1	Omega-3 fatty acid: 1.2 g/day of eicosapentaenoic acid (E-EPA)+ 0.9 g/day of decosahexaenoic acid (DHA) N = 22*
	2	Placebo: capsules contained 99% corn oil and a 1% E-EPA + DHA mixture, ensuring blindness by also causing ‘fishy breath’, the most frequent side-effect of the active drug N = 27* Concomitant psychotherapy: Patients actually receiving psychotherapy were not eligible for study participation Concomitant pharmacotherapy: Patients could continue to receive standard psychiatric care and had changes to their psychotropic medication as prescribed (53.1% of participants actually did). Patients with changes to or introduction of psychotropic medication during the 6 weeks prior to screening were not eligible
Outcomes	Suicidal ideation: Number of patients with OAS-M-suicidality subscale score of 1 or higher, indication at least slight suicidal tendencies Self-mutilating behaviour: Number of patients with episodes of self-harm during treatment Depression: Number of patients with at least 50% and 70% reduction of depressive pathology as assessed by both BDI and Ham-D	

Attrition: Number of non-completers

Notes

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Quote: "computer-generated list" (p. 119)
Allocation concealment?	Yes	Quote: "An independent colleague dispensed either active or placebo capsules according to a computer-generated list."
Blinding? self-rated outcomes	Yes	Quote: "Participants were prescribed four identical capsules of either active agent or placebo [...] Placebo ensured a degree of equality in the incidence of 'fishy breath', the most frequent side-effect of taking active treatment." (p. 119)
Blinding? observer-rated outcomes	Yes	Quote: "identical capsules [...] Placebo ensured a degree of equality in the incidence of 'fishy breath' [...] An independent colleague dispensed [...] capsules according to a computer-generated list. The code was only revealed to the researchers once data collection was complete." (p. 119)
Incomplete outcome data addressed? All outcomes	Unclear	Comment: LOCF used, reasons for early termination specified (p. 120) Of the 49 patients enrolled, 39 completed treatment (19 of the 22 allocated to active treatment, 20 of the 27 allocated to placebo) Reasons for early termination: Left district: 1 in active group, 2 in placebo group Lost to follow-up: 2 in active group, 2 in placebo group Admitted to psychiatric hospital: 0 in active group, 2 in placebo group Refused to continue treatment: 0 in active group, 1 in placebo group Dichotomous outcomes calculated on basis of the ITT sample
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	No	Quote: "Patients had changes to their psychotropic medication as prescribed by their treating agency." (p. 118)
Bias due to sponsoring improbable?	Yes	Quote: "Pronova (now Epax) AS, Lysaker, Norway, provided the active preparation and placebo but were not otherwise involved in the study." (p. 118) Quote: "B.H. [i.e. first author] received salary support from the Department of Psychiatry, University of Illinois at Chicago, USA." (p. 122)

Hollander 2001

Methods

Design: RCT
Allocation: Randomised, no further details
Blinding: Double-blind; treating psychiatrist was kept blind to patient medication, blood valproate levels were read and dose adjustments to both valproate semisodium and placebo were determined by a psychiatrist not seeing patients for this study

Duration: 10 weeks (no washout reported) Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-IV; SCID-II) Age: Mean 38.6 years (SD = 10.37, range 18 - 62) Sex: 11 F, 10 M Exclusions: Current suicidal ideation, current substance abuse, current major depression, bipolar disorder type I or II, psychotic disorders, medical or neurologic illness, pregnancy
Interventions	1 Valproate semisodium: dose sufficient to maintain blood valproate level at 80 micrograms/mL or the highest tolerated dose; mean endpoint blood valproate level 64.57 micrograms/mL (SD 15.21, range 47-85 micrograms/mL) N = 12** 2 Placebo: no further details N = 4** Concomitant psychotherapy: Not specified Concomitant pharmacotherapy: Not specified
Outcomes	Impulsivity: AQ, OAS-M-aggression Anger: OAS-M-irritability Suicidal behaviour: OAS-M-suicidality Depression: BDI Mental health status: non-responders (CGI-I score of 3 or more) Attrition
Notes	**Initially 21 subjects entered the study, only 16 were randomised to a treatment group without giving reasons Continuous outcomes based on ITT (LOCF) Of the 16 patients randomised, 6 completed treatment (6 in valproate semisodium group, 0 in placebo group) Reasons for early termination: All patients dropped out owing to either lack of efficacy or impulsive decisions, none dropped out owing to side effects

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Patients were randomly assigned [...] at an approximate ratio of 2:1" (p. 201)
Allocation concealment?	No	Quote: "Although the planned patient assignment ratio was 2:1 [...], the ratio was actually 3:1" (p.202) Comment: First, the authors say that there was an approximate ratio of 2:1 randomisation was planned. However, it remains unclear why the actual ratio turned out to be 3:1, even if taking the small number (16) of participants into account. A 2:1 ratio assignment would have been feasible
Blinding? self-rated outcomes	Unclear	Quote: "The treating psychiatrist was kept blind to patient medication, blood valproate levels were read and dose adjustments to both valproate semisodium and placebo were determined by a psychiatrist not seeing patients for this study." (p. 201) Comment: No information given if opaque capsules were used, and if the placebo pseudo-dose was also "adjusted"
Blinding? observer-rated outcomes	Unclear	Quote: "clinician-rated outcome measures [...] based on the average of the ratings of the treating psychiatrist and independent evaluator (a psychologist blind to side effects as well as to medication group)" (p. 201) Comment: No information given on who exactly assessed observer-rated outcomes
Incomplete outcome data addressed? All outcomes	No	Quote: "Patients taking valproate semisodium had a 50% dropout rate [...] versus 100% dropout in the placebo group. [...] No patients dropped out owing to side effects; all dropped out owing to either lack of efficacy or impulsive decisions. [...]" (p. 201) Comment: LOCF used (p. 202) Initially 21 subjects entered the study, only 16 were randomised to a treatment group without giving reasons Of the 16 patients

		randomised, 6 completed treatment (6 in divalproex group, 0 in placebo group) Reasons for early termination: "No patients dropped out owing to side effects; all dropped out owing to either lack of efficacy or impulsive decisions. [...]" (p. 201)
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Unclear	Comment: Not specified if there was a washout-period preceding the trial or if concomitant psychotropic medication was allowed
Bias due to sponsoring improbable?	No	"Supported in part by grants from the National Institute of Mental Health (1 RO3 MH58168-01A1), Richville, Md. (Dr. Hollander); Abbott Laboratories, Abbott Park, Ill. (Dr. Hollander); the National Center for Research Resources, National Institutes of Health (5 MO1 RR00071), Rockville, Md., for the Mount Sinai General Clinical Research Center; and the Seaver Foundation and the PBO Foundation, New York, N.Y." (p. 199)

Leone 1982

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Drugs were supplied in identical opaque capsules Duration: 6 weeks Setting: Outpatient
Participants	Diagnosis: BPD (DSM-III, no further details) Age: Loxapine group: mean 29.5 years (range 16-54 years), chlorpromazine group: mean 32 years (range 16-59 years) Sex: 48 F, 32 M Exclusions: Using sedatives or tranquilisers, having been treated with psychotropic drugs within 48 hours of beginning treatment with study drugs, allergy/hypersensitivity to study drugs, organic brain syndrome, mental retardation, severe medical disease
Interventions	- Loxapine: capsules of 5 mg, starting dose one or two capsules daily, increased based on symptom severity and tolerance; maximum dose 12 capsules/d; mean final dose 13.5 mg/day, overall mean daily dose 14.4 mg N = 40* - Chlorpromazine: capsules of 50 mg, starting dose one or two capsules daily, increased based on symptom severity and tolerance; maximum dose 12 capsules daily; mean final dose 105 mg/day, overall mean daily dose 110 mg N = 40* Concomitant psychotherapy: Not specified Concomitant pharmacotherapy: Patients did not receive any other psychotropic medication during the study; nighttime sedatives were limited to flurazepam and chloral hydrate
Outcomes	BPD severity Affective instability: POMS Psychotic symptoms: BPRS Mental health status: CGI, Systematic Nurses' Observation of Psychopathology (SNOOP) Attrition Adverse effects: Recorded upon appearance in terms of data of onset, intensity, duration, and any remedial action Unable to use outcome data (except for Attrition)
Notes	*As randomised

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Matched groups [...] Subjects [...] were selected randomly to receive loxapine or chlorpromazine. [...] There were 24 women and 16 men in each treatment group." (p. 148) Comment: probably matching procedure used
Allocation concealment?	Unclear	Comment: No information given.
Blinding? self-rated outcomes	Yes	Quote: "drugs were supplied in identical opaque capsules" (p. 148) Comment: No self-rated outcomes used.
Blinding? observer-rated outcomes	Unclear	Comment: No information given. Within this review, only the outcomes of Attrition and adverse effects, that were "recorded upon appearance" (p. 148), were used. For these, the review authors assume the risk of bias as moderately
Incomplete outcome data addressed? All outcomes	Yes	Continuous outcomes based on available cases Of the 80 patients enrolled, 69 completed at least 3 weeks of treatment and were included (34 in loxapine group, 35 in placebo group) Reasons for early termination: Did not follow study procedures: 4 in loxapine group, 4 in chlorpromazine group Had to be admitted to hospital within 3 days: 2 in loxapine group, 1 in chlorpromazine group Comment: Only dichotomous outcomes used here, for which dropped-out patients were imputed as having the negative outcome
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Unclear	Quote: "Patients did not receive any other psychotropic medication during the study; nighttime sedatives were limited to flu-raxepam and chloral hydrate." (p. 148) Comment: Actually, patient could thus receive concomitant sedatives, but it is not specified how many did
Bias due to sponsoring improbable?	No	Quote: "This study was supported by a grant from Lederle Laboratories, Pear River, New York." (p. 148)

Linehan 2008

Methods	Design: RCT Allocation: Content of tablets was determined by a random number sequence Blinding: Patients, psychotherapists, pharmacotherapist, and assessment interviewers were kept naive to medication assignment Duration: 24 weeks, last assessment after week 21, however Setting: Outpatient
Participants	Diagnosis: BPD according to DSM-IV (SCID-II, PDE) + BPD criterion for inappropriate anger met + score of 6 or higher on the irritability scale of the OAS-M Age: Overall mean age 36.8 years (SD=9.0) Sex: 24 F

Exclusions: Episode of self-inflicted self-injury including suicide attempts during 8 weeks prior to screening, current diagnosis of schizophrenia, bipolar I disorder, schizoaffective disorder, major depressive disorder with psychotic features or other psychotic disorder, substance dependence during last 6 months, mental retardation, seizure disorder, pregnant women or planning to be, breastfeeding

Interventions	1	DBT + olanzapine (allowed dosage range: 2.5 to 15 mg/day; mean daily dose 4.46 mg/day, SD 1.16) N = 12*
	2	DBT + placebo (dose was adjusted in response to perceived response and side effects, no further details) N = 12*
		Data refer to the intention to treat sample Concomitant psychotherapy: All participants received DBT Concomitant pharmacotherapy: n.s.
Outcomes		Suicidal ideation: Number of patients with high suicidality score on the OAS-M suicidality subscale Self-mutilating behaviour: number of patients with self-injury Depression: Ham-D Attrition Adverse effects: Weight gain (lb), remaining data on adverse effects not usable
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Quote: "random number sequence" (p. e2)
Allocation concealment?	Yes	Quote: "each tablet contained either 5 mg of olanzapine or matching inert placebo as determined by a random number sequence" (p. e2)
Blinding? self-rated outcomes	Yes	Quote: "Patients, psychotherapists, pharmacotherapist, and assessment interviewers were kept naive to medication assignment. (p. e2)
Blinding? observer-rated outcomes	Yes	Quote: "Patients, psychotherapists, pharmacotherapist, and assessment interviewers were kept naive to medication assignment. At the end of the study, the pharmacotherapist and interviewers were unable to guess group assignment above chance." (p. e2)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "Outcomes were intent-to-treat analyses" (p. e3) Comment: Reasons for early termination specified (p. e4); dropped-out patients were imputed as having the negative outcome
Free of selective reporting?	Unclear	Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Unclear	Quote: "To enhance compliance, tablets were given in [...] prescription bottles programmed to sound a sequence of alarms when medications were due, terminating only when the medication top was removed." Comment: Compliance was thus controlled for Comment: Not specified if concomitant medication was allowed or not
Bias due to sponsoring improbable?	No	Quote: This research was supported by a grant from Eli Lilly and Co., Protocol F1D-US-X173, to Dr. Linehan; by Remind Rx Medication Compliance Systems; and by a contribution of electronic pill bottles from IBV Technologies, Seattle, Wash. [...] Dr. Linehan is a consultant for, has received grant/research support and honoraria from, and is a member of the speakers/ advisory board from Eli Lilly. Drs. McDavid, Brown, Savrs, and Gallop report no additional financial or

other relationships relevant to the subject of this article." (p. 999)

Loew 2006

Methods	Design: RCT Allocation: Randomisation was carried out confidentially by the clinic administration, tablets were supplied in numbered boxes Blinding: Double-blind, blind medication which constituted either the active drug or placebo Duration: 10 weeks Setting: Outpatient
Participants	Diagnosis: BPD (SCID-II) Age: Mean age active drug group: 24.9 (SD = 5.3), placebo group 25.6 (SD = 5.7) Sex: 56 F Exclusions: Currently suicidal patients, abusing alcohol or drugs, schizophrenia, severe somatic illness, current use of topiramate or other psychotropic medication, or psychotherapy
Interventions	1 Topiramate: 200 mg/day N = 28* 2 Placebo: analogous pseudo-dose N = 28* Concomitant psychotherapy: Not allowed Concomitant pharmacotherapy: Any other psychotropic medication not allowed
Outcomes	Interpersonal problems: SCL-90-R-INT Anger: SCL-90-R-HOS Psychotic symptoms: SCL-90-R-PAR, SCL-90-R-PSY Depression: SCL-90-R-DEP Anxiety: SCL-90-R-ANX General psychiatric pathology: SCL-90-R-GSI Attrition Adverse effects: non-structured questionnaire

Notes

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Randomization was carried out confidentially by the clinic administration" (Loew 2006, p. 63)
Allocation concealment?	Yes	Quote: "Tablets were supplied in numbered boxes." (Loew 2006, p. 63)
Blinding? self-rated outcomes	Yes	Quote: "blinded medication" (Loew 2006, p. 63), "subjects [...] were blinded regarding [...] assignment" (Loew 2006, p. 63)
Blinding? observer-rated outcomes	Yes	Quote: "blinded medication" (Loew 2006, p. 63), "clinicians were blinded regarding [...] assignment" (Loew 2006, p. 63)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "Fifty-nine subjects were eligible to take part in the study [...] 56 patients were required [...] randomization was carried out [...] with a 1:1 assignment to the active drug (N = 28) and placebo (N = 28)" (Loew 2006, p. 63) Of the 56 patients enrolled, 52 completed treatment (27 in topiramate group, 25 in placebo group) Reasons for early termination: Absent more than twice for weekly evaluation: 1 in the topiramate group, 3 in the placebo group

		LOCF used, reasons for early termination specified (Loew 2006, p. 63) Comment: Not clear, why or how the 56 participants were finally chosen out of the 59 potential participants
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "exclusion criteria included [...] the current use of topiramate or other psychotropic medication." (p. 62)
Bias due to sponsoring improbable?	Unclear	Quote: "The study was planned and conducted independently] of any institutional influence and approved by the clinic's ethics committee in accordance with the Declaration of Helsinki and ethical laws pertaining to the medical professions." (Loew 2006, p. 63)

Montgomery 1979/82

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double-blind, flupenthixol decanoate and placebo drawn from identical matching ampoules Duration: 24 weeks Setting: Outpatient	
Participants	Diagnosis: Patients admitted following a suicidal act, having a history of 2 or more previous documented suicidal acts; more than 75% BPD (23 out of 30*; DSM-III, clinical interview)* Age: Flupenthixol group: 38.2 years (SD = 15.53), placebo group: 31.9 (SD = 11.0)* Sex: 21 F, 9 M* Exclusions: Overt schizophrenia or depression, organic illness	
Interventions	1	Flupenthixol decanoate intramuscular: 20 mg every four weeks N= 14
	2	Placebo: drawn from identical ampoules N= 16 Concomitant psychotherapy: All patients attended the special crisis intervention clinic within two weeks of the index suicidal act Concomitant pharmacotherapy: Not specified
Outcomes	Suicidal behaviour: Number of participants in each group with/without suicidal act within the 6 months of treatment Adverse effects: Assessed by standard reporting form	
Notes	*Only reported for the completers	

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Patients were randomly allocated" (Montgomery 1979, p. 227)
Allocation concealment?	Unclear	Comment: No information given
Blinding? self-rated outcomes	Yes	Quote: "intramuscular flupenthixol decanoate or placebo drawn from identical matching ampoules" (Montgomery 1979, p. 227)
Blinding? observer-rated outcomes	Yes	Quote: "intramuscular flupenthixol decanoate or placebo drawn from identical matching ampoules" (Montgomery 1979, p. 227)

Incomplete outcome data addressed? All outcomes	Yes	Quote: "To preserve blindness patients with significant Parkinsonian side effects were removed from the trial and counted as drop outs." (Montgomery 1979, p. 227) Reported dichotomous outcomes based on the completer sample (no further details on drop-out patients concerning diagnosis, sex, and age) Of the 37 patients enrolled, 30 completed treatment (4 drop-outs in the active group leaving 14 completers, 3 drop-outs in the placebo group leaving 16 completers) Reasons for early termination: Parkinsonian side effects: 2 in flupenthixol group/0 in placebo group No reason given: 2/3 Comment: Only dichotomous data used in this review, drop-outs were imputed as having the negative outcome
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Unclear	Comment: Not specified if there was a washout-period or if concomitant psychotropic medication was allowed
Bias due to sponsoring improbable?	Unclear	No details about funding/sponsoring provided.

Montgomery 81/82/83

Methods	Design: RCT Allocation: Randomised Blinding: No further details Duration: 6 months Setting: Outpatient	
Participants	Diagnosis: Patients admitted following a suicidal act, having a history of 2 or more previous documented suicidal acts; more than 75% BPD (30 out of 38*; DSM-III, clinical interview) Age: Mianserin group: mean age 35.1 (SD = 12.24), placebo group: mean age 36.2 (SD = 13.38)* Sex: 26 F, 12 M* Exclusions: Overt schizophrenia or depression, organic illness	
Interventions	1	Mianserin (30 mg nightly) N = 17 completers of N = 29 allocated to mianserin
	2	Placebo N = 21 completers of N = 29 allocated to placebo Concomitant psychotherapy: Patients were followed up in a clinic with back up from social workers, community nurses and a crisis intervention team Concomitant pharmacotherapy: Not specified
Outcomes	Suicidal behaviour: Number of participants in each group with/without act of self-harm within the 6 months of treatment Attrition	
Notes	*Only reported for the completers	
<i>Risk of bias</i>		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "randomly allocated" (Montgomery 1981, p. 787)
Allocation concealment?	Unclear	Comment: No information given

Blinding? self-rated outcomes	Unclear	Quote: "double-blind conditions" (Montgomery 1981, p. 787)
Blinding? observer-rated outcomes	Unclear	Quote: "double-blind conditions" (Montgomery 1981, p. 787)
Incomplete outcome data addressed? All outcomes	Yes	Dichotomous outcomes used here are based on the ITT sample, dropped-out patients were imputed as having the negative outcome Comment: High drop-out rate (20 out of 58; Montgomery 1981, p. 787), but reasons not specified, nor to which treatment group the lost patients belonged. Therefore, drop-outs could not be imputed in categorial outcomes as having the negative outcome
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Unclear	Quote: "Compliance was checked by tablet count." (Montgomery 1983, p. 184S) Comment: Not specified if there was a washout-period or if concomitant psychotropic medication was allowed
Bias due to sponsoring improbable?	Unclear	No details about funding/sponsoring provided.

Nickel 2004

Methods	Design: RCT Allocation: Randomisation was carried out confidentially by the clinic administration and arranged so that twice as many subjects would be treated with the active drug as with placebo Blinding: Tablets were supplied in numbered boxes Duration: 8 weeks Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-IV, SCID-II) Age: Topiramate: mean age 25.5 years, placebo 26.6 years (no further details) Sex: 31 F Exclusions: Actively suicidal patients, abusing alcohol or drugs, major depression, schizophrenia, bipolar disorder, current use of topiramate or other psychotropic medication, psychotherapy, pregnant or planning to become, somatically ill	
Interventions	1	Topiramate: 250 mg/day N = 21*
	2	Placebo: Matching N = 10* Concomitant psychotherapy: Not allowed Concomitant pharmacotherapy: Not allowed
Outcomes	Impulsivity: STAXI-anger-out Anger: STAXI-trait anger Attrition Adverse effects: Non-structured questionnaire	
Notes		
<i>Risk of bias</i>		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Randomization was carried out confidentially by the clinic administration. " (Nickel 2004, p. 1516)
Allocation concealment?	Yes	Quote: "Tablets were supplied in numbered boxes." (Nickel 2004, p. 1516)

Blinding? self-rated outcomes	Yes	Quote: "blinded medication" (Nickel 2004, p. 1516), "subjects [...] were blinded regarding [...] assignment" (Nickel 2004, p. 1516)
Blinding? observer-rated outcomes	Yes	Quote: "blinded medication" (Nickel 2004, p. 1516), "clinicians [...] were blinded regarding [...] assignment" (Nickel 2004, p. 1516)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "Two subjects, who failed to appear 2 to 3 times for the weekly evaluations, dropped out of the study, and their data were not further analyzed. Finally, data from 29 women [...] were evaluated." (Nickel 2004, p. 1516). Continuous outcomes based on available case analysis Of the 31 patients enrolled, 29 completed treatment Reasons for early termination: Failed to appear at least 2 times for weekly evaluation, no further details: 2 in topiramate group/0 in placebo group
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "exclusion [...] current use of topi-ramate or other psychotropic medication"
Bias due to sponsoring improbable?	Unclear	Quote: "The authors report no financial affiliation or other relationship relevant to the subject matter of this article." (Nickel 2004, p. 1515)

Nickel 2005

Methods	Design: RCT Allocation: Randomisation was carried out confidentially by the clinic administration, tablets were supplied in numbered boxes Blinding: Double-blind (both clinician and subjects were blinded) Duration: 8 weeks Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-IV, SCID-II) Age: Mean age 29.1 years Sex: 44 M Exclusions: Actively suicidal, currently fulfilling criteria for an addictive illness, severe major depression, acute psychosis, bipolar disorder, current use of topiramate or other psychotropic medication, current psychotherapy, somatically ill	
Interventions	1	Topiramate: 250 mg/day N = 22*
	2	Placebo: matching N = 22* Concomitant psychotherapy: Not allowed Concomitant pharmacotherapy: Psychotropic medication not allowed
Outcomes	Impulsivity: STAXI-anger out Anger: STAXI-trait anger Attrition Adverse effects: Weight	
Notes		
<i>Risk of bias</i>		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Randomization was carried out confidentially by the clinic administration. " (Nickel 2005, p. 496)

Allocation concealment?	Yes	Quote: "Tablets were supplied in numbered boxes." (Nickel 2005, p. 496)
Blinding? self-rated outcomes	Yes	Quote: "blinded medication" (Nickel 2005, p. 496) "subjects [...] were blinded regarding [...] assignment" (Nickel 2005, p. 496)
Blinding? observer-rated outcomes	Yes	Quote: "blinded medication" (Nickel 2005, p. 496), "clinicians [...] were blinded regarding [...] assignment" (Nickel 2005, p. 496)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "Forty-eight subjects were eligible to take part in the study [...] 44 patients were required [...] randomization was carried out confidentially by the clinical administration [...] 1:1 randomisation ratio for topiramate (TG, N = 22) versus placebo treatment (N = 22)" (Nickel 2005, p. 496) Comment: Unclear, why or how the 44 participants were finally chosen out of the 47 potential participants Quote: "Two subjects from the placebo group failed to appear appear more than twice for the weekly evaluations and dropped out of the study; their data were not further analyzed. Thus, data from 42 men (42 out of 44) were evaluated." (Nickel 2004, p. 1516). Comment: Reasons for early termination not further specified. Continuous outcomes based on available case analysis. For dichotomous data, drop-outs were imputed as having the negative outcome Of the 44 patients enrolled, 42 completed treatment (22 in the active group, 20 in the placebo group) Reasons for early termination: Failed to appear more than twice for weekly evaluation, no further reasons given: 0 in the active group, 2 in the placebo group
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "reasons for exclusion were [...] the current use of topiramate or other psychotropic medication." (Nickel 2004, p. 495)
Bias due to sponsoring improbable?	Unclear	Quote: "The study was planned and conducted in accordance with the Declaration of Helsinki and ethical laws pertaining to the medical professions and its design approved by Ethikkommission der ROMED Kliniken KG. All subjects gave written informed consent. The study was conducted independent of any institutional influence and was not funded, and there were no conflicts of interest." (Nickel 2004, p. 496)

Nickel 2006

Methods	Design: RCT Allocation: Randomisation was carried out confidentially by the clinic administration and arranged so that twice as many subjects would be treated with the active drug as with placebo Blinding: Tablets were supplied in numbered boxes, both patients and clinicians were blinded Duration: 8 weeks Setting: Outpatient
Participants	Diagnosis: BPD (DSM-IV, SCID-II) Age: Aripiprazole group: mean age 22.1 years (SD = 3.4), Placebo group: mean age 21.2 years (SD = 4.6) Sex: 43 F, 9 M Exclusions: Current suicidal ideation, schizophrenia, current use of aripiprazole or another psychotropic medication, current psychotherapy, pregnancy, planned pregnancy or sexual activity without contraception, severe somatic illness
Interventions	1 Aripiprazole: 15 mg/day

- N = 26*
- 2 Placebo: matching dose
- N = 26*
- Concomitant psychotherapy: Not allowed
- Concomitant pharmacotherapy: Not allowed

Outcomes	Interpersonal problems: SCL-90-R-INT (t-value transformed) Impulsivity: STAXI-anger out Self-mutilating behaviour: Number of patients with/without self-injury during the 8 week treatment Anger: SCL-90-R-HOS (t-value transformed), STAXI-trait anger Psychotic symptoms: SCL-90-R-PAR, SCL-90-R-PSY (both t-value transformed) Depression: SCL-90-R-DEP (t-value transformed), Ham-D Anxiety: SCL-90-R-ANX (t-value transformed), HARS General psychiatric pathology: SCL-90-R-GSI (t-value transformed) Adverse effects: Serious side effects, suicidal acts	
Notes	*As randomised	
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "The random assignment was carried out confidentially by the clinic administration." (Nickel 2006, p. 835)
Allocation concealment?	Yes	Quote: "Tablets were supplied in numbered boxes." (Nickel 2006, p. 835)
Blinding? self-rated outcomes	Yes	Quote: "subjects received medication in a blinded manner" (Nickel 2006, p. 835), "the subjects [...] were blinded regarding the assignment (Nickel 2006, p. 835)
Blinding? observer-rated outcomes	Yes	Quote: "subjects received medication in a blinded manner" (Nickel 2006, p. 835), "the clinicians [...] were blinded regarding the assignment (Nickel 2006, p. 835)
Incomplete outcome data addressed? All outcomes	Yes	Of the 52 patients enrolled, 47 completed treatment Quote: "Five subjects who missed more than two weekly evaluations dropped out. " (Nickel 2006, p. 835) Comment: Reasons for drop-out not further specified. Quote: "according to the intent-to-treat principle performed with the last observation carried forward" (Nickel 2007, p. 1025) Continuous outcomes based on ITT sample (LOCF) Dichotomous outcomes based on ITT sample Reasons for early termination: Failed to appear more than twice for weekly evaluation, no further reasons given: 5 subjects, no further details
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "Criteria for exclusion [...] current use of aripiprazole or another psychotropic medication" (p. 8349
Bias due to sponsoring improbable?	Unclear	Quote: "The study was planned and conducted in accordance with the Declaration of Helsinki and ethical laws pertaining to the medical profession, and its design was approved by the clinic's ethics committee. The study was conducted independently of any institutional influence an was not funded." (Nickel 2006, p. 835)

Pascual 2008

Methods	Design: RCT
---------	-------------

Allocation: Randomised, randomisation was performed by blocks of 4 generated using the SPSS software package (SPSS Inc., Chicago, Ill) Blinding: Double, no further details Duration: 12 weeks, following a 2-week baseline period Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-IV, SCID-II, DIB-R) Age: Mean 29.2 years Sex: 49 F, 11 M Exclusions: Schizophrenia, alcohol or other substance dependence, current major depressive episode, bipolar disorder, drug-induced psychosis, organic brain syndrome, mental retardation
Interventions	1 Ziprasidone: flexible dose 40 to 200 mg/day, mean dose 84.1 mg/day (SD 54.4, range 40 - 200 mg/day) N = 30* 2 Placebo: No further details N = 30* Concomitant psychotherapy: Patients participated in weekly, 2-hour, non-specific group psychotherapy sessions Concomitant pharmacotherapy: Allowed to continue with benzodiazepine (max. 40 mg/day), antidepressants, mood stabilisers if initiated prior to inclusion; doses could not be modified
Outcomes	BPD severity: CGI-BPD-global Avoidance of abandonment: CGI-BPD-abandonment Interpersonal problems: CGI-BPD-unstable relations Identity disturbance: CGI-BPD-identity Impulsivity: CGI-BPD-impulsivity, BIS Suicidal ideation: CGI-BPD-suicide Affective instability: CGI-BPD-affect instability Feelings of emptiness: CGI-BPD-emptiness Anger: CGI-BPD-anger Psychotic paranoid symptoms: CGI-BPD-paranoid ideation, BPRS Depression: Ham-D-17, BDI Anxiety: HARS General psychiatric pathology: SCL-90-R-GSI Attrition Adverse effects: treatment-emergent adverse events, EKG, laboratory assessment, UKU Side Effect Rating Scale for extrapyramidal side effects
Notes	*As randomised

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Quote: "Randomization was performed by blocks of 4 generated using the SPSS software package" (p. 604)
Allocation concealment?	Unclear	Comment: No information given
Blinding? self-rated outcomes	Unclear	Quote: "double-blind" (p. 604) Comment: No further information given
Blinding? observer-rated outcomes	Unclear	Quote: "double-blind" (p. 604) Comment: No further information given
Incomplete outcome data addressed? All outcomes	No	Quote: "All analyses were conducted on an intent-to-treat basis. [...] Patients were included in the analyses only if they had a baseline measure and at least 1 postbaseline measure. [...] The end point was based on a last-observation-carried-forward (LOCF) strategy." (p. 604 et seq.) Comment: intent-to-treat data refer to all participants that were randomly assigned and initiated the experimental phase (p. 605) However, it remains unclear for which reason 5 out of the 65 eligible subjects "dropped out during the selection phase" (p. 605) Reasons for drop-out specified and balanced across the two groups, including withdrawal due to "clinician decision/insufficient treatment effect (p. 605 et seq.) Continuous data based on LOCF data of the ITT sample

		Dichotomous data based on ITT sample Of the 60 patients enrolled, 29 completed the full 12 weeks of the trial (13 in ziprasidone group, 16 in placebo group) Reasons for early termination: Need of psychiatric hospitalization: 4 in ziprasidone group/3 in placebo group, Adverse events/patient decision: 9/4, Clinician decision/insufficient treatment effect 3/7, Other reasons: 1/0
Free of selective reporting?	Yes	Comment: The study protocol is available and all of the study's pre-specified primary and secondary outcomes that are of interest in the review are reported in the pre-specified way
Free of other bias?	Unclear	Quote: "Compliance was assessed by direct questioning of patients and by counting the capsules returned at follow-up visits." Quote: "patients were allowed to continue with benzodiazepines [max. 40 mg/day], antidepressants, and mood stabilisers if they had been initiated prior to inclusion, but doses could not be modified during the study." (p. 604)
Bias due to sponsoring improbable?	No	Quote: "This study was supported by grants from the Fondo de Investigacion Sanitaria (Ministry of Health, Spain), the REM-TAP Network, and Pfizer, Madrid, Spain. The authors report no additional financial or other relationships relevant to the subject of this article." (p. 603)

Reich 2009

Methods	Design: RCT Allocation: Randomised (prearranged random number sequence) Blinding: Double Duration: 12 weeks Setting: Outpatient
Participants	Diagnosis: BPD (DSM-IV, DIB-R ≥ 8) Age: Mean 31.2 years Sex: 24 F, 3 M Exclusions: Diagnosis of dementia, psychiatric disorder secondary to a general medical condition, bipolar disorder, or psychotic disorder (schizophrenia, schizoaffective disorder, or mood disorder with psychotic features); diagnosis of substance dependence (active within last 60 days); currently being hospitalized; unstable general medical condition; previous treatment with lamotrigine for 1 week or more; enrollment in a drug study within last 60 days; enrollment in psychotherapy in the last 30 days; active suicidal or homicidal ideation; pregnancy or nursing
Interventions	1 Lamotrigine: flexible dose 25 to 275 mg/day, mean final dose 106.7 mg (range 25–225 mg/day) N = 15* 2 Placebo: No further details N = 12 (one patient of the 13 assigned to placebo was disqualified because of failure to adhere to the study protocol and not included in analyses) Concomitant psychotherapy: Patients enrolled in psychotherapy in the last 30 days were not eligible Concomitant pharmacotherapy: Patients could be taking one antidepressant, but had to have been on a stable dose of this medication for 1 month

Outcomes
 BPD severity: ZAN-BPD total score
 Impulsivity: ZAN-BPD-impulsivity score
 Affective instability: ZAN-BPD-affective instability score, ALS
 Attrition
 Adverse effects: rash

Notes

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Quote: "patients were randomized [...] in a 1:1 manner. This was determined by a prearranged random number sequence." (p. e-3) Twenty-eight patients completed all aspects of assessment before randomization. Fifteen patients were randomized to receive lamotrigine, and 13 patients were assigned to receive placebo." (p. e-3)
Allocation concealment?	Unclear	Insufficient information to permit judgement of 'Yes' or 'No' (unclear, if the number sequence was kept confidentially or if enrolling investigators could possibly foresee assignment)
Blinding? self-rated outcomes	Yes	No self-rated outcomes used for this review
Blinding? observer-rated outcomes	Yes	Quote: "double-blind placebo-controlled study" (e.g. p. e-1); "double-distinction between "prescribing psychiatrist (D.B.R.)" who fixed the dose and "study staff" who made assessments (p. 3)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "One patient in the placebo group was disqualified because of failure to adhere to the study protocol." (p. e-3) Not clear if the reported mean changes are based on the ITT sample or completers only
Free of selective reporting?	Yes	Comment: The study protocol is available and all of the study's pre-specified primary and secondary outcomes that are of interest in the review are reported in the pre-specified way
Free of other bias?	Unclear	Quote: "Patients could be taking one antidepressant, but had to have been on a stable dose of this medication for 1 month." (p. e-2)
Bias due to sponsoring improbable?	No	Quote: "The study was supported by a grant from GlaxoSmithKline." (p. e-5)

Rinne 2002

Methods	Design: RCT, followed by single-blind half crossover and an open treatment phase; only the first RCT phase will be regarded in the following Allocation: Randomised, no further details Blinding: Double-blind, no further details Duration: 6 weeks, patients had to be medication free for at least 2 weeks before entering the trial Setting: Outpatients
Participants	Diagnosis: BPD (DSM-IV, SCID-II) + score of 110 or more on the borderline trait and distress scale of a self-report screener for personality disorders (ADP-IV) + score of 20 or more on the BPDSI Age: 29.2 (SD = 7.6) Sex: 38 F

Exclusions: n.s.		
Interventions	1	Fluvoxamine: 150 mg/day N = 20*
	2	Placebo: No further details N = 18*
		Concomitant psychotherapy: Two patients who began psychotherapy dropped-out the study; thus, psychotherapeutic treatment is likely to not have been allowed
		Concomitant pharmacotherapy: Patients had to stop taking all psychoactive drugs and be medication free for at least 2 weeks before entering the trial (6 weeks for fluoxetine)
Outcomes		Impulsivity: BPDSI-impulsivity Affective instability: BPDSI-rapid mood shifts Anger: BPDSI-anger Attrition Adverse effects: Any, number of subjects experiencing specific adverse events (not used here as data refer to intermediate assessment, whereas post-treatment data are not available)
Notes		*As randomised
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "randomized trial" (p. 2049)
Allocation concealment?	Unclear	Comment: No information given
Blinding? self-rated outcomes	Unclear	Quote: "double-blind" (p. 2049) Comment: No further information given
Blinding? observer-rated outcomes	Unclear	Quote: "double-blind" (p. 2049) Comment: No further information given
Incomplete outcome data addressed? All outcomes	Yes	Quote: "The final study group comprised the 38 subjects eligible for participation" (p. 2049), "an intent-to-treat analysis was performed" (p. 2050) Continuous outcomes based on ITT, BMDP imputation technique used for drop-outs Of the 38 patients enrolled, 35 completed the RCT phase (19 in active drug group, 16 in placebo group) Reasons for early termination: Serious aggravation of self-damaging behaviours: 0 in the fluvoxamine group, 2 in the placebo group Severe side effects: 1/0
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "The participants had to stop taking all psychoactive drugs [...] and they all had to be medication free for at least 2 weeks before entering the trial; the medication -free interval was 6 weeks for fluoxetine." (p. 2049)
Bias due to sponsoring improbable?	No	Quote: "Supported by the De Geestgronden Institute of Mental Health Care, by Stichting tot Steun of Vereniging Bennekom, by national Fund for Menal Health grant 4820, and by Solvay Pharma." (p. 2053)
Salzman 1995		
Methods	Design: RCT Allocation: Randomised, no further details	

	Blinding: Double-blind, active drug and placebo administered in identical capsules Duration: 12 weeks (after 1 week placebo run-in) Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-III-R; DIB-R, SCID-II, clinical interview) Age: Mean age of fluoxetine group: 37.0 (no further details), placebo group: 35.6 (no further details)* Sex: 14 F, 8 M* Exclusions: Self-mutilating behaviours during the past 4 years, recent suicidal behaviour, current suicidal or aggressive behaviour, current substance abuse or excessive daily alcohol use (> 2 drinks/day), history of psychiatric hospitalization, concurrent secondary axis II disorder, major depression or other axis I disorder	
Interventions	1	Fluoxetine: maximum of 60 mg/day, according to needs of the patient and in accordance with package insert guidelines; mean daily dose 40 mg/day Completers: N =13
	2	Placebo: No further details Completers: N = 9 Data are only reported for treatment completers Concomitant psychotherapy: Not specified Concomitant pharmacotherapy: Not allowed
Outcomes	Anger: PDRS-anger, POMS-anger, OAS-M-anger against objects Depression: Ham-D, PDRS-depression, POMS-depression Mental health status: GAS	
Notes	* Completers only	

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "random-assignment comparison" (p. 24)
Allocation concealment?	Unclear	Comment: No information given
Blinding? self-rated outcomes	Yes	Comment: No self-rated outcomes used within this review
Blinding? observer-rated outcomes	Yes	Quote: "Subjects were evaluated by independent observers" (p. 24)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "Thirty-one subjects met criteria for this study; four decided not to enroll and were lost to follow-up. Of 27 subjects who enrolled in the study, 22 completed the trial. One subject dropped out because she wanted assurance that she would be in the medication group; four others dropped out without explanation and were lost to follow-up." (p. 24) Of the 27 patients enrolled, 22 completed treatment Reasons for early termination: Wanted assurance to be in the active drug group: 1 (not specified, which group) Dropped out without explanation: 4 (not specified, which group) Comment: Continuous outcomes based on completer analysis
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "Subjects were not included [...] if they were taking any other psychotropic medication"; "1-week placebo run-in" (p. 24),
Bias due to sponsoring improbable?	Unclear	No details provided.

Schulz 2007

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double-blind; olanzapine started at 2.5 or 5.0 mg/d at investigator's discretion, flexible dose thereafter Duration:12 weeks (after screening period of 2 - 14 days) Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-IV; DIPD-IV) + ZAN-BPD total score of 9 or higher Age: Olanzapine group mean age 31.79 (SD = 9.54), placebo group mean age 31.83 (SD = 9.62) Sex: 223 F, 91 M Exclusions: Bipolar disorder, schizophrenia, major depressive disorder or substance dependence within last 3 months, current PTSD, panic disorder, or obsessive-compulsive disorder	
Interventions	1	Olanzapine: flexible dose, 2.5 to 20 mg/day, mean modal dose 7.09 mg/day N = 150
	2	Placebo: No further details N= 155 Concomitant psychotherapy: Not specified Concomitant pharmacotherapy: No medications with primarily CNS activity (except for protocol-specified benzodiazepines and hypnotics)
Outcomes	BPD severity: number of patients in each group with response/no response, i.e. 50% reduction at least in ZAN-BPD total score Avoidance of abandonment: ZAN-BPD-frantic efforts to avoid abandonment Interpersonal problems: ZAN-BPD unstable interpersonal relationships Identity disturbance: ZAN-BPD-identity disturbance Impulsivity: ZAN-BPD-impulsivity, OAS-M-aggression Suicidal ideation: OAS-M-suicidal ideation Suicidal behaviour: ZAN-BPD-suicidal or self-mutilating behaviour Affective instability: ZAN-BPD-affective instability Feelings of emptiness: ZAN-BPD-chronic feelings of emptiness Anger: ZAN-BPD-intense anger, OAS-M-irritability, SCL-90-R-HOS Dissociative symptoms: ZAN-BPD-paranoid ideation of disassociation Depression: MADRS General psychiatric pathology: SCL-90-R GSI Mental health status: Sheehan Disability Scale-total, GAF Attrition Adverse effects: weight, Simpson-Angus Scale, BARS, AIMS	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Quote: "patients [...] were randomly assigned to treatment" (Eli Lilly, 2008, p. 15) , "All participants, study site personnel and investigators were masked to randomisation codes." (Schulz 2008, p. e1) Comment: Randomisation conducted centrally
Allocation concealment?	Yes	Quote: "All participants, study site personnel and investigators were masked to randomisation codes." (Schulz 2008, p. e1)
Blinding? self-rated outcomes	Yes	Quote: "All participants, study site personnel and investigators were masked to randomisation codes." (Schulz 2008, p. e1)
Blinding? observer-rated outcomes	Yes	Quote: "All participants, study site personnel and investigators were masked to randomisation codes." (Schulz 2008, p. e1)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "Analyses were done on an intent-to-treat basis [...] In general, LOCF mean change analyses" (Eli Lilly, 2008, p. 5)

		Quote: "Of the 314 randomized patients, 305 had both a baseline and a non-missing post-baseline observation and were thus qualified for the primary efficacy analysis." (Eli Lilly, p. 16) Comment: Unclear, what "non-missing post-baseline observation" exactly means. However, discontinuing participants were enclosed in the 305 participants whose results were analysed using LOCF Continuous outcomes based on LOCF/ITT 314 patients were enrolled and randomly allocated. Outcomes refer partly to all of them, partly to 310 or 305 patients. No further details given
Free of selective reporting?	No	Comment: Several outcome measures (secondary and adverse events) are reported that were not pre-specified according to the study protocol
Free of other bias?	Unclear	Quote: "No medication with primarily CNS activity (except for protocol-specified benzodiazepines and hypnotics)" (p. EliLilly, p. 4)
Bias due to sponsoring improbable?	No	Quote: "This study was sponsored by Eli Lilly. S.C.S. has received honorarium from Eli Lilly, AstraZeneca and Bristol-Meyers Squibb; grant fees from Eli Lilly, AstraZeneca, Abbott, MIND Institute and the NIMH; and consultation fees from Eli Lilly, AstraZeneca and Vanda. H.C.D., Q. T., Y.T., D.L. and S.C. are employed by Lilly Research Laboratories." (p. e-1)

Simpson 2004

Methods	Design: RCT Allocation: Randomised block assignment, equal number of patients with major depressive disorder, PTSD, or both were assigned to each treatment condition in order to minimize the possible confound to treatment response Blinding: Double-blind, a non-treating study psychiatrist was available to break the blind in the event of a clinical emergency, but didn't occur Duration: 12 weeks (after a 1-week placebo run-in) Setting: Partial hospitalization
Participants	Diagnosis: BPD (DSM-IV, SCID-II), patients had to meet at least one BPD criterion pertaining to affective instability or anger and one pertaining to impulsivity Age: Fluoxetine completers mean age 39.78 (SD = 9.81), placebo completers mean age 32.73 (SD = 10.76) Sex: 25 F Exclusions: Primary diagnosis of substance dependence, seizure disorder, unstable medical conditions, lifetime history of schizophrenia or bipolar disorder, MAOI treatment in the prior 2 weeks, previous adequate trial of fluoxetine, pregnancy, lactating women, unwillingness to use effective contraception
Interventions	1 DBT (weekly 1-hour sessions of individual DBT, weekly 2-hour skills group, round-the-clock emergency consultation availability) + fluoxetine 40 mg/day N= 12 2 DBT (weekly 1 -hour sessions of individual DBT, weekly 2-hour skills group, round-the-clock emergency consultation availability) + placebo (no further details) N= 13 Data are only available for the 20 completers (fluoxetine N = 9, placebo N = 11) Concomitant psychotherapy: Patients were recruited from a partial hospital program, all received DBT as depicted above Concomitant pharmacotherapy: Only other psychotropic allowed was 50 to 100 mg/day trazodone for insomnia

Outcomes
 Impulsivity: OAS-M-aggression, STAXI-anger out
 Suicidal ideation: OAS-M-suicidality
 Self-mutilating behaviour: OAS-M-assault against self
 Psychotic symptoms/dissociation: DES
 Depression: BDI
 Anxiety: STAI-trait
 Mental health status: GAF
 Attrition: number of patients lost after randomisation

Notes

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "randomized block assignment minimized the possible confound of co-morbid axis-I presentations expected to response to fluoxetine by assignment of an equal number of patients with major depressive disorder, posttraumatic stress disorder, or both to each treatment condition." (p. 380)
Allocation concealment?	Unclear	Comment: No information given.
Blinding? self-rated outcomes	Unclear	Quote: "This study was double-blind" (p. 380) Comment: No information given how blinding of participants was attempted, especially in light of the day-clinic setting with possibly shared group therapy
Blinding? observer-rated outcomes	Yes	Quote: "This study was double-blind" (p. 380), "A non-treating study psychiatrist was available to break the blind in event of a clinical emergency." (p. 381) Comment: In contrast, the treating clinician was probably blind
Incomplete outcome data addressed? All outcomes	No	Of the 25 patients enrolled, 12 were randomised to fluoxetine and 13 to placebo. 20 completed treatment (9 in fluoxetine group, 11 in placebo group) Reasons for early termination: Negative experience of the placebo washout period, which led to a reversal of their willingness to tolerate a potential assignment to the placebo condition: 3 in fluoxetine group, 0 in placebo group Sought hospitalization at another facility: 0/ Intolerable lack of improvement: 0/1 Comment: Reasons for early termination specified (p. 381) Continuous outcomes are only reported for study completers, while drop-outs could be imputed as having the negative outcome for dichotomous data
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "Diary card records of pill ingestion were reviewed, and pill counts were made as a compliance measure." (p. 381) Quote: "1-week placebo run-in" (p. 380), "the only other medication allowed was 50 to 100 mg/day of trazodone for insomnia." (p. 381)
Bias due to sponsoring improbable?	No	Quote: "Support for this study was provided by the Department of Psychiatry and Human Behaviour at Brown Medical School and Eli Lilly."

Soler 2005

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double-blind, no further details Duration: 12 weeks (after a 4 weeks selection phase during which the pre-intervention baseline was established but no therapeutic intervention was given) Setting: Outpatient
---------	---

Participants	Diagnosis: BPD (DSM-IV, SCID-II and DIB-R), CGI-S score of at least 4 Age: DBT + olanzapine group mean Age: 27.57 (SD = 6.3), DBT + placebo group mean Age: 26.33 (SD = 5.4) Sex: 52 F, 8 M Exclusions: Comorbid unstable axis I disorder, women not using medically accepted contraception	
Interventions	1	DBT + olanzapine: weekly 150-minute skills training group sessions, phone calls + olanzapine flexible dose between 5 to 20 mg/day (mean dose 8.83 mg/day, SD = 3.8) N = 30*
	2	DBT + placebo: weekly 150-minute skills training group sessions, phone calls + placebo (no further details) N = 30*
		Concomitant psychotherapy: All patients received DBT as depicted above Concomitant pharmacotherapy: Subjects could continue treatment with benzodi-azepines, antidepressants, and mood stabilisers, but doses could not be modified
Outcomes	Impulsivity: Behavioural biweekly reports of episodes of impulsivity/aggressive behaviour Suicidal behaviour/self-mutilating behaviour: behavioural biweekly reports of episodes of self-injuring behaviour/suicide attempts Depression: Ham-D Anxiety: HARS Mental health status: CGI-S Attrition Adverse effects: As reported by patients, scales assessing extrapyramidal side effects, weight, cholesterol levels	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "randomly assigned to receive dialectical behaviour therapy plus either olanzapine or placebo on a 1:1 ratio" (p. 1222)
Allocation concealment?	Unclear	Comment: No information given.
Blinding? self-rated outcomes	Unclear	Quote: "double blind [...] study
Blinding? observer-rated outcomes	Unclear	Quote: "double blind [...] study
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "All analyses were conducted on an intent-to-treat basis. The endpoint was based on a last-observation-carried-for-ward strategy. Patients were included in the analyses only if they had a baseline measure and at least one post-baseline measure." (p. 1222) "Quote: Sixty subjects were randomly assigned to dialectical behaviour therapy plus olanzapine or placebo and started the experimental phase; 42 subjects (70%) completed the study. There were no be-tween-group differences regarding demographic variables or concomitant treatments at baseline. Neither dialectical behaviour therapy intervention time nor dropout rates differed significantly between the two groups (eight of the 30 patients who received olanzapine versus 10 of the 30 who received placebo dropped out before the end of the study." (p. 1222 et seq.) Comment: reasons for drop-out given; numbers balanced across groups Continuous outcomes based on ITT (LOCF) Of the 60 patients enrolled, 42 completed treatment (22 in active drug group, 20 in placebo group) Reasons for early termination: No reasons given

Free of selective reporting?	Yes	Comment: The study protocol is available and all of the study's pre-specified primary and secondary outcomes that are of interest in the review are reported in the pre-specified way
Free of other bias?	Unclear	Quote: "Patients could continue treatment with benzodiazepines, antidepressants, and mood stabilisers, but doses could not be modified." (p. 1222)
Bias due to sponsoring improbable?	No	Quote: "Supported by grants from the Fondo de Investigacion Sanitaria (Ministry of Health, Spain) and from Eli Lilly and Co. Madrid." (p. 1223)

Soloff 1989

Methods	Design: RCT Allocation: Randomised Blinding: Double-blind, no further details Duration: 5 weeks (after 1-week washout) Setting: Inpatient (after 3 weeks some allowed to complete as outpatients)	
Participants	Diagnosis: BPD (DSM-III, DIB), GAS score of 50 or less and either score of 17 or higher on the Ham-D or 66 or greater on the IMPS Age: Mean 25.1 years, no further details Sex: 68 F, 22 M Exclusions: Schizophrenia, mania, related disorders, chronicity of illness, organicity	
Interventions	1	Amitriptyline: Mean dose after 3 weeks of treatment: 149.1 mg/day, plasma levels of 240.4 ng/mL amitriptyline + nortriptyline (SD = 99.4) N = 29 2
	2	Haloperidol: Mean dose after 3 weeks of treatment 4.8 mg/day, plasma level of 8.66 ng /mL (SD = 3.7) N = 28 3
	3	Placebo: No further details N = 28 Concomitant psychotherapy: Patients were treated as psychiatric inpatients for at least 3 weeks, no further details Concomitant pharmacotherapy: Biperiden hydrochloride (2 mg) was allowed as needed for extrapyramidal reactions
Outcomes	Interpersonal problems: SCL-90-INT Impulsivity: Ward Scale of Impulsive Action Patterns, BIS, STIC Anger: SCL-90-HOS, BDHI, BDH Psychotic symptoms: SCL-90-PAR, SCL-90-PSY, IMPS, SSI Depression: SCL-90-DEP, Ham-D, BDI Anxiety: SCL-90-ANX General psychiatric pathology: SCL-90-GSI Mental health status: GAS Attrition Adverse effects	
Notes		
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Patients were randomly assigned" (Soloff 1986, p. 692)
Allocation concealment?	Yes	Quote: "Numbered tablets [...] were given" (Soloff 1986, p. 692)

Blinding? self-rated outcomes	Unclear	Quote: "Double-blind [...] trial" (Soloff 1989, p. 239)
Blinding? observer-rated outcomes	Yes	Quote: "Weekly ratings by two 'blind investigators', an onward psychiatrist serving as the nonblind psychiatrist (for safety)." (Soloff 1986, p. 693)
Incomplete outcome data addressed? All outcomes	Yes	Quote: "Five patients failed to complete the minimum two weeks on medication needed for inclusion in outcome analysis, one taking amitriptyline, three taking haloperidol, and one taking placebo." (Soloff 1989, p. 242) Continuous outcomes based on LOCF/ITT A minimum of 2 weeks receiving medication was required to include data for end-point analysis Of the 90 patients enrolled, 85 completed treatment (29 in amitriptyline group, 28 in haloperidol group, 28 in placebo group) Reasons for early termination: Failed to complete the minimum 2 weeks on medication needed for inclusion in outcome analysis (1 in amitriptyline group, 3 in haloperidol group, 1 in placebo group) Comment: Reasons for drop-out not further specified. Total number of drop-outs small, though, and balanced across groups
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "Patients were kept free of all medication for 7 days" (Soloff 1989, p. 239) , "Biperiden hydrochloride (2 mg) was allowed as needed for extrapyramidal reactions" (Soloff 1986, p. 692)
Bias due to sponsoring improbable?	Yes	Quote: "This work was supported by NIMH grants 35392, MHCRC 30915, and MH00658." (p. 245)

Soloff 1993

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double-blind, no further details Duration: 5 weeks (after 1-week washout) Setting: Patient in the hospital for a minimum of 2 weeks and after discharge were seen weekly as outpatients	
Participants	Diagnosis: BPD (DSM-III-R, DIB), GAS score of 50 or less and either score of 17 or higher on the Ham-D or 66 or greater on the IMPS Age: Mean 26.7 years (SD=7.2) Sex: 82 F, 26 M Exclusions: Drug- and/or alcohol-related deficits or physical dependence, evidence of central nervous system disease, physical disorders of known psychiatric consequence, borderline mental retardation	
Interventions	1	Haloperidol up to 6 mg/day (six tablets); average dose after 3 weeks of medication 3.93 mg/day (SD = 0.65); mean plasma level by 4 weeks 5.29 ng/mL (SD = 4.04) N = 30
	2	Phenelzine sulfate up to 90 mg/day (six tablets); average dose after 3 weeks of medication 60.45 mg/day (SD = 9.55); mean plasma level by 3 weeks 77.54% platelet MAO inhibition (SD = 16.97) N = 34
	3	Placebo up to six tablets; average dose after 3 weeks of medication 4.31 tablets/day (SD = 0.6) N = 28 Concomitant psychotherapy: Not specified, patients were inpatients, some were allowed after 2 weeks to complete as outpatients

Concomitant pharmacotherapy: Patients were at the start kept free of medication for at least 7 days in order to washout street drugs or prescribed medications

Outcomes	BPD severity: Borderline Syndrome Index Interpersonal problems: ADDS - rejection sensitivity Impulsivity: Ward Scale of Impulsive Action Patterns, BIS, STIC Anger: SCL-90-HOS, BDHI, BDHI, ADDS-reactivity Psychotic symptoms: SCL-90-PAR, SCL-90-PSY, IMPS, SSI Depression: SCL-90-DEP, Ham-D, BDI Anxiety: SCL-90-ANX General psychiatric pathology: SCL-90-GSI Mental health status: GAS Attrition Adverse effects: Weight gain	
Notes	*Completers only	
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Patients were randomly assigned. " (p. 378)
Allocation concealment?	Unclear	Quote: "double-blind [...] trial" (p. 377)
Blinding? self-rated outcomes	Yes	Average daily doses of medication, including placebo pseudo-dose, are given (p. 380) Comment: The measures undertaken to ensure blinding seem elaborated and are described in detail, so the blinding of participants seems to have been thoroughly ensured
Blinding? observer-rated outcomes	Yes	Quote: "Medication could be increased up to six tablets (haloperidol, 6 mg; phenelzine sulfate, 90 mg; placebo, six tablets)" (p. 378) Average daily doses of medication, including placebo pseudo-dose, are given (p. 380) Comment: The measures undertaken to ensure blinding seem elaborated and are described in detail, so the blinding of the rating study personnel seems to have been thoroughly ensured
Incomplete outcome data addressed? All outcomes	Yes	Quote: "Sixteen patients failed to complete the minimum 3 weeks of medication required for end-point analysis (p. 380) Comment: Reasons for these drop-outs not further specified. Total number of dropouts small, though, and balanced across groups Continuous outcomes based on all cases with a minimum of3 weeks of medication exposure Of the 108 patients enrolled, 92 completed treatment (30 in haloperidol group, 34 in phenelzine group, 28 in placebo group) Reasons for early termination: Relating to medication assignment (e.g. side effects), clinical worsening, factors unrelated to the protocol; not specified by group Patients failing to complete the minimum 3 weeks of medication required for end-point analysis: 6 in the haloperidol group, 4 in the phenelzine group, 6 in the placebo group
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "Patients were at the start kept free of medication for at least 7 days to [...] washout street drugs or prescribed medications." (p. 378)
Bias due to sponsoring improbable?	Yes	Quote: "This study was supported by National Institute of Mental Health grant MH35392 and by Clinical Research Center grant MH30915." (p. 697)

Tritt 2005

Methods	Design: RCT Allocation: Randomization in secrecy by the clinic administration so that twice as many subjects would be treated with the active drug compared to placebo Blinding: Double-blind, tablets were supplied in numbered bottles Duration: 8 weeks Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-IV, SCID-II) Age: Lamotrigine group mean age 29.4, no further details; placebo group mean age 28.9, no further details Sex: 27 F Exclusions: Actively suicidal, abusing alcohol or drugs, major depression, bipolar disorder, schizophrenia, current use of lamotrigine or other psychotropic medication, psychotherapy, pregnant or planning to be or not using contraception, somatically ill	
Interventions	1	Lamotrigine: final dose 200 mg/day (one blinded capsule medication daily) N= 18 2
		Placebo, one blinded capsule medication daily N=9 Concomitant psychotherapy: Other psychotropic medication not allowed Concomitant pharmacotherapy: Not allowed
Outcomes	Impulsivity: STAXI-anger out Anger: STAXI-trait Attrition Adverse effects: Non-structured questionnaire, patients were asked to note down any new symptoms, weight	
Notes	Continuous outcomes based on ITT data (LOCF)	
Risk of bias		
Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "Randomization was carried out confidentially in secrecy by the clinic administration section" (Tritt 2005, p. 288)
Allocation concealment?	Yes	Quote: "Tablets were supplied in numbered boxes." (Tritt 2005, p. 288)
Blinding? self-rated outcomes	Yes	Quote: "Each individual received one blinded capsule medication daily [...] Both subjects and clinicians were blinded regarding assignment." (Tritt 2005, p. 288)
Blinding? observer-rated outcomes	Yes	Quote: "Each individual received one blinded capsule medication daily [...] Both subjects and clinicians were blinded regarding assignment." (Tritt 2005, p. 288)
Incomplete outcome data addressed? All outcomes	No	Quote: "Thirty-eight subjects were eligible to take part in the study [...] The necessary sample size was calculated [...] This resulted in a group size of n = 27 patients [...] active drug (n=18) compared to placebo (n = 9)" (Tritt 2005, p. 288) Comment: Not clear, why or how the 27 participants were finally chosen out of the 38 potential participants
Free of selective reporting?	No	Comment: All outcomes (i.e. one assessment instrument) reported as planned to be assessed are also reported. However, it seems implausible to use only one assessment instrument in such a complex trial. There is no protocol available to check the pre-defined outcome measure(s)
Free of other bias?	Yes	Quote: "reasons for exclusion were [...] current use of lamotrigine or other psychotropic medication " (p. 288)

Bias due to sponsoring improbable?	Yes	Quote: "The study was conducted independently of any institutional influence and was not funded." (p. 288)
------------------------------------	-----	--

Zanarini 2001

Methods	Design: RCT Allocation: Numbered bottles containing drug or placebo as determined by a random number sequence Blinding: Tablets were supplied in numbered bottles containing drug or placebo Duration: 6 months Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-IV; DIB-R) Age: Olanzapine group mean age 27.6 years (SD = 7.7), placebo group mean age 25.8 years (SD = 4.5) Sex: 28 F Exclusions: Actively abusing alcohol or drugs, acutely suicidal, current or lifetime schizophrenia, schizoaffective disorder, bipolar disorder, medically ill, seizure disorder, pregnant or planning to be, breastfeeding, not using reliable forms of contraception, having been treated with olanzapine, being prescribed any psychotropic medication that patients thought was helpful	
Interventions	1	Olanzapine 2.5 mg/day at beginning, adjusted according to perceived response and side effects, mean dose at endpoint 5.33 mg/day (SD = 3.43); endpoint mean number of tablets/day 1.1 (SD = 0.68) N= 19
	2	Placebo: Endpoint mean number of tablets/day 1.2 (SD = 0.75) N = 9 Concomitant psychotherapy: Not specified Concomitant pharmacotherapy: No other psychotropic medication allowed
Outcomes	BPD severity Avoidance of abandonment Interpersonal problems: SCL-90-INT Identity disturbance Impulsivity Suicidal ideation Suicidal behaviour Self-mutilating behaviour Affective instability Feelings of emptiness Anger: SCL-90-HOS Psychotic symptoms: SCL-90-PAR, SCL-90-PSY, PANSS Dissociative symptoms: DES Depression: SCL-90-DEP, Ham-D, Anxiety: SCL-90-ANX General psychiatric pathology: SCL-90-GSI Mental health status: GAF Attrition Adverse effects: Weight, Simpson-Angus Scale, BARS, AIMS, structured questionnaire	
Notes		
<i>Risk of bias</i>		
Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Quote: "random number sequence" (p. 850)
Allocation concealment?	Yes	Quote: "Tablets were supplied in numbered bottles containing drug or placebo as determined by a random number sequence. " (p. 850)
Blinding? self-rated outcomes	Yes	Quote: "Each tablet contained either 2. 5 mg of olanzapine or matching inert placebo. [...] Both subjects

		and clinicians were blinded to olanzapine/placebo assignment. The blind was broken after the acquisition of all endpoint data for all subjects." (p. 850)
Blinding? observer-rated outcomes	Yes	Quote: "Each tablet contained either 2.5 mg of olanzapine or matching inert placebo. [...] Both subjects and clinicians were blinded to olanzapine/placebo assignment. The blind was broken after the acquisition of all endpoint data for all subjects." (p. 850)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "Thirty subjects completed all aspects of pre-randomization assessment. However, 2 of these subjects were excluded [...] because it was determined that they were responding well to a selective serotonin reuptake inhibitor. Twenty-eight subjects entered the trial and were randomly assigned [...] All [...] completed at least 2 post-baseline visits and were included in all subsequent analyses." (p. 851) Of the 28 patients enrolled, 9 completed treatment (8 in olanzapine group, 1 in placebo group) Reasons for early termination: Sedation: 1 in olanzapine group, 0 in placebo group Increased anxiety or depression: 3/2 Perceived weight gain: 2/0 Lost to follow-up: 5/6 Continuous outcomes based on ITT sample (LOCF) Comment: Overall high drop-out rate but adequately addressed
Free of selective reporting?	No	Quote: "Due to the small number of subjects, results pertaining to secondary outcome measures will not be reported." (p. 851)
Free of other bias?	Yes	Quote: "excluded if [...] currently were being prescribed any psychotropic medication that they thought was helping" (p. 850)
Bias due to sponsoring improbable?	No	Quote: "Supported, in part, by a grant from Eli Lilly." (p. 849)

Zanarini 2003

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double-blind Duration: 8 weeks Setting: Outpatient	
Participants	Diagnosis: BPD (DSM-IV; DIB-R) Age: Mean age 26.3 years (SD = 6.2) Sex: 30 F Exclusions: Major depressive episode, current or lifetime schizophrenia, schizoaffective disorder, bipolar I or bipolar II disorder	
Interventions	1	Ethyl-eicosapentaenoic acid (E-EPA): 1 g/day N = 20
	2	Placebo: Mineral oil N= 10 Concomitant psychotherapy: Not specified Concomitant pharmacotherapy: Not specified
Outcomes	Impulsivity: MOAS Depression: MADRS Attrition Adverse effects: Structured questionnaire	
Notes		

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "randomly assigned [...] 2:1 randomization ratio." (p. 168)
Allocation concealment?	Unclear	Comment: No information given.
Blinding? self-rated outcomes	Unclear	Quote: "double-blind study" (p. 167), "subjects received two capsules per day [...] each contained either [...] E-EPA or placebo" (p. 167)
Blinding? observer-rated outcomes	Unclear	Quote: "double-blind study" (p. 167), "subjects received two capsules per day [...] each contained either [...] E-EPA or placebo" (p. 167)
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "The three subjects who discontinued their participation (two taking E-EPA and one taking placebo) did so because of life events unrelated to the study." (p. 168) of the 30 patients enrolled, 27 completed treatment (18 in E-EPA group, 9 in placebo group) Reasons for early termination: life events unrelated to the study: 2 in E-EPA group, 1 in placebo group Comment: Continuous outcomes are based on completers only.
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "Potential subjects were excluded if they were [...] currently being prescribed any psychotropic medication" (p. 167)
Bias due to sponsoring improbable?	No	Quote: "Capsules were supplied by Laxdale Pharmaceuticals (Stirling, U.K.)" (p. 167), "Supported by an Independent Investigator Award from the National Alliance for Research on Schizophrenia and Depression to Dr. Zanarini." (p. 169)

Zanarini 2004

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double-blind, dose was adjusted by an unblinded psychiatrist according to perceived response and side effects Duration: 8 weeks Setting: Outpatient
Participants	Diagnosis: BPD (DSM-IV, DIB-R) Age: Mean age 23 years (SD = 5.7) Sex: 45 F Exclusions: Current major depression, current or lifetime schizophrenia, schizoaffective disorder, bipolar disorder
Interventions	1 Fluoxetine: Mean dose at endpoint 15.0 mg/day (SD = 6.5, range 10.0 - 30.0 mg/day) N = 14 2 Olanzapine: Mean dose at endpoint 3.3 mg/day (SD = 1.8, range 2.5 - 7.5 mg/day) N = 16 3 Fluoxetine + olanzapine: Mean dose at endpoint 12.7 mg/day fluoxetine (SD = 5.9, range 10.0 - 30.0 mg/day) and 3.2 mg/day olanzapine (SD = 1.5, range 2.5 - 7.5 mg/day) N = 15 Concomitant psychotherapy: Not specified

Concomitant pharmacotherapy: Not specified

Outcomes	Impulsivity: OAS-M total Depression: MADRS Attrition Adverse effects: Weight, Simpson-Angus Rating Scale, BARS, AIMS, structured questionnaire
----------	--

Notes

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Unclear	Quote: "The randomization procedure was designed to assign equal numbers of subjects to the 3 treatment groups." (p. 904)
Allocation concealment?	No	Quote: "Forty-five subjects entered the trial and were randomized to fluoxetine (N = 14), olanzapine (N = 16), or OFC (N = 15)." (p. 905) Comment: Allocation probably not adequately concealed, since equal numbers were intended (cf. to item above), but group sizes differed, actually
Blinding? self-rated outcomes	Yes	Quote: "Dose was adjusted by an un-blinded psychiatrist according to perceived response and side effects. Both subjects and raters were blinded to study assignment. The blind was broken after acquisition of all endpoint data for all subjects." (p. 904)
Blinding? observer-rated outcomes	Yes	Quote: "Dose was adjusted by an un-blinded psychiatrist according to perceived response and side effects. Both subjects and raters were blinded to study assignment. The blind was broken after acquisition of all endpoint data for all subjects." (p. 904)
Incomplete outcome data addressed? All outcomes	Unclear	Comment: Reasons for drop-out specified (p. 905), but outcome data were only reported for completers Of the 45 patients enrolled, 42 completed treatment (13 in fluoxetine group, 16 in olanzapine group, 13 in fluoxetine + olanzapine group) Reasons for early termination: Onset of a number of psychosocial stressors culminating in a suicide gesture: 1/0/0 Dizziness and headaches: 0/0/1 Lost to follow-up: 0/0/1
Free of selective reporting?	Unclear	Comment: Insufficient information to permit judgment of 'Yes' or 'No'
Free of other bias?	Yes	Quote: "Potential subjects were excluded if they [...] were currently being prescribed any psychotropic medication" (p. 904)
Bias due to sponsoring improbable?	No	Quote: "Supported by a grant from Eli Lilly, Indianapolis, Ind." (p. 903)

Zanarini 2007

Methods	Design: RCT Allocation: Randomised, no further details Blinding: Double-blind, no further details Duration: 12 weeks (after a 2 weeks screening period) Setting: Outpatient
Participants	Diagnosis: BPD (DSM-IV-TR, DIPD-IV), Zan-BPD total score of 9 or more Age: Mean 32.98 (SD = 10.83) Sex: 332 F, 119 M Exclusions: Bipolar disorder, schizophrenia, major depressive disorder within last 3 months, substance dependence within last 3 months, current PTSD, current panic disorder, current obsessive-compulsive disorder, comorbid Cluster A Axis II PD, active suicidality, pregnancy

Interventions	1 Olanzapine: 2.5 mg/day N = 150 2 Olanzapine: 5 to 10 mg/day, mean dose 6.66 mg/day (SD = 2.91) N = 148 (106 F, 42 M) 3 Placebo: Daily capsules N = 153 (117 F, 36 M)
Outcomes	BPD severity: Number of patients in each group with response/no response, i.e. 50% reduction at least in ZAN-BPD total score Avoidance of abandonment: ZAN-BPD-frantic efforts to avoid abandonment Interpersonal problems: ZAN-BPD unstable interpersonal relationships, SCL-90-R-INT Identity disturbance: ZAN-BPD-identity disturbance Impulsivity: ZAN-BPD-impulsivity, OAS-M-aggression Suicidal ideation: OAS-M-suicidal ideation Suicidal behaviour: ZAN-BPD-suicidal or self-mutilating behaviour Affective instability: ZAN-BPD-affective instability Feelings of emptiness: ZAN-BPD-chronic feelings of emptiness Anger: ZAN-BPD-intense anger, OAS-M-irritability, SCL-90-R-HOS Dissociative symptoms: ZAN-BPD-paranoid ideation of disassociation, SCL-90-R-PAR Depression: MADRS, SCL-90-R-DEP Anxiety: SCL-90-R-ANX General psychiatric pathology: SCL-90-R GSI Mental health status: Sheehan Disability Scale-total, GAF Attrition Adverse effects: Weight, Simpson-Angus Scale, BARS, AIMS
Notes	*As randomised

Risk of bias

Item	Authors' judgement	Description
Adequate sequence generation?	Yes	Quote: "patients were randomised to 1 of 3 treatment groups" (Eli Lilly, 2008, p. 3) Comment: randomisation conducted centrally
Allocation concealment?	Yes	Comment: Probably done, since this RCT was conducted in parallel with the olanzapine flexible-dose trial (Schulz 2008), where a randomisation code was used
Blinding? self-rated outcomes	Yes	Quote: "Double-blind treatment" (Eli Lilly, 2008, p. 1) Comment: Probably equally managed as in Schulz 2008
Blinding? observer-rated outcomes	Yes	Quote: "Double-blind treatment" (Eli Lilly, 2008, p. 1) Comment: Probably equally managed as in Schulz 2008
Incomplete outcome data addressed? All outcomes	Unclear	Quote: "451 were randomly assigned (148 [] to olanzapine 5-to-10 mg/day treatment group, 150 [] to olanzapine 2.5-mg/day treatment group, and 153 to placebo)" (Eli Lilly 2008, p. 14), "last observation carried forward" (Eli Lilly 2008, p. 3) Continuous data based on LOCF data Dichotomous data based on ITT sample Of the 451 patients enrolled, 294 completed the full 12 weeks of the double-blind treatment phase (97/103/94) Comment: In this review, only the groups receiving olanzapine 5 to 10 mg/day or placebo group were included
Free of selective reporting?	Yes	Comment: The study protocol is available and all of the study's pre-specified (primary and secondary) outcomes that are of interest in the review are reported in the pre-specified way
Free of other bias?	Unclear	Comment: Not specified if there was a washout period or if concomitant psychotropic medication was allowed. Probably, medication with primarily CNS

		activity was not allowed (except for protocol-specified benzodiazepines and hypnotics), as was the case for the Schulz 2007 trial
Bias due to sponsoring improbable?	No	Eli Lilly was the study sponsor. Most study results used here are from the company's study report (the remaining references were either clinical trial register entries or congress abstracts and did not provide detailed data)

ADDS - Atypical Depression Inventory
 ADP-IV - Assessment of DSM-IV Personality Disorders
 AIAQ - Anger, Irritability, and Assault Questionnaire
 AIMS - Abnormal Involuntary Movement Scale
 ALS - Affective Lability Scale
 AQ - Assault Questionnaire
 BARS - Barnes Akathisia Rating Scale
 BDHI - Buss-Durkee Hostility Inventory
 BID - Beck Depression Inventory
 BIS - Barratt Impulsiveness Scale
 BPDSI - Borderline Personality Disorder Severity Index
 BPRS - Brief Psychiatric Rating Scale
 CGI - Clinical Global Impressions Scale
 CGI-BPD - Clinical Global Impressions Scale modified for borderline personality disorder
 CGI-I Clinical Global Impressions Scale-global improvement
 CGI-S Clinical Global Impressions Scale-Severity of Illness
 DBT - Dialectical Behaviour Therapy
 DES - Dissociative Experiences Scale
 DIB - Diagnostic Interview for Borderline Patients
 DIB-R - Diagnostic Interview for Borderline Patients-Revised
 GAF - Global Assessment of Functioning
 GAS - Global Assessment Scale
 Ham-D - Hamilton Rating Scale for Depression
 HARS - Hamilton Anxiety Rating Scale
 HDRS-24 - Hamilton's 24-item Depression Rating Scale
 HSCL - Hopkins Symptom Checklist
 HSCL-DEP - depression
 HSCL-HOS - anger-hostility
 HSCL-INT - interpersonal sensitivity
 HSCL-PSY - psychotic
 IMPS - Inpatient Multidimensional Rating Scale
 ITT - intention to treat
 LOCF - Last observation carried forward
 MADRS - Montgomery-Asberg Depression Rating Scale
 MAOI - monoamine oxidase inhibitor agents
 MOAS - Modified Overt Aggression Scale
 OAS-M - Overt Aggression Scale-Modified
 PANSS - Positive and Negative Syndrome Scale

PDRS - Personality Disorder Rating Scale
 POMS - Profile of Mood States
 SCID-II - Structured Clinical Interview for DSM Axis II Personality Disorders
 SCL-90 - Symptom Checklist-90
 SCL-90-R - Symptom Checklist-90 - Revised
 SCL-90-ANX - anxiety
 SCL-90-DEP - depression
 SCL-90-GSI - global severity index
 SCL-90-HOS - hostility
 SCL-90-INT - interpersonal sensitivity
 SCL-90-PAR - paranoid ideation
 SCL-90-PST - positive symptom total
 SCL-90-PSY - psychoticism
 SIB - Schedule for Interviewing Borderlines
 SSI - Schizotypal Symptom Inventory
 SSRI - Selective Serotonin Reuptake Inhibitor
 STAXI - State-Trait Anger Expression Inventory
 STIC - Self Report Test of Impulse Control
 TCA - tricyclic antidepressant
 ZAN-BPD - Zanarini Rating Scale for Borderline Personality Disorder
 *
 as randomised

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Bellino 2005	Allocation: not randomised; open trial
Bellino 2006a	Allocation: not randomised; open trial
Bellino 2006b	Comparison: fluoxetine vs. fluoxetine + Interpersonal Therapy; testing the effects of additional psychotherapy as compared to pharmacotherapy alone
Benedetti 1998	Allocation: not randomised; open trial
Bohus 1999	Allocation: not randomised; open trial
Chengappa 1999	Allocation: not randomised; retrospective chart review
Coccaro 1997	Patients: less than 70% BPD
Cornelius 1990	Allocation: not randomised; open trial
Cornelius 1993	Investigates the effects of continuation therapy in patients of the Soloff 1989 trial, but only responders were allowed to enter this study
Cowdry 1988	Randomised cross-over trial, no separate data first period available
Frankenburg 1993	Allocation: not randomised; open trial
Hilger 2003	Allocation: not randomised; case series
Hollander 2005	Data not sufficient for effect size calculation concerning any outcome of interest
Koenigsberg 2003	Participants: less than 70% BPD
La Malfa 2003	Participants: less than 70% BPD; separate data on BPD patients available but not sufficient for effect size calculation

Links 1990	Participants: BPD characteristics, mean DIB score 9.47 (SD = 0.75); exact number of BPD patients unclear Data: no separate data for first arm of cross-over trial (randomised cross-over trial)
Markovitz 1991	Allocation: not randomised; open trial
Markovitz 1995a	Data: not sufficient for effect size calculation
Markovitz 1995b	Allocation: not randomised; open trial
Norden 1989	Allocation: not randomised; open trial
Parsons 1989	Participants: less than 75% BPD; no separate data available
Philipsen 2004a	Intervention: administration in acute dissociative states only, not for continuous treatment
Philipsen 2004b	Allocation: not randomised; open trial Intervention: administered in acute states of aversive inner tension only, not for continuous treatment
Rocca 2002	Allocation: not randomised; open trial
Russell 2003	Participants: PD but not BPD patients
Schulz 1999	Allocation: not randomised; open trial
Serban 1984	Participants: less than 70% BPD
Soloff 1986	Midpoint analysis of the Soloff 1989 trial which has been included
Soloff 1987	Compares haloperidol responders to patients receiving placebo of the Soloff 1989 trial
Verkes 1998	Participants: suicide attempt repeaters, not clear how many patients actually had a BPD
Ziegenhorn 2009	Within-subject crossover-design, no separate data for first study periods available

Characteristics of ongoing studies [ordered by study ID]

AstraZen NCT00254748

Trial name or title	The effect of quetiapine on psychotic-like symptoms in borderline personality disordered patients: a randomised placebo-controlled trial
Methods	RCT
Participants	BPD according to DSM-IV, including criterion 9; 18-55 years, in- or outpatients
Interventions	8 weeks of quetiapine (flexible dose between 200 mg/day and 600 mg/day) vs. placebo
Outcomes	Psychotic-like symptoms, severity of psychiatric symptoms; mood, anger, impulsiveness, hostility, anxiety
Starting date	June 2004
Contact information	AstraZeneca Netherlands
Notes	

Bohus NCT00124839

Trial name or title	Evaluation of the efficacy of the opioid antagonist naltrexone on the incidence and intensity of flashbacks and dissociative states in patients with borderline personality disorder
Methods	RCT
Participants	BPD according to DSM-IV, Dissociation Experience Scale (DES) score 25 at least, not actively abusing opiates, no other psychopharmacological treatment for at least two weeks

Interventions	Naltrexone vs. placebo
Outcomes	Reduction of dissociative symptoms, flashbacks, self-injurious behaviour, psychopathology (depression, anxiety, anger, borderline symptoms), safety
Starting date	October 2005
Contact information	Bohus M, Central Institute of Mental Health, Mannheim, Germany
Notes	This study has been terminated in April 2008. (Difficulties in recruiting enough subjects)

Casas NCT00437099

Trial name or title	Efficacy of omega-3 fatty acids on borderline personality disorder: a randomised, double-blind, clinical trial
Methods	RCT
Participants	BPD according to DSM-IV, CGI-S (BPD) > 3, 18-65 years
Interventions	12 weeks of cognitive behaviour therapy (CBT) + placebo vs. CBT + Omacor (R) 1680 mg/day vs. CBT + Omacor (R) 3360 mg/day
Outcomes	Depression, manic symptoms, impulsivity, aggression, anger, psychotic symptoms, anxiety, suicidal and para-suicidal behaviour, adverse events
Starting date	Unknown
Contact information	Casa M, Hospital Univesitari Vall d'Hebron, Barcelona, Spain
Notes	

Goodman NCT00255554

Trial name or title	Effects of Dialectical Behavioural Therapy and escitalopram on impulsive aggression, affective instability and cognitive processing in borderline personality disorder
Methods	RCT
Participants	BPD, 18-60 years, off psychotropic medication for at least 2 weeks
Interventions	Six months of DBT + escitalopram vs. DBT + placebo
Outcomes	Impulsivity, aggression, affective impulsivity, immediate and delayed memory, cognitive processing
Starting date	November 2005
Contact information	Goodman M, Bronx VA Medical Center, New York, USA
Notes	

Malev ISRCTN11135486

Trial name or title	Quetiapine versus sertraline as the pharmacological component in a standardised psychopharmacological and psychotherapeutic treatment of borderline personality disorder: a randomised, rater-blinded study
Methods	RCT
Participants	BPD according to DSM-IV, females, at least 18 years of age

Interventions	24 weeks of quetiapine (50-800 mg/day) vs. sertraline (25-200 mg/day)
Outcomes	Anger, hostility, severity of affective symptoms, anxiety, depression, psychotic symptoms, interpersonal problems, duration of hospitalisation, co-medication
Starting date	October 2006
Contact information	Malevani J, University of Dusseldorf, Germany
Notes	

Ralevski NCT00463775

Trial name or title	Topiramate for treatment of patients with borderline personality disorder and alcohol dependence
Methods	RCT
Participants	BPD and alcohol dependence, 21-60 years
Interventions	8 weeks of topiramate (250 mg/day) vs. placebo
Outcomes	Hostility, aggression, drinking, craving, side effects
Starting date	March 2007
Contact information	Ralevski E, Yale University, USA
Notes	

Schulz NCT00222482

Trial name or title	A double-blind and placebo controlled assessment of Depakote ER in borderline personality disorder
Methods	RCT
Participants	BPD, 21-55 years
Interventions	12 weeks of DBT + Depakote ER vs. DBT + placebo
Outcomes	SCL-90, impulsivity
Starting date	March 2003
Contact information	Schulz SC, University of Minnesota Medical School, USA
Notes	

DATA AND ANALYSES

Comparison 1 Active drug versus placebo: BPD severity

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	1	58	Std. Mean Difference (IV, Random, 95% CI)	0.30 [-0.22, 0.82]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.2 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.47 [-0.98, 0.05]
1.3 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	-0.15 [-0.65, 0.35]
2 SMD on basis of post-means and pre-SDs	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 First-generation antipsychotics: Thiothixene	1	50	Std. Mean Difference (IV, Random, 95% CI)	0.28 [-0.28, 0.83]
3 SMD on basis of change from baseline scores	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
3.1 Second-generation antipsychotics: Olanzapine	2	596	Std. Mean Difference (IV, Random, 95% CI)	-0.15 [-0.41, 0.10]
3.2 Mood stabilizers: Lamotrigine	1	27	Std. Mean Difference (IV, Random, 95% CI)	-0.43 [-1.20, 0.34]

Comparison 2

Active drug versus placebo: Avoidance of abandonment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.08 [-0.58, 0.43]
2 SMD on basis of change from baseline scores	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Second-generation antipsychotics: Olanzapine	3	631	Std. Mean Difference (IV, Random, 95% CI)	-0.01 [-0.22, 0.21]

Comparison 3

Active drug versus placebo: Interpersonal problems

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	7		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	2	114	Std. Mean Difference (IV, Random, 95% CI)	Not estimable
1.2 Second-generation antipsychotics: Aripiprazole	1	52	Std. Mean Difference (IV, Random, 95% CI)	-0.77 [-1.33, -0.20]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.3 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.12 [-0.63, 0.38]
1.4 Mood stabiliser: Carbamazepine	1	19	Std. Mean Difference (IV, Random, 95% CI)	-0.54 [-1.46, 0.38]
1.5 Mood stabiliser: Valproate semisodium	1	30	Std. Mean Difference (IV, Random, 95% CI)	-1.04 [-1.85, -0.23]
1.6 Mood stabiliser: Topiramate	1	56	Std. Mean Difference (IV, Random, 95% CI)	-0.91 [-1.46, -0.35]
1.7 Antidepressants: Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.39 [-0.92, 0.13]
1.8 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	0.24 [-0.26, 0.74]
2 SMD on basis of post-means and pre-SDs	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 First-generation antipsychotics: Thiothixene	1	50	Std. Mean Difference (IV, Random, 95% CI)	-0.18 [-0.74, 0.37]
3 SMD on basis of change from baseline scores	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
3.1 Second-generation antipsychotics: Olanzapine	2	340	Std. Mean Difference (IV, Random, 95% CI)	-0.10 [-0.31, 0.12]
4 Mean Change Difference	1		MCD (Random, 95% CI)	Subtotals only
4.1 Second-generation antipsychotics: Olanzapine	1	291	MCD (Random, 95% CI)	-0.2 [-0.62, 0.22]

Comparison 4 Active drug versus placebo: Identity disturbance

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.38 [-0.90, 0.13]
2 SMD on basis of change from baseline scores	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Second-generation antipsychotics: Olanzapine	3	631	Std. Mean Difference (IV, Random, 95% CI)	-0.06 [-0.21, 0.10]

Comparison 5
Active drug versus placebo: Impulsivity

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	13		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	2	114	Std. Mean Difference (IV, Random, 95% CI)	0.07 [-0.30, 0.43]
1.2 Second-generation antipsychotics: Aripiprazole	1	52	Std. Mean Difference (IV, Random, 95% CI)	-1.84 [-2.49, -1.18]
1.3 Second-generation antipsychotics: Olanzapine	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.04 [-0.54, 0.47]
1.4 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	0.03 [-0.48, 0.53]
1.5 Mood stabiliser: Valproate semisodium	2	46	Std. Mean Difference (IV, Random, 95% CI)	-0.62 [-1.48, 0.24]
1.6 Mood stabiliser: Lamotrigine	1	27	Std. Mean Difference (IV, Random, 95% CI)	-1.62 [-2.54, -0.69]
1.7 Mood stabiliser: Topiramate	2	71	Std. Mean Difference (IV, Random, 95% CI)	-3.36 [-4.44, -2.27]
1.8 Antidepressants: Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.12 [-0.64, 0.40]
1.9 Antidepressants: Fluoxetine	1	20	Std. Mean Difference (IV, Random, 95% CI)	-0.59 [-1.50, 0.31]
1.10 Antidepressants: Fluvoxamine	1	38	Std. Mean Difference (IV, Random, 95% CI)	-0.05 [-0.68, 0.59]
1.11 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	-0.00 [-0.50, 0.50]
1.12 Miscellaneous: Omega-3 fatty acids	1	27	Std. Mean Difference (IV, Random, 95% CI)	-0.47 [-1.28, 0.34]
2 SMD on basis of change from baseline scores	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Second-generation antipsychotics: Olanzapine	2	340	Std. Mean Difference (IV, Random, 95% CI)	-0.18 [-0.40, 0.03]
2.2 Mood stabilizers: Lamotrigine	1	27	Std. Mean Difference (IV, Random, 95% CI)	-1.41 [-2.27, -0.55]
3 Mean Change Difference	1		MCD (Random, 95% CI)	Subtotals only
3.1 Second-generation antipsychotics: Olanzapine	1	291	MCD (Random, 95% CI)	-0.10 [-0.40, 0.20]
4 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
4.1 Mood stabiliser: Carbamazepine	1	20	Risk Ratio (M-H, Random, 95% CI)	0.88 [0.53, 1.46]

Comparison 6

Active drug versus placebo: Suicidal ideation

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.27 [-0.78, 0.23]
1.2 Mood stabiliser: Valproate semisodium	1	16	Std. Mean Difference (IV, Random, 95% CI)	0.52 [-0.63, 1.67]
1.3 Antidepressants: Fluoxetine	1	20	Std. Mean Difference (IV, Random, 95% CI)	0.44 [-0.46, 1.33]
2 SMD on basis of change from baseline scores	2		Std. Mean Difference (IV, Fixed, 95% CI)	Subtotals only
2.1 Second-generation antipsychotics: Olanzapine	2	340	Std. Mean Difference (IV, Fixed, 95% CI)	0.29 [0.07, 0.50]
3 Mean Change Difference	1		MCD (Random, 95% CI)	Subtotals only
3.1 Second-generation antipsychotics: Olanzapine	1	291	MCD (Random, 95% CI)	-0.1 [-0.20, -0.00]
4 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
4.1 Second-generation antipsychotics versus placebo: Olanzapine	1	24	Risk Ratio (M-H, Random, 95% CI)	1.2 [0.50, 2.88]
4.2 Miscellaneous: Omega-3 fatty acids	1	49	Risk Ratio (M-H, Random, 95% CI)	0.52 [0.28, 0.95]

Comparison 7

Active drug versus placebo: Suicidal behaviour

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Flupenthixol decanoate	1	37	Risk Ratio (M-H, Random, 95% CI)	0.49 [0.26, 0.92]
1.2 Antidepressants: Mianserin	1	58	Risk Ratio (M-H, Random, 95% CI)	1.0 [0.71, 1.41]
2 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.1 Second-generation antipsychotics: Olanzapine	1	60	Std. Mean Difference (IV, Random, 95% CI)	0.15 [-0.36, 0.65]

Comparison 8 Active drug versus placebo: Self-mutilating behaviour

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Aripiprazole	1	52	Risk Ratio (M-H, Random, 95% CI)	0.29 [0.07, 1.25]
1.2 Second-generation antipsychotics: Olanzapine	1	24	Risk Ratio (M-H, Random, 95% CI)	1.2 [0.50, 2.88]
1.3 Miscellaneous: Omega-3 fatty acids	1	49	Risk Ratio (M-H, Random, 95% CI)	1.23 [0.51, 2.97]
2 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Antidepressants: Fluoxetine	1	20	Std. Mean Difference (IV, Random, 95% CI)	0.03 [-0.85, 0.92]

Comparison 9 Active drug versus placebo: Affective instability

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.10 [-0.61, 0.41]
1.2 Antidepressants: Fluvoxamine	1	38	Std. Mean Difference (IV, Random, 95% CI)	-0.64 [-1.30, 0.01]
2 SMD on basis of change from baseline scores	4		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Second-generation antipsychotics: Olanzapine	3	631	Std. Mean Difference (IV, Random, 95% CI)	-0.16 [-0.32, -0.01]
2.2 Mood stabilizers: Lamotrigine	1	27	Std. Mean Difference (IV, Random, 95% CI)	-0.61 [-1.39, 0.17]

Comparison 10
Active drug versus placebo: Chronic feelings of
emptiness

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	0.18 [-0.32, 0.69]
2 SMD on basis of change from baseline scores	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Second-generation antipsychotics: Olanzapine	3	631	Std. Mean Difference (IV, Random, 95% CI)	-0.03 [-0.22, 0.16]

Comparison 11
Active drug versus placebo: Anger

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	13		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	2	114	Std. Mean Difference (IV, Random, 95% CI)	-0.46 [-0.84, -0.09]
1.2 Second-generation antipsychotics: Aripiprazole	1	52	Std. Mean Difference (IV, Random, 95% CI)	-1.14 [-1.73, -0.55]
1.3 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	0.08 [-0.43, 0.58]
1.4 Mood stabiliser: Carbamazepine	1	19	Std. Mean Difference (IV, Random, 95% CI)	-0.34 [-1.25, 0.57]
1.5 Mood stabiliser: Valproate semisodium	1	30	Std. Mean Difference (IV, Random, 95% CI)	-0.15 [-0.91, 0.61]
1.6 Mood stabiliser: Valproate semisodium	1	16	Std. Mean Difference (IV, Random, 95% CI)	-1.83 [-3.17, -0.48]
1.7 Mood stabiliser: Lamotrigine	1	27	Std. Mean Difference (IV, Random, 95% CI)	-1.69 [-2.62, -0.75]
1.8 Mood stabiliser: Topiramate (females)		85	Std. Mean Difference (IV, Random, 95% CI)	-1.00 [-3.64, -2.36]
1.9 Mood stabiliser: Topiramate (males)	1	42	Std. Mean Difference (IV, Random, 95% CI)	-0.65 [-1.27, -0.03]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.10 Antidepressants: Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.26 [-0.78, 0.26]
1.11 Antidepressants: Fluoxetine	1	22	Std. Mean Difference (IV, Random, 95% CI)	-0.65 [-1.53, 0.22]
1.12 Antidepressants: Fluvoxamine	1	38	Std. Mean Difference (IV, Random, 95% CI)	-0.37 [-1.01, 0.28]
1.13 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	-0.34 [-0.84, 0.17]
2 SMD on basis of post-means and pre-SDs	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 First-generation antipsychotics: Thiothixene	1	50	Std. Mean Difference (IV, Random, 95% CI)	-0.07 [-0.63, 0.48]
3 SMD on basis of change from baseline scores	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
3.1 Second-generation antipsychotics: Olanzapine	3	631	Std. Mean Difference (IV, Random, 95% CI)	-0.27 [-0.43, -0.12]

Comparison 12
Active drug versus placebo: Psychotic symptoms

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	6		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	2	114	Std. Mean Difference (IV, Random, 95% CI)	-0.44 [-1.09, 0.20]
1.2 Second-generation antipsychotics: Aripiprazole	1	52	Std. Mean Difference (IV, Random, 95% CI)	-1.05 [-1.64, -0.47]
1.3 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.23 [-0.74, 0.28]
1.4 Mood stabiliser: Carbamazepine	1	19	Std. Mean Difference (IV, Random, 95% CI)	-0.58 [-1.50, 0.35]
1.5 Mood stabiliser: Topiramate	1	56	Std. Mean Difference (IV, Random, 95% CI)	-0.49 [-1.02, 0.05]
1.6 Antidepressants: Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.43 [-0.96, 0.09]
1.7 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	-0.28 [-0.78, 0.22]
2 SMD on basis of post-means and pre-SDs	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.1 First-generation antipsychotics: Thiothixene	1	50	Std. Mean Difference (IV, Random, 95% CI)	0.19 [-0.37, 0.75]
3 SMD on basis of change from baseline scores	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
3.1 Second-generation antipsychotics: Olanzapine	3	631	Std. Mean Difference (IV, Random, 95% CI)	-0.18 [-0.34, -0.03]

Comparison 13 Active drug versus placebo: Dissociation

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Antidepressants: Fluoxetine	1	20	Std. Mean Difference (IV, Random, 95% CI)	0.42 [-0.47, 1.32]

Comparison 14 Active drug versus placebo: Depression

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	13		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	2	114	Std. Mean Difference (IV, Random, 95% CI)	-0.09 [-0.87, 0.68]
1.2 Second-generation antipsychotics: Aripiprazole	1	52	Std. Mean Difference (IV, Random, 95% CI)	-1.25 [-1.85, -0.65]
1.3 Second-generation antipsychotics: Olanzapine	2	84	Std. Mean Difference (IV, Random, 95% CI)	-0.37 [-0.80, 0.07]
1.4 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.30 [-0.81, 0.21]
1.5 Mood stabiliser: Carbamazepine	1	19	Std. Mean Difference (IV, Random, 95% CI)	-0.66 [-1.59, 0.27]
1.6 Mood stabiliser: Valproate semisodium	2	46	Std. Mean Difference (IV, Random, 95% CI)	-0.66 [-1.31, -0.01]
1.7 Mood stabiliser: Topiramate	1	56	Std. Mean Difference (IV, Random, 95% CI)	-0.51 [-1.04, 0.02]
1.8 Antidepressants: Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.59 [-1.12, -0.06]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.9 Antidepressant: Fluoxetine	2	42	Std. Mean Difference (IV, Random, 95% CI)	0.12 [-1.13, 1.36]
1.10 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	-0.34 [-0.84, 0.16]
1.11 Miscellaneous: Omega-3 fatty acids	1	27	Std. Mean Difference (IV, Random, 95% CI)	-0.34 [-1.15, 0.46]
2 SMD on basis of post-means and pre-SDs	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 First-generation antipsychotics: Thiothixene	1	50	Std. Mean Difference (IV, Random, 95% CI)	0.12 [-0.43, 0.68]
3 Mean Change Difference	2		MCD (Random, 95% CI)	Subtotals only
3.1 Second-generation antipsychotics: Olanzapine	2	596	MCD (Random, 95% CI)	0.39 [-0.20, 0.97]
4 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
4.1 Miscellaneous: Omega-3 fatty acid	1	49	Risk Ratio (M-H, Random, 95% CI)	0.48 [0.28, 0.81]

Comparison 15 Active drug versus placebo: Anxiety

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	8		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	2	114	Std. Mean Difference (IV, Random, 95% CI)	0.06 [-0.68, 0.79]
1.2 Second-generation antipsychotics: Aripiprazole	1	52	Std. Mean Difference (IV, Random, 95% CI)	-0.73 [-1.29, -0.17]
1.3 Second-generation antipsychotics: Olanzapine	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.23 [-0.74, 0.28]
1.4 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.12 [-0.63, 0.39]
1.5 Mood stabiliser: Carbamazepine	1	19	Std. Mean Difference (IV, Random, 95% CI)	-0.51 [-1.43, 0.41]
1.6 Mood stabiliser: Topiramate	1	56	Std. Mean Difference (IV, Random, 95% CI)	-1.40 [-1.99, -0.81]
1.7 Antidepressants: Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.15 [-0.67, 0.37]
1.8 Antidepressants: Fluoxetine	1	20	Std. Mean Difference (IV, Random, 95% CI)	0.15 [-0.73, 1.03]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.9 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	-0.14 [-0.65, 0.36]
2 Mean Change Difference	1		MCD (Random, 95% CI)	Subtotals only
2.1 Second-generation antipsychotics: Olanzapine	1	274	MCD (Random, 95% CI)	-0.22 [-0.41, -0.03]

Comparison 16
Active drug versus placebo: General psychiatric pathology

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	6		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	2	114	Std. Mean Difference (IV, Random, 95% CI)	-0.08 [-0.71, 0.54]
1.2 Second-generation antipsychotics: Aripiprazole	1	52	Std. Mean Difference (IV, Random, 95% CI)	-1.27 [-1.87, -0.67]
1.3 Second-generation antipsychotics: Ziprasidone	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.41 [-0.92, 0.10]
1.4 Mood stabiliser: Carbamazepine	1	19	Std. Mean Difference (IV, Random, 95% CI)	-0.57 [-1.49, 0.36]
1.5 Mood stabiliser: Topiramate	1	56	Std. Mean Difference (IV, Random, 95% CI)	-1.19 [-1.76, -0.61]
1.6 Antidepressants: Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.34 [-0.87, 0.18]
1.7 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	-0.23 [-0.73, 0.27]
2 SMD on basis of change from baseline scores	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Second-generation antipsychotics: Olanzapine	2	557	Std. Mean Difference (IV, Random, 95% CI)	-0.21 [-0.53, 0.10]

Comparison 17
Active drug versus placebo: Mental health status

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	6		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 First-generation antipsychotics: Haloperidol	2	114	Std. Mean Difference (IV, Random, 95% CI)	0.16 [-0.77, 1.08]
1.2 Second-generation antipsychotics: Olanzapine	1	60	Std. Mean Difference (IV, Random, 95% CI)	-0.03 [-0.53, 0.48]
1.3 Mood stabiliser: Carbamazepine	1	19	Std. Mean Difference (IV, Random, 95% CI)	0.34 [-0.57, 1.25]
1.4 Antidepressants: Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	0.27 [-0.25, 0.79]
1.5 Antidepressants: Fluoxetine	2	42	Std. Mean Difference (IV, Random, 95% CI)	0.40 [-0.27, 1.07]
1.6 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	0.14 [-0.36, 0.64]
2 SMD on basis of post-means and pre-SDs	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 First-generation antipsychotics: Thiothixene	1	50	Std. Mean Difference (IV, Random, 95% CI)	0.06 [-0.50, 0.61]
3 Mean Change Difference	2		MCD (Random, 95% CI)	Subtotals only
3.1 Second-generation antipsychotics: Olanzapine	2	596	MCD (Random, 95% CI)	1.52 [-0.75, 3.79]
4 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
4.1 Mood stabiliser: Valproate semisodium	1	16	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.37, 1.11]

Comparison 18
Active drug versus placebo: Attrition/leaving the study early for any reason

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	24		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Flupenthixol decanoate	1	37	Risk Ratio (M-H, Random, 95% CI)	1.41 [0.36, 5.43]
1.2 First-generation antipsychotics versus placebo: Haloperidol	2	130	Risk Ratio (M-H, Random, 95% CI)	1.15 [0.45, 2.92]
1.3 First-generation antipsychotics versus placebo: Thiothixene	1	50	Risk Ratio (M-H, Random, 95% CI)	2.53 [0.74, 8.68]
1.4 Second-generation antipsychotics: Olanzapine	6	767	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.72, 1.29]
1.5 Second-generation antipsychotics: Ziprasidone	1	60	Risk Ratio (M-H, Random, 95% CI)	1.21 [0.74, 1.99]
1.6 Mood stabiliser: Carbamazepine	1	20	Risk Ratio (M-H, Random, 95% CI)	5.0 [0.27, 92.62]
1.7 Mood stabiliser: Valproate semisodium	2	46	Risk Ratio (M-H, Random, 95% CI)	0.78 [0.40, 1.53]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.8 Mood stabiliser: Lamotrigine	2	55	Risk Ratio (M-H, Random, 95% CI)	0.74 [0.22, 2.48]
1.9 Mood stabiliser: Topiramate	3	133	Risk Ratio (M-H, Random, 95% CI)	0.55 [0.14, 2.16]
1.10 Antidepressants: Amitriptyline	1	59	Risk Ratio (M-H, Random, 95% CI)	0.97 [0.06, 14.74]
1.11 Antidepressants: Fluoxetine	1	25	Risk Ratio (M-H, Random, 95% CI)	1.63 [0.33, 8.11]
1.12 Antidepressants: Fluvoxamine	1	38	Risk Ratio (M-H, Random, 95% CI)	0.45 [0.04, 4.55]
1.13 Antidepressants: Mianserin	1	58	Risk Ratio (M-H, Random, 95% CI)	1.5 [0.72, 3.12]
1.14 Antidepressants: Phenelzine sulfate	1	72	Risk Ratio (M-H, Random, 95% CI)	0.60 [0.18, 1.94]
1.15 Miscellaneous: Omega-3 fatty acids	2	79	Risk Ratio (M-H, Random, 95% CI)	0.61 [0.21, 1.79]

Comparison 19 Active drug versus placebo: AE - body weight change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	12		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 First-generation antipsychotics: Haloperidol	1	58	Std. Mean Difference (IV, Random, 95% CI)	-0.18 [-0.70, 0.34]
1.2 Second-generation antipsychotics: Olanzapine	6	752	Std. Mean Difference (IV, Random, 95% CI)	1.05 [0.90, 1.20]
1.3 Mood stabiliser: Valproate semisodium	1	30	Std. Mean Difference (IV, Random, 95% CI)	0.68 [-0.10, 1.47]
1.4 Mood stabiliser: Lamotrigine	1	27	Std. Mean Difference (IV, Random, 95% CI)	-0.13 [-0.93, 0.67]
1.5 Mood stabilizer: Topiramate	3	127	Std. Mean Difference (IV, Random, 95% CI)	-0.55 [-0.91, -0.19]
1.6 Antidepressants: Phenelzine sulfate	1	62	Std. Mean Difference (IV, Random, 95% CI)	0.11 [-0.39, 0.61]

Comparison 20 Active drug versus placebo: AE - any AE

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics versus placebo: Olanzapine	2	615	Risk Ratio (M-H, Random, 95% CI)	1.13 [1.00, 1.28]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.2 Second-generation antipsychotics versus placebo: Ziprasidone	1	60	Risk Ratio (M-H, Random, 95% CI)	2.75 [0.99, 7.68]

Comparison 21
Active drug versus placebo: AE - increased appetite

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	2	615	Risk Ratio (M-H, Random, 95% CI)	2.76 [1.75, 4.34]

Comparison 22
Active drug versus placebo: AE - paraesthesia

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Mood stabiliser: Topiramate	1	56	Risk Ratio (M-H, Random, 95% CI)	3.00 [0.33, 27.12]

Comparison 23
Active drug versus placebo: AE - headache

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	2	615	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.43, 1.92]
1.2 Mood stabiliser: Topiramate	1	56	Risk Ratio (M-H, Random, 95% CI)	1.0 [0.15, 6.61]

Comparison 24
Active drug versus placebo: AE - dizziness

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 Second-generation antipsychotics versus placebo: Ziprasidone	1	60	Risk Ratio (M-H, Random, 95% CI)	9.00 [0.51, 160.17]
1.2 Mood stabiliser: Topiramate	1	56	Risk Ratio (M-H, Random, 95% CI)	1.5 [0.27, 8.30]

Comparison 25
Active drug versus placebo: AE - disturbance in attention

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	1	301	Risk Ratio (M-H, Random, 95% CI)	1.37 [0.63, 20 3.81]
1.2 Mood stabiliser: Topiramate	1	56	Risk Ratio (M-H, Random, 95% CI)	2.0 [0.55, 7.22]

Comparison 26
Active drug versus placebo: AE - memory problems

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Mood stabiliser: Topiramate	1	56	Risk Ratio (M-H, Random, 95% CI)	2.0 [0.55, 7.22]

Comparison 27
Active drug versus placebo: AE - fatigue

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	2	615	Risk Ratio (M-H, Random, 95% CI)	2.04 [0.79, 5.23]
1.2 Mood stabiliser: Topiramate	1	56	Risk Ratio (M-H, Random, 95% CI)	2.00 [0.40, 10.05]

Comparison 28
Active drug versus placebo: AE - somnolence

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	2	615	Risk Ratio (M-H, Random, 95% CI)	2.97 [1.75, 5.03]

Comparison 29
Active drug versus placebo: AE - sedation

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	3		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine (1)	1	314	Risk Ratio (M-H, Random, 95% CI)	9.23 [2.18, 39.12]
1.2 Second-generation antipsychotics: Olanzapine (2)	1	28	Risk Ratio (M-H, Random, 95% CI)	1.26 [0.44, 3.66]
1.3 Second-generation antipsychotics versus placebo: Ziprasidone	1	60	Risk Ratio (M-H, Random, 95% CI)	6.0 [0.77, 46.87]

Comparison 30
Active drug versus placebo: AE - insomnia

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	2	615	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.33, 1.37]

Comparison 31
Active drug versus placebo: AE - anxiety

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	1	314	Risk Ratio (M-H, Random, 95% CI)	0.90 [0.33, 2.42]

Comparison 32
Active drug versus placebo: AE - nausea

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	2	615	Risk Ratio (M-H, Random, 95% CI)	0.83 [0.43, 1.59]

Comparison 33
Active drug versus placebo: AE - uneasy feeling

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics versus placebo: Ziprasidone	1	60	Risk Ratio (M-H, Random, 95% CI)	7.0 [0.38, 129.93]

Comparison 34
Active drug versus placebo: AE - constipation

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics versus placebo: Olanzapine	1	28	Risk Ratio (M-H, Random, 95% CI)	6.5 [0.41, 104.20]

Comparison 35
Active drug versus placebo: AE - dry mouth

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	2	615	Risk Ratio (M-H, Random, 95% CI)	2.24 [1.08, 4.67]

Comparison 36
Active drug versus placebo: AE - nasopharyngitis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	1	301	Risk Ratio (M-H, Random, 95% CI)	0.62 [0.23, 1.66]

Comparison 37
Active drug versus placebo: AE - menstrual pain

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Risk Ratio	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Mood stabiliser: Topiramate	1	56	Risk Ratio (M-H, Random, 95% CI)	1.67 [0.44, 6.31]

Comparison 38
Active drug versus placebo: AE - liver function: AST/SGOT baseline to endpoint mean change (U/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	2	526	Std. Mean Difference (IV, Random, 95% CI)	0.35 [0.18, 0.52]

Comparison 39
Active drug versus placebo: AE - liver function: ALT/SGPT baseline to endpoint mean change (U/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotics: Olanzapine	2	530	Std. Mean Difference (IV, Random, 95% CI)	0.46 [0.29, 0.63]

Comparison 40
Active drug versus placebo: AE - liver function: GGT
(GGPT/SGGT/YGGT) baseline to endpoint mean
change (U/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	268	Std. Mean Difference (IV, Random, 95% CI)	0.26 [0.02, 0.50]

Comparison 41
Active drug versus placebo: AE - liver function: total
bilirubin baseline to endpoint mean change (μ mol/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	264	Std. Mean Difference (IV, Random, 95% CI)	-0.29 [-0.53, -0.05]

Comparison 42
Active drug versus placebo: AE - liver function: direct
bilirubin baseline to endpoint mean change (μ mol/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	258	Std. Mean Difference (IV, Random, 95% CI)	-0.35 [-0.60, -0.11]

Comparison 43
Active drug versus placebo: AE - lipids: total
cholesterol baseline to endpoint change (mmol/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation	2	327	Std. Mean Difference (IV, Random, 95% CI)	0.42 [0.20, 0.64]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
antipsychotics: Olanzapine				

Comparison 44
Active drug versus placebo: AE - lipids: LDL
cholesterol baseline to endpoint mean change (mmol/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	259	Std. Mean Difference (IV, Random, 95% CI)	0.35 [0.10, 0.59]

Comparison 45
Active drug versus placebo: AE - lipids: HDL
cholesterol (dextran precip.) baseline to endpoint mean
change (mmol/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	269	Std. Mean Difference (IV, Random, 95% CI)	-0.28 [-0.52, -0.04]

Comparison 46
Active drug versus placebo: AE - lipids: triglycerides,
fasting, baseline to endpoint mean change (mmol/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	203	Std. Mean Difference (IV, Random, 95% CI)	0.37 [0.09, 0.64]

Comparison 47
Active drug versus placebo: AE - prolactin: baseline to endpoint mean change (µg/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	2	528	Std. Mean Difference (IV, Random, 95% CI)	0.41 [0.23, 0.59]

Comparison 48
Active drug versus placebo: AE - platelet count baseline to endpoint mean change (GI/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	2		Std. Mean Difference (IV, Random, 95% CI)	Totals not selected
1.1 Second-generation antipsychotic: Olanzapine	2		Std. Mean Difference (IV, Random, 95% CI)	Not estimable

Comparison 49
Active drug versus placebo: AE - erythrocyte count baseline to endpoint mean change (TI/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	262	Std. Mean Difference (IV, Random, 95% CI)	-0.18 [-0.42, 0.06]

Comparison 50
Active drug versus placebo: AE - leukocyte count baseline to endpoint mean change (GI/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation	1	262	Std. Mean Difference (IV, Random, 95% CI)	-0.40 [-0.65, -0.16]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
antipsychotic: Olanzapine				

Comparison 51
Active drug versus placebo: AE - neutrophils,
segmented, baseline to endpoint mean change (GI/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	262	Std. Mean Difference (IV, Random, 95% CI)	-0.39 [-0.63, -0.14]

Comparison 52
Active drug versus placebo: AE - basophils baseline to
endpoint mean change (GI/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	262	Std. Mean Difference (IV, Random, 95% CI)	-0.28 [-0.53, -0.04]

Comparison 53
Active drug versus placebo: AE - monocytes baseline to
endpoint mean change (GI/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	262	Std. Mean Difference (IV, Random, 95% CI)	-0.28 [-0.53, -0.04]

Comparison 54
Active drug versus placebo: AE - haemoglobin baseline to endpoint mean change (mmol/L-F)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	262	Std. Mean Difference (IV, Random, 95% CI)	-0.21 [-0.45, 0.03]

Comparison 55
Active drug versus placebo: AE - mean cell haemoglobin concentration (MCHC) baseline to endpoint mean change (mmol/L-F)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	260	Std. Mean Difference (IV, Random, 95% CI)	0.03 [-0.22, 0.27]

Comparison 56
Active drug versus placebo: AE - calcium baseline to endpoint mean change (mmol/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	268	Std. Mean Difference (IV, Random, 95% CI)	-0.33 [-0.57, -0.09]

Comparison 57
Active drug versus placebo: AE - albumin baseline to endpoint mean change (g/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation	1	269	Std. Mean Difference (IV, Random, 95% CI)	-0.21 [-0.45, 0.03]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
antipsychotic: Olanzapine				

Comparison 58
Active drug versus placebo: AE - creatine
phosphokinase baseline to endpoint mean change (U/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	268	Std. Mean Difference (IV, Random, 95% CI)	-0.21 [-0.45, 0.03]

Comparison 59
Active drug versus placebo: AE - urea nitrogen baseline
to endpoint mean change (mmol/L)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	269	Std. Mean Difference (IV, Random, 95% CI)	-0.14 [-0.38, 0.10]

Comparison 60
Active drug versus placebo: AE - pulse, standing,
baseline to endpoint mean change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	290	Std. Mean Difference (IV, Random, 95% CI)	0.08 [-0.15, 0.31]

Comparison 61
Active drug versus placebo: AE - pulse, supine, baseline to endpoint mean change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	290	Std. Mean Difference (IV, Random, 95% CI)	0.02 [-0.21, 0.25]

Comparison 62
Active drug versus placebo: AE - diastolic blood pressure, standing, baseline to endpoint mean change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	290	Std. Mean Difference (IV, Random, 95% CI)	-0.03 [-0.26, 0.20]

Comparison 63
Active drug versus placebo: AE - diastolic blood pressure, supine, baseline to endpoint mean change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	290	Std. Mean Difference (IV, Random, 95% CI)	-0.01 [-0.24, 0.22]

Comparison 64
Active drug versus placebo: AE - systolic blood pressure, standing, baseline to endpoint mean change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation	1	290	Std. Mean Difference (IV, Random, 95% CI)	0.03 [-0.20, 0.26]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
antipsychotic: Olanzapine				

Comparison 65
Active drug versus placebo: AE - systolic blood pressure, supine, baseline to endpoint mean change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 SMD	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Second-generation antipsychotic: Olanzapine	1	290	Std. Mean Difference (IV, Random, 95% CI)	-0.04 [-0.27, 0.19]

Comparison 66
Drug versus drug: BPD severity

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	0.46 [-0.03, 0.96]

Comparison 67
Drug versus drug: Interpersonal problems

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol vs. Amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.14 [-0.66, 0.38]
1.2 Haloperidol vs. Phenelzine Sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	-0.46 [-0.96, 0.04]

Comparison 68
Drug versus drug: Impulsivity

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 Haloperidol versus amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	0.20 [-0.32, 0.72]
1.2 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	0.09 [-0.40, 0.58]
2 Second-generation antipsychotic versus antidepressant	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Olanzapine versus fluoxetine	1	29	Std. Mean Difference (IV, Random, 95% CI)	-0.20 [-0.93, 0.53]

Comparison 69 Drug versus drug: Anger

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol versus amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.36 [-0.89, 0.16]
1.2 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	0.08 [-0.41, 0.57]

Comparison 70 Drug versus drug: Psychotic symptoms

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol versus amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.35 [-0.87, 0.18]
1.2 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	0.15 [-0.34, 0.64]

Comparison 71 Drug versus drug: Depression

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol versus amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	0.08 [-0.44, 0.59]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.2 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	0.68 [0.17, 1.19]
2 Second-generation antipsychotic versus antidepressant	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Olanzapine versus fluoxetine	1	29	Std. Mean Difference (IV, Random, 95% CI)	-0.73 [-1.49, 0.03]

Comparison 72 Drug versus drug: Anxiety

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol versus amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.18 [-0.70, 0.34]
1.2 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	0.66 [0.15, 1.16]

Comparison 73 Drug versus drug: General psychiatric pathology

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol versus amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	-0.07 [-0.59, 0.45]
1.2 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	0.53 [0.03, 1.03]

Comparison 74 Drug versus drug: Mental health status

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol versus amitriptyline	1	57	Std. Mean Difference (IV, Random, 95% CI)	0.29 [-0.23, 0.81]
1.2 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	-0.51 [-1.01, -0.01]

Comparison 75
Drug versus drug: AE - attrition

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus first-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Loxapine versus chlorpromazine	1	80	Risk Ratio (M-H, Random, 95% CI)	1.14 [0.46, 2.85]
2 First-generation antipsychotic versus antidepressant	2		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 Haloperidol versus amitriptyline	1	61	Risk Ratio (M-H, Random, 95% CI)	2.90 [0.32, 26.38]
2.2 Haloperidol versus phenelzine sulfate	1	74	Risk Ratio (M-H, Random, 95% CI)	1.58 [0.49, 5.15]
3 Second-generation antipsychotic versus antidepressant	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
3.1 Olanzapine versus fluoxetine	1	30	Risk Ratio (M-H, Random, 95% CI)	0.29 [0.01, 6.69]

Comparison 76
Drug versus drug: AE - body weight change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus antidepressant	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Haloperidol versus phenelzine sulfate	1	64	Std. Mean Difference (IV, Random, 95% CI)	-0.29 [-0.78, 0.21]
2 Second-generation antipsychotic versus antidepressant	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Olanzapine versus fluoxetine	1	29	Std. Mean Difference (IV, Random, 95% CI)	0.98 [0.20, 1.76]

Comparison 77
Drug versus drug: AE - any AE

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus first-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Loxapine versus chlorpromazine	1	80	Risk Ratio (M-H, Random, 95% CI)	1.27 [0.66, 2.45]

Comparison 78
Drug versus drug: AE - sedation

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Second-generation antipsychotic versus antidepressant	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Olanzapine versus fluoxetine	1	30	Risk Ratio (M-H, Random, 95% CI)	3.5 [1.23, 9.92]

Comparison 79
Drug versus drug: AE - sleepiness/drowsiness

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus first-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Loxapine versus chlorpromazine	1	80	Risk Ratio (M-H, Random, 95% CI)	0.8 [0.23, 2.76]

Comparison 80
Drug versus drug: restlessness

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus first-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Loxapine versus chlorpromazine	1	80	Risk Ratio (M-H, Random, 95% CI)	1.5 [0.26, 8.50]
2 Second-generation antipsychotic versus antidepressant	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 Olanzapine versus fluoxetine	1	30	Risk Ratio (M-H, Random, 95% CI)	0.7 [0.23, 2.11]

Comparison 81
Drug versus drug: AE - muscle spasms

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus first-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Loxapine versus chlorpromazine	1	80	Risk Ratio (M-H, Random, 95% CI)	3.00 [0.33, 27.63]

Comparison 82
Drug versus drug: AE - fainting spells

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 First-generation antipsychotic versus first-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Loxapine versus chlorpromazine	1	80	Risk Ratio (M-H, Random, 95% CI)	0.14 [0.01, 2.68]

Comparison 83
Drug versus combination of drugs: Impulsivity

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Olanzapine versus olanzapine + fluoxetine	1	29	Std. Mean Difference (IV, Random, 95% CI)	0.02 [-0.71, 0.76]
2 Antidepressant versus antidepressant + second-generation antipsychotic	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Fluoxetine versus fluoxetine + olanzapine	1	26	Std. Mean Difference (IV, Random, 95% CI)	0.25 [-0.53, 1.02]

Comparison 84
Drug versus combination of drugs: Depression

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Olanzapine versus olanzapine + fluoxetine	1	29	Std. Mean Difference (IV, Random, 95% CI)	-0.26 [1.00, 0.47]
2 Antidepressant versus antidepressant + second-generation antipsychotic	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Fluoxetine versus fluoxetine + olanzapine	1	26	Std. Mean Difference (IV, Random, 95% CI)	0.54 [-0.24, 1.33]

Comparison 85
Drug versus combination of drugs: AE - attrition

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Olanzapine versus olanzapine + fluoxetine	1	31	Risk Ratio (M-H, Random, 95% CI)	0.19 [0.01, 3.63]
2 Antidepressant versus antidepressant + second-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 Fluoxetine versus fluoxetine + olanzapine	1	29	Risk Ratio (M-H, Random, 95% CI)	0.54 [0.05, 5.28]

Comparison 86
Drug versus combination of drugs: AE - body weight change

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
1.1 Olanzapine versus olanzapine + fluoxetine	1	29	Std. Mean Difference (IV, Random, 95% CI)	0.70 [-0.05, 1.46]
2 Antidepressant versus antidepressant + second-generation antipsychotic	1		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Fluoxetine versus fluoxetine + olanzapine	1	26	Std. Mean Difference (IV, Random, 95% CI)	-0.54 [-1.32, 0.25]

Comparison 87
Drug versus combination of drugs: AE - sedation

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Olanzapine versus olanzapine + fluoxetine	1	31	Risk Ratio (M-H, Random, 95% CI)	1.61 [0.87, 2.96]
2 Antidepressant versus antidepressant + second-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 Fluoxetine versus fluoxetine + olanzapine	1	29	Risk Ratio (M-H, Random, 95% CI)	0.46 [0.15, 1.44]

Comparison 88

Drug versus combination of drugs: AE - akathisia

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
1.1 Olanzapine versus olanzapine + fluoxetine	1	31	Risk Ratio (M-H, Random, 95% CI)	0.75 [0.25, 2.28]
2 Antidepressant versus antidepressant + second-generation antipsychotic	1		Risk Ratio (M-H, Random, 95% CI)	Subtotals only
2.1 Fluoxetine versus fluoxetine + olanzapine	1	29	Risk Ratio (M-H, Random, 95% CI)	1.07 [0.39, 2.92]


Analysis 1.1

Comparison 1 Active drug versus placebo: BPD severity, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 1 Active drug versus placebo: BPD severity

Outcome: 1 SMD




Analysis 1.2

Comparison 1 Active drug versus placebo: BPD severity, Outcome 2 SMD on basis of postmeans and pre-SDs

Review: Pharmacological interventions for borderline personality disorder

Comparison: 1 Active drug versus placebo: BPD severity

Outcome: 2 SMD on basis of post-means and pre-SDs




Analysis 1.3

Comparison 1 Active drug versus placebo: BPD severity, Outcome 3 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder

Comparison: 1 Active drug versus placebo: BPD severity

Outcome: 3 SMD on basis of change from baseline scores




Analysis 2.1

Comparison 2 Active drug versus placebo: Avoidance of abandonment, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 2 Active drug versus placebo: Avoidance of abandonment

Outcome: 1 SMD




Analysis 2.2

Comparison 2 Active drug versus placebo: Avoidance of abandonment, Outcome 2 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder

Comparison: 2 Active drug versus placebo: Avoidance of abandonment

Outcome: 2 SMD on basis of change from baseline scores




Analysis 3.1

Comparison 3 Active drug versus placebo:

Interpersonal problems, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 3 Active drug versus placebo: Interpersonal problems
 Outcome: 1 SMD




Analysis 3.2

Comparison 3 Active drug versus placebo: Interpersonal problems, Outcome 2 SMD on basis of post-means and pre-SDs

Review: Pharmacological interventions for borderline personality disorder

Comparison: 3 Active drug versus placebo: Interpersonal problems

Outcome: 2 SMD on basis of post-means and pre-SDs




Analysis 3.3

Comparison 3 Active drug versus placebo: Interpersonal problems, Outcome 3 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder

Comparison: 3 Active drug versus placebo: Interpersonal problems


Outcome: 3 SMD on basis of change from baseline scores



Analysis 3.4

Comparison 3 Active drug versus placebo: Interpersonal problems, Outcome 4 Mean Change Difference


Review: Pharmacological interventions for borderline personality disorder
Comparison: 3 Active drug versus placebo: Interpersonal problems
Outcome: 4 Mean Change Difference



Analysis 4.1

Comparison 4 Active drug versus placebo: Identity disturbance, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder
Comparison: 4 Active drug versus placebo: Identity disturbance
Outcome: 1 SMD




Analysis 4.2

Comparison 4 Active drug versus placebo: Identity disturbance, Outcome 2 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder

Comparison: 4 Active drug versus placebo: Identity disturbance

Outcome: 2 SMD on basis of change from baseline scores




Analysis 5.1


Comparison 5 Active drug versus placebo: Impulsivity, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 5 Active drug versus placebo: Impulsivity

Outcome: 1 SMD






Analysis 5.2

Comparison 5 Active drug versus placebo: Impulsivity, Outcome 2 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder
Comparison: 5 Active drug versus placebo: Impulsivity
Outcome: 2 SMD on basis of change from baseline scores



Analysis 5.3


Comparison 5 Active drug versus placebo: Impulsivity,

Outcome 3 Mean Change Difference

Review: Pharmacological interventions for borderline personality disorder

Comparison: 5 Active drug versus placebo: Impulsivity

Outcome: 3 Mean Change Difference



Analysis 5.4


Comparison 5 Active drug versus placebo: Impulsivity,

Outcome 4 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 5 Active drug versus placebo: Impulsivity

Outcome: 4 Risk Ratio




Analysis 6.1

Comparison 6 Active drug versus placebo: Suicidal ideation, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 6 Active drug versus placebo: Suicidal ideation

Outcome: 1 SMD




Analysis 6.2

Comparison 6 Active drug versus placebo: Suicidal ideation, Outcome 2 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder

Comparison: 6 Active drug versus placebo: Suicidal ideation

Outcome: 2 SMD on basis of change from baseline scores




Analysis 6.3

Comparison 6 Active drug versus placebo: Suicidal ideation, Outcome 3 Mean Change Difference

Review: Pharmacological interventions for borderline personality disorder

Comparison: 6 Active drug versus placebo: Suicidal ideation

Outcome: 3 Mean Change Difference




Analysis 6.4

Comparison 6 Active drug versus placebo: Suicidal ideation, Outcome 4 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 6 Active drug versus placebo: Suicidal ideation

Outcome: 4 Risk Ratio




Analysis 7.1

Comparison 7 Active drug versus placebo: Suicidal behaviour, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 7 Active drug versus placebo: Suicidal behaviour

Outcome: 1 Risk Ratio




Analysis 7.2

Comparison 7 Active drug versus placebo: Suicidal behaviour, Outcome 2 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 7 Active drug versus placebo: Suicidal behaviour


Outcome: 2 SMD



Analysis 8.1

Comparison 8 Active drug versus placebo: Self-mutilating behaviour, Outcome 1 Risk Ratio


Review: Pharmacological interventions for borderline personality disorder
 Comparison: 8 Active drug versus placebo: Self-mutilating behaviour
 Outcome: 1 Risk Ratio



Analysis 8.2

Comparison 8 Active drug versus placebo: Self-mutilating behaviour, Outcome 2 SMD

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 8 Active drug versus placebo: Self-mutilating behaviour
 Outcome: 2 SMD




Analysis 9.1

Comparison 9 Active drug versus placebo: Affective instability, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 9 Active drug versus placebo: Affective instability

Outcome: 1 SMD




Analysis 9.2

Comparison 9 Active drug versus placebo: Affective instability, Outcome 2 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder

Comparison: 9 Active drug versus placebo: Affective instability


Outcome: 2 SMD on basis of change from baseline scores



Analysis 10.1

Comparison 10 Active drug versus placebo: Chronic feelings of emptiness, Outcome 1 SMD


Review: Pharmacological interventions for borderline personality disorder
 Comparison: 10 Active drug versus placebo: Chronic feelings of emptiness
 Outcome: 1 SMD



Analysis 10.2

Comparison 10 Active drug versus placebo: Chronic feelings of emptiness, Outcome 2 SM

on basis of change from baseline scores.
 Review: Pharmacological interventions for borderline personality disorder
 Comparison: 10 Active drug versus placebo: Chronic feelings of emptiness
 Outcome: 2 SMD on basis of change from baseline scores



Analysis 11.1


Comparison 11 Active drug versus placebo: Anger,


Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 11 Active drug versus placebo: Anger

Outcome: 1 SMD





Heterogeneity: not applicable
Test for overall effect: $Z = 1.31$ ($P = 0.19$)

-4 -2 0 2 4
Favours experimental Favours control

(1) Frankenburg 2002 and Hollander 2001 were not pooled, as heterogeneity seemed considerable (I^2 78%), and could not definitely be explained.

(2) cf. to (1)

(3) For Topiramate, data were analysed for male and female samples separately (I^2 of all three estimates 93%).

(4) cf. to (3)

(5) cf. to (3)

Analysis 11.2


Comparison 11 Active drug versus placebo: Anger,

Outcome 2 SMD on basis of post-means and pre-SDs

Review: Pharmacological interventions for borderline personality disorder

Comparison: 11 Active drug versus placebo: Anger

Outcome: 2 SMD on basis of post-means and pre-SDs



Analysis 11.3

Comparison 11 Active drug versus placebo: Anger,


Outcome 3 SMD on basis of change from baseline

scores

Review: Pharmacological interventions for borderline personality disorder

Comparison: 11 Active drug versus placebo: Anger

Outcome: 3 SMD on basis of change from baseline scores




Analysis 12.1

Comparison 12 Active drug versus placebo: Psychotic symptoms, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 12 Active drug versus placebo: Psychotic symptoms

Outcome: 1 SMD




Analysis 12.2

Comparison 12 Active drug versus placebo: Psychotic symptoms, Outcome 2 SMD on basis of post-means and pre-SDs

Review: Pharmacological interventions for borderline personality disorder

Comparison: 12 Active drug versus placebo: Psychotic symptoms

Outcome: 2 SMD on basis of post-means and pre-SDs




Analysis 12.3

Comparison 12 Active drug versus placebo: Psychotic symptoms, Outcome 3 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder

Comparison: 12 Active drug versus placebo: Psychotic symptoms

Outcome: 3 SMD on basis of change from baseline scores



Analysis 13.1


Comparison 13 Active drug versus placebo:

Dissociation, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 13 Active drug versus placebo: Dissociation

Outcome: 1 SMD



Analysis 14.1


Comparison 14 Active drug versus placebo: Depression,


Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 14 Active drug versus placebo: Depression

Outcome: 1 SMD






Analysis 14.2

Comparison 14 Active drug versus placebo: Depression, Outcome 2 SMD on basis of postmeans and pre-SDs

Review: Pharmacological interventions for borderline personality disorder
Comparison: 14 Active drug versus placebo: Depression
Outcome: 2 SMD on basis of post-means and pre-SDs




Analysis 14.3

Comparison 14 Active drug versus placebo: Depression, Outcome 3 Mean Change Difference

Review: Pharmacological interventions for borderline personality disorder

Comparison: 14 Active drug versus placebo: Depression

Outcome: 3 Mean Change Difference




Analysis 14.4

Comparison 14 Active drug versus placebo: Depression, Outcome 4 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 14 Active drug versus placebo: Depression

Outcome: 4 Risk Ratio



Analysis 15.1 **Comparison 15 Active drug versus placebo: Anxiety,** **Outcome 1 SMD**

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 15 Active drug versus placebo: Anxiety
 Outcome: 1 SMD




Analysis 15.2

Comparison 15 Active drug versus placebo: Anxiety, Outcome 2 Mean Change Difference

Review: Pharmacological interventions for borderline personality disorder

Comparison: 15 Active drug versus placebo: Anxiety

Outcome: 2 Mean Change Difference




Analysis 16.1


Comparison 16 Active drug versus placebo: General psychiatric pathology, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 16 Active drug versus placebo: General psychiatric pathology

Outcome: 1 SMD






Analysis 16.2


Comparison 16 Active drug versus placebo: General psychiatric pathology, Outcome 2 SMD on basis of change from baseline scores

Review: Pharmacological interventions for borderline personality disorder
Comparison: 16 Active drug versus placebo: General psychiatric pathology
Outcome: 2 SMD on basis of change from baseline scores



Analysis 17.1 **Comparison 17 Active drug versus placebo: Mental** **health status, Outcome 1 SMD**

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 17 Active drug versus placebo: Mental health status
 Outcome: 1 SMD




(1) Despite of the high heterogeneity (τ^2 84%), we decided to pool estimates, as clinical heterogeneity was quite low, and the same pattern of more favourable results in the

(2) cf. to (1)

Analysis 17.2

Comparison 17 Active drug versus placebo: Mental health status, Outcome 2 SMD on basis of post-means and pre-SDs


Review: Pharmacological interventions for borderline personality disorder
 Comparison: 17 Active drug versus placebo: Mental health status
 Outcome: 2 SMD on basis of post-means and pre-SDs



Analysis 17.3

Comparison 17 Active drug versus placebo: Mental health status, Outcome 3 Mean Change Difference

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 17 Active drug versus placebo: Mental health status
 Outcome: 3 Mean Change Difference




Analysis 17.4 **Comparison 17 Active drug versus placebo: Mental health status, Outcome 4 Risk Ratio**

Review: Pharmacological interventions for borderline personality disorder

Comparison: 17 Active drug versus placebo: Mental health status


Outcome: 4 Risk Ratio




Analysis 18.1

Comparison 18 Active drug versus placebo: Attrition/leaving the study early for any reason, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 18 Active drug versus placebo: Attrition/leaving the study early for any reason
 Outcome: 1 Risk Ratio






Analysis 19.1

Comparison 19 Active drug versus placebo: AE - body weight change, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 19 Active drug versus placebo: AE - body weight change
 Outcome: 1 SMD




Analysis 20.1

Comparison 20 Active drug versus placebo: AE - any AE, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 20 Active drug versus placebo: AE - any AE

Outcome: 1 Risk Ratio



(1) event: reporting any adverse event

(2) cf. to (1)

(3) cf. to (1)


Analysis 21.1

Comparison 21 Active drug versus placebo: AE - increased appetite, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 21 Active drug versus placebo: AE - increased appetite

Outcome: 1 Risk Ratio




Analysis 22.1

Comparison 22 Active drug versus placebo: AE - paraesthesia, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 22 Active drug versus placebo: AE - paraesthesia

Outcome: 1 Risk Ratio




Analysis 23.1

Comparison 23 Active drug versus placebo: AE - headache, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 23 Active drug versus placebo: AE - headache

Outcome: 1 Risk Ratio




Analysis 24.1

Comparison 24 Active drug versus placebo: AE - dizziness, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 24 Active drug versus placebo: AE - dizziness

Outcome: 1 Risk Ratio



Analysis 25.1


Comparison 25 Active drug versus placebo: AE - disturbance in attention, Outcome 1 Risk Ratio

Ratio.

Review: Pharmacological interventions for borderline personality disorder

Comparison: 25 Active drug versus placebo: AE - disturbance in attention

Outcome: 1 Risk Ratio




Analysis 26.1

Comparison 26 Active drug versus placebo: AE - memory problems, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 26 Active drug versus placebo: AE - memory problems

Outcome: 1 Risk Ratio




Analysis 27.1

Comparison 27 Active drug versus placebo: AE - fatigue, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 27 Active drug versus placebo: AE - fatigue

Outcome: 1 Risk Ratio




Analysis 28.1

Comparison 28 Active drug versus placebo: AE - somnolence, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 28 Active drug versus placebo: AE - somnolence

Outcome: 1 Risk Ratio




Analysis 29.1

Comparison 29 Active drug versus placebo: AE - sedation, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 29 Active drug versus placebo: AE - sedation

Outcome: 1 Risk Ratio




Analysis 30.1

Comparison 30 Active drug versus placebo: AE - insomnia, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 30 Active drug versus placebo: AE - insomnia

Outcome: 1 Risk Ratio




Analysis 31.1

Comparison 31 Active drug versus placebo: AE - anxiety, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 31 Active drug versus placebo: AE - anxiety

Outcome: 1 Risk Ratio




Analysis 32.1

Comparison 32 Active drug versus placebo: AE - nausea, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 32 Active drug versus placebo: AE - nausea

Outcome: 1 Risk Ratio




Analysis 33.1

Comparison 33 Active drug versus placebo: AE - uneasy feeling, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 33 Active drug versus placebo: AE - uneasy feeling

Outcome: 1 Risk Ratio




Analysis 34.1

Comparison 34 Active drug versus placebo: AE - constipation, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 34 Active drug versus placebo: AE - constipation

Outcome: 1 Risk Ratio




Analysis 35.1

Comparison 35 Active drug versus placebo: AE - dry mouth, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 35 Active drug versus placebo: AE - dry mouth

Outcome: 1 Risk Ratio




Analysis 36.1

Comparison 36 Active drug versus placebo: AE - nasopharyngitis, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 36 Active drug versus placebo: AE - nasopharyngitis

Outcome: 1 Risk Ratio




Analysis 37.1

Comparison 37 Active drug versus placebo: AE - menstrual pain, Outcome 1 Risk Ratio

Review: Pharmacological interventions for borderline personality disorder

Comparison: 37 Active drug versus placebo: AE - menstrual pain

Outcome: 1 Risk Ratio




Analysis 38.1

Comparison 38 Active drug versus placebo: AE - liver function: AST/SGOT baseline to endpoint mean change (U/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 38 Active drug versus placebo: AE - liver function: AST/SGOT baseline to endpoint mean change (U/L)

Outcome: 1 SMD




Analysis 39.1

Comparison 39 Active drug versus placebo: AE - liver function: ALT/SGPT baseline to endpoint mean change (U/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 39 Active drug versus placebo: AE - liver function: ALT/SGPT baseline to endpoint mean change (U/L)

Outcome: 1 SMD




Analysis 40.1
Comparison 40 Active drug versus placebo: AE - liver
function: GGT (GGPT/SGGT/YGGT) baseline to
endpoint mean change (U/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 40 Active drug versus placebo: AE - liver function: GGT (GGPT/SGGT/YGGT) baseline to endpoint mean change (U/L)

Outcome: 1 SMD




Analysis 41.1
Comparison 41 Active drug versus placebo: AE - liver
function: total bilirubin baseline to endpoint mean
change (µmol/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 41 Active drug versus placebo: AE - liver function: total bilirubin baseline to endpoint mean change (µmol/L)

Outcome: 1 SMD




Analysis 42.1

Comparison 42 Active drug versus placebo: AE - liver function: direct bilirubin baseline to endpoint mean change ($\mu\text{mol/L}$), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 42 Active drug versus placebo: AE - liver function: direct bilirubin baseline to endpoint mean change ($\mu\text{mol/L}$)

Outcome: 1 SMD




Analysis 43.1

Comparison 43 Active drug versus placebo: AE - lipids: total cholesterol baseline to endpoint change (mmol/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 43 Active drug versus placebo: AE - lipids: total cholesterol baseline to endpoint change (mmol/L)

Outcome: 1 SMD




Analysis 44.1

Comparison 44 Active drug versus placebo: AE - lipids: LDL cholesterol baseline to endpoint mean change (mmol/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 44 Active drug versus placebo: AE - lipids: LDL cholesterol baseline to endpoint mean change (mmol/L)

Outcome: 1 SMD




Analysis 45.1

Comparison 45 Active drug versus placebo: AE - lipids: HDL cholesterol (dextran precip.) baseline to endpoint mean change (mmol/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 45 Active drug versus placebo: AE - lipids: HDL cholesterol (dextran precip.) baseline to endpoint mean change (mmol/L)

Outcome: 1 SMD




Analysis 46.1

Comparison 46 Active drug versus placebo: AE - lipids: triglycerides, fasting, baseline to endpoint mean change (mmol/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 46 Active drug versus placebo: AE - lipids: triglycerides, fasting, baseline to endpoint mean change (mmol/L)

Outcome: 1 SMD




Analysis 47.1

Comparison 47 Active drug versus placebo: AE - prolactin: baseline to endpoint mean change (µg/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 47 Active drug versus placebo: AE - prolactin: baseline to endpoint mean change (µg/L)

Outcome: 1 SMD




Analysis 48.1

Comparison 48 Active drug versus placebo: AE - platelet count baseline to endpoint mean change (GI/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 48 Active drug versus placebo: AE - platelet count baseline to endpoint mean change (GI/L)

Outcome: 1 SMD




Analysis 49.1

Comparison 49 Active drug versus placebo: AE - erythrocyte count baseline to endpoint mean change (TI/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 49 Active drug versus placebo: AE - erythrocyte count baseline to endpoint mean change (TI/L)


Outcome: 1 SMD



Analysis 50.1

Comparison 50 Active drug versus placebo: AE - leukocyte count baseline to endpoint mean change (GI/L), Outcome 1 SMD


Review: Pharmacological interventions for borderline personality disorder
 Comparison: 50 Active drug versus placebo: AE - leukocyte count baseline to endpoint mean change (GI/L)
 Outcome: 1 SMD



Analysis 51.1

Comparison 51 Active drug versus placebo: AE - neutrophils, segmented, baseline to endpoint mean change (GI/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 51 Active drug versus placebo: AE - neutrophils, segmented, baseline to endpoint mean change (GI/L)
 Outcome: 1 SMD




Analysis 52.1

Comparison 52 Active drug versus placebo: AE - basophils baseline to endpoint mean change (GI/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 52 Active drug versus placebo: AE - basophils baseline to endpoint mean change (GI/L)

Outcome:




Analysis 53.1

Comparison 53 Active drug versus placebo: AE - monocytes baseline to endpoint mean change (GI/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 53 Active drug versus placebo: AE - monocytes baseline to endpoint mean change (GI/L)

Outcome: 1 SMD




Analysis 54.1

Comparison 54 Active drug versus placebo: AE - haemoglobin baseline to endpoint mean change (mml/L-F), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 54 Active drug versus placebo: AE - haemoglobin baseline to endpoint mean change (mml/L-F)

Outcome: 1 SMD




Analysis 55.1

Comparison 55 Active drug versus placebo: AE - mean cell haemoglobin concentration (MCHC) baseline to endpoint mean change (mml/L-F), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 55 Active drug versus placebo: AE - mean cell haemoglobin concentration (MCHC) baseline to endpoint mean change (mml/L-F)

Outcome: 1 SMD




Analysis 56.1

Comparison 56 Active drug versus placebo: AE - calcium baseline to endpoint mean change (mmol/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 56 Active drug versus placebo: AE - calcium baseline to endpoint mean change (mmol/L)

Outcome: 1 SMD




Analysis 57.1

Comparison 57 Active drug versus placebo: AE - albumin baseline to endpoint mean change (g/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 57 Active drug versus placebo: AE - albumin baseline to endpoint mean change (g/L)

Outcome: 1 SMD




Analysis 58.1

Comparison 58 Active drug versus placebo: AE - creatine phosphokinase baseline to endpoint mean change (U/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 58 Active drug versus placebo: AE - creatine phosphokinase baseline to endpoint mean change (U/L)

Outcome: 1 SMD




Analysis 59.1

Comparison 59 Active drug versus placebo: AE - urea nitrogen baseline to endpoint mean change (mmol/L), Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 59 Active drug versus placebo: AE - urea nitrogen baseline to endpoint mean change (mmol/L)


Outcome: 1 SMD



Analysis 60.1

Comparison 60 Active drug versus placebo: AE - pulse, standing, baseline to endpoint mean change, Outcome 1 SMD


Review: Pharmacological interventions for borderline personality disorder
 Comparison: 60 Active drug versus placebo: AE - pulse, standing, baseline to endpoint mean change
 Outcome: 1 SMD



Analysis 61.1

Comparison 61 Active drug versus placebo: AE - pulse, supine, baseline to endpoint mean change, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder
 Comparison: 61 Active drug versus placebo: AE - pulse, supine, baseline to endpoint mean change
 Outcome: 1 SMD




Analysis 62.1

Comparison 62 Active drug versus placebo: AE - diastolic blood pressure, standing, baseline to endpoint mean change, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 62 Active drug versus placebo: AE - diastolic blood pressure, standing, baseline to endpoint mean change

Outcome: 1 SMD




Analysis 63.1

Comparison 63 Active drug versus placebo: AE - diastolic blood pressure, supine, baseline to endpoint mean change, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 63 Active drug versus placebo: AE - diastolic blood pressure, supine, baseline to endpoint mean change

Outcome: 1 SMD




Analysis 64.1

Comparison 64 Active drug versus placebo: AE - systolic blood pressure, standing, baseline to endpoint mean change, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 64 Active drug versus placebo: AE - systolic blood pressure, standing, baseline to endpoint mean change

Outcome: 1 SMD




Analysis 65.1

Comparison 65 Active drug versus placebo: AE - systolic blood pressure, supine, baseline to endpoint mean change, Outcome 1 SMD

Review: Pharmacological interventions for borderline personality disorder

Comparison: 65 Active drug versus placebo: AE - systolic blood pressure, supine, baseline to endpoint mean change

Outcome: 1 SMD




Analysis 66.1

Comparison 66 Drug versus drug: BPD severity, Outcome 1 First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 66 Drug versus drug: BPD severity

Outcome: 1 First-generation antipsychotic versus antidepressant




Analysis 67.1

Comparison 67 Drug versus drug: Interpersonal problems, Outcome 1 First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 67 Drug versus drug: Interpersonal problems

Outcome: 1 First-generation antipsychotic versus antidepressant



Analysis 68.1

Comparison 68 Drug versus drug: Impulsivity,


Outcome 1 First-generation antipsychotic versus

antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 68 Drug versus drug: Impulsivity

Outcome: 1 First-generation antipsychotic versus antidepressant



Analysis 68.2

Comparison 68 Drug versus drug: Impulsivity,


Outcome 2 Second-generation antipsychotic versus

antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 68 Drug versus drug: Impulsivity

Outcome: 2 Second-generation antipsychotic versus antidepressant



Analysis 69.1


Comparison 69 Drug versus drug: Anger, Outcome 1

First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 69 Drug versus drug: Anger

Outcome: 1 First-generation antipsychotic versus antidepressant




Analysis 70.1

Comparison 70 Drug versus drug: Psychotic symptoms, Outcome 1 First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 70 Drug versus drug: Psychotic symptoms

Outcome: 1 First-generation antipsychotic versus antidepressant




Analysis 71.1

Comparison 71 Drug versus drug: Depression, Outcome 1 First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 71 Drug versus drug: Depression

Outcome: 1 First-generation antipsychotic versus antidepressant




Analysis 71.2

Comparison 71 Drug versus drug: Depression, Outcome 2 Second-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 71 Drug versus drug: Depression

Outcome: 2 Second-generation antipsychotic versus antidepressant



Analysis 72.1


Comparison 72 Drug versus drug: Anxiety, Outcome 1

First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 72 Drug versus drug: Anxiety

Outcome: 1 First-generation antipsychotic versus antidepressant




Analysis 73.1

Comparison 73 Drug versus drug: General psychiatric pathology, Outcome 1 First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 73 Drug versus drug: General psychiatric pathology

Outcome: 1 First-generation antipsychotic versus antidepressant



Analysis 74.1

Comparison 74 Drug versus drug: Mental health status,


Outcome 1 First-generation antipsychotic versus

antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 74 Drug versus drug: Mental health status

Outcome: 1 First-generation antipsychotic versus antidepressant



Analysis 75.1

Comparison 75 Drug versus drug: AE - attrition,


Outcome 1 First-generation antipsychotic versus first-

generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 75 Drug versus drug: AE - attrition

Outcome: 1 First-generation antipsychotic versus first-generation antipsychotic




Analysis 75.2

Comparison 75 Drug versus drug: AE - attrition, Outcome 2 First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 75 Drug versus drug: AE - attrition

Outcome: 2 First-generation antipsychotic versus antidepressant




Analysis 75.3

Comparison 75 Drug versus drug: AE - attrition, Outcome 3 Second-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 75 Drug versus drug: AE - attrition

Outcome: 3 Second-generation antipsychotic versus antidepressant




Analysis 76.1

Comparison 76 Drug versus drug: AE - body weight change, Outcome 1 First-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 76 Drug versus drug: AE - body weight change

Outcome: 1 First-generation antipsychotic versus antidepressant




Analysis 76.2

Comparison 76 Drug versus drug: AE - body weight change, Outcome 2 Second-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 76 Drug versus drug: AE - body weight change

Outcome: 2 Second-generation antipsychotic versus antidepressant




Analysis 77.1
Comparison 77 Drug versus drug: AE - any AE,
Outcome 1 First-generation antipsychotic versus first-
generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 77 Drug versus drug: AE - any AE

Outcome: 1 First-generation antipsychotic versus first-generation antipsychotic




Analysis 78.1
Comparison 78 Drug versus drug: AE - sedation,
Outcome 1 Second-generation antipsychotic versus
antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 78 Drug versus drug: AE - sedation

Outcome: 1 Second-generation antipsychotic versus antidepressant




Analysis 79.1

Comparison 79 Drug versus drug: AE - sleepiness/drowsiness, Outcome 1 First-generation antipsychotic versus first-generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 79 Drug versus drug: AE - sleepiness/drowsiness

Outcome: 1 First-generation antipsychotic versus first-generation antipsychotic




Analysis 80.1

Comparison 80 Drug versus drug: AE - restlessness, Outcome 1 First-generation antipsychotic versus first-generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 80 Drug versus drug: AE - restlessness

Outcome: 1 First-generation antipsychotic versus first-generation antipsychotic




Analysis 80.2

Comparison 80 Drug versus drug: AE - restlessness, Outcome 2 Second-generation antipsychotic versus antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 80 Drug versus drug: AE - restlessness

Outcome: 2 Second-generation antipsychotic versus antidepressant



(1) event: akathisia


Analysis 81.1

Comparison 81 Drug versus drug: AE - muscle spasms, Outcome 1 First-generation antipsychotic versus first- generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 81 Drug versus drug: AE - muscle spasms

Outcome: 1 First-generation antipsychotic versus first-generation antipsychotic




Analysis 82.1

Comparison 82 Drug versus drug: AE - fainting spells, Outcome 1 First-generation antipsychotic versus first- generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 82 Drug versus drug: AE - fainting spells

Outcome: 1 First-generation antipsychotic versus first-generation antipsychotic




Analysis 83.1

Comparison 83 Drug versus combination of drugs: Impulsivity, Outcome 1 Secondgeneration antipsychotic versus second-generation antipsychotic + antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 83 Drug versus combination of drugs: Impulsivity

Outcome: 1 Second-generation antipsychotic versus second-generation antipsychotic +
antidepressant




Analysis 83.2
Comparison 83 Drug versus combination of drugs:
Impulsivity, Outcome 2 Antidepressant versus
antidepressant + second-generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 83 Drug versus combination of drugs: Impulsivity

Outcome: 2 Antidepressant versus antidepressant + second-generation antipsychotic




Analysis 84.1
Comparison 84 Drug versus combination of drugs:
Depression, Outcome 1 Secondgeneration antipsychotic
versus second-generation antipsychotic +
antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 84 Drug versus combination of drugs: Depression

Outcome: 1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant




Analysis 84.2
Comparison 84 Drug versus combination of drugs:
Depression, Outcome 2 Antidepressant versus
antidepressant + second-generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 84 Drug versus combination of drugs: Depression

Outcome: 2 Antidepressant versus antidepressant + second-generation antipsychotic




Analysis 85.1
Comparison 85 Drug versus combination of drugs: AE -
attrition, Outcome 1 Secondgeneration antipsychotic
versus second-generation antipsychotic +
antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 85 Drug versus combination of drugs: AE - attrition

Outcome: 1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant




Analysis 85.2

Comparison 85 Drug versus combination of drugs: AE - attrition, Outcome 2 Antidepressant versus antidepressant + second-generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 85 Drug versus combination of drugs: AE - attrition

Outcome: 2 Antidepressant versus antidepressant + second-generation antipsychotic




Analysis 86.1

Comparison 86 Drug versus combination of drugs: AE - body weight change, Outcome 1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 86 Drug versus combination of drugs: AE - body weight change

Outcome: 1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant




Analysis 86.2**Comparison 86 Drug versus combination of drugs: AE - body weight change, Outcome 2 Antidepressant versus antidepressant + second-generation antipsychotic**

Review: Pharmacological interventions for borderline personality disorder

Comparison: 86 Drug versus combination of drugs: AE - body weight change


Outcome: 2 Antidepressant versus antidepressant + second-generation antipsychotic

**Analysis 87.1****Comparison 87 Drug versus combination of drugs: AE - sedation, Outcome 1 Secondgeneration antipsychotic versus second-generation antipsychotic + antidepressant**

Review: Pharmacological interventions for borderline personality disorder

Comparison: 87 Drug versus combination of drugs: AE - sedation

Outcome: 1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant




Analysis 87.2

Comparison 87 Drug versus combination of drugs: AE - sedation, Outcome 2 Antidepressant versus antidepressant + second-generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 87 Drug versus combination of drugs: AE - sedation

Outcome: 2 Antidepressant versus antidepressant + second-generation antipsychotic




Analysis 88.1

Comparison 88 Drug versus combination of drugs: AE - akathisia, Outcome 1 Secondgeneration antipsychotic versus second-generation antipsychotic + antidepressant

Review: Pharmacological interventions for borderline personality disorder

Comparison: 88 Drug versus combination of drugs: AE - akathisia

Outcome: 1 Second-generation antipsychotic versus second-generation antipsychotic + antidepressant




>Analysis 88.2
Comparison 88 Drug versus combination of drugs: AE -
akathisia, Outcome 2 Antidepressant versus
antidepressant + second-generation antipsychotic

Review: Pharmacological interventions for borderline personality disorder

Comparison: 88 Drug versus combination of drugs: AE - akathisia

Outcome: 2 Antidepressant versus antidepressant + second-generation antipsychotic



ADDITIONAL TABLES

Table 1
First-generation antipsychotics vs. placebo: outcome
scales

	Haloperidol		Flupenthixol decanoate	Thiothixene
	Soloff 1989	Soloff 1993	Montgomery 1979/82	Goldberg 1986
BPD severity	-	BSI	-	SIB-borderline score
avoidance of abandonment	-	-	-	-
interpersonal problems	SCL-90-INT	ADS-rejection sensitivity	-	HSCL-INT
identity disturbance	-	-	-	-
impulsivity	BIS, Ward Scale of Impulse Action Patterns, STIC BIS used because of width of use and self-reporting format	BIS, Ward Scale of Impulse Action Patterns, STIC BIS used because of width of use and self-reporting format		
suicidal ideation	-	-	-	-
suicidal behaviour	-	-	number of patients with suicidal act during treatment (6 months period)	-
self-mutilating behaviour	-	-	-	-
affective instability	-	-	-	-
feelings of emptiness	-	-	-	-

	Haloperidol		Flupenthixol decanoate	Thiothixene
	Soloff 1989	Soloff 1993	Montgomery 1979/82	Goldberg 1986
anger	SCL-90-HOS, BDHI SCL-90-HOS used because of greater sensitivity to change	SCL-90-HOS, BDHI, ADS-reactivity SCL-90-HOS used because of greater sensitivity to change than BDHI and greater width of than ADS scale		HSCL-HOS
psychotic/paranoid symptoms	SCL-90-PAR, SCL-90-PSY, SSI SCL-90-PAR used because regarded as most adequately reflecting BPD relevant pathology	SCL-90-PAR, SCL-90-PSY, SSI SCL-90-PAR used because regarded as most adequately reflecting BPD relevant pathology		SIB-suspicious/paranoid subscale, HSCL-90-PSY SIB-suspicious/paranoid subscale used because regarded as most adequately reflecting BPD-relevant pathology
dissociative symptoms	-	-	-	
depression	BDI, SCL-90-DEP, Ham-D BDI used because of width of use and self-report format	BDI, SCL-90-DEP, Ham-D, ADS total BDI used because of width of use and self-report format		HSCL-DEP
anxiety	SCL-90-ANX	SCL-90-ANX	-	-
general psychiatric pathology	SCL-90-GSI	SCL-90-GSI	-	-
mental health status	GAS	GAS	-	GAS
attrition	number of patients lost after randomisation	number of patients lost after randomisation	number of patients lost after randomisation	number of patients lost after randomisation
adverse events	-	ADS-weight gain	-	-

Table 2
Second-generation antipsychotics vs. placebo: outcome scales (part 1)

	Aripiprazole	Ziprasidone
	Nickel 2006	Pascual 2008
BPD severity	-	CGI-BPD-global
avoidance of abandonment	-	CGI-BPD-abandonment
interpersonal problems	SCL-90-R-INT (t-transformed)	CGI-BPD-unstable relations
identity disturbance	-	CGI-BPD-identity
impulsivity	STAXI-OUT	CGI-BPD-impulsivity
suicidal ideation		CGI-BPD-suicide
self-mutilating behaviour	number of patients with self-injury during treatment (8 weeks period)	-
affective instability	-	CGI-BPD-affect instability

	Aripiprazole	Ziprasidone
	Nickel 2006	Pascual 2008
feelings of emptiness	-	CGI-BPD-emptiness
anger	SCL-90-R-HOS (t-transformed), STAXI-trait, STAXI-state, STAXI-anger in SCL-90-R-HOS used because regarded as most comprehensive measure	CGI-BPD-anger
psychotic/paranoid symptoms	SCL-90-R-PAR, SCL-90-R-PSY SCL-90-R-PAR used because considered as most adequately reflecting BPD relevant pathology	CGI-BPD-paranoid ideation; BPRS CGI-BPD used because specific for assessment in BPD patients
dissociative symptoms		
depression	Ham-D, SCL-90-R-DEP Ham-D used because also reported by other trials within this comparison category	Ham-D-17; BDI Ham-D used because also reported by other trials within this comparison category
anxiety	HARS; SCL-90-R-ANX HARS used because also reported by other trials within this comparison category	HARS
general psychiatric pathology	SCL-90-R-GSI	SCL-90-R-GSI
adverse effects	-	attrition
patient reported adverse events	-	minor sedation, dizziness, uneasy feeling

Table 3
Second-generation antipsychotics vs. placebo: outcome scales (part 2)

	Olanzapine			
	Bogenschutz 2004	Linehan 2007	Soler 2005	Zanarini 2001
BPD severity	-	-	-	-
avoidance of abandonment	CGI-abandonment	-	-	-
interpersonal problems	CGI-unstable relationships, SCL-90-R-INT CGI-unstable relationships used because specific for assessment in BPD patients			
identity disturbance	CGI-identity disturbance	-	-	-
impulsivity	CGI-impulsivity	-	Behavioural reports of numbers of episodes of impulsivity/aggressive behaviour	-
suicidal ideation	CGI-recurrent suicidal ideation	number of patients with high suicidality scores on OAS-M-suicidality sub-scale (i.e., reporting frequent		

Olanzapine				
	Bogenschutz 2004	Linehan 2007	Soler 2005	Zanarini 2001
		suicide ideation and/or planning or behaviour)		
suicidal behaviour	-		Behavioural reports of numbers of episodes of self-injuring behaviour/ suicide attempts	-
self-mutilating behaviour	-	number of patients with self-injury during treatment	-	-
affective instability	CGI-affective instability	-	-	-
feelings of emptiness	CGI-chronic feelings of emptiness	-	-	-
anger	CGI-inappropriate anger, OAS-M, AIAQ, SCL-90-HOS CGI-inap- propriate anger used because specific for assessment in BPD patients			
psychotic/paranoid symptoms	CGI-transient paranoia or dissociation, SCL-90-PSY, SCL-90-PAR CGI-transient paranoia or dissociation used because specific for assessment in BPD patients			
depression	-	Ham-D	Ham-D	-
anxiety	-		HARS	-
general psychiatric - - pathology				
mental health status	-		CGI-S	-
attrition	number of patients lost after randomisation	number of patients lost after randomisation	number of patients lost after randomisation	number of patients lost after randomisation
adverse effects	baseline to endpoint weight change (kg)	baseline to endpoint weight change (kg)	baseline to end-point weight change (kg) , baseline to endpoint increase in cholesterol levels (mg/dl)	baseline to endpoint weight change (kg)
patient reported adverse events	-	-	-	constipation, sedation

Table 4
Second-generation antipsychotics vs. placebo: outcome
scales (part 3)

Olanzapine		
	Schulz 2007	Zanarini 2007
BPD severity	Zan-BPD-total	Zan-BPD-total
avoidance of abandonment	Zan-BPD-frantic efforts to avoid abandonment	Zan-BPD-frantic efforts to avoid abandonment
interpersonal problems	ZAN-BPD-unstable interpersonal relationships	ZAN-BPD-unstable interpersonal relationships, SCL-90-R-INT ZAN-BPD-unstable relationships used because specific for assessment in BPD patients
identity disturbance	ZAN-BPD-identity disturbance	ZAN-BPD-identity disturbance
impulsivity	ZAN-BPD-impulsivity that are self-damaging, OAS-M-aggression ZAN-BPD-impulsivity used because specific for assessment in BPD patients	ZAN-BPD-impulsivity that are self-damaging, OAS-M-aggression ZAN-BPD-impulsivity used because specific for assessment in BPD patients
suicidal ideation	ZAN-BPD-suicidal or self-mutilating behaviour	ZAN-BPD-suicidal or self-mutilating behaviour, OAS-M-suicidality ZAN-BPD subscale used because specific for assessment in BPD patients
suicidal behaviour		
self-mutilating behaviour		
affective instability	ZAN-BPD-affective instability	ZAN-BPD-affective instability
chronic feelings of emptiness	ZAN-BPD-chronic feelings of emptiness	ZAN-BPD-chronic feelings of emptiness
anger	ZAN-BPD-intense anger	ZAN-BPD-intense anger, OAS-M-irritability, SCL-90-R-HOS ZAN-BPD subscale used because specific for assessment in BPD patients
psychotic/paranoid symptoms	ZAN-BPD-paranoid ideation of dissociation	ZAN-BPD-paranoid ideation of dissociation, SCL-90-R-PAR ZAN-BPD subscale used because specific for assessment in BPD patients
depression	MADRS	MADRS, SCL-90-R-DEP MADRS used because also available for Schulz 2007
anxiety	-	SCL-90-R-ANX
general psychiatric pathology	SCL-90-R-GSI	SCL-90-R-GSI
mental health status	GAF, Sheehan Scale GAF used because of width of use	GAF, Sheehan Scale GAF used because of width of use
attrition	number of patients lost after randomisation	number of patients lost after randomisation
adverse effects	baseline to endpoint weight change (kg)	baseline to endpoint weight change (kg)

Olanzapine		
	Schulz 2007	Zanarini 2007
patient-reported adverse events	anxiety, dry mouth, fatigue, headache, increased appetite, insomnia, nausea, number of patients experiencing any AE, sedation, somnolence all used	disturbed attention, dry mouth, fatigue, headache, increased appetite, insomnia, nausea, number of patients experiencing any AE, somnolence
laboratory values	lipids (baseline to endpoint mean changes): LDL cholesterol (mmol/L), total cholesterol (mmol/L) liver function values (baseline to endpoint mean changes): ALT/SGPT (U/L), AST/SGOT (U/L), total bilirubin (umol/L), direct bilirubin (umol/L) prolactin (micrograms/L)	lipids (baseline to endpoint mean changes): HDL cholesterol (mmol/L), total cholesterol (mmol/L), triglycerides fasting (mmol/L) liver function values (baseline to endpoint mean changes): GGT (GGPT/SGGT/YGGT; U/L), ALT/SGPT (U/L), AST/SGOT (U/L) prolactin baseline to endpoint mean change (micrograms/L) blood values (baseline to endpoint mean changes): leukocyte count (GI/L), monocytes (GI/L), neutrophils segmented (GI/L), platelet count (GI/L)

Table 5
Mood stabiliser vs. placebo: outcome scales (part 1)

	Carbamazepine	Valproate Semisodium	
	De la Fuente 1993	Frankenburg 2002	Hollander 2001
BPD severity	-	-	-
avoidance of abandonment	-	-	-
interpersonal problems	SCL-90-INT	SCL-90-INT	
identity disturbance	-	-	-
impulsivity	Acting-out Scale: number of patients worsened or unimproved as compared to baseline	MOAS	OAS-M-aggression (not used: Assault Questionnaire, because of close affinity of OAS-M with the MOAS scale as used by the Frankenburg trial)
suicidal ideation	-	-	OAS-M-suicidality
suicidal behaviour	-	-	-
self-mutilating behaviour	-	-	-
affective instability	-	-	-
feelings of emptiness	-	-	-
anger	SCL-90-HOS	SCL-90-HOS	OAS-M-irritability
psychotic/paranoid symptoms	SCL-90-PAR (not used: BPRS as the SCL-scale had also been used by another trial within this comparison category; SCL-90-PSY as the PAR subscale reflects BPD relevant pathology more adequately)		

	Carbamazepine	Valproate Semisodium	
dissociative symptoms	-	-	-
depression	SCL-90-DEP (not used: Ham-D as the SCL-90-DEP subscale was also available from other trials within this comparison category)	SCL-90-DEP	BDI
anxiety	SCL-90-ANX	-	-
general psychiatric pathology	SCL-90-total	-	-
mental health status	-	-	CGI-I: number of patients with an CGI-I of 3 or more (i.e., minimally improved to very much worse)
attrition	number of patients lost after randomisation	number of patients lost after randomisation	number of patients lost after randomisation
adverse effects: weight gain	-	weight gain (kg; derived out of lb data)	-

Table 6
Mood stabilisers vs. placebo: outcome scales (part 2)

	Lamotrigine	
	Tritt 2005	Reich 2009
BPD severity	-	ZAN-BPD-total
avoidance of abandonment	-	-
interpersonal problems	-	-
identity disturbance	-	-
impulsivity	STAXI-anger out (not used: STAXI-anger in, STAXI-control)	ZAN-BPD-impulsivity
suicidal ideation	-	-
suicidal behaviour	-	-
self-mutilating behaviour	-	-
ffective instability	-	ZAN-BPD-affective instability (not used: Affective Liability Scale)
feelings of emptiness		-
anger	STAXI-anger trait (not used: STAXI-anger state, as this subscale refers only to the intensity of angry feelings at the time of testing)	-
psychotic/paranoid symptoms		-
dissociative symptoms		-
depression		-
anxiety		-
general psychiatric pathology		-
mental health status		-

Lamotrigine		
attrition	number of patients lost after randomisation	number of patients lost after randomisation
adverse effects: absolute weight	weight (kg)	-
patient-reported adverse events	-	number of patients with rash

Table 7
Mood stabilisers vs. placebo: outcome scales (part 3)

Topiramate			
	Loew 2006	Nickel 2004	Nickel 2005
BPD severity	-	-	-
avoidance of abandonment	-	-	-
interpersonal problems	SCL-90-R-INT (t-transformed)	-	-
identity disturbance	-	-	-
impulsivity	-	STAXI-anger out (not used: STAXI-anger in, STAXI-control)	STAXI-anger out (not used: STAXI-anger in, STAXI-control)
suicidal ideation	-	-	-
suicidal behaviour			
self-mutilating behaviour	-	-	-
affective instability	-	-	-
feelings of emptiness	-	-	-
anger	SCL-90-R-HOS (t-transformed)	STAXI-anger trait (not used: STAXI-anger state, as this sub-scale refers only to the intensity of angry feelings at the time of testing)	STAXI-anger trait (not used: STAXI-anger state, as this sub-scale refers only to the intensity of angry feelings at the time of testing)
psychotic/paranoid symptoms	SCL-90-R-PAR (t-transformed) (not used: SCL-90-R-PSY, as the SCL-90-R-PAR subscale reflects BPD relevant pathology more adequately)		
dissociative symptoms	-	-	-
depression	SCL-90-R-DEP (t-transformed)	-	-
anxiety	SCL-90-R-ANX (t-transformed)	-	-
general psychiatric pathology	SCL-90-R-GSI (t-transformed)	-	-
mental health status	-	-	-
attrition	number of patients lost after randomisation	number of patients lost after randomisation	number of patients lost after randomisation
adverse effects: absolute weight	weight (kg)	weight (kg)	weight (kg)
patient-reported adverse events	memory problems, troubles in concentrating, headache, fatigue, dizziness, menstrual pain, paresthesia	-	-

Table 8
Antidepressants vs. placebo: outcome scales (part 1)

	Amitriptyline	Fluoxetine	
	Soloff 1989	Salzman 1995	Simpson 2004
BPD severity	-	-	-
avoidance of abandonment	-	-	-
interpersonal problems	SCL-90-INT	-	-
identity disturbance	-	-	-
impulsivity	BIS; not used: Ward Scale of Impulsive Action, Self-Report Test of Impulse Control total because BIS is a self-report measure and broadly used		OAS-M-aggression; not used: STAXI-anger-out because other OAS-M-sub-scales were also used for several outcomes (s. below)
suicidal ideation	-	-	OAS-M-suicidality
suicidal behaviour	-	-	-
self-mutilating behaviour	-	-	OAS-M-assault against self
ffective instability			
feelings of emptiness	-	-	-
anger	SCL-90-HOS not used: BDHI, ADDS-reactivity)	POMS-anger; not used: OAS-M-anger against objects (only one spectrum of anger entailed), PDRS-anger (only based on one interviewer-rated item)	
psychotic/paranoid symptoms	SCL-90-PAR (not used: SCL-90-PSY, IMPS, SSI because SCL-90-PAR is a self-rated measure most adequately reflecting BPD relevant pathology	-	-
dissociative symptoms	-	-	DES
depression	BDI; not used: SCL-90-DEP, Ham-D-17, Ham-D-24, ADS, as all but the SCL-90-DEP scale are observer rated, and the BDI is more commonly used in the	Ham-D; not used: POMS-dep because the Ham-D was also used by other studies within this comparison category	BDI
	assessment of depression than the SCL-90-DEP		
anxiety	SCL-90-ANX	-	STAI
general psychiatric pathology	SCL-90-GSI	-	-
mental health status	GAS	GAS	GAF
attrition	number of non-completers	-	number of non-completers
adverse effects		-	-

Table 9
Antidepressants vs. placebo: outcome scales (part 2)

	SSRI: Fluvoxamine	MAOI: Phenelzine	Mianserin
	Rinne 2002	Soloff 1993	Montgomery 81/82/83
BPD severity		BSI	-
avoidance of abandonment			-
dysfunctional interpersonal patterns		ADI-rejection sensitivity	-
identity disturbance			-
impulsivity	BPDSI-impulsivity	BIS; not used: Ward Scale of Impulsive Action, Self-Report Test of Impulse Control total because BIS is a self-report measure and broadly used	
suicidal ideation			-
suicidal behaviour			-
self-mutilating behaviour	-	-	number of patients with self-harming behaviour during 6 months of treatment
affective instability	BPDSI-rapid mood shifts	-	-
feelings of emptiness			-
anger	BPDSI-anger	SCL-90-HOS; not used: BDHI, ADDS-reactivity)	-
psychotic/paranoid ideation		SCL-90-PAR; not used: SCL-90-PSY, IMPS, SSI because SCL-90-PAR is a self-rated measure most adequately reflecting BPD relevant pathology	
dissociative symptoms	-	-	-
depression		BDI; not used: SCL-90-DEP, Ham-D-17, Ham-D-24, ADS, as all but the SCL-90-DEP scale are observer rated, and the BDI is more commonly used in the assessment of depression than the SCL-90-DEP	
anxiety	-	SCL-90-ANX	-
general psychiatric pathology	-	SCL-90-GSI	-
mental health status	-	GAS	-
attrition	number of patients not completing the study protocol	number of patients not completing the study protocol	-
adverse effects	-	ADI-weight gain	

Table 10
Miscellaneous active agents vs. placebo: outcome scales

Omega-3 fatty acid		
	Hallahan 2007	Zanarini 2003
BPD severity	-	-
avoidance of abandonment	-	-
interpersonal problems	-	-
identity disturbance	-	-
impulsivity	-	MOAS
suicidal ideation	OAS-M-suicidality: number of patients with suicidality subscale score >1 (i.e., at least slight suicidal tendency)	-
suicidal behaviour	-	-
self-mutilating behaviour	number of patients with self-harm episodes during treatment	-
affective instability	-	-
feelings of emptiness	-	-
anger	-	-
psychotic/paranoid symptoms	-	-
dissociative symptoms	-	-
depression	number of patients not experiencing at least a 50% or 70% reduction of depressive pathology as assessed both by BDI and HAM-D BDI used as it shows high concurrent validity with MADRS, whereas HAM-D mainly fo-cusses on somatic depressive symptoms 50% cut-off data used as this is more sensitive to change, as are the continuous data reported by the other relevant trial in this comparison category	MADRS
anxiety	-	-
general psychiatric pathology	-	-
mental health status	-	-
attrition	number of patients lost after randomisation	number of patients lost after randomisation
adverse effects	-	-

Table 11
First-generation antipsychotic vs. first generation antipsychotic: outcome scales

Loxapine vs. Chlorpromazine	
	Leone 1982
BPD severity	-
avoidance of abandonment	-
interpersonal problems	-
identity disturbance	-

Loxapine vs. Chlorpromazine	
Leone 1982	
impulsivity	-
suicidal ideation	-
suicidal behaviour	-
self-mutilating behaviour	-
affective instability	-
feelings of emptiness	-
anger	-
psychotic/paranoid symptoms	-
dissociative symptoms	-
depression	-
anxiety	-
general psychiatric pathology	-
mental health status	-
attrition	number of patients lost after randomisation
patient-reported adverse events	number of patients experiencing anyAE, sleepiness/drowsiness, restlessness, muscle spasms, fainting spells

Table 12
First-generation antipsychotics vs. antidepressants:
outcome scales

Haloperidol vs. Amitriptyline		Haloperidol vs. Phenelzine Sulfate
Soloff 1989		Soloff 1993
BPD severity	-	BSI
avoidance of abandonment	-	-
interpersonal problems	SCL-90-INT	ADS-rejection sensitivity
identity disturbance	-	-
impulsivity	BIS, Ward Scale of Impulse Action Patterns, STIC BIS used because of width of use and self-reporting format	BIS, Ward Scale of Impulse Action Patterns, STIC BIS used because of width of use and self-reporting format
suicidal ideation	-	-
suicidal behaviour	-	-
self-mutilating behaviour	-	-
affective instability	-	-
feelings of emptiness	-	-
anger	SCL-90-HOS, BDHI SCL-90-HOS used because of greater sensitivity to change	SCL-90-HOS, BDHI, ADS-reactivity SCL-90-HOS used because of greater sensitivity to change than BDHI and greater width of than ADS scale
psychotic/paranoid symptoms	SCL-90-PAR, SCL-90-PSY, SSI SCL-90-PAR used because regarded as most adequately reflecting BPD relevant pathology	SCL-90-PAR, SCL-90-PSY, SSI

	Haloperidol vs. Amitriptyline	Haloperidol vs. Phenelzine Sulfate
	Soloff 1989	Soloff 1993
		SCL-90-PAR used because regarded as most adequately reflecting BPD relevant pathology
dissociative symptoms	-	-
depression	BDI, SCL-90-DEP, Ham-D BDI used because of width of use and self-report format	BDI, SCL-90-DEP, Ham-D, ADS total BDI used because of width of use and self-report format
anxiety	SCL-90-ANX	SCL-90-ANX
general psychiatric pathology	SCL-90-GSI	SCL-90-GSI
mental health status	GAS	GAS
attrition	number of patients lost after randomisation	number of patients lost after randomisation
adverse events	-	ADS-weight gain

Table 13
Second-generation antipsychotics vs. antidepressants:
outcome scales

	Olanzapine vs. Fluoxetine
	Zanarini 2004
BPD severity	-
avoidance of abandonment	-
interpersonal problems	-
identity disturbance	-
impulsivity	OAS-M
suicidal ideation	-
suicidal behaviour	-
self-mutilating behaviour	-
affective instability	-
feelings of emptiness	-
anger	-
psychotic/paranoid symptoms	-
dissociative symptoms	-
depression	MADRS
anxiety	-
general psychiatric pathology	-
mental health status	-
attrition	number of patients lost after randomisation
Adverse effects	baseline to endpoint weight change (kg)
patient-reported adverse events	number of patients experiencing mild sedation, akathisia

Table 14
Second-generation antipsychotics vs. second-generation antipsychotics plus antidepressants: outcome scales

Olanzapine vs. Olanzapine + Fluoxetine	
Zanarini 2004	
BPD severity	-
avoidance of abandonment	
interpersonal problems	-
identity disturbance	-
impulsivity	OAS-M
suicidal ideation	-
suicidal behaviour	-
self-mutilating behaviour	-
affective instability	-
feelings of emptiness	-
anger	-
psychotic/paranoid symptoms	-
dissociative symptoms	-
depression	MADRS
anxiety	-
general psychiatric pathology	-
mental health status	-
attrition	number of patients lost after randomisation
Adverse effects	baseline to endpoint weight change (kg)
patient-reported adverse events	number of patients experiencing mild sedation, akathisia

Table 15
Antidepressants vs. antidepressants plus second-generation antipsychotics: outcome scales

Fluoxetine vs. Fluoxetine + Olanzapine	
Zanarini 2004	
BPD severity	-
avoidance of abandonment	-
interpersonal problems	-
identity disturbance	-
impulsivity	OAS-M
suicidal ideation	-
suicidal behaviour	-
self-mutilating behaviour	-

Fluoxetine vs. Fluoxetine + Olanzapine	
Zanarini 2004	
affective instability	-
feelings of emptiness	-
anger	-
psychotic/paranoid symptoms	-
dissociative symptoms	-
depression	MADRS
anxiety	-
general psychiatric pathology	-
mental health status	-
attrition	number of patients lost after randomisation
Adverse effects	baseline to endpoint weight change (kg)
patient-reported adverse events	number of patients experiencing mild sedation, akathisia

Appendix 1. ASSIA search strategy

We searched Applied Social Sciences Index and Abstracts (1987 to September 2009):
 ((personality near disorder*) or ((antisocial* near disorder*) or (avoidant* near disorder*) or (borderline* near disorder*)) or ((dependent* near disorder*) or (histrionic* near disorder*) or (narcissistic* near disorder*)) or ((obsessive* near disorder*) or (compulsive* near disorder*) or (paranoid* near disorder*)) or (((passive* near disorder*) or (aggress* near disorder*) or (somasochistic* near disorder*)) or (schizo* near disorder*)) or (((passive* and disorder*) or (aggress* and disorder*) or (somasochistic* and disorder*)) or ((schizo* and disorder*) or (paranoid* and disorder*) or (compulsive* and disorder*)) or ((obsessive* and disorder*) or (narcissistic and disorder*) or (histrionic* and disorder*))) or (((personality and disorder*) or (antisocial* and disorder*) or (avoidant* and disorder*)) or ((borderline* and disorder*) or (dependent* and disorder*))) and ((AB=randomi* or TI=randomi*) or (DE=(randomi?ed controlled trials) or AB=(double* blind*) or TI=(double* blind*)) or (DE=(double blind studies) or (single* near blind*)))

Appendix 2. BIOSIS search strategy

We searched BIOSIS (1985 to 16 September 2009) using the phrase: (((al: ((personality and disorder))) or al: ((antisocial and behaviour))) or al: ((antisocial and behavior)) or (((al: ((self and defeating))) or al: ((parano* and person*))) or al: ((gender and identity)) or (al: ((asocial or antisocial* or dissocial* or psychopath* or sadist* or sociopath*))) and al: ((person*)) and or (al: ((moral and insanity)) or ((al: ((psychopath* or sociopath* or dissocial* or sadist* or schizotypal self-defeating or borderline or avoidant or dependent or depressive))) and al: (person*) or ((al: ((histrionic or multi-impulsive or multiple or narcissistic or passive-aggressive))) and al: (person*) and ((al: ((randomi* or crossover or random-assignment))) or al: (((singl* or doubl* or tripl* or trebl*) and (mask* or blind*))))

Appendix 3. COPAC search strategy

We searched the Consortium of University Research Libraries joint catalogue in September 2009 using the phrase: randomi* OR ((double OR single OR triple OR treble) and blind) OR prospective OR (clinical and trial) We then downloaded results into a Procite5 database and searched again using the terms: (antisocial* OR asocial* OR avoidant OR borderline OR dependent OR depressive OR dissocial OR dissocial* OR histrionic OR moral OR multi-impulsive OR multiple* OR narcissistic OR parano* OR passive-aggressive OR psychopath* OR sadis* OR schizotypal OR self-defeating OR sociopath*)

Appendix 4. CENTRAL search strategy

We searched CENTRAL (Cochrane library) 2009, issue 3, using the phrase: [(antisocial-personality-disorder*:me OR personality-disorders*:me OR sexual-and-gender-disorders*:me OR multiple-personality-disorder*:me OR paraphilias*:me) OR (multi-impulsive and personality) OR (parano* NEAR person*) OR (asocial* NEAR person) OR (dissocial* NEAR person) OR (psychopath* NEAR person) OR (sadist* NEAR person) OR (sociopath* NEAR person*) OR (moral NEAR insanity) OR ((personality and disorder*) and (((avoidant OR multiimpulsive) OR narcissistic) OR self-defeating) OR personality)]

Appendix 5. CINAHL search strategy

We searched CINAHL 1982 to September 2009 using the phrase:

1. exp Personality Disorders/
2. exp Antisocial Personality Disorder/
3. exp Borderline Personality Disorder/
4. exp Compulsive Personality Disorder/
5. exp Dependent Personality Disorder/
6. exp Impulse Control Disorders/
7. exp Passive-Aggressive Personality Disorder/
8. (histrionic\$ adj2 person\$).tw.
9. (parano\$ adj2 person\$).tw.
10. (schizo\$ adj3 person\$).tw.
11. ((asocial\$ or antisocial\$ or dissocial\$ or psychopath\$ or sadist\$ or sociopath\$) adj2 person\$).tw.
12. psychopath.tw.
13. sociopath.tw.
14. (moral adj2 insanity).tw.
15. dyssocial.tw.

16. (DSM and (Axis and II)).tw.
17. or/1-16
18. randomi\$.mp. [mp=title, subject heading word, abstract, instrumentation]
19. clin\$.mp. [mp=title, subject heading word, abstract, instrumentation]
20. trial\$.mp. [mp=title, subject heading word, abstract, instrumentation]
21. (clin\$ adj3 trial\$).mp. [mp=title, subject heading word, abstract, instrumentation]
22. singl\$.mp. [mp=title, subject heading word, abstract, instrumentation]
23. doubl\$.mp. [mp=title, subject heading word, abstract, instrumentation]
24. tripl\$.mp. [mp=title, subject heading word, abstract, instrumentation]
25. trebl\$.mp. [mp=title, subject heading word, abstract, instrumentation]
26. mask\$.mp. [mp=title, subject heading word, abstract, instrumentation]
27. blind\$.mp. [mp=title, subject heading word, abstract, instrumentation]
28. (22 or 23 or 24 or 25) and (26 or 27)
29. crossover.mp. [mp=title, subject heading word, abstract, instrumentation]
30. random\$.mp. [mp=title, subject heading word, abstract, instrumentation]
31. allocate\$.mp. [mp=title, subject heading word, abstract, instrumentation]
32. assign\$.mp. [mp=title, subject heading word, abstract, instrumentation]
33. (random\$ adj3 (allocate\$ or assign\$)).mp.
34. Random Assignment/
35. exp Clinical Trials/
36. exp Meta Analysis/
37. 33 or 29 or 28 or 21 or 18 or 34 or 35 or 36
38. 17 and 37

Appendix 6. DISSERTATION ABSTRACTS INTERNATIONAL search strategy

We searched Dissertation Abstracts International using the term “Borderline Personality Disorder” (1861 to September 2009) and then read through the titles of the dissertations that were found.

Appendix 7. EMBASE search strategy

We searched EMBASE (1980 to 37th week 2009) using the phrase:

1. exp Personality Disorder/
2. exp Borderline State/

3. exp Character Disorder/
4. exp Compulsive Personality Disorder/
5. exp DELUSION/
6. exp Dependent Personality Disorder/
7. exp DEPERSONALIZATION/
8. exp JEALOUSY/
9. exp KLEPTOMANIA/
10. exp Multiple Personality/
11. exp NARCISSISM/
12. exp PSYCHOPATHY/
13. exp SCHIZOIDISM/
14. exp SOCIOPATHY/
15. (antisoci\$ adj2 person\$).tw.
16. (aggres\$ adj2 person\$).tw.
17. (border\$ adj2 person\$).tw.
18. histrion\$ person\$.tw.
19. paranoid person\$.tw.
20. (passive adj2 aggressive).tw.
21. ((asocial\$ or antisocial\$ or dissocial\$ or psychopath\$ or sadist\$ or sociopath\$) adj person\$).tw.
22. (moral adj2 insan\$).tw.
23. dyssocial.tw.
24. (DSM and (Axis and II)).tw.
25. or/1-24
26. clin\$.tw.
27. trial\$.tw.
28. (clin\$ adj3 trial\$).tw.
29. singl\$.tw.
30. doubl\$.tw.
31. trebl\$.tw.
32. tripl\$.tw.
33. blind\$.tw.

34. mask\$.tw.
35. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj3 (blind\$ or mask\$)).tw.
36. randomi\$.tw.
37. random\$.tw.
38. allocat\$.tw.
39. assign\$.tw.
40. (random\$ adj3 (allocat\$ or assign\$)).tw.
41. crossover.tw.
42. 41 or 40 or 36 or 35 or 28
43. exp Randomized Controlled Trial/
44. exp Double Blind Procedure/
45. exp Crossover Procedure/
46. exp Single Blind Procedure/
47. exp RANDOMIZATION/
48. 43 or 44 or 45 or 46 or 47 or 42
49. 25 and 48

Appendix 8. MEDLINE search strategy

We searched Medline 1966 to 11 September 2009 using the phrase:

1. exp Personality Disorders/
2. exp Antisocial Personality Disorder/
3. exp Borderline Personality Disorder/
4. exp Compulsive Personality Disorder/
5. exp Dependent Personality Disorder/
6. exp Histrionic Personality Disorder/
7. exp Hysteria/
8. exp Paranoid Personality Disorder/
9. exp Passive-Aggressive Personality Disorder/
10. exp Schizoid Personality Disorder/
11. exp Schizotypal Personality Disorder/
12. ((asocial\$ or antisocial\$ or dissocial\$ or psychopath\$ or sadist\$ or sociopath\$) adj2 person\$).tw.

13. psychopath.tw.
14. sociopath\$.tw.
15. (moral adj2 insanity).tw.
16. (DSM and (axis and II)).tw.
17. or/1-16
18. randomized controlled trial.pt.
19. controlled clinical trial.pt.
20. randomized controlled trials.sh.
21. random allocation.sh.
22. double blind method.sh.
23. single-blind method.sh.
24. or/18-23
25. (animal not human).sh.
26. 24 not 25
27. clinical trial.pt.
28. exp clinical trials/
29. (clin\$ adj25 trial\$).ti,ab.
30. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj25 (blind\$ or mask\$)).ti,ab.
31. Placebos.sh.
32. placebo\$.ti,ab.
33. random\$.ti,ab.
34. research design.sh.
35. or/27-34
36. 35 not 25
37. 36 not 26
38. comparative study.sh.
39. exp evaluation studies/
40. follow up studies.sh.
41. prospective studies.sh.
42. (control\$ or prospectiv\$ or volunteer\$).ti,ab.
43. or/38-42
44. 43 not 25

45. 44 not (26 or 37)

46. 26 or 37 or 45

47. 17 and 46

Appendix 9. NATIONAL CRIMINAL JUSTICE REFERENCE SERVICE ABSTRACTS search strategy

We searched NCJRS 1970 to July 2008 using the phrase:

(randomi* OR double blind) and (antisocial* OR asocial* OR avoidant OR borderline OR dependent OR depressive OR dissocial OR dissocial* OR histrionic OR moral OR multiimpulsive OR multiple* OR narcissistic OR parano* OR passiveaggressive OR psychopath* OR sadis* OR schizotypal OR self defeating OR sociopath*)

Appendix 10. PsycINFO search strategy

We searched PsycINFO 1872 to 2nd week September 2009 using the phrase:

1. Personality Disorders/
2. exp Antisocial Personality Disorder/
3. exp Avoidant Personality Disorder/
4. exp Borderline Personality Disorder/
5. exp Dependent Personality Disorder/
6. exp Histrionic Personality Disorder/
7. exp Narcissistic Personality Disorder/
8. exp Obsessive Compulsive Personality Disorder/
9. exp Paranoid Personality Disorder/
10. exp Passive Aggressive Personality Disorder/
11. exp Sadomasochistic Personality/
12. exp Schizoid Personality Disorder/
13. exp Schizotypal Personality Disorder/
14. (personality adj disorders).tw.
15. (antisocial adj personality).tw.
16. (avoidant adj personality).tw.
17. (borderline adj personality).tw.
18. (dependent adj personality).tw.
19. (histrionic adj (personality and disorder)).tw.

20. (narcissistic adj personality).tw.
21. (obsessive adj (compulsive and personality)).tw.
22. (paranoid adj personality).tw.
23. (passive adj (aggressive and personality)).tw.
24. (somasochistic adj personality).tw.
25. (schizoid adj personality).tw.
26. (schizotypal adj personality).tw.
27. or/1-26
28. randomi\$.tw.
29. singl\$.tw.
30. doubl\$.tw.
31. trebl\$.tw.
32. tripl\$.tw.
33. blind\$.tw.
34. mask\$.tw.
35. (or/29-32) adj3 (or/33-34)
36. clin\$.tw.
37. trial\$.tw.
38. (clin\$ adj3 trial\$).tw.
39. placebo\$.tw.
40. exp PLACEBO/
41. crossover.tw.
42. exp Treatment Effectiveness Evaluation/
43. exp Mental Health Program Evaluation/
44. random\$.tw.
45. assign\$.tw.
46. allocate\$.tw.
47. (random\$ adj3 (assign\$ or allocate\$)).tw.
48. 27 or 35 or 38 or 39 or 40 or 41 or 42 or 43 or 47
49. 27 and 48

Appendix 11. SIGLE search strategy

We searched SIGLE 1980 to April 2006 using the phrase: ((randomisation) OR (randomised) OR (randomisee) OR (randomises) OR (randomize) OR (randomized) OR (randomly) OR ((double AND blind) OR double-blind OR double* blind* OR randomi?ed controlled trials)) AND ((psychopath* OR sociopath* OR dissocial OR sadis* OR schizotypal OR selfdefeating OR borderline OR avoidant OR dependent OR depressive OR histrionic OR multi-impulsive OR multiple OR narcissistic OR passive-aggressive) AND (person*) OR (antisocial AND behaviour) OR (personality AND disorder*) OR (gender AND identity) OR (parano* AND person*) OR (self AND defeating) OR ((asocial* OR antisocial* OR dissocial* OR psychopath* OR sadist* OR sociopath*) AND person*) OR (moral AND insanity))

Appendix 12. SOCIOLOGICAL ABSTRACTS search strategy

We searched SOCIOLOGICAL ABSTRACTS 1963 to September 2009 using the phrase: (((personality near disorder*) or ((antisocial* near disorder*) or (avoidant* near disorder*) or (borderline* near disorder*)) or ((dependent* near disorder*) or (histrionic* near disorder*)) or (narcissistic* near disorder*)) or ((obsessive* near disorder*) or (compulsive* near disorder*) or (paranoid* near disorder*)) or (((passive* near disorder*) or (aggress* near disorder*) or (somasochistic* near disorder*)) or (schizo* near disorder*)) or (((passive* and disorder*) or (aggress* and disorder*) or (somasochistic* and disorder*)) or ((schizo* and disorder*) or (paranoid* and disorder*) or (compulsive* and disorder*)) or ((obsessive* and disorder*) or (narcissistic and disorder*) or (histrionic* and disorder*))) or (((personality and disorder*) or (antisocial* and disorder*) or (avoidant* and disorder*)) or (borderline* and disorder*)) and ((AB=randomi* or TI=randomi*) or (DE=(randomi?ed controlled trials) or AB=(double* blind*) or TI=(double* blind*)) or (DE=(double blind studies) or (single* near blind*)))

Appendix 13. WEB OF SCIENCE search strategy

We searched the Web of Science 1981 to 12 September 2009 using the phrase: (double blind OR randomi*) AND ((passive-aggressive OR psychopath* OR sociopath* OR dissocial OR sadis* OR schizotypal OR self-defeating OR borderline OR avoidant OR dependent OR depressive OR parano* OR asocial* OR antisocial* OR dissocial* OR psychopath* OR sadist* OR sociopath* OR histrionic OR multi-impulsive OR multiple* OR narcissistic) AND personality*) OR ((moral AND insanity) OR (self AND defeating) OR (gender AND identity) OR (personality AND disorder) OR (antisocial AND behaviour))

Appendix 14. Summary of findings: haloperidol vs. placebo

Haloperidol for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: haloperidol						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Haloperidol				
BPD severity Borderline Syndrome Index (BSI) Follow-up: mean 5 weeks	The mean BPD severity in the control groups was 20.08 ¹ points	The mean BPD severity in the intervention groups was 0.30 standard deviations higher (0.22 lower to 0.82 higher)		58 (1 study)	++OO low ^{2,3}	
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems outcome was measured on different scales in different studies Follow-up: mean 5 weeks	The mean interpersonal problems in the control groups was 0	The mean interpersonal problems in the intervention groups was 0.37 standard deviations lower (0.74 lower to 0 higher)		114 (2 studies)	++OO low ²⁴	
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity Barrett Impulsiveness Scale (BIS) Follow-up: mean 5 weeks	The mean impulsivity ranged across control groups from 103.06 to 237.74 ¹ points	The mean impulsivity in the intervention groups was 0.07 standard deviations higher (0.3 lower to 0.43 higher)		114 (2 studies)	++OO low ^{2,4}	
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Haloperidol for borderline personality disorder**Patient or population:** patients with borderline personality disorder**Settings:****Intervention:** haloperidol

Anger SCL-90-R-hostility. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean anger ranged across control groups from 0.97 to 1.05 points	The mean anger in the intervention groups was 0.46 standard deviations lower (0.84 to 0.09 lower)	114 (2 studies)	++OO low ^{2,4}
Psychotic symptoms SCL-90-paranoid ideation. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean psychotic symptoms ranged across control groups from 0.98 to 1.00 points	The mean psychotic symptoms in the intervention groups was 0.44 standard deviations lower (1.09 lower to 0.2 higher)	114 (2 studies)	++OO low ²⁴
Dissociation - not measured	See comment	See comment	Not estimable	- See comment Not measured
Depression BDI. Scale from: 0 to 63. Follow-up: mean 5 weeks	The mean depression ranged across control groups from 13.04 to 14.88 points	The mean depression in the intervention groups was 0.09 standard deviations lower (0.87 lower to 0.68 higher)	114 (2 studies)	++OO low ²⁴
Anxiety SCL-90-R-anxiety. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean anxiety ranged across control groups from 1.00 to 1.07 points	The mean anxiety in the intervention groups was 0.06 standard deviations higher (0.68 lower to 0.79 higher)	114 (2 studies)	++OO low ^{2,4}
General psychiatric pathology SCL-90-R global severity index. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean general psychiatric pathology ranged across control groups from 0.84 to 0.87 points	The mean general psychiatric pathology in the intervention groups was 0.08 standard deviations lower (0.71 lower to 0.54 higher)	114 (2 studies)	++OO low ^{2,4}
Mental health status GAS. Scale from: 0 to 100. Follow-up: mean 5 weeks	The mean mental health status ranged across control groups from 48.16 to 58.43 points	The mean mental health status in the intervention groups was 0.16 standard deviations higher (0.77 lower to 1.08 higher)	114 (2 studies)	++OO low ^{2,4}
Attrition lost after randomisation Follow-up: mean 5 weeks	111 per 1000	128 per 1000 (50 to 324)	RR 1.15 (0.45 to 2.92)	130 (2 studies) ++OO low ^{2,4}

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

⁴ Two study effects only, publication bias cannot be excluded

Appendix 15. Summary of findings: thiothixene vs. placebo

Thiothixene for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: Thiothixene						
Outcomes	Illustrative comparative risks [*] (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Haloperidol				
BPD severity Schedule of Interviewing Schizotypal Personalities, borderline score Follow-up: mean 12 weeks	The mean BPD severity in the control groups was 4.1812₁ points	The mean BPD severity in the intervention groups was 0.28 standard deviations higher (0.28 lower to 0.83 higher) ²		50 (1 study)	+000 _{3,4,5} very low	
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems HSCL-90-interpersonal sensitivity. Scale from: 0 to 4. Follow-up: mean 12 weeks	The mean interpersonal problems in the control groups was 0.6209 points₁	The mean interpersonal problems in the intervention groups was 0.18 standard deviations lower (0.74 lower to 0.37 higher) ²		50 (1 study)	+000 _{3,4,5} very low	
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Thiothixene for borderline personality disorder**Patient or population: patients with borderline personality disorder****Settings:****Intervention: Thiothixene**

Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger SCL-90-hostility. Scale from: 0 to 4. Follow-up: mean 12 weeks	The mean anger in the control groups was 0.6769¹ points	The mean anger in the intervention groups was 0.07 standard deviations lower (0.63 lower to 0.48 higher) ²		50 (1 study)	+000 very low ^{3,4,5}	
Psychotic symptoms Schedule of Interviewing Schizotypal Personalities (SIB)-suspicious/paranoid score Follow-up: mean 5 weeks	The mean psychotic symptoms in the control groups was 1.8654¹ points	The mean psychotic symptoms in the intervention groups was 0.19 standard deviations higher (0.37 lower to 0.75 higher) ²		50 (1 study)	+000 very low ^{3,4,5}	
Dissociation not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression SCL-90-depression. Scale from: 0 to 4. Follow-up: mean 12 weeks	The mean depression in the control groups was 1.5065¹ points	The mean depression in the intervention groups was 0.12 standard deviations higher (0.43 lower to 0.68 higher) ²		50 (1 study) ^{3,4,5}	+000 very low ^{3,4,5}	
Anxiety - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
General psychiatric pathology - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Mental health status GAS. Scale from: 0 to 100. Follow-up: mean 12 weeks	The mean mental health status in the control groups was 71.92¹ points	The mean mental health status in the intervention groups was 0.06 standard deviations higher (0.5 lower to 0.61 higher) ²		50 (1 study)	+000 very very low ^{3,4,5}	
Attrition attrition Follow-up: mean 12 weeks	115 per 1000	291 per 1000 (85 to 998)	RR2.53 (0.74 to 8.68)	50 (1 study)	+000 low ^{3,5}	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.**Moderate quality:** Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.**Low quality:** Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.**Very low quality:** We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls² SMD calculated on basis of post mean and pre mean SD, might overestimate the treatment effect³ Small sample size

⁴ Effect estimate had to be calculated by using the SD of the pre-treatment means. This may have led to an overestimation of effect sizes, as the pre-SDs are commonly smaller than post-SDs.

⁵ Single study effect only, publication bias cannot be excluded

Appendix 16. Summary of findings: flupenthixol decanoate vs. placebo

Haloperidol for borderline personality disorder					
Patient or population: patients with borderline personality disorder					
Settings:					
Intervention: flupenthixol decanoate					
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)
	Assumed risk	Corresponding risk			
	Control	Flupenthixol decanoate			
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment
Interpersonal problems - not measured	See comment	See comment	Not estimable	-	See comment
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment
Impulsivity - not measured	See comment	See comment	Not estimable	-	See comment
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment
Suicidal behaviour suicidal act during treatment Follow-up: mean 6 months	789 per 1000	387 per 1000 (205 to 726)	RR 0.49 (0.26 to 0.92)	37 (1 study)	++OO low ^{1,2}
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment
Anger - not measured	See comment	See comment	Not estimable	-	See comment
Psychotic symptoms - not measured	See comment	See comment	Not estimable	-	See comment
Dissociation - not measured	See comment	See comment	Not estimable	-	See comment

Haloperidol for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: flupenthixol decanoate						
Depression - not measured	See comment	See comment	Not estimable	-		See comment
Anxiety - not measured	See comment	See comment	Not estimable	-		See comment
General psychiatric pathology - not measured	See comment	See comment	Not estimable	-		See comment
Mental health status - not measured	See comment	See comment	Not estimable			See comment
Attrition lost after randomisation Follow-up: mean 6 months	158 per 1000	223 per 1000 (57 to 858)	RR 1.41 (0.36 to 5.43)	37 (1 study)	++oo low ^{1,2}	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Small sample size

² Single study effect only, publication bias cannot be excluded

Appendix 17. Summary of findings: aripiprazole vs. placebo

Aripiprazole for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: aripiprazole						
Outcomes	Illustrative comparative risks [*] (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	aripiprazole				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	

Aripiprazole for borderline personality disorder**Patient or population:** patients with borderline personality disorder**Settings:****Intervention:** aripiprazole

Interpersonal problems SCL-90-R-in-terpersonal sensitivity Follow-up: mean 8 weeks	The mean interpersonal problems in the control groups was 64.2 points (Z-transformed score)	The mean inter-personal problems in the intervention groups was 0.77 standard deviations lower (1.33 to 0.2 lower)	52 (1 study)	++OO low ^{2,3}	
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment
Impulsivity STAXI-anger out Follow-up: mean 8 weeks	The mean impulsivity in the control groups was 20.7 points (Z-transformed score)	The mean impulsivity in the intervention groups was 1.84 standard deviations lower (2.49 to 1.18 lower)	52 (1 study)	++OO low ^{2,3}	
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment
Self-mutilating behaviour self-injury during treatment Follow-up: mean 8 weeks	269 per 1000	78 per 1000 (19 to 336)	RR 0.29 (0.07 to 1.25)	52 (1 study)	++OO low ^{2,3}
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment
Anger SCL-90-R-hostility Follow-up: mean 8 weeks	The mean anger in the control groups was 73.1 points (Z-transformed score)	The mean anger in the intervention groups was 1.14 standard deviations lower (1.73 to 0.55 lower)	52 (1 study)	++OO low ^{2,3}	
Psychotic symptoms SCL-90-R-paranoid ideation Follow-up: mean 8 weeks	The mean psychotic symptoms in the control groups was 68.3 points (Z-transformed score)	The mean psychotic symptoms in the intervention groups was 1.05 standard deviations lower (1.64 to 0.47 lower)	52 (1 study)	++OO low ^{2,3}	
Dissociation not measured	See comment	See comment	Not estimable	-	See comment
Depression Ham-D. Scale from: 0 to 50. Follow-up: mean 8 weeks	The mean depression in the control groups was 18.8 points	The mean depression in the intervention groups was 1.25 standard deviations lower (1.85 to 0.65 lower)	52 (1 study)	++OO low ^{2,3}	

Aripiprazole for borderline personality disorder

Patient or population: patients with borderline personality disorder

Settings:

Intervention: aripiprazole

Anxiety HARS. Scale from: 0 to 56. Follow-up: mean 8 weeks	The mean anxiety in the control groups was 19.5 points ¹	The mean anxiety in the intervention groups was 0.73 standard deviations lower (1.29 to 0.17 lower)	52 (1 study)	++OO low ^{2,3}
General psychiatric pathology SCL-90-R-global severity index Follow-up: mean 8 weeks	The mean general psychiatric pathology in the control groups was 69.4 points (Z-transformed score) ¹	The mean general psychiatric pathology in the intervention groups was 1.27 standard deviations lower (1.87 to 0.67 lower)	52 (1 study)	++OO low ^{2,3}
Mental health status - not measured	See comment	See comment	Not estimable	- See comment
Attrition - not reported	See comment	See comment	Not estimable	- See comment

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

Appendix 18. Summary of findings: olanzapine vs. placebo

olanzapine compared to for borderline personality disorder

Patient or population: patients with borderline personality disorder

Settings:

Intervention: olanzapine

Comparison:

Outcomes	Illustrative comparative risks [*] (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Olanzapine				
BPD severity Zan-BPD-total score.	The mean BPD severity ranged across	The mean BPD severity in the intervention		596 (2 studies)	++++ high	

olanzapine compared to for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: olanzapine						
Comparison:						
Scale from: 0 to 36. Follow-up: mean 12 weeks	control groups from -6.19 to -6.69 points ¹	groups was 0.15 standard deviations lower (0.41 lower to 0.1 higher) ²				
Avoidance of abandonment outcome was measured on different scales in different studies. Follow-up: mean 12 weeks		The mean avoidance of abandonment in the intervention groups was 0.01 standard deviations lower (0.22 lower to 0.21 higher) ²	631 (3 studies)	++++ high		
Interpersonal problems I outcome was measured on different scales in different studies. Follow-up: mean 12 weeks		The mean interpersonal problems I in the intervention groups was 0.10 standard deviations lower (0.31 lower to 0.12 higher) ²	340 (2 studies)	+++O moderate ³		
Interpersonal problems II Zan-BPD-unstable interpersonal relationships. Scale from: 0 to 4. Follow-up: mean 12 weeks	See comment 1	See comment	291 (1 study)	+++O moderate ³		Mean baseline change difference between groups was 0.2 (greater change in controls, 95% CI -0.31 to 0.12)
Identity disturbance outcome was measured on different scales in different studies. Follow-up: mean 12 weeks		The mean identity disturbance in the intervention groups was 0.06 standard deviations lower (0.21 lower to 0.1 higher) ²	631 (3 studies)	++++ high		
Impulsivity I behavioural reports of numbers of episodes of impulsivity/aggressive behaviour. Follow-up: mean 12 weeks	The mean impulsivity I in the control groups was 6.12 points ⁴	The mean impulsivity I in the intervention groups was 0.04 standard deviations lower (0.54 lower to 0.47 higher)	60 (1 study)	+++O moderate ³		
Impulsivity II	See comment	See comment	340 (2 studies)	+++O moderate ³		Mean baseline change

olanzapine compared to for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: olanzapine						
Comparison:						
outcome was measured on different scales in different studies Follow-up: mean 12 weeks						difference between groups was $\text{MD} = -0.18$ (95% CI $\text{MD} = -0.40$ to 0.03)
Impulsivity III Zan-BPD-impulsivity that are self-damaging. Scale from: 0 to 4. Follow-up: median 12 weeks	See comment ¹	See comment		291 (1 study)	+++O moderate ³	Mean baseline change difference between groups was $\text{MD} = -0.10$ (95% CI $\text{MD} = -0.40$ to 0.20)
Suicidal ideation I outcome was measured on different scales in different studies Follow-up: mean 12 weeks		The mean suicidal ideation I in the intervention groups was 0.29 standard deviations higher (0.07 to 0.5 higher) ²		340 (2 studies)	+++O moderate ³	
Suicidal ideation II high suicidality according to OAS-M Follow-up: mean 21 weeks	417 per 1000	500 per 1000 (209 to 1000)	RR 1.20 (0.5 to 2.88)	24 (1 study)	+++O moderate ³	
Suicidal behaviour I behavioural reports of numbers of episodes of self-injuring behaviour/ suicide attempts Follow-up: mean 12 weeks	The mean suicidal behaviour i in the control groups was 0.88 points ⁴	The mean suicidal behaviour I in the intervention groups was 0.15 standard deviations higher (0.36 lower to 0.65 higher)		60 (1 study)	+++O moderate ³	
Suicidal behaviour II Zan-BPD-suicidal or self-mutilating behaviour. Scale from: 0 to 4. Follow-up: mean 12 weeks	See comment ¹	See comment		291 (1 study)	+++O moderate ³	Mean baseline change difference between groups was -0.2 (95% CI -0.20 to 0.00)
Suicidal behaviour III	417 per 1000	500 per 1000 (209 to 1000)	RR 1.20 (0.5 to 2.88)	24 (1 study)	+++O moderate ³	

olanzapine compared to for borderline personality disorder

Patient or population: patients with borderline personality disorder**Settings:****Intervention:** olanzapine**Comparison:**

 high
 suicidality
 according to
 OAS-M
 Follow-up:
 mean 21
 weeks

Affective instability outcome was measured on different scales in different studies Follow-up: mean 12 weeks	The mean affective instability in the intervention groups was 0.16 standard deviations lower (0.32 to 0.01 lower) ²	631 (3 studies)	++++	high	
Chronic feelings of emptiness outcome was measured on different scales in different studies Follow-up: mean 12 weeks	The mean chronic feelings of emptiness in the intervention groups was 0.03 standard deviations lower (0.22 lower to 0.16 higher) ²	631 (3 studies)	++++	high	
Anger outcome was measured on different scales in different studies Follow-up: mean 12 weeks	The mean anger in the intervention groups was 0.27 standard deviations lower (0.43 to 0.12 lower) ²	631 (3 studies)	++++	high	
Psychotic symptoms outcome was measured on different scales in different studies Follow-up: mean 12 weeks	The mean psychotic symptoms in the intervention groups was 0.18 standard deviations lower (0.34 to 0.03 lower) ²	631 (3 studies)	++++	high	
Dissociation - not measured	See comment	See comment	Not estimable	-	See comment Not measured
Depression I Ham-D. Scale from: 0 to 50. Follow-up: mean 12 weeks	The mean depression I ranged across control groups from 15.4 to 15.8 points ⁴	The mean depression I in the intervention groups was 0.37 standard deviations lower (0.8 lower to 0.07 higher)	84 (2 studies)	+++O	moderate ³
Depression II	See comment ¹	See comment	596 (2 studies)	++++	high Mean baseline

olanzapine compared to for borderline personality disorder**Patient or population:** patients with borderline personality disorder**Settings:****Intervention:** olanzapine**Comparison:**

MADRS. Scale from: 0 to 60. Follow-up: mean 12 weeks						change difference was 0.39 (95% CI -0.20 to 0.97)
Anxiety I HARS. Scale from: 0 to 56. Follow-up: mean 12 weeks	The mean anxiety I in the control groups was 19.93 points ⁴	The mean anxiety I in the intervention groups was 0. 23 standard deviations lower (0.74 lower to 0. 28 higher)	60 (1 study)	+++O moderate ³		
Anxiety II SCL-90-R-anxiety. Scale from: 0 to 4. Follow-up: mean 12 weeks	See comment ¹	See comment	274 (1 study)	+++O moderate ³		Mean baseline change difference was -0.22 (95% CI -0.41 to -0.03)
General psychiatric pathology SCL-90-R-global severity index. Scale from: 0 to 4. Follow-up: mean 12 weeks	The mean general psychiatric pathology ranged across control groups from -0.53 to -0.56 points ¹	The mean general psychiatric pathology in the intervention groups was 0. 21 standard deviations lower (0.53 lower to 0. 1 higher) ¹	557 (2 studies)	++++ high		
Mental health status I CGI-S. Scale from: 0 to 7. Follow-up: mean 12 weeks	The mean mental health status I in the control groups was 3.97 points ⁴	The mean mental health status I in the intervention groups was 0. 03 standard deviations lower (0.53 lower to 0. 48 higher)	60 (1 study)	+++O moderate ³		
Mental health status II GAF. Scale from: 0 to 100. Follow-up: mean 12 weeks	See comment ¹	See comment	596 (2 studies)	++++ high		Mean baseline change difference was 1.52 (95% CI -0.75 to 3.79)
Attrition lost after randomisation Follow-up: mean 16 weeks	389 per 1000	377 per 1000 (280 to 502)	RR 0.97 (0.72 to 1.29)	767 (6 studies)	++++ high	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.**Moderate quality:** Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Change value in controls

² SMD on basis of change scores

³ Small sample size

⁴ Final value in controls

Appendix 19. Summary of findings: ziprasidone vs. placebo

Ziprasidone for borderline personality disorder					
Patient or population: patients with borderline personality disorder					
Settings:					
Intervention: haloperidol					
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)
	Assumed risk	Ziprasidone			
	Control	Haloperidol			
BPD severity Clinical Global Impression scale for use in borderline patients-global. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean BPD severity in the control groups was 4.3 points ¹	The mean BPD severity in the intervention groups was 0.47 standard deviations lower (0.98 lower to 0.05 higher)		60 (1 study)	+OOO _{2,3,4} very low
Avoidance of abandonment CGI-BPD-abandonment. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean avoidance of abandonment in the control groups was 4.53 points ¹	The mean avoidance of abandonment in the intervention groups was 0.08 standard deviations lower (0.58 lower to 0.43 higher)		60 (1 study)	+OOO _{2,3,4} very low
Interpersonal problems CGI-BPD- unstable relationships. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean interpersonal problems in the control groups was 4.5 points ¹	The mean interpersonal problems in the intervention groups was 0.12 standard deviations lower (0.63 lower to 0.38 higher)		60 (1 study)	+OOO _{2,3,4} very low
Identity disturbance CGI-BPD-identity disturbance. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean identity disturbance in the control groups was 5.03 points ¹	The mean identity disturbance in the intervention groups was		60 (1 study)	+OOO _{2,3,4} very low

Ziprasidone for borderline personality disorder

Patient or population: patients with borderline personality disorder

Settings:

Intervention: haloperidol

								0.38 standard deviations lower (0.9 lower to 0. 13 higher)
Impulsivity CGI-BPD-impulsivity. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean impulsivity in the control groups was ¹ 3.96 points	The mean impulsivity in the intervention groups was 0.03 standard deviations higher (0.48 lower to 0. 53 higher)		60 (1 study)		+000 ^{2,3,4} very low		
Suicidal ideation CGI-BPD- recurrent suicidal ideation. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean suicidal ideation in the control groups was ¹ 3.13 points	The mean suicidal ideation in the intervention groups was 0.27 standard deviations lower (0.78 lower to 0. 23 higher)		60 (1 study)		+000 ^{2,3,4} very low		
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-		See comment		Not measured
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-		See comment		Not measured
Affective instability CGI-BPD- affective instability. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean affective instability in the control groups was 30.30	The mean affective instability in the intervention groups was 0.10 standard deviations lower (0.61 lower to 0. 41 higher)		60 (1 study)		+000 ^{2,3,4} very very low		
Chronic feelings of emptiness CGI-BPD- chronic feelings of emptiness. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean chronic feelings of emptiness in the control groups was 4.4 points ¹	The mean chronic feelings of emptiness in the intervention groups was 0.18 standard deviations higher (0.32 lower to 0. 69 higher)		60 (1 study)		+000 ^{2,3,4} very low		
Anger CGI-BPD-inap-proprieate anger. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean anger in the control groups was ¹ 3.56 points	The mean anger in the intervention groups was 0.08 standard		60 (1 study)		+000 ^{2,3,4} very low		

Ziprasidone for borderline personality disorder**Patient or population:** patients with borderline personality disorder**Settings:****Intervention:** haloperidol

							deviations higher (0.43 lower to 0.58 higher)
Psychotic symptoms CGI-BPD-transient paranoia or dissociation. Scale from: 1 to 7. Follow-up: mean 12 weeks	The mean psychotic symptoms in the control groups was 2.23 points ¹	The mean psychotic symptoms in the intervention groups was 0.23 standard deviations lower (0.74 lower to 0.28 higher)		60 (1 study)	+000, very low ^{2,3,4}		
Dissociation not measured	See comment	See comment	Not estimable	-	See comment	Not measured	
Depression Ham-D. Scale from: 0 to 50. Follow-up: mean 12 weeks	The mean depression in the control groups was 16.07 points ¹	The mean depression in the intervention groups was 0.30 standard deviations lower (0.81 lower to 0.21 higher)		60 (1 study)	+000, very low ^{2,3,4}		
Anxiety HARS. Scale from: 0 to 56. Follow-up: mean 12 weeks	The mean anxiety in the control groups was 16.53 points ¹	The mean anxiety in the intervention groups was 0.12 standard deviations lower (0.63 lower to 0.39 higher)		60 (1 study)	+000, very low ^{2,3,4}		
General psychiatric pathology SCL-90-R-GSI. Scale from: 0 to 4. Follow-up: mean 12 weeks	The mean general psychiatric pathology in the control groups was 2.39 points ¹	The mean general psychiatric pathology in the intervention groups was 0.41 standard deviations lower (0.92 lower to 0.1 higher)		60 (1 study)	+000, very low ^{2,3,4}		
Mental health status - not measured	See comment	See comment	Not estimable	-	See comment	Not measured	
Attrition lost after randomisation Follow-up: mean 12 weeks	467 per 1000	565 per 1000 (346 to 929)	RR 1.21 (0.74 to 1.99)	60 (1 study)	+000 low ^{3,4}		

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls

² Very high attrition (>50%), unclear how participants were selected from eligible patients

³ Small sample size

⁴ Single study effect only, publication bias cannot be excluded

Appendix 20. Summary of findings: carbamazepine vs. placebo

Carbamazepine for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: carbamazepine						
Outcomes	Illustrative comparative risks [*] (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Carbamazepine				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems SCL-90- interpersonal sensitivity. Scale from: 0 to 4. Follow-up: mean 32 days	The mean interpersonal problems in the control groups was 11.0 points ¹	The mean interpersonal problems in the intervention groups was 0.54 standard deviations lower (1.46 lower to 0.38 higher)		19 (1 study)	++OO low ^{2,3}	
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity status quo or worsened after treatment according to Acting out scale Follow-up: mean 32 days	800 per 1000	704 per 1000 (424 to 1000)	RR 0.88 (0.53 to 1.46)	20 (1 study)	++OO low ^{2,3}	
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Carbamazepine for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: carbamazepine						
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger SCL-90-hostility Follow-up: mean 32 days	The mean anger in the control groups was 8.33 points ¹	The mean anger in the intervention groups was 0.34 standard deviations lower (1.25 lower to 0.57 higher)		19 (1 study)	++OO low ^{2,3}	
Psychotic symptoms SCL-90-paranoid ideation Follow-up: mean 32 days	The mean psychotic symptoms in the control groups was 9.22 points ¹	The mean psychotic symptoms in the intervention groups was 0.58 standard deviations lower (1.5 lower to 0.35 higher)		19 (1 study)	++OO low ^{2,3}	
Dissociation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression SCL-90-depression Follow-up: mean 32 days	The mean depression in the control groups was 22.6 points ¹	The mean depression in the intervention groups was 0.66 standard deviations lower (1.59 lower to 0.27 higher)		19 (1 study)	++OO low ^{2,3}	
Anxiety SCL-90-anxiety Follow-up: mean 32 days	The mean anxiety in the control groups was 17.44 points ¹	The mean anxiety in the intervention groups was 0.51 standard deviations lower (1.43 lower to 0.41 higher)		19 (1 study)	++OO low ^{2,3}	
General psychiatric pathology SCL-90-GSI Follow-up: mean 32 days	The mean general psychiatric pathology in the control groups was 125.11 points ¹	The mean general psychiatric pathology in the intervention groups was 0.57 standard deviations lower (1.49 lower to 0.36 higher)		19 (1 study)	++OO low ^{2,3}	
Mental health status GAS. Scale from: 0 to 100. Follow-up: mean 32 days	The mean mental health status in the control groups was 60.1 points ¹	The mean mental health status in the intervention groups was 0.34 standard deviations higher (0.57 lower to 1.25 higher)		19 (1 study)	++OO low ^{2,3}	

Carbamazepine for borderline personality disorder

Patient or population: patients with borderline personality disorder

Settings:

Intervention: carbamazepine

Attrition lost after randomisation Follow-up: mean 32 days	RR 5 (0.27 to 92.62)	20 (1 study)	++oo low ^{2,3}
---	--------------------------------	--------------	--------------------------------

CI: Confidence interval; **RR:** Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

Appendix 21. Summary of findings: valproate semisodium vs. placebo

Valproate semisodium for borderline personality disorder

Patient or population: patients with borderline personality disorder

Settings:

Intervention: valproate semisodium

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Valproate semisodium				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems SCL-90-R-interpersonal sensitivity. Scale from: 0 to 4. Follow-up: mean 24 weeks	The mean interpersonal problems in the control groups was 2.2 points ¹	The mean interpersonal problems in the intervention groups was 1.04 standard deviations lower (1.85 to 0.23 lower)		30 (1 study)	++oo low ^{2,3}	
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Valproate semisodium for borderline personality disorder**Patient or population:** patients with borderline personality disorder**Settings:****Intervention:** valproate semisodium

Impulsivity outcome was measured on different scales in different studies Follow-up: 10 to 24 weeks		The mean impulsivity in the intervention groups was 0.62 standard deviations lower (1.48 lower to 0.24 higher)	46 (2 studies)	++OO low ^{2,4}		
Suicidal ideation Overt Aggression Scale-Modified (OAS-M)-suicidality Follow-up: mean 10 weeks	The mean suicidal ideation in the control groups was 0.3 points	The mean suicidal ideation in the intervention groups was 0.52 standard deviations higher (0.63 lower to 1.67 higher)	16 (1 study)	+OOO very low ^{2,35}		
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger I OAS-M-irritability. Scale from: 0 to 10. Follow-up: mean 10 weeks	See comment	See comment	Not estimable	16 (1 study)	+OOO very low ^{2,35}	
Anger II SCL-90-R-hostility. Scale from: 0 to 4. Follow-up: mean 24 weeks	See comment	See comment	Not estimable	30 (1 study)	++OO low ^{2,3}	
Psychotic symptoms - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Dissociation not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression outcome was measured on different scales in different studies Follow-up: 10 to 24 weeks		The mean depression in the intervention groups was 0.66 standard deviations lower (1.31 to 0.01 lower)	46 (2 studies)	++OO low ^{2,4}		
Anxiety - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
General psychiatric pathology - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Mental health status CGI-I score of 3 or more (i. e. minimally improved to very much worse) Follow-up: mean 10 weeks	1000 per 1000	640 per 1000 (370 to 1000)	RR 0.64 (0.37 to 1.11)	16 (1 study)	+OOO very low ^{2,35}	
Attrition lost after randomisation Follow-up: 10 to 24 weeks	714 per 1000	557 per 1000 (286 to 1000)	RR 0.78 (0.4 to 1.53)	46 (2 studies)	++OO low ^{2,4}	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

⁴ Two study effects only, publication bias cannot be excluded

⁵ Unclear how study participants were chosen out of eligible patients; very high attrition rate (> 50%)

Appendix 22. Summary of findings: lamotrigine vs. placebo

Lamotrigine for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: lamotrigine						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Lamotrigine				
BPD severity Zan-BPD total score. Scale from: 0 to 36. Follow-up: mean 12 weeks	The mean BPD severity in the control groups was ¹ -6.6 points	The mean BPD severity in the intervention groups was 0.43 standard deviations lower (1.2 lower to 0.34 higher)		27 (1 study)	++OO low ²³	
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity I STAXI-anger out. Scale from: 8 to 32. Follow-up: mean 8 weeks	The mean impulsivity I in the control groups was ⁴ 23.2 points	The mean impulsivity I in the intervention groups was 1.62 standard deviations lower (2.54 to 0.69 lower)		27 (1 study)	++OO low ²³	
Impulsivity II Zan-BPD-impulsivity that are self-damaging. Scale from: 0 to 4. Follow-up: mean 12 weeks	The mean impulsivity II in the control groups was ¹ -0.1 points	The mean impulsivity II in the intervention groups was 1.41 standard deviations lower (2.27 to 0.55 lower) ⁵		27 (1 study)	++OO low ²³	
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Lamotrigine for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: lamotrigine						
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Affective instability Zan-BPD-affective instability. Scale from: 0 to 4. Follow-up: mean 12 weeks	The mean affective instability in the control groups was -0.8 points ¹	The mean affective instability in the intervention groups was 0.61 standard deviations lower (1.39 lower to 0.17 higher) ⁵		27 (1 study)	++OO low ^{2,3}	
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger STAXI- trait anger. Scale from: 10 to 40. Follow-up: mean 8 weeks	The mean anger in the control groups was 27.9 points ⁴	The mean anger in the intervention groups was 1.69 standard deviations lower (2.62 to 0.75 lower)		27 (1 study)	++OO low ^{2,3}	
Psychotic symptoms - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Dissociation not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anxiety - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
General psychiatric pathology - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Mental health status - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Attrition lost after randomisation Follow-up: 8 to 12 weeks	333 per 1000	246 per 1000 (73 to 826)	RR 0.74 (0.22 to 2.48)	54 (2 studies)	++OO low ^{2,6}	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Change score in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

⁴ Final value in controls

⁵ SMD on basis of change scores

⁶Two study effects only, publication bias cannot be excluded

Appendix 23. Summary of findings: topiramate vs. placebo

Topiramate for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: topiramate						
Outcomes	Illustrative comparative risks [*] (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Topiramate				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems SCL-90-R- interpersonal sensitivity Follow-up: mean 10 weeks	The mean interpersonal problems in the control groups was 66.9 points (Z- trans-formed score) ¹	The mean inter- personal problems in the intervention groups was 0.91 standard deviations lower (1.46 to 0.35 lower)		56 (1 study)	++OO low ²³	
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity STAXI-anger- out Follow- up: mean 8 weeks	The mean impulsivity ranged across control groups from 22.6 to 24.8 points (Z- trans-formed scores) ⁴	The mean impulsivity in the intervention groups was 3.36 standard deviations lower (4.44 to 2.27 lower)		71 (2 studies)	++OO low ¹²	
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self- mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger I outcome was measured on different scales in		The mean anger I in the intervention groups was 3.00 standard deviations lower		85 (2 studies)	++OO low ¹²	

Topiramate for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: topiramate						
different studies		(3.64 to 2.36 lower)				
Follow-up: 8 to 10 weeks						
Anger II STAXI- trait anger. Scale from: 10 to 40. Follow-up: mean 8 weeks	The mean anger II in the control groups was 27.4 points	The mean anger II in the intervention groups was 0. 65 standard deviations lower (1.27 to 0.03 lower)	42 (1 study)		++OO low ²³	
Psychotic symptoms SCL-90-R-paranoid ideation Follow-up: mean 10 weeks	The mean psychotic symptoms in the control groups was 71.1 points (Z-transformed score)	The mean psychotic symptoms in the intervention groups was 0. 49 standard deviations lower (1.02 lower to 0. 05 higher)	56 (1 study)		++OO low ²³	
Dissociation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression SCL-90-R-depression Follow-up: mean 10 weeks	The mean depression in the control groups was 72.7 points (Z-transformed)	The mean depression in the intervention groups was 0. 51 standard deviations lower (1.04 lower to 0. 02 higher)	56 (1 study)		++OO low ²³	
Anxiety SCL-90-R-anxiety Follow-up: mean 10 weeks	The mean anxiety in the control groups was 69.8 points (Z-transformed score)	The mean anxiety in the intervention groups was 1. 40 standard deviations lower (1.99 to 0.81 lower)	56 (1 study)		++OO low ²³	
General psychiatric pathology SCL-90-R-global severity index Follow-up: mean 10 weeks	The mean general psychiatric pathology in the control groups was 70.1 points (Z-transformed score)	The mean general psychiatric pathology in the intervention groups was 1. 19 standard deviations lower (1.76 to 0.61 lower)	56 (1 study)		++OO low ²³	
Mental health status - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Attrition lost after randomisation Follow-up: 8 to 10 weeks	98 per 1000	54 per 1000 (14 to 212)	RR 0.55 (0.14 to 2.16)	133 (3 studies)	+++O moderate ²	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

¹ Two study effects only, publication bias cannot be excluded

² Small sample size

³ Single study effect only, publication bias cannot be excluded

⁴ Final value in controls

Appendix 24. Summary of findings: amitriptyline vs. placebo

Amitriptyline for borderline personality disorder						
Patient or population: patients with borderline personality disorder Settings: Intervention: amitriptyline						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Amitriptyline				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems SCL-90-interpersonal sensitivity. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean interpersonal problems in the control groups was 1.74 points ¹	The mean inter-personal problems in the intervention groups was 0.39 standard deviations lower (0.92 lower to 0.13 higher)		57 (1 study)	++OO low ²³	
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity Barratt Impulsiveness Scale. Scale from: 30 to 120. Follow-up: mean 5 weeks	The mean impulsivity in the control groups was 103.06 points ¹	The mean impulsivity in the intervention groups was 0.12 standard deviations lower (0.64 lower to 0.4 higher)		57 (1 study)	++OO low ²³	
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger SCL-90-hostility. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean anger in the control	The mean anger in the intervention groups was 0.26		57 (1 study)	++OO low ²³	

Amitriptyline for borderline personality disorder**Patient or population:** patients with borderline personality disorder **Settings:** Intervention: amitriptyline

	groups was 1.39 points ¹	standard deviations lower (0.78 lower to 0. 26 higher)				
Psychotic symptoms SCL-90-paranoid ideation. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean psychotic symptoms in the control groups was 1.44 points ¹	The mean psychotic symptoms in the intervention groups was 0.43 standard deviations lower (0.96 lower to 0. 09 higher)	57 (1 study)	++OO low ²³		
Dissociation not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression BDI. Scale from: 0 to 63. Follow-up: mean 5 weeks	The mean depression in the control groups was 23.04 points ¹	The mean depression in the intervention groups was 0.59 standard deviations lower (1.12 to 0.06 lower)	57 (1 study)	++OO low ²³		
Anxiety SCL- 90-anxiety. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean anxiety in the control groups was 1.54 points ¹	The mean anxiety in the intervention groups was 0.15 standard deviations lower (0.67 lower to 0. 37 higher)	57 (1 study)	++OO low ²³		
General psychiatric pathology SCL- 90-global severity index. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean general psychiatric pathology in the control groups was 1.35 points ¹	The mean general psychiatric pathology in the intervention groups was 0.34 standard deviations lower (0.87 lower to 0. 18 higher)	57 (1 study)	++OO low ²³		
Mental health status GAS. Scale from: 0 to 100. Follow-up: mean 5 weeks	The mean mental health status in the control groups was 48.16 points ¹	The mean mental health status in the intervention groups was 0.27 standard deviations higher (0.25 lower to 0. 79 higher)	57 (1 study)	++OO low ²³		
Attrition lost after randomisation Follow-up: mean 5 weeks	34 per 1000	33 per 1000 (2 to 501)	RR 0.97 (0.06 to 14.74)	59 (1 study)	++OO low ²³	

CI: Confidence interval; **RR:** Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.**Moderate quality:** Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.**Low quality:** Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.**Very low quality:** We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

Appendix 25. Summary of findings: fluoxetine vs. placebo

Fluoxetine for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: fluoxetine						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Fluoxetine				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity OAS-M- aggression Follow-up: mean 12 weeks	The mean impulsivity in the control groups was 7.45 points ¹	The mean impulsivity in the intervention groups was 0.59 standard deviations lower (1.5 lower to 0.31 higher)		20 (1 study)	+0.00 very low ^{2,3,4}	
Suicidal ideation OAS-M- suicidality Follow-up: mean 12 weeks	The mean suicidal ideation in the control groups was 1 points ¹	The mean suicidal ideation in the intervention groups was 0.44 standard deviations higher (0.46 lower to 1.33 higher)		20 (1 study)	+0.00 very low ^{2,3,4}	
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self- mutilating behaviour OAS-M- autoaggression Follow-up: mean 12 weeks	The mean self- mutilating behaviour in the control groups was 6.55 points ¹	The mean self- mutilating behaviour in the intervention groups was 0.03 standard deviations higher (0.85 lower to 0.92 higher)		20 (1 study)	+0.00 very low ^{2,3,4}	
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Fluoxetine for borderline personality disorder							
Patient or population: patients with borderline personality disorder							
Settings:							
Intervention: fluoxetine							
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured	
Anger Profile of Mood States (POMS)-anger Follow-up: mean 12 weeks	The mean anger in the control groups was 44.89 points ¹	The mean anger in the intervention groups was 0. 65 standard deviations lower (1.53 lower to 0. 22 higher)		22 (1 study)	++OO low ³⁴		
Psychotic symptoms - not measured	See comment	See comment	Not estimable	-	See comment	Not measured	
Dissociation dissociative experiences scales Follow-up: mean 12 weeks	The mean dissociation in the control groups was 12.66 points ¹	The mean dissociation in the intervention groups was 0. 42 standard deviations higher (0.47 lower to 1. 32 higher)		20 (1 study)	+OOO very low ²³⁴		
Depression BDI. Scale from: 0 to 63. Follow-up: mean 12 weeks	The mean depression ranged across control groups from 6 to 13.91 points	The mean depression in the intervention groups was 0. 12 standard deviations higher (1.13 lower to 1. 36 higher)		42 (2 studies)	++OO low ³⁴		
Anxiety STAI-trait Follow-up: mean 12 weeks	The mean anxiety in the control groups was 96.18 points	The mean anxiety in the intervention groups was 0. 15 standard deviations higher (0.73 lower to 1. 03 higher)		20 (1 study)	+OOO very low ²³⁴		
General psychiatric pathology - not measured	See comment	See comment	Not estimable	-	See comment	Not measured	
Mental health status GAS/GAF. Scale from: 0 to 100. Follow-up: mean 12 weeks	The mean mental health status ranged across control groups from 59.3 to 82.6 points ¹	The mean mental health status in the intervention groups was 0. 40 standard deviations higher (0.27 lower to 1. 07 higher)		42 (2 studies)	++OO low ³⁵		
Attrition lost after randomisation Follow-up: mean 12 weeks	154 per 1000	251 per 1000 (51 to 1000)	RR 1.63 (0.33 to 8.11)	25 (1 study)	++OO low ³⁴		

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls

² High overall attrition rate (20%), analysis refers to completers only

³ Small sample size

⁴ Single study effect only, publication bias cannot be excluded

⁵ Effect based on a two study estimates only

Appendix 26. Summary of findings: fluvoxamine vs. placebo

Fluvoxamine for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: fluvoxamine						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Fluvoxamine				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Avoidance of abandonment - not measure	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity Borderline Personality Disorder Severity Index-impulsivity Follow-up: 12 weeks	The mean impulsivity in the control groups was 0.73 points ¹	The mean impulsivity in the intervention groups was 0.05 standard deviations lower (0.68 lower to 0.59 higher)		38 (1 study)	++OO low ²³	
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Fluvoxamine for borderline personality disorder						
Affective instability BPDSI-rapid mood shifts	The mean affective instability in the control groups was	The mean affective instability in the intervention groups was		38 (1 study)	++OO low ²³	
Follow-up: mean 6 weeks	5.83 points ¹	0.64 standard deviations lower (1.3 lower to 0.01 higher)				
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger BPDSI-anger Follow-up: mean 6 weeks	The mean anger in the control groups was 2.49 points ¹	The mean anger in the intervention groups was 0.37 standard deviations lower (1.01 lower to 0.28 higher)		38 (1 study)	++OO low ²³	
Psychotic symptoms - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Dissociation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anxiety - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
General psychiatric pathology - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Mental health status - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Attrition lost after randomisation Follow-up: mean 6 weeks	111 per 1000	50 per 1000 (4 to 505)	RR 0.45 (0.04 to 4.55)	38 (1 study)	++OO low ²³	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

¹ Final value in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

Appendix 27. Summary of findings: phenelzine sulfate vs. placebo

Phenelzine sulfate for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: phenelzine sulfate						
Outcomes	Illustrative comparative risks [*] (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk					
	Control	Phenelzine sulfate				
BPD severity Borderline Syndrome Index Follow-up: mean 5 weeks	The mean BPD severity in the control groups was 20.08 ¹ points	The mean BPD severity in the intervention groups was 0. 15 standard deviations lower (0.65 lower to 0. 35 higher)		62 (1 study)	++O_{2,3} low	
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems Atypical Depression Inventory (ADDS) - rejection sensitivity Follow-up: mean 5 weeks	The mean interpersonal problems in the control groups was 0.73 ¹ points	The mean inter-personal problems in the intervention groups was 0. 24 standard deviations higher (0.26 lower to 0. 74 higher)		62 (1 study)	++O_{2,3} low	
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity Barrett Impulsiveness Scale (BIS) Follow-up: mean 5 weeks	The mean impulsivity in the control groups was 237.74 ¹ points	The mean impulsivity in the intervention groups was 0. 00 standard deviations lower (0.5 lower to 0.5 higher)		62 (1 study)	++O_{2,3} low	
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self-mutilating behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Phenelzine sulfate for borderline personality disorder**Patient or population:** patients with borderline personality disorder**Settings:****Intervention:** phenelzine sulfate

Anger SCL-90-hostility. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean anger in the control groups was 1.04 points ¹	The mean anger in the intervention groups was 0.34 standard deviations lower (0.84 lower to 0.17 higher)	62 (1 study)	++OQ _{2,3} low	
Psychotic symptoms SCL-90-paranoid ideation. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean psychotic symptoms in the control groups was 1.18 points ¹	The mean psychotic symptoms in the intervention groups was 0.28 standard deviations lower (0.78 lower to 0.22 higher)	62 (1 study)	++OQ _{1,2} low	
Dissociation - not measured	See comment	See comment	Not estimable	-	See comment Not measured
Depression BDI. Scale from: 0 to 63. Follow-up: mean 5 weeks	The mean depression in the control groups was 19.54 points ¹	The mean depression in the intervention groups was 0.34 standard deviations lower (0.84 lower to 0.16 higher)	62 (1 study)	++OQ _{2,3} low	
Anxiety SCL-90-anxiety. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean anxiety in the control groups was 1.18 points ¹	The mean anxiety in the intervention groups was 0.14 standard deviations lower (0.65 lower to 0.36 higher)	62 (1 study)	++OQ _{2,3} low	
General psychiatric pathology SCL- 90-global severity index. Scale from: 0 to 4. Follow-up: mean 5 weeks	The mean general psychiatric pathology in the control groups was 1.16 points ¹	The mean general psychiatric pathology in the intervention groups was 0.23 standard deviations lower (0.73 lower to 0.27 higher)	62 (1 study)	++OQ _{2,3} low	
Mental health status GAS. Scale from: 0 to 100. Follow-up: mean 5 weeks	The mean mental health status in the control groups was 58.43 points ¹	The mean mental health status in the intervention groups was 0.14 standard deviations higher (0.36 lower to 0.64 higher)	62 (1 study)	++OQ _{2,3} low	
Attrition lost after randomisation Follow-up: mean 5 weeks	176 per 1000	106 per 1000 (32 to 341)	RR 0.60 (0.18 to 1.94)	72 (1 study)	++OQ _{2,3} low

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

¹ Final value in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

Appendix 28. Summary of findings: mianserin vs. placebo

Mianserin for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: mianserin						
Outcomes	Illustrative comparative risks [*] (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Mianserin				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal ideation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self- mutilating behaviour self-harm during treatment Follow-up: mean 6 months	690 per 1000	690 per 1000 (490 to 973)	RR 1.00 (0.71 to 1.41)	58 (1 study)	++O₂ low ^{1,2}	
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured

Mianserin for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: mianserin						
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Psychotic symptoms - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Dissociation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anxiety - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
General psychiatric pathology - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Mental health status - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Attrition lost after randomisation Follow-up: mean 6 months	276 per 1000	414 per 1000 (199 to 861)	RR 1.50 (0.72 to 3.12)	58 (1 study)	++O₂ low	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

¹ Small sample size

² Single study effect only, publication bias cannot be excluded

Appendix 29. Summary of findings: omega-3 fatty acids vs. placebo

Omega-3 fatty acids for borderline personality disorder						
Patient or population: patients with borderline personality disorder						
Settings:						
Intervention: omega-3 fatty acids						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Omega-3 fatty acids				
BPD severity - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Avoidance of abandonment - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Interpersonal problems - not reported	See comment	See comment	Not estimable	-	See comment	Not measured
Identity disturbance - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Impulsivity Modified Overt Aggression Scale (MOAS), total Follow-up: mean 8 weeks	The mean impulsivity in the control groups was 12.9 final value in controls¹	The mean impulsivity in the intervention groups was 0.47 standard deviations lower (1.28 lower to 0.34 higher)		27 (1 study)	++O₃ low^{2,3}	
Suicidal ideation OAS-M- suicidality subscale score of 1 or higher (at least slight suicidal tendencies) Follow-up: mean 12 weeks	704 per 1000	366 per 1000 (197 to 669)	RR 0.52 (0.28 to 0.95)	49 (1 study)	+OOO very low^{2,3,4}	
Suicidal behaviour - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Self-mutilating behaviour self-harm during treatment Follow-up: mean 12 weeks	259 per 1000	319 per 1000 Follow-up: mean 12 weeks (132 to 769)	RR 1.23 Follow-up: mean 12 weeks (0.51 to 2.97)	49 (1 study)	+OOO very low^{2,3,5}	
Affective instability - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Chronic feelings of emptiness - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Anger - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Psychotic symptoms - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Dissociation - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Depression no remission (i. e. at least 50%)	852 per 1000	409 per 1000 (239 to 690)	RR 0.48 (0.28 to 0.81)	49 (1 study)	++O₃ low^{2,3}	

Omega-3 fatty acids for borderline personality disorder

Patient or population: patients with borderline personality disorder

Settings:

Intervention: omega-3 fatty acids

reduction of depressive pathology as assessed by BDI Follow-up: mean 12 weeks

Depression MADRS. Scale from: 0 to 60. Follow-up: mean 8 weeks	The mean depression in the control groups was 8 points ¹	The mean depression in the intervention groups was 0.34 standard deviations lower (1.15 lower to 0.46 higher)	2746 higher)(1 study)	+OOOvery low ^{2,3,4}		
Anxiety - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
General psychiatric pathology - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Mental health status - not measured	See comment	See comment	Not estimable	-	See comment	Not measured
Attrition lost after randomisation Follow-up: 8 to 12 weeks	216 per 1000	132 per 1000 (45 to 387)	RR 0.61 (0.21 to 1.79)	79 (2 studies)	++OOOlow ^{2,5}	

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

* The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

¹ Final value in controls

² Small sample size

³ Single study effect only, publication bias cannot be excluded

⁴ Concomitant psychotropic treatment was allowed without any restrictions and could be changed anytime

⁵ Findings based on two study estimates only

WHAT'S NEW

Last assessed as up-to-date: 24 January 2010.

Date	Event	Description
10 May 2010	New citation required but conclusions have not changed	Substantive amendment undertaken by a new author team
10 May 2010	New search has been performed	This is an update of the review of the same title published first in 2006

HISTORY

Date	Event	Description
30 April 2008	New citation required and conclusions have changed	Substantive amendment

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Because of the deficiency in numbers of study effects per comparison, we did not perform sensitivity analyses as planned, nor was it possible to draw funnel plots.

References to studies included in this review

- Bogenschutz 2004 {published data only} . *Bogenschutz MP, Nurnberg H. Olanzapine versus placebo in the treatment of borderline personality disorder. *Journal of Clinical Psychiatry*. 2004; 65(1):104–9. [PubMed: 14744178]
- Nurnberg, HG.; Bogenschutz, MP. Atypical antipsychotic treatment of BPD: a 12-week, double-blind, placebo-control trial with olanzapine; 156th annual meeting of the American Psychiatric Association; San Francisco. 2003 May 17-22;
- De la Fuente 1994 {published data only} . *De la Fuente JM, Lotstra F. A trial of carbamazepine in borderline personality disorder. *European Neuropsychopharmacology*. 1994; 4:479–86. [PubMed: 7894258]
- De la Fuente JM, Lotstra F. Carbamazepine in borderline personality disorder. *European Neuropsychopharmacology*. 1993; 3:350.
- Frankenburg 2002 {published data only} . Frankenburg FR, Zanarini MC. Divalproex sodium treatment of women with borderline personality disorder and bipolar II disorder: a double-blind placebo-controlled pilot study. *Journal of Clinical Psychiatry*. 2002; 63(5):442–6. [PubMed: 12019669]
- Goldberg 1986 {published data only} . Goldberg SC, Schulz SC, Schulz PM, Resnick RJ, Hamer RM, Friedel RO. Borderline and schizotypal personality disorders treated with low-dose thiothixene vs placebo. *Archives of General Psychiatry*. 1986; 43(7):680–6. [PubMed: 3521531]
- Hallahan 2007 {published data only} . Hallahan B, Hibbeln JR, Davis JM, Garland MR. Omega-3 fatty acid supplementation in patients with recurrent self-harm. *British Journal of Psychiatry*. 2007; 190:118–22. [PubMed: 17267927]
- Hollander 2001 {published data only} . Hollander E, Allen A, Lopez RP, Bienstock CA, Grossman R, Siever LJ, et al. A preliminary double-blind, placebocontrolled trial of divalproex sodium in borderline personality disorder. *Journal of Clinical Psychiatry*. 2001; 62(3):199–203. [PubMed: 11305707]
- Leone 1982 {published data only} . Leone NF. Response of borderline patients to loxapine and chlorpromazine. *Journal of Clinical Psychiatry*. 1982; 43(4):148–50. [PubMed: 7040351]
- Linehan 2008 {published data only} . Linehan, M. Update on dialectical behavioral therapy for BPD; 158th Annual Meeting of the American Psychiatric Association; Atlanta, GA. 2005 May 21-26;

- Linehan MM, McDavid JD, Brown MZ, Sayrs JHR, Gallop RJ. Olanzapine plus dialectical behavior therapy for women with high irritability who meet criteria for borderline personality disorder: a double-blind, placebocontrolled pilot study. *Journal of Clinical Psychiatry*. 2008; 69(9):999–1005. [PubMed: 18466045]
- *Linehan MM, McDavid JD, Brown MZ, Sayrs JHR, Gallop RJ. Olanzapine plus dialectical behavior therapy for women with high irritability who meet criteria for borderline personality disorder: a double-blind, placebocontrolled pilot study. published ahead of print: *Journal of Clinical Psychiatry*. May 6th.2008 :e1–e7. pii: ej07m03158.
- Loew 2006 {published data only} . Loew TH, Nickel MK. Topiramate treatment of women with borderline personality disorder, part II: an open 18-month follow-up. *Journal of Clinical Psychopharmacology*. 2008; 3:355–7. [PubMed: 18480701]
- *Loew TH, Nickel MK, Muehlbacher M, Kaplan P, Nickel C, Kettler C, et al. Topiramate treatment for women with borderline personality disorder: a double-blind, placebocontrolled study. *Journal of Clinical Psychopharmacology*. 2006; 26(1):61–6. [PubMed: 16415708]
- Montgomery 1979/82 {published data only} . *Montgomery SA, Montgomery D. Pharmacological prevention of suicidal behaviour. *Journal of Affective Disorders*. 1982; 4:291–8. [PubMed: 6131083]
- Montgomery SA, Montgomery DB, Janyanthi Rani S, Roy DH, Shaw PJ, McAuley R. Maintenance therapy in repeat suicidal behaviour: a placebo controlled trial. *Proceedings of the 10th international congress for suicide prevention and crisis intervention*. Ottawa. 1979:227–9.
- Montgomery 81/82/83 {published data only} . Montgomery D, Roy D, Montgomery S. Mianserin in the prophylaxis of suicidal behaviour: a double-blind placebo controlled trial. *Dépression et Suicide*. 1981; 1981:786–90.
- *Montgomery SA, Montgomery D. Pharmacological prevention of suicidal behaviour. *Journal of Affective Disorders*. 1982; 4:291–8. [PubMed: 6131083]
- Montgomery SA, Roy D, Montgomery DB. The prevention of recurrent suicidal acts. *British Journal of Clinical Pharmacology*. 1983; 15:183–85.
- Nickel 2004 {published data only} . Nickel M. Topiramate reduced aggression in female patients with borderline personality disorder. *European Archives of Psychiatry & Clinical Neuroscience*. 2007; 257(7):432–3. [PubMed: 17902007]
- *Nickel MK, Nickel C, Mitterlehner FO, Tritt K, Lahmann C, Leiberich PK, et al. Topiramate treatment of aggression in female borderline personality disorder patients: a double-blind, placebo-controlled study. *The Journal of Clinical Psychiatry*. 2004; 65(11):1515–9. [PubMed: 15554765]
- Nickel 2005 {published data only} . Nickel MK. Topiramate treatment of aggression in male borderline patients. *Australian and New Zealand Journal of Psychiatry*. 2007; 41(5):461–2. [PubMed: 17464743]
- Nickel MK, Loew TH. Treatment of aggression with topiramate in male borderline patients, part II: 18-month follow-up. *European Psychiatry*. 2008; 23(2):115–7. [PubMed: 18024088]
- *Nickel MK, Nickel C, Kaplan P, Lahmann C, Muehlbacher M, Tritt K, et al. Treatment of aggression with topiramate in male borderline patients: a double-blind, placebocontrolled study. *Biological Psychiatry*. 2005; 57(5):495–9. [PubMed: 15737664]
- Nickel 2006 {published data only} . Nickel MK. Aripiprazole treatment of patients with borderline personality disorder. *Journal of Clinical Psychiatry*. 2007; 68(11):1815–6. [PubMed: 18052583]
- Nickel MK, Loew TH, Pedrosa Gil F. Aripiprazole in treatment of borderline patients, part II: an 18-month follow-up. *Psychopharmacology*. 2007; 191:1023–6. [PubMed: 17318503]
- *Nickel MK, Mühlbacher M, Nickel C, Kettler C, Pedrosa Gil F, Bachler E, et al. Aripiprazole in the treatment of patients with borderline personality disorder: a double-blind, placebo-controlled study. *American Journal of Psychiatry*. 2006; 163:833–48. [PubMed: 16648324]
- Pascual 2008 {published data only} . [accessed 9 September 2009] Fundacio Institut de Recerca de l'Hospital de la Santa Creu i Sant Pau, Fondo de Investigación Sanitaria (Ministry of Health, Spain), Pfizer. Ziprasidone in the treatment of borderline personality disorder. www.clinicaltrials.gov/NCT00635921

- *Pascual JC, Soler J, Puigdemont D, Perez-Egea R, Tiana T, Alvarez E, et al. Ziprasidone in the treatment of borderline personality disorder: a double-blind, placebo-controlled, randomized study. *Journal of Clinical Psychiatry*. 2008; 69(4):603–8. [PubMed: 18251623]
- Reich 2009 {published data only} . McLean Hospital, GlaxoSmithKline. [accessed 8 September 2009] Study of lamotrigine treatment of affective instability in borderline personality disorder. www.clinicaltrials.gov/NCT00634062
- *Reich DB, Zanarini MC, Bieri KA. A preliminary study of lamotrigine in the treatment of affective instability in borderline personality disorder. *International Clinical Psychopharmacology*. 2009 **e-pub ahead of print**. [DOI: 10.1097/YIC.0b013e32832d6c2f].
- Rinne 2002 {published data only} . Rinne T, van den Brink W, Wouters L, van Dyck R. SSRI treatment of borderline personality disorder: a randomized, placebo-controlled clinical trial for female patients with borderline personality disorder. *American Journal of Psychiatry*. 2002; 159(12):2048–54. [PubMed: 12450955]
- Salzman 1995 {published data only} . Salzman C, Wolfson AN, Schatzberg A, Looper J, Henke R, Albanese M, et al. Effect of fluoxetine on anger in symptomatic volunteers with borderline personality disorder. *Journal of Clinical Psychopharmacology*. 1995; 15(1):23–9. [PubMed: 7714224]
- Schulz 2007 {published data only} . *Eli Lilly, Company. [accessed 14 February 2008] Efficacy and safety of olanzapine in patients with borderline personality disorder: a randomized, flexible-dose, double-blind comparison with placebo [Summary ID #6257. Clinical Study Summary: F1D-MC-HGKL]. www.lillytrials.com/resultsfiles/zyprexa/zyprexasummary6257
- Eli Lilly, Company. [accessed 14 February 2008] Olanzapine in patients with borderline personality disorder. www.clinicaltrials.gov/NCT00091650
- Eli Lilly, Company. [accessed 14 February 2008] Olanzapine in patients with borderline personality disorder. www.who.int/trialsearch
- Schulz SC, Zanarini M, Detke H, Trzaskoma Q, Lin D, DeBerdt W. Olanzapine for the treatment of borderline personality disorder: a flexible-dose 12-week randomized double-blind placebo-controlled study. *International Journal of Neuropsychopharmacology*. 2006; 9(Suppl.1):S191.
- Schulz SC, Zanarini MC, Bateman A, Bohus M, Detke HC, Trzaskoma Q, et al. Olanzapine for the treatment of borderline personality disorder: variable dose, 12-week, randomised double-blind placebo-controlled study. *The British Journal of Psychiatry*. 2008; 193:485–92. [PubMed: 19043153]
- Schulz, SC.; Zanarini, MC.; Detke, HC.; Trzaskoma, Q.; Lin, D.; DeBerdt, W., et al. Olanzapine for the treatment of borderline personality disorder: A flexible-dose, 12-week, randomized, double-blind, placebo-controlled study; Proceedings of the Association of European Psychiatrists 15th European Congress of Psychiatry; Madrid (Spain). 2007 March 17–21;
- Schulz SC, Zanarini MC, Detke HC, Trzaskoma Q, Lin D, DeBerdt W, et al. Olanzapine for the treatment of borderline personality disorder: a flexible-dose 12-week randomized double-blind placebo-controlled study. *European Psychiatry*. 2007; 22:S172.
- Zanarini MC, Schulz SC, Detke HC, Tanaka Y, Zhao F, Trzaskoma Q, et al. Olanzapine for the treatment of borderline personality disorder: two 12-week randomized double-blind placebo-controlled trials. *Neuropsychopharmacology*. 2006; 31:S229–30.
- Simpson 2004 {published data only} . Simpson EB, Yen S, Costello E, Rosen K, Begin A, Pistorello J, et al. Combined dialectical behavior therapy and fluoxetine in the treatment of borderline personality disorder. *Journal of Clinical Psychiatry*. 2004; 65(3):379–85. [PubMed: 15096078]
- Soler 2005 {published data only} . Pascual, JC.; Soler, J.; Barrachina, J.; Campins, J.; Puigdemont, D.; Alvarez, E., et al. Olanzapine plus dialectical behavior therapy of patients with BPD; Proceedings of the 157th annual meeting of the American Psychiatric Association; 2004 May 1–6;
- *Soler J, Pascual JC, Campins J, Barrachina J, Puigdemont D, Alvarez E, et al. Double-blind, placebo-controlled study of dialectical behavior therapy plus olanzapine for borderline personality disorder. *American Journal of Psychiatry*. 2005; 162:1221–4. [PubMed: 15930077]

- Soloff 1989 {published data only} . Soloff PH, George A, Nathan RS, Schulz PM, Ulrich RF, Perel JM. Progress in pharmacotherapy of borderline disorders. *Archives of General Psychiatry*. 1986; 43:691–7. [PubMed: 3521532]
- *Soloff PH, George A, Nathan S, Schulz PM, Cornelius JR, Herring J, et al. Amitriptyline versus haloperidol in borderlines: final outcomes and predictors of response. *Journal of Clinical Psychopharmacology*. 1989; 9(4):238–46. MEDLINE: 24. [PubMed: 2768542]
- Soloff 1993 {published data only} . Soloff PH, Cornelius J, George A, Nathan S, Perel JM, Ulrich RF. Efficacy of phenelzine and haloperidol in borderline personality disorder. *Archives of General Psychiatry*. 1993; 50(5):377–85. [PubMed: 8489326]
- Tritt 2005 {published data only} . Leiberich P, Nickel MK, Tritt K, Pedrosa Gil F. Lamotrigine treatment of aggression in female borderline patients, part II: an 18-month follow-up. *Journal of Psychopharmacology*. 2008; 22(7):805–8. [PubMed: 18308777]
- *Tritt K, Nickel C, Lahmann C, Leiberich PK, Rother WK, Loew TH, et al. Lamotrigine treatment of aggression in female borderline-patients: a randomized, double-blind, placebo-controlled study. *Journal of Psychopharmacology*. 2005; 19(3):287–91. [PubMed: 15888514]
- Zanarini 2001 {published data only} . Zanarini MC, Frankenburg FR. Olanzapine treatment of female borderline personality disorder patients: a double-blind, placebo-controlled pilot study. *Journal of Clinical Psychiatry*. 2001; 62(11):849–54. [PubMed: 11775043]
- Zanarini 2003 {published data only} . Zanarini MC, Frankenburg FR. Omega-3 Fatty acid treatment of women with borderline personality disorder: a double-blind, placebo-controlled pilot study. *American Journal of Psychiatry*. 2003; 160(1):167–9. [PubMed: 12505817]
- Zanarini MC, Frankenburg FR, Parachini EA. A preliminary, randomized trial of fluoxetine, olanzapine, and the olanzapine-fluoxetine combination in women with borderline personality disorder. *Journal of Clinical Psychiatry*. 2004; 65(7):903–7. [PubMed: 15291677]
- Zanarini 2007 {published data only} . *Eli Lilly, Company. [accessed 14 February, 2008] Efficacy and safety of olanzapine in patients with borderline personality disorder. www.who.int/trialsearch/NCT00088036
- Eli Lilly, Company. [accessed 14 February 2008] Efficacy and safety of olanzapine in patients with borderline personality disorder: a randomized double-blind comparison with placebo [Summary ID # 6253. Clinical Study Summary: Study F1D-MC-HGKK]. www.lillytrials.com/resultsfiles/zyprexa/zyprexasummary6523/
- Eli Lilly, Company. [accessed 28 July 2007] Efficacy and safety of olanzapine in patients with borderline personality disorder: a randomized double-blind comparison with placebo. www.clinicaltrials.gov/
- Zanarini, MC.; Schulz, SC.; Detke, HC.; Tanaka, Y.; Zhao, F.; Lin, D., et al. A dose comparison of olanzapine for the treatment of borderline personality disorder: a 12-week randomized, double-blind, placebo-controlled study; Proceedings of the Association of European Psychiatrists 15th European Congress of Psychiatry; Madrid (Spain). 2007 March 17–21;
- Zanarini MC, Schulz SC, Detke HC, Tanaka Y, Zhao F, Lin D, et al. A dose comparison of olanzapine for the treatment of borderline personality disorder: a 12-week randomized double-blind placebo controlled study. *European Psychiatry*. 2007; 22:S172–3.
- Zanarini MC, Schulz SC, Detke HC, Tanaka Y, Zhao F, Lin D, et al. A dose comparison of olanzapine for the treatment of borderline personality disorder: a 12-week randomized double-blind placebo-controlled study. *International Journal of Neuropsychopharmacology*. 2006; 9(Suppl.1):S191.
- Zanarini MC, Schulz SC, Detke HC, Tanaka Y, Zhao F, Trzaskoma Q, et al. Olanzapine for the treatment of borderline personality disorder: two 12-week randomized double-blind placebo-controlled trials. *Neuropsychopharmacology*. 2006; 31:S229–30.

References to studies excluded from this review

- Bellino 2005 {published data only} . Bellino S, Paradiso E, Bogetto F. Oxcarbazepine in the treatment of borderline personality disorder: a pilot study. *Journal of Clinical Psychiatry*. 2005; 66(9):1111–5. [PubMed: 16187767]

- Bellino 2006a {published data only} . Bellino S, Paradiso E, Bogetto F. Efficacy and tolerability of quetiapine in the treatment of borderline personality disorder: A pilot study. *Journal of Clinical Psychiatry*. 2006; 67(7):1042–6. [PubMed: 16889446]
- Bellino 2006b {published data only} . Bellino S, Zizza M, Rinaldi C, Bogetto F. Combined treatment of major depression in patients with borderline personality disorder: a comparison with pharmacotherapy. *Canadian Journal of Psychiatry*. 2006; 51(7):453–60.
- Benedetti 1998 {published data only} . Benedetti F, Sforzini L, Colombo C, Maffei C, Smeraldi E. Low-dose clozapine in acute and continuation treatment of severe borderline personality disorder. *Journal of Clinical Psychiatry*. 1998; 59(3):103–7. [PubMed: 9541151]
- Bohus 1999 {published data only} . Bohus MJ, Landwehrmeyer GB, Stiglmayr CE, Limberger MF, Bohme R, Schmahl CG. Naltrexone in the treatment of dissociative symptoms in patients with borderline personality disorder: an open-label trial. *Journal of Clinical Psychiatry*. 1999; 60(9):598–603. [PubMed: 10520978]
- Chengappa 1999 {published data only} . Chengappa KN, Ebeling T, Kang JS, Levine J, Parepally H. Clozapine reduces severe self-mutilation and aggression in psychotic patients with borderline personality disorder. *Journal of Clinical Psychiatry*. 1999; 60(7):477–84. [PubMed: 10453803]
- Coccaro 1997 {published data only} . Coccaro EF, Kavoussi RJ. Fluoxetine and impulsive aggressive behavior in personality-disordered subjects. *Archives of General Psychiatry*. 1997; 54(12):1081–8. [PubMed: 9400343]
- Cornelius 1990 {published data only} . Cornelius JR, Soloff PH, Perel JM, Ulrich RF. Fluoxetine trial in borderline personality disorder. *Psychopharmacology Bulletin*. 1990; 26(1):151–4. [PubMed: 2371370]
- Cornelius 1993 {published data only} . *Cornelius JR, Soloff PH, George A, Ulrich RF, Perel JM. Haloperidol vs. phenelzine in continuation therapy of borderline disorder. *Psychopharmacology Bulletin*. 1993; 29(2):333–7. [PubMed: 8290683]
- Cornelius JR, Soloff PH, Perel JM, Ulrich RF. Continuation pharmacotherapy of borderline personality disorder with haloperidol and phenelzine. *American Journal of Psychiatry*. 1993; 150(12):1843–8. [PubMed: 8238640]
- Cowdry 1988 {published data only} . Cowdry RW, Gardner DL. Pharmacotherapy of borderline personality disorder. Alprazolam, carbamazepine, trifluoperazine, and tranylcypromine. *Archives of General Psychiatry*. 1988; 45(2):111–9. [PubMed: 3276280]
- Frankenburg 1993 {published data only} . Frankenburg FR, Zanarini MC. Clozapine treatment of borderline patients: a preliminary study. *Comprehensive Psychiatry*. 1993; 34(6):402–5. [PubMed: 8131384]
- Hilger 2003 {published data only} . Hilger E, Barnas C, Kasper S. Quetiapine in the treatment of borderline personality disorder. *World Journal of Biological Psychiatry*. 2003; 4(1):42–4. [PubMed: 12582977]
- Hollander 2005 {published data only} . *Hollander E, Swann AC, Coccaro EF, Jiang P, Smith TB. Impact of trait impulsivity and state aggression on divalproex versus placebo response in borderline personality disorder. *American Journal of Psychiatry*. 2005; 162(3):621–4. [PubMed: 15741486]
- Hollander E, Tracy KA, Swann AC, Coccaro EF, McElroy SL, Wozniak P, et al. Divalproex in the treatment of impulsive aggression: efficacy in cluster B personality disorders. *Neuropsychopharmacology*. 2003; 28:1186–97. [PubMed: 12700713]
- Koenigsberg 2003 {published data only} . Koenigsberg HW, Reynolds D, Goodman M, New AS, Mitropoulou V, Trestman RL, et al. Risperidone in the treatment of schizotypal personality disorder. *Journal of Clinical Psychiatry*. 2003; 64:628–34. [PubMed: 12823075]
- La Malfa 2003 {published data only} . La Malfa G, Lampronti L, Bertelli M, Conte M, Cabras P. Aggressiveness and personality disorders: Evaluation of efficacy and existence of predictors in the treatment with fluvoxamine and lithium] Aggressività e disturbi della personalità. *Minerva Psichiatrica*. 2003; 44:75–81.
- Links 1990 {published data only} . Links PS, Steiner M, Boiago I, Irwin D. Lithium therapy for borderline patients: Preliminary findings. *Journal of Personality Disorders*. 1990; 4(2):173–81.

- Markovitz 1991 {published data only} . Markovitz PJ, Calabrese JR, Schulz SC, Meltzer HY. Fluoxetine in the treatment of borderline and schizotypal personality disorders. *American Journal of Psychiatry*. 1991; 148(8):1064–7. [PubMed: 1853957]
- Markovitz 1995a {published data only} . Markovitz, PJ. Pharmacotherapy of impulsivity, aggression, and related disorders. In: Hollander, E.; Stein, DJ., editors. *Impulsivity and Aggression*. Wiley; New York, NY: 1995. p. 263–87.
- Markovitz 1995b {published data only} . Markovitz PJ, Wagner SC. Venlafaxine in the treatment of borderline personality disorder. *Psychopharmacology Bulletin*. 1995; 31(4):773–7. [PubMed: 8851652]
- Norden 1989 {published data only} . Norden MJ. Fluoxetine in borderline personality disorder. *Progress in Neuropsychopharmacology Biology & Psychiatry*. 1989; 13(6):885–93.
- Parsons 1989 {published data only} . Parsons B, Quitkin FM, McGrath PJ, Stewart JW, Tricamo E, Ocepek-Welickson K, et al. Phenelzine, imipramine, and placebo in borderline patients meeting criteria for atypical depression. *Psychopharmacology Bulletin*. 1989; 25(4):524–34. [PubMed: 2698483]
- Philipsen 2004a {published data only} . Philipsen A, Schmahl C, Lieb K. Naloxone in the treatment of acute dissociative states in female patients with borderline personality disorder. *Pharmacopsychiatry*. 2004; 37(5):196–9. [PubMed: 15470797]
- Philipsen 2004b {published data only} . Philipsen A, Richter H, Schmahl C, Peters J, Rusch N, Bohus M, et al. Clonidine in acute aversive inner tension and self-injurious behavior in female patients with borderline personality disorder. *Journal of Clinical Psychiatry*. 2004; 65(10):1414–9. [PubMed: 15491247]
- Rocca 2002 {published data only} . Rocca P, Marchiaro L, Cocuzza E, Bogetto F. Treatment of borderline personality disorder with risperidone.[see comment]. *The Journal of Clinical Psychiatry*. 2002; 63:241–4. [PubMed: 11926724]
- Russell 2003 {published data only} . Russell JM, Kornstein SG, Shea M, McCullough JP, Harrison WM, Hirschfeld RM, et al. Chronic depression and comorbid personality disorders: Response to sertraline versus imipramine. *Journal of Clinical Psychiatry*. 2003; 64(5):554–61. [PubMed: 12755659]
- Schulz 1999 {published data only} . Schulz SC, Camlin KL, Berry SA, Jesberger JA. Olanzapine safety and efficacy in patients with borderline personality disorder and comorbid dysthymia. *Biological Psychiatry*. 1999; 46(10):1429–35. [PubMed: 10578457]
- Serban 1984 {published data only} . Serban G, Siegel S. Response of borderline and schizotypal patients to small doses of thiothixene and haloperidol. *American Journal of Psychiatry*. 1984; 141(11):1455–8. [PubMed: 6388363]
- Soloff 1986 {published data only} . *Soloff PH, George A, Nathan RS, Schulz PM, Ulrich RF, Perel JM. Progress in pharmacotherapy of borderline disorders. A double-blind study of amitriptyline, haloperidol, and placebo. *Archives of General Psychiatry*. 1986; 43(7):691–7. [PubMed: 3521532]
- Soloff PH, George A, Nathan S, Schulz PM, Ulrich RF, Perel JM. Amitriptyline and haloperidol in unstable and schizotypal borderline disorders. *Psychopharmacology Bulletin*. 1986; 22(1):177–82. [PubMed: 3523575]
- Soloff 1987 {published data only} . *Soloff PH, George A, Nathan RS, Schulz PM, Perel JM. Behavioral dyscontrol in borderline patients treated with amitriptyline. *Psychopharmacology Bulletin*. 1987; 23(1):177–81. [PubMed: 3602316]
- Soloff PH, George A, RS Nathan, Schulz PM, Perel JM. Paradoxical effects of amitriptyline on borderline patients. *American Journal of Psychiatry*. 1986; 143(12):1603–5. [PubMed: 3538914]
- Verkes 1998 {published data only} . Verkes RJ, Van der Mast RC, Hengeveld MW, Tuyl JP, Zwinderman AH, Van Kempen GM. Reduction by paroxetine of suicidal behavior in patients with repeated suicide attempts but not major depression. *American Journal of Psychiatry*. 1998; 155(4):543–7. [PubMed: 9546002]
- Ziegenhorn 2009 {published data only} . Ziegenhorn AA, Roepke S, Schommer NC, Merkl A, Danker-hopfe H, Perschel FH, et al. Clonidine improves hyperarousal in borderline personality

disorder with or without comorbid posttraumatic stress disorder. *Journal of Clinical Psychopharmacology*. 2009; 29(2):170–3. [PubMed: 19512980]

References to ongoing studies

- AstraZen NCT00254748 {published data only} . *AstraZeneca Netherlands Medical Director. [accessed 28 July 2007] The effect of quetiapine on psychotic-like symptoms in borderline personality disordered patients: a randomised placebo-controlled trial. www.clinicaltrials.gov/NCT00254748
- Bohus NCT00124839 {published data only} . *Bohus, M. [accessed 27 July 2007] Evaluation of the efficacy of the opioid antagonist naltrexone on the incidence and intensity of flashbacks and dissociative states in patients with borderline personality disorder. www.clinicaltrials.gov/NCT00124839
- Casas NCT00437099 {published data only} . *Casas, M. [accessed 28 July 2007] Efficacy of omega-3 fatty acids on borderline personality disorder: a randomised, double blind clinical trial. www.clinicaltrials.gov/NCT00437099
- Goodman NCT00255554 {published data only} . Goodman, M. [accessed 28 July 2007] Effects of Dialectical Behavioral Therapy and escitalopram on impulsive aggression, affective instability and cognitive processing in Borderline Personality Disorder. www.clinicaltrials.gov/NCT00255554
- Goodman, M.; New, AS.; Koenigsberg, HW.; Hazlett, E.; Flory, J.; Siever, L. Psychotherapy and combined treatment in BPD: possible effects; 158th Annual Meeting of the American Psychiatric Association; Atlanta, GA. 2005 May 21-26;
- Malev ISRCTN11135486 {published data only} . *Malevani J. Quetiapine versus sertraline as the pharmacological component in a standardised psychopharmacological and psychotherapeutic treatment of borderline personality disorder: a randomised, rater-blinded study. www.controlled-trials.com/ISRCTN11135486.
- Ralevski NCT00463775 {published data only} . *Ralevski, E. [accessed 28 July 2007] Topiramate for treatment of patients with borderline personality disorder and alcohol dependence. www.clinicaltrials.gov/NCT00463775
- Schulz NCT00222482 {published data only} . *Schulz SC. A double-blind and placebo controlled assessment of Depakote ER in borderline personality disorder. www.clinicaltrials.gov/NCT00222482.

Additional references

- Akiskal 2004 . Akiskal HS. Demystifying borderline personality: critique of the concept and unorthodox reflections on its natural kinship with the bipolar spectrum. *Acta Psychiatrica Scandinavica*. 2004; 110(6):401–7. [PubMed: 15521823]
- APA 1980 . American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 3rd Edition. American Psychiatric Association; Washington, DC: 1980.
- APA 1994 . American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th Edition. American Psychiatric Association; Washington, DC: 1994.
- APA 2000a . American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4th edition. American Psychiatric Association; Washington, DC: 2000. text revision
- APA 2000b . Task force for the Handbook of Psychiatric Measures. Handbook of Psychiatric Measures. 1st Edition. American Psychiatric Association; Washington (DC): 2000.
- APA 2001 . American Psychiatric Association. Practice guideline for the treatment of patients with borderline personality disorder. *American Journal of Psychiatry*. 2001; 158(10 Suppl):1–52.
- APA 2005 . Oldham, JM. Guideline Watch: Practice guideline for the treatment of patients with borderline personality disorder. American Psychiatric Association; Arlington, VA: 2005. Available online at <http://www.psych.org/psychpract/treatg/pg/pracguide.cfm>

- Arntz 2003 . Arntz A, van den Hoorn M, Cornelis J, Verheul R, van den Bosch W, de Bie AJHT. Reliability and validity of the Borderline Personality Disorder Severity Index. *Journal of Personality Disorders*. 2003; 17(1):45–59. [PubMed: 12659546]
- Binks 2006 . Binks CA, Fenton M, McCarthy L, Lee T, Adams CE, Duggan C. Pharmacological interventions for people with borderline personality disorder. *Cochrane Database of Systematic Reviews*. 2006; (1) DOI: 10.1002/14651858. CD005653.
- Black 2004 . Black DW, Blum N, Pfohl B, Hale N. Suicidal behavior in borderline personality disorder: prevalence, risk factors, prediction, and prevention. *Journal of Personality Disorders*. 2004; 18(3):226–39. [PubMed: 15237043]
- Duggan 2008 . Duggan C, Huband N, Smailagic N, Ferriter M, Adams C. The use of pharmacological treatments for people with personality disorder: a systematic review of randomized controlled trials. *Personality and Mental Health*. 2008; 2:119–70.
- Elbourne 2002 . Elbourne DR, Altman DG, Higgins JPT, Curtin F, Worthington HV, Vail A. Meta-analyses involving cross-over trials: methodological issues. *International Journal of Epidemiology*. 2002; 31:140–9. [PubMed: 11914310]
- Endicott 1976 . Endicott J, Spitzer RL, Fleiss JL, Cohen J. The Global Assessment Scale: A procedure for measuring overall severity of psychiatric disturbance. *Archives of General Psychiatry*. 1976; 33:766–71. [PubMed: 938196]
- Franke 2002 . Franke, GH. SCL-90-R. Symptom-Checkliste von L.R. Derogatis. Göttingen; Beltz: 2002.
- Gunderson 1981 . Gunderson JG, Kolb JE, Austin V. The diagnostic interview for borderline patients. *American Journal of Psychiatry*. 1981; 138:896–903. [PubMed: 7258348]
- Guy 1976 . Guy, W. Psychopharmacology Research Branch, Division of Extramural Research Programs. U.S. Department of Health, Education, and Welfare, Public Health Service, Alcohol, Drug Abuse, and Mental Health Administration, NIMH; Rockville, MD: 1976. ECDEU Assessment Manual for Psychopharmacology - Revised (DHEW Publ No ADM 76-338); p. 218-222.
- Herpertz 2007 . Herpertz SC, Zanarini M, Schulz CS, Siever L, Lieb K, Möller H-J, et al. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for biological treatment of personality disorders. *The World Journal of Biological Psychiatry*. 2007; 8(4):212–44. [PubMed: 17963189]
- Higgins 2003 . Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003; 327(7414):557–60. [PubMed: 12958120]
- Higgins 2008 . Higgins, JPT.; Green, S., editors. *Cochrane Handbook for Systematic Review of Interventions* Version 5.0.0 [updated February 2008]. The Cochrane Collaboration; 2008. Available from www.cochrane-handbook.org. The Cochrane Collaboration
- Ingenhoven 2010 . Ingenhoven T, Lafay P, Rinne T, Passchier J, Duivenvoorden H. Effectiveness of pharmacotherapy for severe personality disorders: meta-analyses of randomized controlled trials. *Journal of Clinical Psychiatry*. 2010; 71(1):14–25. [PubMed: 19778496]
- Kernberg 1975 . Kernberg, OF. Borderline conditions and primary narcissism. Jason Aronson; New York: 1975.
- Lenzenweger 2007 . Lenzenweger MF, Lane M, Loranger AW, Kessler RC. DSM-IV personality disorders in the National Comorbidity Survey Replication (NCS-R). *Biological Psychiatry*. 2007; 62:553–64. [PubMed: 17217923]
- Lieb 2004 . Lieb K, Zanarini MC, Schmahl C, Linehan MM, Bohus M. Borderline personality disorder. *Lancet*. 2004; 364(9432):453–61. [PubMed: 15288745]
- Linehan 1997 . Linehan MM. Behavioral treatments of suicidal behaviors. Definitional obfuscation and treatment outcomes. *Annals of the New York Academy of Sciences*. 1997; 836:302–28. [PubMed: 9616806]
- Links 2009 . Links, PS.; Kolla, N. Assessing and managing suicide risk. In: Oldham, JM.; Skodol, AE.; Bender, DS., editors. *Essentials of personality disorders*. 1st Edition. American Psychiatric Publishing; Washington, DC: 2009.
- McGlashan 2000 . McGlashan TH, Grilo CM, Skodol AE, Gunderson JG, Shea MT, Morey LC, et al. The Collaborative Longitudinal Personality Disorders Study: baseline Axis I/II and II/II

- diagnostic co-occurrence. *Acta Psychiatrica Scandinavica*. 2000; 102(4):256–64. [PubMed: 11089725]
- Moher 1999 . Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, Stroup DF. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. *Quality of Reporting of Meta-analyses*. *Lancet*. 1999; 354(9193):1896–900. [PubMed: 10584742]
- NICE 2009 . National Collaborating Centre for Mental Health (Commissioned by the National Institute for Health and Clinical Excellence). Borderline Personality Disorder: treatment and management. National Clinical Practice Guideline Number 78. The British Psychological Society and The Royal College of Psychiatrists; <http://www.nice.org.uk/nicemedia/pdf/Borderline%20personality%20disorder%20full%20guideline-published.pdf> [accessed 12 March 2010]
- NICE 2010 . National Institute for Health and Clinical Excellence. [accessed 12 March 2010] <http://www.nice.org.uk/aboutnice/whatwedo/niceandthenhs/UsingNICEGuidanceToCutCostsInTheDownturn.jsp>
- Nosè 2006 . Nosè M, Cipriani A, Biancosino B, Grassi L, Barbui C. Efficacy of pharmacotherapy against core traits of borderline personality disorder: meta-analysis of randomized controlled trials. *International Clinical Psychopharmacology*. 2006; 21:345–53. [PubMed: 17012981]
- Oldham 2004 . Oldham JM, Bender DS, Skodol AE, Dyk IR, Sanislow CA, Yen S, et al. Testing an APA practice guideline: symptom-targeted medication utilization for patients with borderline personality disorder. *Journal of Psychiatric Practice*. 2004; 10(3):156–61. [PubMed: 15330221]
- Paris 2007 . Paris J, Gunderson J, Weinberg I. The interface between borderline personality disorder and bipolar spectrum disorders. *Comprehensive Psychiatry*. 2007; 48(2):145–54. [PubMed: 17292705]
- Perez 2007 . Perez V, Barrachina J, Soler J, Pascual JC, Campins MJ, Puigdemont D, et al. The clinical global impression scale for borderline personality disorder patients (CGI-BPD): a scale sensible to detect changes [Impresión clínica global para pacientes con trastorno límite de la personalidad (ICG-TLP): una escala sensible al cambio]. *Actas Españolas de Psiquiatría*. 2007; 35(4):229–35. [PubMed: 17592784]
- RevMan 2008 . The Nordic Cochrane Centre. The Cochrane Collaboration. Review Manager (RevMan). 5.0. The Nordic Cochrane Centre, The Cochrane Collaboration; Copenhagen: 2008.
- Skodol 2002 . Skodol AE, Gunderson JG, McGlashan TH, Dyck IR, Stout RL, Bender DS, et al. Functional impairment in patients with schizotypal, borderline, avoidant, or obsessive-compulsive personality disorder. *American Journal of Psychiatry*. 2002; 159(2):276–83. [PubMed: 11823271]
- Soloff 1998 . Soloff PH. Algorithms for pharmacological treatment of personality dimensions: Symptom-specific treatments for cognitive-perceptual, affective, and impulsive-behavioral dysregulation. *Bulletin of the Menninger Clinic*. 1998; 62:195–214. [PubMed: 9604516]
- Soloff 2002 . Soloff PH, Lynch KG, Kelly TM. Childhood abuse as a risk factor for suicidal behavior in borderline personality disorder. *Journal of Personality Disorder*. 2002; 16(3):201–14.
- Stern 1938 . Stern A. Psychoanalytic investigation of and therapy in the borderline group of neuroses. *Psychoanalytic Quarterly*. 1938; 7:467–89.
- Stone 1993 . Stone MH. Long-term outcome in personality disorders. *British Journal of Psychiatry*. 1993; 162:299–313. [PubMed: 8453424]
- Torgersen 2005 . Torgersen S. Prevalence, sociodemographics, and functional impairment. In: Oldham, JM.; Skodol, AE.; Bender, DS., editors. *Essentials of Personality Disorders*. 1st Edition. American Psychiatric Publishing; Arlington, VA: 2009. p. 83-102.
- WHO 1993 . World Health Organisation. The ICD-10 Classification of Mental and Behavioral Disorders: Diagnostic Criteria for Research. World Health Organization; Geneva: 1993.
- Zanarini 2003a . Zanarini MC. Zanarini Rating Scale for Borderline Personality Disorder (Zan-BPD): a continuous measure of DSM-IV borderline psychopathology. *Journal of Personality Disorders*. 2003; 17(3):233–42. [PubMed: 12839102]
- Zanarini 2004a . Zanarini MC, Frankenburg FR, Hennen J, Silk KR. Mental health service utilization by borderline personality disorder patients and Axis II comparison subjects followed prospectively for 6 years. *Journal of Clinical Psychiatry*. 2004; 65(1):28–36. [PubMed: 14744165]

- Zanarini 2004b . Zanarini MC, Frankenburg FR, Hennen J, Reich DB, Silk KR. Axis I comorbidity in patients with borderline personality disorder: 6-year follow-up and prediction of time to remission. *American Journal of Psychiatry*. 2004; 161:2108–14. [PubMed: 15514413]
- Zanarini 2005 . Zanarini MC, Frankenburg FR, Hennen J, Reich DB, Silk KR. The McLean Study of Adult Development (MSAD): overview and implications of the first six years of prospective follow-up. *Journal of Personality Disorders*. 2005; 19(5):505–23. [PubMed: 16274279]
- Zanarini 2007 . Zanarini MC, Frankenburg FR, Reich DB, Silk KR, Hudson JI, McSweeney LB. The subsyndromal phenomenology of borderline personality disorder: a 10-year follow-up study. *American Journal of Psychiatry*. 2007; 164(6):929–35. [PubMed: 17541053]

References to other published versions of this review

- Lieb 2010 . Lieb K, Völm B, Rücker G, Timmer A, Stoffers JM. Pharmacotherapy for borderline personality disorder: Cochrane systematic review of randomised trials. *British Journal of Psychiatry*. 2010; 196:4–12. [PubMed: 20044651]

* *Indicates the major publication for the study*

PLAIN LANGUAGE SUMMARY

Drug treatment for borderline personality disorder

Many people with borderline personality disorder (BPD) receive medical treatment. However, there are no drugs available for BPD treatment specifically. A certain drug is most often chosen because of its known properties in the treatment of associated disorders, or BPD symptoms that are also known to be present in other conditions, such as depressive, psychotic, or anxious disorders. BPD itself is characterised by a pervasive pattern of instability in affect regulation (with symptoms such as inappropriate anger, chronic feelings of emptiness, and affective instability), impulse control (symptoms: self-mutilating or suicidal behaviour, ideation, or suicidal threats to others), interpersonal problems (symptoms: frantic efforts to avoid abandonment, patterns of unstable relationships with idealization and depreciation of others), and cognitive-perceptual problems (symptoms: identity disturbance in terms of self perception, transient paranoid thoughts or feelings of dissociation in stressful situations). This review aimed to summarise the current evidence of drug treatment effects in BPD from high-quality randomised trials.

Available studies tested the effects of antipsychotic, antidepressant and mood stabiliser treatment in BPD. In addition, the dietary supplement omega-3 fatty acid (commonly derived from fish) which is supposed to have mood stabilising effects was tested. Twentyeight studies covering 1742 study participants were included.

The findings tended to suggest a benefit from using second-generation antipsychotics, mood stabilisers, and omega-3 fatty acids, but most effect estimates were based on single study effects so repeat studies would be useful. Moreover, the long-term use of these drugs has not been assessed. The small amount of available information for individual comparisons indicated marginal effects for first-generation antipsychotics and antidepressants.

The data also indicated that there may be an increase in self-harming behaviour in patients treated with olanzapine. In general, attention must be paid to adverse effects. Most trials did not provide detailed data of adverse effects and thus could not be considered within this review. We assumed their effects were similar to those experienced by patients with other conditions. Available data of the studies included here suggested adverse effects included weight gain, sedation and change of haemogram parameters with olanzapine treatment, and weight loss with topiramate. Very few beneficial effects were identified for first-generation antipsychotics and antidepressants. However, they may be helpful in the presence of comorbid problems that are not part of BPD core pathology, but can often be found in BPD patients.

There are only few study results per drug comparison, with small numbers of included participants. Thus, current findings of trials and this review are not robust and can easily be changed by future research endeavours. In addition, the studies may not adequately reflect several characteristics of clinical settings (among others, patients' characteristics and duration of interventions and observation periods).

	Adequate sequence generation?	Allocation concealment?	Blinding? (self-rated outcomes)	Blinding? (observer-rated outcomes)	Incomplete outcome data addressed?	Free of selective reporting?	Free of other bias?	Bias due to sponsoring improbable?
Bogenschutz 2004	?	?	?	?	?	?	+	+
De la Fuente 1994	?	?	?	+	?	?	+	?
Frankenburg 2002	+	+	+	?	?	?	+	+
Goldberg 1986	?	?	+	+	+	?	+	?
Hallahan 2007	+	+	+	+	?	?	+	+
Hollander 2001	?	+	?	?	+	?	?	+
Leone 1982	?	?	+	?	+	?	?	+
Linehan 2008	+	+	+	+	?	?	?	+
Loew 2006	?	+	+	+	?	?	+	?
Montgomery 1979/82	?	?	+	+	+	?	?	?
Montgomery 81/82/83	?	?	?	?	+	?	?	?
Nickel 2004	?	+	+	+	?	?	+	?
Nickel 2005	?	+	+	+	?	?	+	?
Nickel 2006	?	+	+	+	+	?	+	?
Pascual 2008	+	?	?	?	+	?	?	+
Reich 2009	+	?	+	?	?	?	?	+
Rinne 2002	?	?	?	?	+	?	+	+
Salzman 1995	?	?	+	+	?	?	+	?
Schulz 2007	+	+	+	+	?	?	?	+
Simpson 2004	?	?	?	+	+	?	+	+
Soler 2005	?	?	?	?	?	+	?	+
Soloff 1989	?	+	?	+	+	?	+	+
Soloff 1993	?	?	+	+	+	?	+	+
Tritt 2005	?	+	+	+	+	+	+	+
Zanarini 2001	+	+	+	+	?	+	+	+
Zanarini 2003	?	?	?	?	?	?	+	+
Zanarini 2004	?	+	+	+	?	?	+	+
Zanarini 2007	+	+	+	+	?	+	?	+

Figure 1. Methodological quality summary: review authors' judgements about each methodological quality item for each included study




Figure 2. Methodological quality graph: review authors' judgements about each methodological quality item presented as percentages across all included studies




Figure 3. Forest plot of comparison: 3.1.1 Active drug versus placebo: Interpersonal problems, SMDs




Figure 4. Forest plot of comparison: 5.1 Active drug versus placebo: Impulsivity, SMDs




Figure 5. Forest plot of comparison: 11.1 Active drug versus placebo: Anger, SMDs





Figure 6. Forest plot of comparison: 12.1 Active drug versus placebo: Psychotic symptoms, SMDs



(1) Despite of the high heterogeneity ($I^2 = 77\%$), we decided to pool estimates, as clinical heterogeneity was quite low, and the same pattern of (2) cf. to (1)

Figure 7. Forest plot of comparison 14.1 Active drug versus placebo: Depression, SMDs




Figure 8. Forest plot of comparison: 15.1 Active drug versus placebo: Anxiety, SMDs





Figure 9. Forest plot of comparison: 16.1 Active drug versus placebo: General psychiatric pathology, SMDs



(1) Despite of the high heterogeneity ($I^2 = 84\%$), we decided to pool estimates, as clinical heterogeneity was quite low, and the same pattern of
(2) cf. to (1)

Figure 10.

Forest plot of comparison: 8.1 Active drug versus placebo: Mental health status, SMDs




Figure 11. Forest plot of comparison: 18.1 Active treatment versus placebo: Attrition, RRs




Figure 12. Funnel plot of comparison: 11.1 Active drug versus placebo: Anger, SMDs