



Distinct Growth Phases in Early Life Associated With the Risk of Type 1 Diabetes: The TEDDY Study

Diabetes Care 2020;43:556–562 | <https://doi.org/10.2337/dc19-1670>

Xiang Liu,¹ Kendra Vehik,¹ Yangxin Huang,² Helena Elding Larsson,³ Jorma Toppari,^{4,5} Anette G. Ziegler,^{6,7,8} Jin-Xiong She,⁹ Marian Rewers,¹⁰ William A. Hagopian,¹¹ Beena Akolkar,¹² Jeffrey P. Krischer,¹ and the TEDDY Study Group*

OBJECTIVE

This study investigates two-phase growth patterns in early life and their association with development of islet autoimmunity (IA) and type 1 diabetes (T1D).

RESEARCH DESIGN AND METHODS

The Environmental Determinants of Diabetes in the Young (TEDDY) study followed 7,522 genetically high-risk children in Sweden, Finland, Germany, and the U.S. from birth for a median of 9.0 years (interquartile range 5.7–10.6) with available growth data. Of these, 761 (10.1%) children developed IA and 290 (3.9%) children were diagnosed with T1D. Bayesian two-phase piecewise linear mixed models with a random change point were used to estimate children's individual growth trajectories. Cox proportional hazards models were used to assess the effects of associated growth parameters on the risks of IA and progression to T1D.

RESULTS

A higher rate of weight gain in infancy was associated with increased IA risk (hazard ratio [HR] 1.09 [95% CI 1.02, 1.17] per 1 kg/year). A height growth pattern with a lower rate in infancy (HR 0.79 [95% CI 0.70, 0.90] per 1 cm/year), higher rate in early childhood (HR 1.48 [95% CI 1.22, 1.79] per 1 cm/year), and younger age at the phase transition (HR 0.76 [95% CI 0.58, 0.99] per 1 month) was associated with increased risk of progression from IA to T1D. A higher rate of weight gain in early childhood was associated with increased risk of progression from IA to T1D (HR 2.57 [95% CI 1.34, 4.91] per 1 kg/year) in children with first-appearing GAD autoantibody only.

CONCLUSIONS

Growth patterns in early life better clarify how specific growth phases are associated with the development of T1D.

Type 1 diabetes (T1D) is a common pediatric chronic disease and is preceded by a preclinical period of islet autoimmunity (IA) in the presence of islet autoantibodies against GAD (GADA), IA2 (IA2A), and/or insulin (IAA). IA commonly develops as early as infancy, with a peak incidence at ~1 to 2 years of age (1–3). The Environmental Determinants of Diabetes in the Young (TEDDY) study reported that HLA-DR-DQ and age are strongly associated with the specific islet autoantibody that appears at the initial seroconversion. The majority of children who seroconvert develop either IAA or GADA as a single first-appearing autoantibody. Over time, these children may remain single autoantibody positive, develop other autoantibody(s), or revert to autoantibody negative. IAA generally appears at an earlier age than GADA, and the order of appearance of the autoantibodies is related to HLA-DR-DQ genotypes (1).

¹Health Informatics Institute, Morsani College of Medicine, University of South Florida, Tampa, FL

²Department of Epidemiology and Biostatistics, College of Public Health, University of South Florida, Tampa, FL

³Department of Clinical Sciences, Lund University/Clinical Research Center, Skane University, Malmö, Sweden

⁴Department of Pediatrics, Turku University Hospital, Turku, Finland

⁵Institute of Biomedicine, Research Centre for Integrative Physiology and Pharmacology, University of Turku, Turku, Finland

⁶Institute of Diabetes Research, Helmholtz Zentrum München (German Research Center for Environmental Health), Munich-Neuherberg, Germany

⁷Forscherguppe Diabetes, Klinikum rechts der Isar, Technical University Munich, Munich, Germany

⁸Forscherguppe Diabetes e.V., Helmholtz Zentrum München (German Research Center for Environmental Health), Munich-Neuherberg, Germany

⁹Center for Biotechnology and Genomic Medicine, Medical College of Georgia, Augusta University, Augusta, GA

¹⁰Barbara Davis Center for Childhood Diabetes, University of Colorado, Aurora, CO

¹¹Pacific Northwest Research Institute, Seattle, WA

¹²National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD

Corresponding author: Xiang Liu, xiang.liu@epi.usf.edu

Received 19 August 2019 and accepted 20 November 2019

This article contains Supplementary Data online at <https://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc19-1670/-/DC1>.

*A complete list of the members of the TEDDY Study Group can be found in the Supplementary Data online.

© 2020 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

Factors in early life likely play an important role in the initiation of IA, and excess growth is a potential candidate, as proposed by the accelerator hypothesis (4) and the overload hypothesis (5). The accelerator hypothesis proposes that excess weight gain and associated insulin resistance accelerate β -cell apoptosis and thereby drive autoimmunity in genetically susceptible children. The overload hypothesis suggests that metabolic overload of the β -cell including insulin resistance might be due to a high growth rate.

Several retrospective studies have reported associations between increased linear growth and weight gain in childhood, particularly in the first 2 years of life, and later onset of T1D (6–8). Longitudinal growth data collected from birth in prospective studies provide an opportunity to study the role of growth in the initiation of IA and the progression to T1D. Several prospective studies reported associations between growth in early life and the risk of IA (9–12). The Australian Baby Diab Study reported that weight SD score (SDS) at 2 years was associated with the risk of IA (9). The U.S. Diabetes Autoimmunity Study in the Young (DAISY) of children aged >2 years showed that height growth velocity, but not weight or BMI growth velocity, was associated with the risks of IA and progression to T1D (10). Moreover, the German BABYDIAB and BABYDIET studies reported that a younger age at infant BMI peak was associated with the risk of IA (13). Previously, the TEDDY study investigated weight and height SDSs at several specific ages during the first 4 years of age and found that the weight SDS at 1 year of age was associated with the risk of IA (12). These findings give evidence that weight gain in very early life and height growth velocity in childhood are associated with the risk of IA. However, there is still no clear picture of whether or how the whole growth trajectory in early life is associated with the development of IA/T1D. This is mainly due to the lack of growth data in infancy and the lack of complex modeling techniques for longitudinal growth data in early life. By modeling of the entire growth trajectory, the growth patterns can be easily identified and help pinpoint the timing of a potential disturbance or exposure.

In this study, we investigated detailed growth data in early life and assessed their relationship with the development

of IA and the progression to T1D in the TEDDY study. We modeled growth trajectories of weight and height in early life in two phases: fast growth in infancy and relatively slow growth subsequently in early childhood (Fig. 1). We used Bayesian two-phase piecewise linear mixed models with a random change point (14) to estimate a child's individual-level random effects (prechange slope, postchange slope, and change point), which correspond to the individual-specific growth parameters (growth rate in infancy, growth rate in early childhood, and age at the phase transition) (Fig. 1B). The estimated change point is the point or time period in which the child transitioned from the phase of rapid infancy growth to another phase of slower childhood growth. Then, we assessed whether these growth parameters were associated with the risk of IA and the risk of progression from IA to T1D.

RESEARCH DESIGN AND METHODS

Participants

TEDDY is a prospective cohort study with the aim to identify environmental causes of T1D. The study included six clinical research centers: three in the U.S. (Colorado, Georgia/Florida, and Washington) and three in Europe (Finland, Germany, and Sweden). Written informed consents were obtained for all study participants from a parent or a primary caretaker, separately, for genetic screening and participation in the prospective follow-up, beginning at birth. The high-risk genotypes selected for inclusion for participants screened from the general population were as follows: DRB1*04-DQA1*03-DQB1*03:02/DRB1*03-DQA1*05-DQB1*02:01 (DR3/4), DRB1*04-DQA1*03-DQB1*03:02/DRB1*04-DQA1*03-DQB1*03:02 (DR4/4), DRB1*04-DQA1*03-DQB1*03:02/DRB1*08-DQA1*04-DQB1*04:02 (DR4/8), and DRB1*03-DQA1*05-DQB1*02:01/DRB1*03-DQA1*




Figure 1—Growth in early life. **A**: Trajectories of body size measurements of 50 randomly selected TEDDY subjects and the mean (loess) curves for weight and height. **B**: Description of the two-phase piecewise linear modeling of growth in early life. **C**: Fitted growth curves of weight and height from the Bayesian two-phase piecewise linear modeling for four randomly selected subjects.

05-DQB1*02:01 (DR3/3). Additional study inclusion genotypes for first-degree relatives of a subject with T1D were: DRB1*04-DQA1*03-DQB1*03:02/DRB1*04-DQA1*03-DQB1*02:02 (DR4/4b), DRB1*04-DQA1*03-DQB1*03:02/DRB1*01-DQA1*01-DQB1*05:01 (DR4/1), DRB1*04-DQA1*03-DQB1*03:02/DRB1*13-DQA1*01-DQB1*06:04 (DR4/13), DRB1*04-DQA1*03-DQB1*03:02/DRB1*09-DQA1*03-DQB1*03:03 (DR4/9), and DRB1*03-DQA1*05-DQB1*02:01/DRB1*09-DQA1*03-DQB1*03:03 (DR3/9) (15). The HLA-DR-DQ genotype abbreviations shown in parentheses will be used throughout this article. A total of 8,676 children were enrolled and followed prospectively from 3 months to 15 years for measurements of weight and height (length before 2 and standing height after 2 years of age), and blood samples were drawn for measurements of islet autoantibodies with study visits every 3 months until 4 years of age and every 3 or 6 months thereafter, depending on autoantibody positivity. All children who were persistently positive for any autoantibody were followed

every 3 months until 15 years of age or onset of T1D. In the analyses, 7,522 children were included after exclusion of 120 children whose HLA eligibility could not be confirmed at a repeated genotyping at 9 months of age, 54 children whose islet autoantibody results were indeterminate, and 980 children with <4 growth measurements in the first 4 years of life. The children were followed from birth for a median of 9.0 years (interquartile range [IQR] 5.7–10.6) (Fig. 2A).

Islet Autoantibodies and T1D

Autoantibodies to IAA, GADA, or IA2A were measured in two laboratories by radio-binding assays (16,17). In the U.S., all sera were assayed at the Barbara Davis Center for Childhood Diabetes at the University of Colorado; in Europe, all sera were assayed at the University of Bristol (Bristol, U.K.). Both laboratories demonstrated high sensitivity and specificity as well as concordance (18). All positive and 5% of negative samples were retested in the

other reference laboratory and deemed confirmed, if concordant. IA was defined as confirmed positive autoantibodies to IAA, GADA, or IA2A in at least two consecutive samples by both laboratories. T1D was diagnosed using American Diabetes Association criteria (19).

Statistical Analyses

The children's weight and height trajectories were analyzed using Bayesian two-phase piecewise linear mixed models with a random change point (14). The models were estimated for girls and boys separately. The estimated individual-level random effects (prechange slope, postchange slope, and change point) (Fig. 1B) correspond to the individual-specific growth parameters (growth rate in infancy, growth rate in early childhood, and age at the phase transition).

Time-to-event analyses using multiple Cox proportional hazards (PH) models were performed to examine the growth parameters (weight and height separately) related to the risk of IA, IAA, or




Figure 2—Study cohort. A: Flow chart of the TEDDY cohort and the study cohort. B: Survival curve of the development of IA. C: Survival curve of progression to T1D from initial seroconversion.

GADA as the first-appearing autoantibody and progression to T1D. The age at the development of IA was the age at the initial of two or more consecutive positive tests. The analyses of progression to T1D included children who developed persistent IA before the onset of T1D. Duration was calculated from the time at first IA positivity to the diagnosis date of T1D or date of last contact to assess T1D status for those who did not develop T1D. The magnitude of the associations was described by hazard ratios (HRs) with 95% CIs. Gestational age-adjusted birth measurement (birth weight in weight models and birth length in height models), HLA-DR-DQ genotype, family history of T1D, duration of exclusive breastfeeding, sex, and country of residence were adjusted for in the Cox PH models. The gestational age-adjusted birth measurements (birth weight or birth length) were the residuals calculated by linear regression of birth measurements on gestational ages for boys and girls separately. In addition, age at IA and type of first-appearing autoantibody (IAA, GADA, or multiple autoantibodies) were adjusted for in the analysis of progression to T1D, in which a small number of children with only IA2A autoantibody were grouped with children with multiple autoantibodies at initial seroconversion. Subgroup analyses stratified by the type of first-appearing autoantibody were also performed.

Sensitivity analyses with additional adjustments for previously identified dietary and genetic risk factors in the TEDDY study (20–24) were performed. Specifically, two dietary factors (early probiotic use and infant formula type in the first 3 months of life), eight T1D-associated non-HLA single nucleotide polymorphisms (SNPs) (rs1004446 in *INS*, rs10517086, rs2476601 in *PTPN22*, rs2816316 in *RGS1*, rs2292239 in *ERBB3*, rs3184504 in *SH2B3*, rs4948088 in *COBL*, and rs12708716 in *CLEC16A*), and two SNPs in complement genes (rs1143678 and rs4597342 in *ITGAM*) were adjusted for in the analysis of IA, while four SNPs (rs1004446 in *INS*, rs10517086, rs1534422, and rs2327832 in *TNFAIP3*) were adjusted for in the analysis of progression to T1D.

The Bayesian two-phase piecewise linear mixed models with a random change point were estimated using the WinBUGS 1.4.3 software (25). All of the other analyses were performed using SAS (version 9.4; SAS Institute, Cary, NC).

RESULTS

As of February 2018, the median (IQR) follow-up time of the last clinic visit for the 7,522 children was 9.0 (5.7–10.6) years. A total of 761 (10.1%) children developed one or more persistent autoantibodies (GADA, IAA, or IA2A) at the median age of 3.0 years (IQR 1.5–6.0), and 3.9% (290/7,522) developed T1D at the median age of 5.5 years (IQR 3.0–8.2) (Fig. 2B). Out of the 761 children with IA, 287 (37.7%) children had IAA only (IAA-first), 329 (43%) children had GADA only (GADA-first), 20 (2.6%) children had IA2A only, and 125 (16.4%) children had multiple autoantibodies at initial seroconversion. Of these children, 35.7% (272 out of 761) progressed to T1D (Fig. 2C). The characteristics of the children by the status of IA (IA negative, any IA, IAA-first IA, and GADA-first IA) and the children diagnosed with T1D are presented in Supplementary Table 1.

Growth Parameters and T1D

The details of the children's growth parameters by the status of IA (IA negative, any IA, IAA-first IA, and GADA-first IA) and T1D are presented in Supplementary Table 1, and additional details are provided in Supplementary Figs. 1–4. The fitted growth curves and measures for four randomly selected subjects are plotted in Fig. 1C.

A higher rate of weight gain in infancy was associated with an increased IA risk adjusted for gestational age-adjusted birth weight, country of residence, sex, duration of exclusive breastfeeding, family history of T1D, and HLA DR-DQ genotype (HR 1.09 [95% CI 1.02, 1.17] per 1 kg/year increase; $P = 0.015$) (Fig. 3A and Supplementary Table 2). The effect of weight growth rate in infancy appeared to be most associated with the risk of GADA-first IA (HR 1.15 [95% CI 1.04, 1.28] per 1 kg/year increase; $P = 0.009$). In addition, gestational age-adjusted birth weight was also positively associated with the risk of IA (HR 1.19 [95% CI 1.02, 1.41] per 1 kg increase; $P = 0.031$) and the risk of GADA-first IA (HR 1.39 [95% CI 1.09, 1.78] per 1 kg increase; $P = 0.008$) (Fig. 3A and Supplementary Table 2).

Height, but not weight, growth parameters were strongly associated with the risk of progression from initial IA seroconversion to T1D, adjusted for birth measurement, country, sex, family history of T1D, HLA DR-DQ genotype, age at IA, and type of first-appearing autoantibody

(Fig. 4 and Supplementary Table 3). A height growth pattern with a lower rate in infancy (HR 0.79 [95% CI 0.70, 0.90] per 1 cm/year increase; $P < 0.001$), a higher rate in early childhood (HR 1.48 [95% CI 1.22, 1.79] per 1 cm/year increase; $P < 0.001$), and a younger age at the phase transition (HR 0.76 [95% CI 0.58, 0.99] per month increase; $P = 0.044$) were associated with an increased risk of progression from IA to T1D (Fig. 4B and Supplementary Table 3).

Subgroup analysis stratified by the type of first-appearing autoantibody showed similar associations between the two height growth rates and the risk of progression to T1D in those who had GADA only or multiple autoantibodies at the time of initial seroconversion (Fig. 4B and Supplementary Table 3). In addition, gestational age-adjusted birth weight and the weight growth rate in early childhood were positively associated with the risk of progression from IA to T1D (HR 2.04 [95% CI 1.03, 4.04] per 1 kg increase, $P = 0.040$; HR 2.57 [95% CI 1.34, 4.91] per 1 kg/year increase, $P = 0.004$) in children who had GADA only at the time of initial seroconversion (Fig. 4A and Supplementary Table 3).

The results were not affected in the sensitivity analyses with the additional adjustments for previously identified risk factors in the TEDDY study (data not shown), in which two dietary factors (early probiotic use and infant formula type in the first 3 months of life), eight T1D-associated non-HLA SNPs (rs1004446 in *INS*, rs10517086, rs2476601 in *PTPN22*, rs2816316 in *RGS1*, rs2292239 in *ERBB3*, rs3184504 in *SH2B3*, rs4948088 in *COBL*, and rs12708716 in *CLEC16A*), and two SNPs in complement genes (rs1143678 and rs4597342 in *ITGAM*) were adjusted for in the analysis of IA, while four SNPs (rs1004446 in *INS*, rs10517086, rs1534422, and rs2327832 in *TNFAIP3*) were adjusted for in the analysis of progression to T1D. Additional analyses by dichotomizing the growth parameters using country- and sex-specific tertiles showed consistent conclusions on the reported associations (data not shown) with an additional association between weight growth rate in early childhood and the risk of progression to T1D overall.

CONCLUSIONS

In this large international multicountry cohort of 7,522 children at high genetic risk for T1D, we modeled growth in early




Figure 3—Forest plots of HR and 95% CI from multiple Cox PH regression analyses of effect of weight (A) and height (B) growth parameters, including gestational age-adjusted birth measurement, on the risk of IA, IAA-first IA, and GADA-first IA. Duration of exclusive breastfeeding, HLA-DR-DQ genotype, family history of T1D, sex, and country of residence were included in the Cox models. mo, months.

life using two distinct phases. Growth transitioned from a fast phase (infancy) to a relatively slow phase (early childhood) at ~7–10 months of age. The weight growth rate in infancy and gestational age-adjusted birth weight were positively associated with the risk of IA and appeared to associate most with the risk of GADA-first IA. Furthermore, the weight growth rate in early childhood and gestational age-adjusted birth weight were

associated with the risk of progression from IA to T1D in children with GADA only at initial seroconversion. A lower height growth rate in early childhood and a higher height growth rate in infancy were associated with the risk of progression from IA to T1D in children who had GADA only or multiple autoantibodies at initial seroconversion.

Our finding that birth weight and the weight growth rate in infancy, but not in early childhood, were positively associated with the risk of IA is consistent with the findings from our previous study (12) and the Australian Baby Diab Study (9), in which weight at a younger age predicted IA more than weight at an older age. Together with our previous finding of a weak association between weight SDS and the risk of IA (12) in the first 4 years of




Figure 4—Forest plots of HR and 95% CI from multiple Cox PH regression analyses of effect of weight (A) and height (B) growth parameters, including gestational age-adjusted birth measurement, on the risk of progression to T1D from IA and from subgroup analysis stratified by the type of first-appearing autoantibody (IAA only, GADA only, or multiple autoantibodies). Duration of exclusive breastfeeding, age at the development of IA, type of first-appearing autoantibody (if applicable), HLA-DR-DQ genotype, family history of T1D, sex, and country of residence were included in the Cox models. mo, months.

designed the study and reviewed and edited the manuscript. X.L. is the guarantor of this work and, as such, had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Prior Presentation. Parts of this study were presented in abstract form at the 79th Scientific Sessions of the American Diabetes Association, San Francisco, CA, 7–11 June 2019.

References

- Krischer JP, Lynch KF, Schatz DA, et al.; TEDDY Study Group. The 6 year incidence of diabetes-associated autoantibodies in genetically at-risk children: the TEDDY study. *Diabetologia* 2015;58:980–987
- Vehik K, Dabelea D. The changing epidemiology of type 1 diabetes: why is it going through the roof? *Diabetes Metab Res Rev* 2011;27:3–13
- Knip M, Veijola R, Virtanen SM, Hyöty H, Vaarala O, Åkerblom HK. Environmental triggers and determinants of type 1 diabetes. *Diabetes* 2005;54(Suppl. 2):S125–S136
- Wilkin TJ. The accelerator hypothesis: weight gain as the missing link between type I and type II diabetes. *Diabetologia* 2001;44:914–922
- Dahlquist G. Can we slow the rising incidence of childhood-onset autoimmune diabetes? The overload hypothesis. *Diabetologia* 2006;49:20–24
- Larsson HE, Hansson G, Carlsson A, et al.; DiPiS Study Group. Children developing type 1 diabetes before 6 years of age have increased linear growth independent of HLA genotypes. *Diabetologia* 2008;51:1623–1630
- EURODIAB Substudy 2 Study Group. Rapid early growth is associated with increased risk of childhood type 1 diabetes in various European populations. *Diabetes Care* 2002;25:1755–1760
- Bruining GJ. Association between infant growth before onset of juvenile type-1 diabetes and autoantibodies to IA-2. Netherlands Kolibriestudy group of childhood diabetes. *Lancet* 2000;356:655–656
- Couper JJ, Beresford S, Hirte C, et al. Weight gain in early life predicts risk of islet autoimmunity in children with a first-degree relative with type 1 diabetes. *Diabetes Care* 2009;32:94–99
- Lamb MM, Yin X, Zerbe GO, et al. Height growth velocity, islet autoimmunity and type 1 diabetes development: the Diabetes Autoimmunity Study in the Young. *Diabetologia* 2009;52:2064–2071
- Beyerlein A, Liu X, Uusitalo UM, et al.; TEDDY Study Group. Dietary intake of soluble fiber and risk of islet autoimmunity by 5 y of age: results from the TEDDY study. *Am J Clin Nutr* 2015;102:345–352
- Elding Larsson H, Vehik K, Haller MJ, et al.; TEDDY Study Group. Growth and risk for islet autoimmunity and progression to type 1 diabetes in early childhood: The Environmental Determinants of Diabetes in the Young study. *Diabetes* 2016;65:1988–1995
- Beyerlein A, Thiering E, Pflueger M, et al. Early infant growth is associated with the risk of islet autoimmunity in genetically susceptible children. *Pediatr Diabetes* 2014;15:534–542
- Brilleman SL, Howe LD, Wolfe R, Tilling K. Bayesian piecewise linear mixed models with a random change point: an application to BMI rebound in childhood. *Epidemiology* 2017;28:827–833
- Hagopian WA, Erlich H, Lernmark A, et al.; TEDDY Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY): genetic criteria and international diabetes risk screening of 421 000 infants. *Pediatr Diabetes* 2011;12:733–743
- TEDDY Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY) study: study design. *Pediatr Diabetes* 2007;8:286–298
- TEDDY Study Group. The Environmental Determinants of Diabetes in the Young (TEDDY) study. *Ann N Y Acad Sci* 2008;1150:1–13
- Bonifacio E, Yu L, Williams AK, et al. Harmonization of glutamic acid decarboxylase and islet antigen-2 autoantibody assays for national institute of diabetes and digestive and kidney diseases consortia. *J Clin Endocrinol Metab* 2010;95:3360–3367
- American Diabetes Association. Standards of medical care in diabetes—2014. *Diabetes Care* 2014;37(Suppl. 1):S14–S80
- Uusitalo U, Liu X, Yang J, et al.; TEDDY Study Group. Association of early exposure of probiotics and islet autoimmunity in the TEDDY study. *JAMA Pediatr* 2016;170:20–28
- Törn C, Hadley D, Lee H-S, et al.; TEDDY Study Group. Role of type 1 diabetes-associated SNPs on risk of autoantibody positivity in the TEDDY study. *Diabetes* 2015;64:1818–1829
- Törn C, Liu X, Hagopian W, et al.; TEDDY Study Group. Complement gene variants in relation to autoantibodies to beta cell specific antigens and type 1 diabetes in the TEDDY Study. *Sci Rep* 2016;6:27887
- Krischer JP, Liu X, Lernmark Å, et al.; TEDDY Study Group. The influence of type 1 diabetes genetic susceptibility regions, age, sex, and family history on the progression from multiple autoantibodies to type 1 diabetes: a TEDDY study report. *Diabetes* 2017;66:3122–3129
- Hummel S, Beyerlein A, Tamura R, et al.; TEDDY Study Group. First infant formula type and risk of islet autoimmunity in The Environmental Determinants of Diabetes in the Young (TEDDY) study. *Diabetes Care* 2017;40:398–404
- Lunn DJ, Thomas A, Best N, Spiegelhalter D. WinBUGS - a Bayesian modelling framework: concepts, structure, and extensibility. *Stat Comput* 2000;10:325–337
- Lifshitz F. Nutrition and growth. *J Clin Res Pediatr Endocrinol* 2009;1:157–163
- Fabricius-Bjerre S, Jensen RB, Færch K, et al. Impact of birth weight and early infant weight gain on insulin resistance and associated cardiovascular risk factors in adolescence. *PLoS One* 2011;6:e20595
- Mendivil CO, Toloza FJ, Ricardo-Silgado ML, et al. Antibodies against glutamic acid decarboxylase and indices of insulin resistance and insulin secretion in nondiabetic adults: a cross-sectional study. *Diabetes Metab Syndr Obes* 2017;10:179–185
- Koskinen MK, Helminen O, Matomäki J, et al. Reduced β -cell function in early preclinical type 1 diabetes. *Eur J Endocrinol* 2016;174:251–259
- Koskinen MK, Lempainen J, Löyttyniemi E, et al. Class II HLA genotype association with first-phase insulin response is explained by islet autoantibodies. *J Clin Endocrinol Metab* 2018;103:2870–2878
- Chen W, Salojin KV, Mi Q-S, et al. Insulin-like growth factor (IGF)-I/IGF-binding protein-3 complex: therapeutic efficacy and mechanism of protection against type 1 diabetes. *Endocrinology* 2004;145:627–638
- Kelley KM, Oh Y, Gargosky SE, et al. Insulin-like growth factor-binding proteins (IGFBPs) and their regulatory dynamics. *Int J Biochem Cell Biol* 1996;28:619–637
- Wen X, Kleinman K, Gillman MW, Rifas-Shiman SL, Taveras EM. Childhood body mass index trajectories: modeling, characterizing, pairwise correlations and socio-demographic predictors of trajectory characteristics. *BMC Med Res Methodol* 2012;12:38