

Classification and Nomenclature of Metacaspases and Paracaspases: No More Confusion with Caspases

Elena A. Minina,^{1,2,*} Jens Staal,³ Vanina E. Alvarez,⁴ John A. Berges,⁵ Ilana Berman-Frank,⁶ Rudi Beyaert,³ Kay D. Bidle,⁷ Frédéric Bornancin,⁸ Magali Casanova,⁹ Juan J. Cazzulo,⁴ Chang Jae Choi,¹⁰ Nuria S. Coll,¹¹ Vishva M. Dixit,¹² Marko Dolinar,¹³ Nicolas Fasel,¹⁴ Christiane Funk,¹⁵ Patrick Gallois,¹⁶ Kris Gevaert,¹⁷ Emilio Gutierrez-Beltran,¹⁸ Stephan Hailfinger,¹⁹ Marina Klemenčič,¹³ Eugene V. Koonin,²⁰ Daniel Krappmann,²¹ Anna Linusson,¹⁵ Maurício F.M. Machado,²² Frank Madeo,²³ Lynn A. Megeney,²⁴ Panagiotis N. Moschou,^{25,26,27} Jeremy C. Mottram,²⁸ Thomas Nyström,²⁹ Heinz D. Osiewacz,³⁰ Christopher M. Overall,³¹ Kailash C. Pandey,³² Jürgen Ruland,^{33,34,35} Guy S. Salvesen,³⁶ Yigong Shi,³⁷ Andrei Smertenko,³⁸ Simon Stael,^{17,39} Jerry Ståhlberg,¹ María Fernanda Suárez,⁴⁰ Margot Thome,¹⁴ Hannele Tuominen,⁴¹ Frank Van Breusegem,³⁹ Renier A.L. van der Hoorn,⁴² Assaf Vardi,⁴³ Boris Zhivotovsky,^{44,45} Eric Lam,⁴⁶ and Peter V. Bozhkov^{1,*}

¹Department of Molecular Sciences, Uppsala BioCenter, Swedish University of Agricultural Sciences and Linnean Center for Plant Biology, Uppsala, Sweden

²COS, Heidelberg University, Heidelberg, Germany

³VIB Center for Inflammation Research, Department of Biomedical Molecular Biology, Ghent University, Ghent, Belgium

⁴Instituto de Investigaciones Biotecnológicas, Universidad Nacional de San Martín, San Martín, Buenos Aires, Argentina

⁵Department of Biological Sciences and School of Freshwater Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI, USA

⁶Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, Israel

⁷Department of Marine and Coastal Sciences, Rutgers University, New Brunswick, NJ, USA

⁸Novartis Institutes for BioMedical Research, Basel, Switzerland

⁹Aix-Marseille Univ, CNRS, LISI, Institut de Microbiologie de la Méditerranée, Marseille, France

¹⁰The University of Texas at Austin, Marine Science Institute, Port Aransas, TX, USA

¹¹Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus UAB, Bellaterra, Barcelona, Spain

¹²Department of Physiological Chemistry, Genentech, South San Francisco, CA, USA

¹³University of Ljubljana, Faculty of Chemistry and Chemical Technology, Ljubljana, Slovenia

¹⁴Department of Biochemistry, University of Lausanne, Epalinges, Switzerland

¹⁵Department of Chemistry, Umeå University, Umeå, Sweden

¹⁶Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK

¹⁷VIB Center for Medical Biotechnology, Department of Biomolecular Medicine, Ghent University, Ghent, Belgium

¹⁸Instituto de Bioquímica Vegetal y Fotosíntesis, Universidad de Sevilla and Consejo Superior de Investigaciones Científicas, Sevilla, Spain

¹⁹Interfaculty Institute for Biochemistry, Eberhard Karls University, Tübingen, Germany

²⁰National Center for Biotechnology Information, National Library of Medicine, Bethesda, MD, USA

²¹Research Unit Cellular Signal Integration, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany

²²Interdisciplinary Center for Biochemical Research, University of Mogi das Cruzes, Mogi das Cruzes, Brazil

²³Institute of Molecular Biosciences, NAWI Graz, University of Graz, BioTechMed Graz, Graz, Austria

(Affiliations continued on next page)

Metacaspases and paracaspases are proteases that were first identified as containing a caspase-like structural fold (Uren et al., 2000). Like caspases, metacaspases and paracaspases are multi-functional proteins regulating diverse biological phenomena, such as aging, immunity, proteostasis, and programmed cell death. The broad phylogenetic distribution of metacaspases and paracaspases across all kingdoms of life and large variation of their biochemical and structural features complicate classification and annotation of the rapidly growing number of identified homologs. Establishment of an adequate classification and

unified nomenclature of metacaspases and paracaspases is especially important to avoid frequent confusion of these proteases with caspases—a tenacious misnomer that unfortunately does not appear to decline with time. This Letter represents a consensus opinion of researchers studying different aspects of caspases, metacaspases, and paracaspases in various organisms, ranging from microbes to plants and animals.

Classification of Metacaspases and Paracaspases

The current classification of proteases provided by the MEROPS database

clusters caspases, metacaspases, and paracaspases to the same family, C14, within the CD clan (<https://www.ebi.ac.uk/merops/>). All members of the C14 family are annotated to possess aspartate P1 cleavage specificity, and the family is further split into two subfamilies: C14A (caspases) and C14B (metacaspases and paracaspases).

Importantly, the MEROPS approach of grouping proteases into families or subfamilies is based on statistically significant similarities of the amino acid sequence within the peptidase domain or part thereof, without considering their biochemical properties (Rawlings et al.,



²⁴Sprott Centre for Stem Cell Research, Ottawa Hospital Research Institute and Departments of Medicine and Cellular and Molecular Medicine, University of Ottawa, Ottawa, ON, Canada

²⁵Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology - Hellas, Heraklion, Greece

²⁶Department of Biology, University of Crete, Greece

²⁷Department of Plant Biology, Uppsala BioCenter, Swedish University of Agricultural Sciences and Linnean Center for Plant Biology, Uppsala, Sweden

²⁸York Biomedical Research Institute, Department of Biology, University of York, York, UK

²⁹Institute for Biomedicine, Sahlgrenska Academy, Centre for Ageing and Health – AgeCap, University of Gothenburg, Gothenburg, Sweden

³⁰Institute for Molecular Biosciences, Faculty of Biosciences, Goethe University, Frankfurt/Main, Germany

³¹Departments of Oral Biological and Medical Sciences and Biochemistry and Molecular Biology, Life Sciences Institute, University of British Columbia, Vancouver, BC, Canada

³²Protein Biochemistry and Engineering Laboratory, ICMR-National Institute of Malaria Research, New Delhi, India

³³Institute of Clinical Chemistry and Pathobiochemistry, School of Medicine, Technical University of Munich, Munich, Germany

³⁴German Cancer Consortium (DKTK), partner site Munich, Germany

³⁵German Center for Infection Research (DZIF), partner site Munich, Germany

³⁶Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA, USA

³⁷School of Life Sciences, Westlake University, Xihu District, Hangzhou Zhejiang Province, China

³⁸Institute of Biological Chemistry, Washington State University, Pullman, WA, USA

³⁹Department of Plant Biotechnology and Bioinformatics, Ghent University, VIB-UGent Center for Plant Systems Biology, Ghent, Belgium

⁴⁰Departamento de Biología Molecular y Bioquímica, Facultad de Ciencias, Universidad de Málaga, Campus de Teatinos, Málaga, Spain

⁴¹Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, Umeå, Sweden

⁴²Department of Plant Sciences, University of Oxford, Oxford, UK

⁴³Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot, Israel

⁴⁴Division of Toxicology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

⁴⁵Faculty of Fundamental Medicine, MV Lomonosov Moscow State University, Moscow, Russia

⁴⁶Department of Plant Biology, Rutgers, The State University of New Jersey, New Brunswick, NJ USA

*Correspondence: alena.minina@slu.se (E.A.M.), peter.bozhkov@slu.se (P.V.B.)

<https://doi.org/10.1016/j.molcel.2019.12.020>

2018). Being valuable for high-throughput protease classification, this approach, however, has substantial drawbacks if implemented without further adjustment. Indeed, in contradiction with the MEROPS description, none of the metacaspases or paracaspases characterized so far cleave after an aspartate residue. Instead, paracaspases are arginine specific (Coornaert et al., 2008; Hachmann et al., 2012; Rebeaud et al., 2008), whereas metacaspases can cleave after either arginine or lysine (Figure S1A; Sundström et al., 2009; Vercammen et al., 2004). Such fundamental differences in the proteolytic specificity between caspases, metacaspases, and paracaspases imply distinct repertoires of new proteoforms that they generate and point to the complex diversification and coevolution of their substrates and downstream pathways. One unfortunate consequence of the current classification is the misuse of caspase-specific probes for studying metacaspases and paracaspases that is commonly found in the literature and leads to false conclusions.

Apart from substrate specificity, caspases, metacaspases, and paracaspases feature other fundamental differences (Figure S1A). For example, active meta-

caspases are monomers and their activation usually requires millimolar concentrations of calcium (Hander et al., 2019; McLuskey et al., 2012; Wong et al., 2012). In contrast, active caspases and paracaspases are calcium-independent dimers (Hachmann et al., 2012; Wiesmann et al., 2012; Yu et al., 2011). This indicates that upstream pathways regulating activation of caspases, metacaspases, and paracaspases are likewise different.

In the past two decades we have learned about important differences between caspases, metacaspases, and paracaspases. Thus, simple extrapolation of features typical for caspases to all other members of the C14 family is not justified anymore. Instead caspases, metacaspases, and paracaspases should be separated into three corresponding groups within the family and each group should be properly annotated by having its key biochemical and structural characteristics provided. We kindly request curators of the MEROPS database to make corresponding changes.

Since the structure and substrate specificity of prokaryotic caspase-like proteases named “orthocaspases” remain largely unknown (Klemenčič et al., 2015), we leave their classification and nomenclature open

until their structural and biochemical properties have been clarified.

Nomenclature of Metacaspases and Paracaspases

The name “caspase” stands for “cysteine-dependent aspartate-specific protease.” Thus, the names “metacaspase” and “paracaspase” imply the wrong substrate specificity for these proteases. However, since these names have been used for two decades, we propose to keep them, provided that caspases, metacaspases, and paracaspases are recognized as three separate groups within the C14 family.

Based on domain composition and arrangement, metacaspases and paracaspases are further subdivided into three and two types, respectively (Figure S1A). For the sake of consistency, we propose to maintain a common nomenclature for the different types of metacaspases and paracaspases using Latin numerals (e.g., type I metacaspases). As for the conserved protein structures, they will be referred to as the p20-like region, the p10-like region, the linker region, and the N-terminal pro-domain, matching the nomenclature of caspases (Figure S1A; Alnemri et al., 1996). The p20, p10, and linker regions have been previously

defined for the caspase group of the C14 family (Fuentes-Prior and Salvesen, 2004) and can be easily identified in metacaspases and paracaspase homologs based on a hidden Markov model (HMM) alignment with the C14 peptidase domain (Figure S1B). Notably, although not always clearly stated in the literature, most known members of the C14 family contain the linker region. Furthermore, type II metacaspases are distinguished by a long linker between the p20 and p10 regions and an additional linker within the p10 region (Figure S1A), which are frequently referred to as a single linker.

We suggest the consideration of the active form of metacaspases or paracaspases as a monomer if it is a cleaved or intact polypeptide chain derived from a single translational event, and a dimer if it comprises uncut or processed products of two translational events.

We propose to establish a unified nomenclature of metacaspases and paracaspases in order to (i) facilitate the comparison of orthologs from different organisms and (ii) make it suitable for annotating homologs of species with partially sequenced genomes. Thus, we suggest using simple root symbols such as MCA for metacaspases and PCA for paracaspases. When naming individual family members, these root symbols will be preceded by the abbreviated Latin name of the species and followed by a hyphen, a Latin number representing the type, and then a small alpha character indicating in alphabetical order the number of the homolog of this type in a given genome (Figure S1C). Proenzymes that require proteolytic processing for activation could be annotated with the prefix “pro-”, e.g. pro-AtMCA-1a for the metacaspase 1 of type I from *A. thaliana*. Splecoforms should be indicated by a decimal number (e.g. AtMCA-1a.1). Please note that these conventions do not consider the letter case, which should conform to gene and protein nomenclature established for a given model organism or taxonomic group.

Importantly, this nomenclature should be used synonymously for metacaspases and paracaspase homologs with well-established names, e.g., human MALT1/HsPCA-1a and *A. thaliana* AtMC1/AtMCA-1a or AtMC4/AtMCA-1la. We encourage all researchers to adopt these recommendations. The new classification and unified nomenclature of metacaspases and paracaspases will facilitate a more comprehensive exchange of relevant findings within the scientific community and help to bridge already existing knowledge with newly discovered homologs, thus promoting mechanistic understanding of these ancient, evolutionarily conserved proteases.

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.molcel.2019.12.020>.

ACKNOWLEDGMENTS

This work was supported by the Knut and Alice Wallenberg Foundation. We apologize to colleagues whose work has not been cited due to space limitations.

REFERENCES

- Alnemri, E.S., Livingston, D.J., Nicholson, D.W., Salvesen, G., Thornberry, N.A., Wong, W.W., and Yuan, J. (1996). Human ICE/CED-3 protease nomenclature. *Cell* 87, 171.
- Coornaert, B., Baens, M., Heyninck, K., Bekaert, T., Haegman, M., Staal, J., Sun, L., Chen, Z.J., Marynen, P., and Beyaert, R. (2008). T cell antigen receptor stimulation induces MALT1 paracaspase-mediated cleavage of the NF- κ B inhibitor A20. *Nat. Immunol.* 9, 263–271.
- Fuentes-Prior, P., and Salvesen, G.S. (2004). The protein structures that shape caspase activity, specificity, activation and inhibition. *Biochem. J.* 384, 201–232.
- Hachmann, J., Snipas, S.J., van Raam, B.J., Cancino, E.M., Houlihan, E.J., Poreba, M., Kasperkiewicz, P., Drag, M., and Salvesen, G.S. (2012). Mechanism and specificity of the human paracaspase MALT1. *Biochem. J.* 443, 287–295.
- Hander, T., Fernández-Fernández, Á.D., Kumpf, R.P., Willems, P., Schatowitz, H., Rombaut, D., Staes, A., Nolf, J., Pottie, R., Yao, P., et al. (2019). Damage on plants activates Ca^{2+} -dependent metacaspases for release of immunomodulatory peptides. *Science* 363, 1–10.
- Klemenčič, M., Novinec, M., and Dolinar, M. (2015). Orthocaspases are proteolytically active prokaryotic caspase homologues: the case of *Microcystis aeruginosa*. *Mol. Microbiol.* 98, 142–150.
- McLuskey, K., Rudolf, J., Proto, W.R., Isaacs, N.W., Coombs, G.H., Moss, C.X., and Mottram, J.C. (2012). Crystal structure of a *Trypanosoma brucei* metacaspase. *Proc. Natl. Acad. Sci. USA* 109, 7469–7474.
- Rawlings, N.D., Barrett, A.J., Thomas, P.D., Huang, X., Bateman, A., and Finn, R.D. (2018). The MEROPS database of proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison with peptidases in the PANTHER database. *Nucleic Acids Res.* 46 (D1), D624–D632.
- Rebeaud, F., Hailfinger, S., Posevitz-Fejfar, A., Tapernoux, M., Moser, R., Rueda, D., Gaide, O., Guzzardi, M., Iancu, E.M., Rufer, N., et al. (2008). The proteolytic activity of the paracaspase MALT1 is key in T cell activation. *Nat. Immunol.* 9, 272–281.
- Sundström, J.F., Vaculova, A., Smertenko, A.P., Savenkov, E.I., Golovko, A., Minina, E., Tiwari, B.S., Rodriguez-Nieto, S., Zamyatin, A.A., Jr., Välineva, T., et al. (2009). Tudor staphylococcal nuclease is an evolutionarily conserved component of the programmed cell death degradome. *Nat. Cell Biol.* 11, 1347–1354.
- Uren, A.G., O'Rourke, K., Aravind, L.A., Pisabarro, M.T., Seshagiri, S., Koonin, E.V., and Dixit, V.M. (2000). Identification of paracaspases and metacaspases: two ancient families of caspase-like proteins, one of which plays a key role in MALT lymphoma. *Mol. Cell* 6, 961–967.
- Vercammen, D., van de Cotte, B., De Jaeger, G., Eeckhout, D., Casteels, P., Vandepoele, K., Vandenberghe, I., Van Beeumen, J., Inzé, D., and Van Breusegem, F. (2004). Type II metacaspases Atmc4 and Atmc9 of *Arabidopsis thaliana* cleave substrates after arginine and lysine. *J. Biol. Chem.* 279, 45329–45336.
- Wiesmann, C., Leder, L., Blank, J., Bernardi, A., Melkko, S., Decock, A., D'Arcy, A., Villard, F., Erbel, P., Hughes, N., et al. (2012). Structural determinants of MALT1 protease activity. *J. Mol. Biol.* 419, 4–21.
- Wong, A.H.H., Yan, C., and Shi, Y. (2012). Crystal structure of the yeast metacaspase Yca1. *J. Biol. Chem.* 287, 29251–29259.
- Yu, J.W., Jeffrey, P.D., Ha, J.Y., Yang, X., and Shi, Y. (2011). Crystal structure of the mucosa-associated lymphoid tissue lymphoma translocation 1 (MALT1) paracaspase region. *Proc. Natl. Acad. Sci. USA* 108, 21004–21009.