

Substrate-dependent CO₂-fixation in heterotrophic bacteria revealed by stable isotope labelling

Running title: Differential CO₂-fixation by *Bacillus subtilis*

Marina Spona-Friedl¹, Alexander Braun¹, Claudia Huber², Wolfgang Eisenreich², Christian Griebler^{1,3},
Andreas Kappler⁴, Martin Elsner^{1,5}

¹ Institute of Groundwater Ecology, Helmholtz Zentrum München, Ingolstädter Landstr. 1, 85764
Neuherberg, Germany

² Chair of Biochemistry, Technische Universität München, Lichtenbergstr. 4, 85747 Garching,
Germany

³ Department of Limnology & Bio-Oceanography, Universität Wien, Althanstr. 14, A-1090 Wien,
Austria

⁴ Geomicrobiology, Eberhard-Karls-University Tuebingen, Sigwartstr. 10, 72076 Tuebingen, Germany

⁵ Chair of Analytical Chemistry and Water Chemistry, Technische Universität München,
Marchioninstr. 17, 81377 München, Germany

Corresponding authors: m.elsner@tum.de; wolfgang.eisenreich@mytum.de

Competing interests: The authors declare that they do not have competing interests.

Abstract

Virtually all heterotrophs incorporate carbon dioxide by anaplerotic fixation. Little explored, however, is the interdependency of pathways and rates of CO₂-fixation on the concurrent usage of organic substrate(s). Potentially, this could reveal which substrates out of a pool of dissolved organic carbon (DOC) are utilised by environmental microorganisms. To explore this possibility, *Bacillus subtilis* W23 was grown in a minimal medium with normalised amounts of either glucose, lactate or malate as only organic substrates, each together with 1 g/L NaH¹³CO₃. Incorporation of H¹³CO₃⁻ was traced by elemental analysis-isotope ratio mass spectrometry (EA-IRMS) of biomass and gas chromatography-mass spectrometry (GC-MS) of protein-derived amino acids. Until the late logarithmic phase, ¹³C-incorporation into the tricarboxylic acid cycle increased with time and occurred *via* [4-¹³C]oxaloacetate formed by carboxylation of pyruvate. The levels of ¹³C-incorporation were highest for growth on glucose and lowest on malate. ¹³C-Incorporation into gluconeogenesis products was mainly detected in the lactate and malate experiment, whereas glucose down-regulated this path. A proof-of-principle study with a natural groundwater community confirmed the ability to determine incorporation from H¹³CO₃⁻ by natural communities leading to specific labelling patterns. This underlines the potential of the labelling approach to characterise carbon sources of heterotrophic microorganisms in their natural environments.

Keywords: Heterotrophic CO₂ Fixation, Isotope Labelling, Organic Substrate Use, Dissolved Organic Matter, Bacterial Substrate Usage, Isotope Analysis

Introduction

Reductive carbon fixation under volcanic conditions played a key role in a potential chemoautotrophic origin of life (1) and CO₂-fixation by autotrophic organisms including plants is among the most important biosynthetic processes in the biosphere (2-5). However, the presence and diversity of carboxylating (CO₂-fixing) enzymes in nature is not restricted to

autotrophs alone. Already 80 years ago, utilisation of CO₂ was reported for heterotrophic bacteria producing propionic acid (*Propionibacteria*) (6). Today, it is recognised that virtually all heterotrophic organisms – from microorganisms to humans – incorporate CO₂ *via* a variety of pathways involving at least 18 different carboxylases in the central and peripheral metabolism (7-10). Among these enzymes, anaplerotic carboxylases incorporate CO₂ into biomass and replenish intermediates of the tricarboxylic acid (TCA) cycle, which are constantly withdrawn for the biosynthesis of amino acids and other metabolic products (10). It is therefore not surprising that carbon from anaplerotic CO₂-incorporation accounts for a significant amount (i.e. 2-8 %) of the cell's biomass carbon abundance (11-16).

Among the protein family of carboxylases, pyruvate carboxylase, an anaplerotic carboxylase, catalyses the bicarbonate (HCO₃⁻)-dependent conversion of pyruvate into oxaloacetate. As a component of a putative ancestral reverse TCA cycle, the enzyme is also present in chemolitho-autotrophic bacteria (17) which are considered as one of the most ancient forms of life. The reaction of pyruvate carboxylase could therefore represent a metabolic feature that goes back to the early evolution of life (18).

The enzyme is widely distributed across the three kingdoms of life and has also been retained in many heterotrophic organisms including the Gram-positive bacterium *Bacillus subtilis* W23. Generally, pyruvate carboxylase occupies a vital position in the central carbon metabolism, since it is located at the “phosphoenolpyruvate-pyruvate-oxaloacetate node” (19-21). This metabolic hub unites structurally entangled reactions that interconnect the major pathways of carbon metabolism, i.e. glycolysis (catabolism), gluconeogenesis (anabolism) and the TCA cycle (energy supply of the cell) (22). However, the direction of the carbon fluxes at this metabolic hub (towards catabolism, anabolism or energy supply) primarily depends on the type of the available dissolved organic carbon (DOC) and it can be expected

that the amount of incorporated CO₂ (or ¹³CO₂/H¹³CO₃⁻ in tracer experiments, respectively) varies even within the same organism depending on the assimilated organic carbon source (12-16). Given a typical metabolic network of a heterotrophic organism capable of carrying out the reaction of pyruvate carboxylase using H¹³CO₃⁻ as a substrate, the following simplified scenarios A – C may be distinguished (Fig. 1):

(A): During growth on carbohydrates like glucose (Glc), the glycolytic flux constantly produces pyruvate (Pyr), which is further oxidised to acetyl coenzyme A (Ac-CoA). Ac-CoA requires oxaloacetate (Oxa) to form citric acid (Cit) in the first reaction of the TCA cycle. The TCA cycle serves, on the one hand, to catabolise substrates to CO₂. On the other hand, intermediates of the TCA cycle are used as building blocks for biosynthesis. Hence, equivalents of Oxa are constantly withdrawn from the TCA cycle for the formation of Asp and related amino acids. Therefore, the pool of Oxa must be replenished to keep the cycle running. To this end, in *B. subtilis* W23, pyruvate carboxylase directly converts Pyr to Oxa *via* the addition of H¹³CO₃⁻. Consequently, TCA cycle metabolites and any products derived thereof, e.g. amino acids like Asp, Lys, Thr, Glu or Pro, are expected to carry this label from H¹³CO₃⁻.

(B): When substrates such as lactate (Lac) enter the metabolic network somewhere between glycolysis and the TCA cycle, Oxa must also be replenished *via* the reaction catalysed by pyruvate carboxylase, in the same way as in the first scenario. In this case, however, also gluconeogenesis *via* Oxa and phosphoenol pyruvate (PEP) would be expected to become active in order to satisfy the need of the organism for glucose and its derivatives (e.g. for building up membranes and the cell wall of the Gram-positive bacterium). As a result, the ¹³C-label from H¹³CO₃⁻ is not only expected in the metabolites and the products of the TCA

cycle (like in the first scenario), but also in those derived from gluconeogenesis or the pentose phosphate pathway, such as Ser, Gly, His, Phe or Tyr.

(C): When cells grow on TCA cycle intermediates like malate (Mal), the substrate directly replenishes the TCA cycle. Hence, Pyr carboxylation seems not to be necessary *per se* and central carbon metabolites are not expected to show ^{13}C -incorporation in labelling experiments with $\text{H}^{13}\text{CO}_3^-$.

However, the assumption that heterotrophic fixation of $\text{CO}_2/\text{HCO}_3^-$ depends on the organic substrate has not been fully exploited yet. This is surprising since this dependency also has the potential to assign the main carbon source which is utilised by heterotrophic microorganisms. Our study aims to close this gap of knowledge by investigating $\text{H}^{13}\text{CO}_3^-$ incorporation into *B. subtilis* W23, a well-known model for a heterotrophic bacterium, during growth on glucose, lactate and malate, respectively. These carbon substrates are indicative of the three different entry points to the central carbon metabolism as depicted in the simplified scenarios in Fig. 1. Additionally, these substrates also represent naturally occurring components in soil and dissolved organic matter (23-25).

To assess and to quantify ^{13}C -incorporation from $\text{H}^{13}\text{CO}_3^-$ in our experiments, we used ^{13}C -based metabolic pathway/flux analysis as a key method (26, 27). Using this technology, carbon from $^{13}\text{CO}_2/\text{H}^{13}\text{CO}_3^-$ can be traced back through the metabolic network of the organism under study. On this basis, mechanisms of CO_2 -fixation, but also downstream fluxes *via* the TCA cycle or gluconeogenesis into metabolic products can be reconstructed on a functional and quantitative basis as shown earlier for plants (28-32) or microorganisms (12, 16, 33-37). Indeed, the latter experiments also pointed at the crucial role of CO_2 -fixation in heterotrophic environmental microbes (24, 33, 38, 39).

In the present study with *B. subtilis* W23 $\text{H}^{13}\text{CO}_3^-$ -labelling experiments were monitored by elemental analysis-isotope ratio mass spectrometry (EA-IRMS) of total biomass and by GC-MS of amino acids to show an example of how to assign main (unlabelled) organic substrates on this basis. Subsequently, $\text{H}^{13}\text{CO}_3^-$ -labelling experiments with a natural microbial community from groundwater were conducted to provide a proof-of-principle that this approach indeed opens a new avenue to elucidate substrate usages in complex environmental samples.

Materials and Methods

Strain and growth conditions

All experiments were performed with *B. subtilis* subsp. *spizizenii* W23 (DSM No.: 6395), a prototrophic derivative of the wild type, obtained from DSMZ (Leibniz Institute DSMZ – German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany). For pre-cultivation, 5 mL of M9 minimal growth medium, supplemented with 1 g/L glucose, 1 g/L lactate or 1.1 g/L malate, respectively, and preheated to 30 °C, were inoculated with 300 µL of a glycerol stock solution of the bacterium. The amounts of organic substrates were chosen in order to ensure that the amount of carbon available to the bacteria (0.4 g/L) was the same in all three setups. The pre-cultures were incubated for 20 h. In order to prevent the formation of biofilms, the culture tubes were shaken vigorously at 300 rpm on an orbital incubation shaker (IKA KS 4000i control, IKA-Werke, Staufen, Germany). Each pre-culture was used to inoculate 195 mL of M9 minimal growth medium, preheated to 30 °C, in 500 mL Schott bottles. The M9 minimal growth medium was a mixture of 165 mL of M9 minimal medium, 20 mL of a 10 g/L glucose, 10 g/L lactate or 11 g/L malate stock solution, respectively, and 10 mL of a 20 g/L sodium bicarbonate stock solution. The bicarbonate was either $\text{NaH}^{13}\text{CO}_3$

(98 atom % ^{13}C , Sigma Aldrich) in the ^{13}C -labelling experiments or unlabelled NaHCO_3 (1.1 % natural ^{13}C -abundance) in the control experiments. The bottles were closed gastight after inoculation to block the release of $^{13}\text{CO}_2$. To avoid depletion of O_2 , an aliquot of fresh air (filter-sterilised using a 0.22 μm syringe filter) that equals the volume of a taken sample, was added at every time point of sampling. The cultivations were performed at 30 °C and 150 rpm on an orbital incubation shaker.

The M9 minimal growth medium consisted of the following components (per litre): 8.5 g of $\text{Na}_2\text{HPO}_4 \cdot 2 \text{H}_2\text{O}$, 3 g of KH_2PO_4 , 1 g of NH_4Cl and 0.5 g of NaCl (=base salts solution). The following components were autoclaved separately before being added to the base salts solution in the given order (per litre): 1 mL of 0.1 M CaCl_2 , 10 mL trace salts stock solution, 1 mL of 1 M MgSO_4 and 1 mL of 50 mM $\text{FeCl}_3 \cdot 6 \text{H}_2\text{O}$ (filter-sterilised using a 0.22 μm syringe filter). The trace salts stock solution contained (per litre): 100 mg of $\text{MnCl}_2 \cdot 4 \text{H}_2\text{O}$, 170 mg of ZnCl_2 , 43 mg of $\text{CuCl}_2 \cdot 2 \text{H}_2\text{O}$, 60 mg of $\text{CoCl}_2 \cdot 6 \text{H}_2\text{O}$ and 60 mg of $\text{Na}_2\text{MoO}_4 \cdot 2 \text{H}_2\text{O}$. The glucose, lactate, malate and sodium bicarbonate stock solutions were filter-sterilised, using a 0.22 μm syringe filter, before being added to the medium. All solutions were prepared using sterilised MilliQ water. All chemicals were purchased from Sigma Aldrich (St. Louis, USA).

^{13}C -Labelling experiments and microbial dry weight

B. subtilis subsp. *spizizenii* W23 was grown in M9 minimal growth medium supplemented with 1 g/L glucose, 1 g/L lactate or 1.1 g/L malate, respectively, and 1 g/L sodium bicarbonate. The ^{13}C -labelling experiments were conducted in triplicates and the control experiments with unlabelled bicarbonate in duplicates. After 10 h of incubation, one control experiment was spiked with sodium ^{13}C -bicarbonate (1 g/L); the second control experiment remained untouched. Bacterial growth was monitored by determining the optical density at

600 nm (OD_{600}). Samples for biomass and amino acid analysis were taken at intervals of 2 h after inoculation. At each of these time points, 20 mL of the bacterial culture were harvested by centrifugation (4 °C, 4000 rpm, 20 min). The supernatant was carefully removed, filter-sterilised, using a 0.22 µm syringe filter, and stored at -20 °C for HPLC analysis (see below). The cell pellet was re-suspended in 2 mL of sterile MilliQ water and transferred into an Eppendorf tube. After this washing step, a second centrifugation step (4 °C, 14000 rpm, 20 min) pelleted the cells again. The supernatant was carefully discarded and the pellet was frozen at -80 °C. The frozen pellets were freeze-dried overnight using a VirTis Sentry 8L benchtop freeze dryer (SP Industries, Warminster, PA, USA). The freeze-dried bacterial pellets were weighed using a high-resolution balance (CP2P, Sartorius AG Göttingen, Germany) to determine the microbial dry weight.

Testing of the $H^{13}CO_3^-$ method with a microbial ground water community

Natural oligotrophic and oxic groundwater samples were collected from a shallow unconsolidated quaternary aquifer composed of fluvio-glacial carbonate gravel and sands at Neuherberg/Munich, Germany. The freshly collected groundwater contained 1.5 ± 0.2 mg/ L DOC, 1.5 ± 0.15 mg/ L nitrate (NO_3^-) and about 15 µg/ L phosphate (PO_4^{3-}), whereas the concentrations of nitrite (NO_2^-) and ammonium (NH_4^+) were below the detection limit of ion chromatography (LOD of 0.1 mg/ L) (Dionex ICS-1100; Thermo Fisher Scientific, USA): 200 µL of the filter-sterilised samples were analysed to determine the concentrations of the nutrients. A stock solution of an organic fertiliser (DOC content 300 mg/ L) was prepared by dissolving the lyophilised substrates at a pH of 9 in ultrapure water, followed by a neutralisation and a centrifugation step. The supernatant was filter-sterilised using a 0.22 µm syringe filter. The organic fertiliser was mainly composed of humic substances of varying molecular size (PhytoGreen®-HumusWP, PhytoSolution).

Each of two bottles with 400 mL groundwater were loaded with 30 mL and 50 mL, respectively, of the organic fertiliser, spiked with 1 g/L ^{13}C -sodium bicarbonate ($\text{NaH}^{13}\text{CO}_3$, 98 atom % ^{13}C , Sigma Aldrich) and closed with a screw cap. The bottles were incubated in the dark at room temperature for 52 days (13th September 2018 – 3rd November 2018) and gently mixed once a week. Growth of the bacterial groundwater community was monitored by optical density measurements at 595 nm.

To enhance biomass yield, the two bottles were combined (on 3rd November 2018), filled with fresh groundwater to a volume of 1000 mL and supplemented with 25 mL of a soil extract, which was prepared from a dark conifer forest soil by solvent extraction: soil from a coniferous wood was extracted with ultra-pure water on a stirrer in darkness at 37 °C overnight. The extract was filter-sterilised using a 0.22 μm syringe filter (Millipore, USA) to remove microbes. The DOC content of the sterile soil extract stock solution was 100 mg/L. On 20th December 2018, the bottle was again loaded with 25 mL sterile soil extract. Bacterial growth was monitored by optical density measurements at 595 nm.

The experiment was ended on 18th January 2019 by sacrificing the whole bottle *via* centrifugation. Biomass was concentrated using centrifugal filter units (30 kDa cut-off, Amicon[®] Ultra-15, Sigma Aldrich) to a final volume of 2 mL. The sample was frozen at -80 °C and freeze-dried overnight to remove the residual water.

Protein hydrolysis and amino acid derivatisation

For protein hydrolysis about 0.5 mg of the freeze-dried bacterial pellet was mixed with 500 μL of 6 M hydrochloric acid and heated at 105 °C for 24 h. After cooling to 70 °C, the residual hydrochloric acid was removed by a constant stream of nitrogen gas. The dried sample was then re-suspended in 50 % glacial acetic acid by sonication for 120 sec. A small column (1 mL pipet tip) of the cation exchanger Dowex 50WX8 [200-400 mesh (=37-74 μm), H^+ form] was prepared and washed with 1 mL of methanol followed by 1 mL of MilliQ

water. After loading the sample onto the column, it was washed twice with 1 mL of MilliQ water. The bound amino acids were then eluted from the column by 1 mL of 4 M ammonium hydroxide. An aliquot of the eluate was dried under a constant stream of nitrogen gas at 70 °C. The dry residue was dissolved in 50 µL of water-free acetonitrile and 50 µL of N-(tert-butyltrimethylsilyl)-N-methyl-trifluoroacetamide containing 1 % tert-butyltrimethylsilylchlorid. This mixture was kept at 70 °C for 30 min. The resulting N-tertbutyltrimethylsilyl-derivatives of the amino acids (TBDMS-amino acid derivatives) were analysed by GC-MS following established protocols (40).

Gas chromatography/mass spectrometry of silylated amino acids

GC-MS analysis was performed using a 7890A GC system (Agilent Technologies, Santa Clara, CA 95051, United States) equipped with a fused silica capillary column (Equity TM-5; 30 m x 0.25 mm, 0.25 µm film thickness; Supelco, Bellefonte, PA 16823, United States). The mass detector worked with electron impact ionisation at 70 eV. An aliquot (1-3 µL) of the solution containing the TBDMS-amino acid derivatives was injected in a 1:10 split mode. The interface temperature was set to 260 °C. The column temperature was held at 140 °C for 3 min, heated with a temperature gradient of 4 °C/ min to 165 °C, heated with a second temperature gradient of 15 °C/ min to 200 °C and heated with a third temperature gradient of 7 °C/ min to 280 °C where the temperature was held for 3 min. Selected ion monitoring data were acquired using a 0.3-sec sampling rate and the samples were analysed three times. Data collection was carried out *via* the GC-MSD Data Analysis software (Agilent Technologies, Santa Clara, CA 95051, United States). The retention times and the detected mass fragments of the amino acids are listed in the supplementary Table T1. ¹³C-Incorporation into amino acids was computed according to Lee et al. (41). The steps include the determination of the contribution of the derivatisation reagent to the observed spectrum of the silylated amino acid and the correction for contribution from ¹³C-carbon natural abundance using multiple linear

regression analysis. The mass isotopomer distribution after this background subtraction provides fractional ^{13}C -excess values for amino acid isotopomers carrying one ^{13}C -carbon atom (M+1), two ^{13}C -carbon atoms (M+2), three ^{13}C -carbon atoms (M+3), and so on, where the sum over all isotopomers [M + (M+1) + (M+2) + (M+3) etc.] is defined as 100 %. As an example, amino acids with an M+1 excess value of 50 % are composed of 50 % unlabelled molecules (M) and 50 % molecules carrying one ^{13}C -carbon (M+1) from the ^{13}C -labelled precursor. Amino acids that carry at least one ^{13}C -carbon atom in excess are termed labelled amino acids in the following.

Carbon isotopic analysis of biomass

Carbon isotopic ratios were determined by an elemental analyser-isotope ratio mass spectrometer (EA-IRMS) consisting of a EuroEA (Euro vector, Milano, Italy) coupled to a FinniganTM MAT253 IRMS (Thermo Fisher Scientific, Bremen, Germany) by a FinniganTM ConFlow III interface (Thermo Fisher Scientific, Bremen, Germany). For EA-IRMS analysis, a small amount of the freeze-dried pellet (100-400 μg) was put into tin capsules (3.3 x 5 mm, IVA Analysentechnik, Meerbusch, Germany) and subjected to elemental analysis by dropping them into a heated reactor which contained silvered cobalt oxide and chromium oxide (IVA Analysentechnik, Meerbusch, Germany and HEKA tech, Wegberg, Germany). The biomass pellets were combusted in a stream of O_2 -containing He at 1000 $^\circ\text{C}$ to produce N_2 , NO_x , H_2O and CO_2 , where NO_x was directly converted to N_2 again in an online reduction reactor filled with metallic copper filings. The gases were subsequently transferred to the isotope ratio mass spectrometer *via* a ConFlow III system using a continuous helium stream of 90 mL/ min. The CO_2 reference gas was provided by CARBO (Bad Hönningen, Germany). The resulting values from EA-IRMS analysis include the natural abundance of ^{13}C -carbon.

Analysis of substrate consumption (HPLC)

The frozen, filter-sterilised supernatant was used for substrate consumption analysis by HPLC. Briefly, glucose, lactate and malate, respectively, were separated and quantified by HPLC using a ligand exchange Aminex HPX 87H column (300 x 7.8 mm) plus precolumn (30 x 4.6 mm) (Bio-Rad Laboratories GmbH, Germany). Aliquots of 20 μ L were injected per run. The column oven was set to 40 °C. The eluent was 5 mM H₂SO₄ at a flow rate of 0.6 mL/ min. Glucose was detected using a RID-10A detector; lactate and malate were detected using the RID-10A and the DAD-SPD-M10Avp detector operating at 210 nm. The retention times of glucose, malate and lactate were 9.1 min, 9.9 min and 12.8 min, respectively.

Statistical Analysis

A two-tailed unpaired Student's t-test was used for the analysis of differences between the mean values of ¹³C-incorporation into selected pairs of amino acids from the experiments with glucose, lactate and malate. Statistical significance is depicted as *ns* = not significant, **p* < 0.05, ***p* < 0.01, or ****p* < 0.001.

Results

Growth of *B. subtilis* in the presence of glucose and H¹³CO₃⁻

The growth experiment with *B. subtilis* W23 in M9 medium containing glucose and H¹³CO₃⁻ displayed the usages of both substrates over time for building up its biomass. The glucose concentration in the medium constantly decreased from 5.6 mM to below the analytical detection limit (3 mM in this setup) at 8 h after inoculation (Fig. 2A). With declining substrate concentration, bacterial biomass increased from 0.03 g/ L to 0.44 g/ L during the experiment

until glucose became limiting. The ^{13}C -abundance of the biomass, as determined by EA-IRMS, steadily rose from 1.1 % (natural abundance of ^{13}C -carbon) to a maximum of 6 % at 6 h after inoculation (Fig. 2B). Then, the ^{13}C -abundance levelled off and stayed constant at about 5 % until the end of the experiment. The control experiment with unlabelled HCO_3^- mirrored the natural abundance of ^{13}C -carbon (1.1 %) in the environment. In the labelling experiment, the maximum of ^{13}C -abundance of 6 % at 6 h and the subsequent decline to 5 % could be explained by the production of unlabelled CO_2 *via* respiration of unlabelled glucose, which led to the formation of unlabelled CO_2 /bicarbonate in the medium as growth occurred. This production of unlabelled bicarbonate led over time to a dilution of the supplied $\text{H}^{13}\text{CO}_3^-$ as indicated by model calculations (see supporting data files F1-F4 for details and supporting Fig. S1). In addition, glucose became limiting so that microbial growth slowed down. Nevertheless, $\text{H}^{13}\text{CO}_3^-$ was still present in the medium and was used by *B. subtilis* W23 for anaplerosis even during the stationary phase of growth finally leading to a stable ^{13}C -abundance of the bacterial biomass till the end of the experiment (Fig. 2B). Specifically, when ^{13}C -bicarbonate was spiked to a non-labelled control after exponential growth in the stationary phase, i.e. after 10 h of inoculation, the ^{13}C -abundance of the biomass still increased up to approximately 2 % (Fig. 2C). This clearly demonstrated that CO_2 -fixation took place even in the absence of cell growth indicating active metabolism during the stationary phase. In similar experiments, we added $\text{H}^{13}\text{CO}_3^-$ to *B. subtilis* W23 during the stationary phase when grown on lactate or malate, respectively (see also below). The ^{13}C -abundance of the respective biomass was again determined by EA-IRMS and accounted for 3 % in the lactate experiment (Fig. S2C) and 2 % in the malate experiment (Fig. S3C). Thus, irrespective of the used carbon substrate and the physiological state (growth phase or stationary phase), metabolic turn-over of oxaloacetate involving the reaction of pyruvate carboxylase remained important, probably to maintain the energy balance also in non-growing *B. subtilis* W23.

Growth of *B. subtilis* in the presence of lactate and $\text{H}^{13}\text{CO}_3^-$

The trends for substrate consumption and biomass production for growth on lactate and $\text{H}^{13}\text{CO}_3^-$ were similar to the experiment with glucose. Briefly, the concentration of lactate decreased from 11.2 mM to 0.3 mM, while the biomass increased from 0.03 g/L to 0.43 g/L during the experiment (Fig. S2A). Again, the formation of unlabelled bicarbonate caused a dilution of the ^{13}C -label at the end of the experiment. EA-IRMS showed that under these conditions *B. subtilis* W23 incorporated 5 % of labelled inorganic carbon into its biomass, which is 1 % less compared to the glucose experiment (Fig. S2B) (see also Fig. 3).

Growth of *B. subtilis* in the presence of malate and $\text{H}^{13}\text{CO}_3^-$

In the third experimental setup, *B. subtilis* W23 was grown in M9 medium supplemented with malate and $\text{H}^{13}\text{CO}_3^-$. Measured substrate consumption demonstrated efficient uptake of malate (42) which was accompanied by an increase of biomass from 0.04 g/L to 0.29 g/L during the experiment. The concentration of malate in the medium decreased from 8.2 mM to below the analytical detection limit of 0.05 mM (Fig. S3A). When using ^{13}C -bicarbonate in the malate medium, the ^{13}C -abundance of the biomass of the bacteria accounted for 3 % under these conditions, as measured by EA-IRMS (Fig. S3B). Notably, this value was significantly lower compared to the glucose and lactate experiments (Fig. 3). Nevertheless, the detection of ^{13}C -incorporation came as a surprise, since exogenous malate could have fully refilled the TCA cycle without the need for anaplerotic replenishment, introducing the ^{13}C -label (see simplified scenario C in Fig. 1).

^{13}C -Labelling patterns of amino acids

The ^{13}C -EA-IRMS results demonstrated the general importance of heterotrophic CO_2 -fixation by pyruvate carboxylase. However, the universal label incorporation under different

conditions makes it difficult to achieve the primary objective of our study, namely to identify the use of different substrates. In a next step, we therefore focused on amino acid-specific incorporation of ^{13}C , which should provide more specific data concerning substrate usage. As an example, amino acids from the TCA cycle (e.g., Asp, Glu) or gluconeogenesis (e.g., Tyr, Phe) were expected to acquire a greater fraction of ^{13}C -carbon as compared to those amino acids derived from pyruvate (e.g., Val, Ala), where the incorporation of ^{13}C -carbon should be low (see Fig. 1). Using established protocols (40), we quantified the ^{13}C -excess (mol-%) in 14 amino acids obtained from acidic hydrolysis of the biomass. Among the labelled amino acids, ^{13}C -excess was found especially for isotopomers carrying one ^{13}C -carbon atom (M+1 isotopomers) as expected for a labelling experiment with $\text{H}^{13}\text{CO}_3^-$.

During growth on glucose, ^{13}C -excess of the M+1 isotopomers of amino acids derived from intermediates of the TCA cycle, such as Asp, Thr, Lys, Glu and Pro (see also Fig. 1) reached values up to 50 % (Fig. 4A). The ^{13}C -excess of the same isotopomers reached values up to 40 % when the bacteria were grown on lactate (Fig. 4B) and values up to 15 % during growth on malate (Fig. 4C). The ^{13}C -excess of the M+1 isotopomers of amino acids derived from gluconeogenic intermediates was low for Ser (about 4 %) and apparently absent for His (derived from the pentose phosphate pathway intermediate, phosphoribosyl pyrophosphate, PRPP) when *B. subtilis* W23 was grown on glucose, lactate or malate, respectively. Glycine, which is also derived from gluconeogenic intermediates, showed a moderate ^{13}C -excess of the M+1 isotopomer under all three conditions (6-8 %). Amino acids derived from pyruvate such as Ala, Val, and Leu received very low ^{13}C -label under all three conditions (< 3 %). Amino acids (Tyr and Phe) that were synthesised from the pentose phosphate pathway intermediate, erythrose-4-phosphate, showed moderate ^{13}C -excess of the respective M+1 isotopomers (about 5 %) when *B. subtilis* W23 was grown on lactate or malate (Fig. 4B and C), and no ^{13}C -excess of the same M+1 isotopomers when grown on glucose (Fig. 4A).

From these comparisons, it becomes already evident that the ^{13}C -patterns in amino acids specifically reflected the (unlabelled) organic carbon substrate used by *B. subtilis* W23 in our model experiments. However, to better visualise the differences in the respective substrate usages, we now compared ratios of ^{13}C -excess of the M+1 isotopomers in specific sets of amino acids (Fig. 5). More specifically, Ala and Val were used as representatives for ^{13}C -incorporation *via* pyruvate (i.e. displaying only very low ^{13}C -excess of the M+1 isotopomers in the experimental settings). Tyr and Phe served as representatives for ^{13}C -bicarbonate incorporation *via* gluconeogenesis and the pentose phosphate pathway, whereas Asp and Glu were used as representatives for ^{13}C -incorporation *via* the TCA cycle (see also Fig. 1). The ratios of the ^{13}C -excess of the M+1 isotopomers at *quasi* steady-state conditions (from 10 h after inoculation till the end of the experiment) gave clear diagnostic trends that uniquely allowed discerning the three different scenarios shown in Fig. 1. When calculating the ratios between the ^{13}C -excess of the M+1 isotopomers in TCA cycle-derived amino acids and those of pyruvate-derived amino acids (i.e., Asp/Val, Asp/Ala, Glu/Val and Glu/Ala), values above 20 were obtained in the experiments with glucose or lactate, where TCA cycle metabolites must be replenished, whereas ratios below 10 were obtained for the same sets of amino acids in the experiments with malate, where anaplerosis is not needed (Fig. 5A). When calculating the ratios between the ^{13}C -excess of the M+1 isotopomers in gluconeogenesis-derived amino acids and those in pyruvate-derived amino acids (i.e., Tyr/Val, Tyr/Ala, Phe/Val and Phe/Ala), ratios above 4 were obtained for experiments with lactate (i.e. under apparently active gluconeogenesis), whereas ratios of 1 or lower were observed for growth on glucose where gluconeogenesis is not needed (Fig. 5B). Thus, ratios of ^{13}C -excess of the M+1 isotopomers between these selected groups of amino acids provided highly selective markers to distinguish the main organic carbon substrates in our model experiments.

Incorporation of ^{13}C -carbon from $\text{H}^{13}\text{CO}_3^-$ by a natural groundwater community

To test whether this approach can also be used to assign unknown substrates for microbes growing in environmental samples, a proof-of-principle experiment was conducted. Specifically, a natural groundwater microbial community was grown for 124 days in water containing organic fertiliser/ soil extract and 1 g/L $\text{H}^{13}\text{CO}_3^-$ (Figs. 7A and B). Indeed, significant ^{13}C -incorporation was found in the biomass of the harvested bacteria (Fig. 7C).

To more specifically assign potential substrates that were used by the microbial community under these conditions, we again analysed amino acids obtained from acidic hydrolysis of the biomass. Overall ^{13}C -incorporation and ^{13}C -excess of the M+1 isotopomers were quantified in 16 amino acids. Meso-diaminopimelic acid (DAP, as a component of peptidoglycan) and lysine reached values up to 12 % ^{13}C -excess of the M+1 isotopomers (Fig. 7C). These amino acids play important roles in cell wall biosynthesis of bacteria (51). DAP is the immediate precursor for lysine (52), which explained the similar ^{13}C -incorporation into these two amino acids. The detected ^{13}C -label of these amino acids showed (i) ^{13}C -incorporation *via* anaplerosis, since DAP and lysine are derived from TCA cycle intermediates and (ii) active cell wall synthesis demonstrating multiplication of the microorganisms under these conditions. Amino acids that are derived from TCA cycle intermediates showed a ^{13}C -excess of the M+1 isotopomers of 3-6%, i.e. Asp (4 %), Thr (3 %), Met (6 %), Glu (4 %) and Pro (4 %).

To check whether it is possible to reveal the type of organic carbon that has been utilised by the groundwater community, we also calculated the ratios of ^{13}C -excess of the M+1 isotopomers between TCA cycle-derived amino acids *vs.* pyruvate-derived amino acids (< 10) (Fig. 7D) and the ratios of ^{13}C -excess of the M+1 isotopomers between gluconeogenesis-derived amino acids *vs.* pyruvate-derived amino acids (< 2) (Fig. 7E). Remarkably, there was

a striking similarity between these values and the corresponding values in the model experiments when *B. subtilis* W23 was grown on malate. Consequently, we hypothesise (also with good statistical significance) that the microorganisms from the groundwater community mainly used organic matter which entered the central carbon metabolism at the stage/level of the TCA cycle – like malate. This result seems reasonable since the utilised organic fertiliser was mostly composed of humic substances, i.e. aromatic compounds, which are degraded to e.g. compounds comprising four carbon atoms, such as succinyl-CoA and succinate. These products could then serve as main substrates entering the central carbon metabolism *via* the TCA cycle such as malate in our model experiment.

Discussion

By means of anaplerotic CO₂-fixation, *B. subtilis* W23 incorporated ¹³C-labelled bicarbonate to a different extent into its biomass depending on the main organic carbon source being present in the minimal medium (i.e. glucose, lactate or malate, respectively). Indeed, the data from EA-IRMS analyses alone could already show significant differences between the three carbon substrates tested in this study, as illustrated in Fig. 3. Incorporation of H¹³CO₃⁻/¹³CO₂ to an extent of 6 % and 5 % into microbial biomass during growth on glucose and lactate, respectively, reflected biomass formation involving anaplerotic carboxylation of pyruvate, which was in some agreement with our simplified scenarios A and B in Fig. 1. Notably, however, in comparison with the glucose experiment, the ¹³C-abundance of the biomass was lower in the lactate experiment (6 % vs. 5 %). At first glance this came as a surprise since we expected the same or an even higher ¹³C-abundance of the bacterial biomass, when lactate was used as the organic substrate. Under lactate conditions, ¹³C-incorporation should also have occurred *via* ¹³C-labelled oxaloacetate into products derived from intermediates of the TCA cycle as well as into those derived from gluconeogenesis and the pentose phosphate

pathway (see scenario B in Fig. 1). The latter routes did not play a major role in the lactate experiment as confirmed by the low levels or the apparent absence of label in His, Ser, Tyr and Phe, respectively. Rather, lactate seemed to be directly channelled *via* pyruvate and Ser into glycerate-3-phosphate and triose phosphates (43) then serving as unlabelled precursors for glucose formation and the pentose phosphate pathway in our experimental setting (23). Following this metabolic flux, cell wall sugars and other gluconeogenic metabolites would not acquire label from $\text{H}^{13}\text{CO}_3^- / ^{13}\text{CO}_2$ *via* ^{13}C -oxaloacetate, thus leading to the observed lower ^{13}C -incorporation. Interestingly, transcriptional, translational and post-translational down-regulation of anaplerotic reactions might be triggered by the presence of exogenous organic acids in the medium (44, 45). Against the background of lactate, as the only organic carbon source in the medium, it seemed safe to assume that anaplerosis was restrained. Similarly, NADH in excess produced by lactate dehydrogenase in the presence of lactate would also down-regulate the TCA cycle and its anaplerotic reactions (46). Together, less ^{13}C -carbon is incorporated into gluconeogenic products and the overall biomass when the bacteria grow on lactate compared to growth on glucose.

The occurrence of the ^{13}C -label in amino acids also clearly assigned pyruvate carboxylase as the $\text{H}^{13}\text{CO}_3^- / ^{13}\text{CO}_2$ -binding enzyme in all of our settings including the experiment with malate (for details, see below). The unexpected incorporation of 3 % ^{13}C -carbon into microbial biomass during growth on malate suggests that, even under this condition, pyruvate carboxylase was still actively transforming pyruvate to oxaloacetate, even though the organism could have directly refilled the oxaloacetate pool of the TCA cycle by taking excess malate from the growth medium (see scenario C in Fig. 1). Thus, although being characterised as non-essential for *B. subtilis* W23 in SubtiWiki (47), the constantly expressed gene for pyruvate carboxylase implies a permanent activity for this enzyme (20, 22) – possibly to be able to quickly react when growth conditions change and anaplerosis becomes necessary for

survival (48, 49). Interestingly, metabolic activity of pyruvate carboxylase could also be detected during the stationary growth phase of *B. subtilis* W23 irrespective of the main organic substrate being used from the medium (Figs. 2C, S2C and S3C). This also indicates the important role of the constantly present enzyme pyruvate carboxylase in the metabolism of *B. subtilis* W23.

EA-IRMS analysis alone could not clearly pinpoint the metabolic history of the incorporated inorganic carbon under the different conditions, nor could it exclude that multiple enzymes contributed to the observed label incorporation. To further confirm that pyruvate carboxylase is responsible for ^{13}C -incorporation into microbial biomass and to follow the labelled inorganic carbon through the metabolic network of the microbial cell, we used GC-MS analysis to reveal information about the carbon positions in amino acids having acquired the label. On the basis of the detected fragmentation patterns of the silylated amino acids (for details, see Table T1), some positional assignments of the ^{13}C -label, especially those involving C-1 of the amino acids, were possible (Fig. 6). As an example, the fragments with m/z of 432 and 286 for Glu and Pro, respectively, contained all five carbon atoms of the original amino acid. When analysing the mass spectra we found that these fragments were accompanied by a high amount (up to 50 %) of the respective M+1 isotopomer (namely m/z 433 for Glu and m/z 287 for Pro). In contrast, the fragments that had lost the C-1 carbon atom of these amino acids (i.e. m/z 404 and 330 for Glu and m/z 258 and 184 for Pro) were not accompanied by a significant excess of the respective M+1 isotopomers (< 1 %). On this basis, it can be safely concluded that Glu and Pro carried the ^{13}C -label at C-1. Similarly, the mass distribution in the fragments observed for Asp, Thr and Lys signalled high amounts of the respective M+1 isotopomers (up to 50 %). Here, the ^{13}C -labelled carbon atom was mainly present at C-4 of these amino acids, as learned from the analysis of the respective fragments. However, lower amounts of ^{13}C -label (1 – 5 %) could also be assigned to C-1 of these amino

acids. As illustrated in Fig. 6, this label distribution can be explained because (i) C-4 of oxaloacetate acquires the ^{13}C -label from $\text{H}^{13}\text{CO}_3^-$ by the reaction of pyruvate carboxylase, (ii) $[4-^{13}\text{C}]$ oxaloacetate is converted into $[1-^{13}\text{C}]\alpha$ -ketoglutarate *via* the oxidative branch of the TCA cycle (leading to the detected $[1-^{13}\text{C}]$ -isotopomers of Glu and Pro), (iii) the biosynthesis of Asp, Thr and Lys is based on the TCA cycle intermediate oxaloacetate (leading to the detected $[4-^{13}\text{C}]$ -isotopomers), and (iv) reversible reactions between oxaloacetate and succinate, in the reductive branch of the TCA cycle, lead to a scrambling of label between C-1 and C-4 of the symmetric intermediates fumarate and succinate. Hence, the ^{13}C -label was transferred also into C-1 of oxaloacetate and its downstream products Asp, Thr and Lys (see also Fig. 6, red half circles). Remarkably, the position-specific incorporation of ^{13}C -carbon at C-1 of Glu and Pro, and C-4 of Asp, Thr and Lys reached values up to 50 % when the bacteria were grown on glucose and as high as 40 % when grown on lactate (Fig. 4A and B). These data demonstrated that the anaplerotic reaction catalysed by pyruvate carboxylase transferred the ^{13}C -label efficiently and quite specifically into C-4 of oxaloacetate and its related downstream products (Fig. 6), and, thereby, gave direct evidence of the heterotrophic CO_2 -fixation. During growth on malate, the ^{13}C -label was found at the same positions, but the ^{13}C -excess of the respective M+1 isotopomers was significantly lower (< 15 %) (Fig. 4C). The amino acids derived from glycolytic precursors, especially Ser and Gly, were characterised by a ^{13}C -incorporation at C-1 of 2.8 %, 3.6 % or 4.5 % (Ser) and 6.4 %, 8.8 % or 7.7 % (Gly), when *B. subtilis* W23 was grown on glucose, lactate or malate, respectively. This ^{13}C -incorporation could again be explained by the equilibrium reactions of the TCA cycle: $[4-^{13}\text{C}]$ oxaloacetate led to $[4-^{13}\text{C}]$ -isotopomers of malate, fumarate and succinate. Since succinate and fumarate are symmetrical intermediates of the TCA cycle, they led in turn to an equal mixture of $[1-^{13}\text{C}]$ - and $[4-^{13}\text{C}]$ oxaloacetate (Fig. 6, red half circles). An active gluconeogenesis could then have transported the ^{13}C -label from the C-1-position of oxaloacetate into the C-1-position of PEP and upstream from there into $[1-^{13}\text{C}]$ Ser and $[1-$

¹³C]Gly. However, *B. subtilis* W23 could also have used the reversible reaction of the Gly cleavage system (50) to synthesise Gly, as shown in the following formula:



The reverse reaction of the glycine cleavage system could have afforded [1-¹³C]Gly which then serves as the precursor for Ser biosynthesis yielding [1-¹³C]Ser without the requirement of an active pyruvate carboxylase (43, 50). Alternatively, [1-¹³C]Gly could be formed by cleavage of 2-aminoacetoacetate (obtained from Thr) whereby C-1 and C-2 of Thr are transformed into C-1 and C-2 of Gly, respectively. Thus, *via* this route the detected (low) label at C-1 of Thr is transferred to C-1 of Gly. Now, one could speculate that these alternative pathways should be active under all the experimental setups and, consequently, the same ¹³C-excess of the respective M+1 isotopomers of Gly and Ser should have resulted in all three cases. This is not true (Fig. 4) and, therefore, the detected differences in ¹³C-incorporation indicate that a significant fraction of Ser and Gly was synthesised *via* [1-¹³C]-PEP. However, we cannot exclude that different Gly biosynthesis pathways are used by *B. subtilis* W23 when growing on different organic substrates.

In summary, our experiments using H¹³CO₃⁻ as a tracer and *B. subtilis* W23 as a model organism, as well as our proof-of-principle experiment with a natural groundwater community, therefore provide solid evidence that EA-IRMS analysis of the biomass in conjunction with GC-MS analysis of protein- and cell wall-derived amino acids (i) reflect the core functional metabolic networks of the organism(s) under study and (ii) can identify the type of the main organic carbon substrate or at least the substrate family being used by the heterotrophic organism or organisms under study.

The general validity of this hypothesis is supported by the fact that almost all heterotrophs need to refill the TCA cycle by anaplerotic CO₂-fixation. For this purpose, heterotrophs either use pyruvate carboxylase which is highly conserved and found in a great variety of organisms including prokaryotes, archaea, yeasts, fungi and higher organisms (e.g. mammals) or phosphoenolpyruvate (PEP) carboxylase which is also widely distributed in bacteria (19, 20, 22, 53). PEP carboxylase serves as another anaplerotic enzyme which catalyses the reaction from phosphoenolpyruvate to oxaloacetate *via* the addition of bicarbonate. Presumably, this results in the same labelling patterns when starting from H¹³CO₃⁻ in comparison to organisms using the pyruvate carboxylase. Thus, labelling experiments using H¹³CO₃⁻ as a tracer bear high potential to generally assign the type of microbial DOC utilisation under various conditions.

Acknowledgments

We thank Ellen Röhm (Geomicrobiology – University of Tübingen) for performing the HPLC measurements and Ramona Brejcha (Institute of Groundwater Ecology, Helmholtz Zentrum München) and Martina Daubmeier (Institute of Groundwater Ecology, Helmholtz Zentrum München) for supporting us in the acquisition of the elemental analyser-isotope ratio mass spectrometer analyses of the bacterial cell pellets. This work was funded by the Helmholtz Association and in part (to W.E.) by grants from the Hans-Fischer-Gesellschaft (Munich, Germany) and the Deutsche Forschungsgemeinschaft DFG (Bonn, Germany) (TRR235).

Competing interests

none

References

1. Huber C, Kraus F, Hanzlik M, Eisenreich W, Wächtershäuser G. Elements of metabolic evolution. *Chemistry*. 2012;18:2063-80.
2. Berg IA. Ecological aspects of the distribution of different autotrophic CO₂ fixation pathways. *Appl Environ Microbiol*. 2011;77(6):1925-36.
3. Fuchs G. Alternative pathways of carbon dioxide fixation: insights into the early evolution of life? *Annu Rev Microbiol*. 2011;65:631-58.
4. Giovannoni SJ, Stingl U. Molecular diversity and ecology of microbial plankton. *Nature*. 2005;437(7057):343-8.
5. Bassham JA, Calvin M. The way of CO₂ in plant photosynthesis. *Comp Biochem Physiol*. 1962;4:187-204.
6. Wood HG, Werkman CH. The utilisation of CO₂ in the dissimilation of glycerol by the propionic acid bacteria. *Biochem J*. 1936;30:48-53.
7. Middelburg JJ. Chemoautotrophy in the ocean. *Geophys Res Lett*. 2011;38:L24604.
8. Baltar F, Arístegui J, Sintes E, Gasol JM, Reinthaler T, Herndl GJ. Significance of non-sinking particulate organic carbon and dark CO₂ fixation to heterotrophic carbon demand in the mesopelagic northeast Atlantic. *Geophysical Research Letters*. 2010;37(9):n/a-n/a.
9. Dijkhuizen L, Harder W. Microbial metabolism of carbon dioxide. *Comprehensive Biotechnology*. 1985;1:409-23.
10. Erb TJ. Carboxylases in natural and synthetic microbial pathways. *Appl Environ Microbiol*. 2011;77(24):8466-77.
11. Herndl GJ, Reinthaler T. Microbial control of the dark end of the biological pump. *Nature Geoscience*. 2013;6:718-24.
12. Miltner A, Richnow H-H, Kopinke F-D, Kästner M. Assimilation of CO₂ by soil microorganisms and transformation into soil organic matter. *Organic Geochemistry*. 2004;35(9):1015-24.
13. Perez RC, Martin A. Carbon dioxide assimilation by *Thiobacillus novellus* under nutrient-limited mixotrophic conditions. *Journal of bacteriology*. 1982;150(1):46-51.
14. Romanenko VI. Heterotrophic CO₂ assimilation by water bacterial flora. *Mikrobiologiya*. 1964;33:679-83.
15. Doronia NV, Trotsenko YA. Levels of carbon dioxide assimilation in bacteria with different pathways of C1 metabolism. *Mikrobiologiya*. 1985;53:885-9.
16. Roslev P, Larsen MB, Jørgensen D, Hesselsoe M. Use of heterotrophic CO₂ assimilation as a measure of metabolic activity in planktonic and sessile bacteria. *Journal of Microbiological Methods*. 2004;59(3):381-93.

17. Giovannelli D, Sievert SM, Hügler M, Markert S, Becher D, Schweder T, et al. Insight into the evolution of microbial metabolism from the deep-branching bacterium, *Thermovibrio ammonificans*. *Elife*. 2017;6.
18. Lombard J, Moreira D. Early evolution of the biotin-dependent carboxylase family. *BMC Evol Biol*. 2011;11:232.
19. Attwood PV. The structure and the mechanism of action of pyruvate carboxylase. *The International Journal of Biochemistry & Cell Biology*. 1995;27(3):231 - 49.
20. Jitrapakdee S, St Maurice M, Rayment I, Cleland WW, Wallace JC, Attwood PV. Structure, mechanism and regulation of pyruvate carboxylase. *Biochem J*. 2008;413:369-87.
21. Owen OE, Kalhan SC, Hanson RW. The key role of anaplerosis and cataplerosis for citric acid cycle function. *Journal of Biological Chemistry*. 2002;277(34):30409-12.
22. Sauer U, Eikmanns BJ. The PEP-pyruvate-oxaloacetate node as the switch point for carbon flux distribution in bacteria. *FEMS Microbiol Rev*. 2005;29(4):765-94.
23. Kleijn RJ, Buescher JM, Le Chat L, Jules M, Aymerich S, Sauer U. Metabolic fluxes during strong carbon catabolite repression by malate in *Bacillus subtilis*. *J Biol Chem*. 2010;285:1587-96.
24. Šantrůčková H, Bird MI, Elhottová D, Novák J, Pícek T, Šimek M, et al. Heterotrophic Fixation of CO₂ in Soil. *Microbial Ecology*. 2005;49(2):218-25.
25. Gleixner G. Soil organic matter dynamics: a biological perspective derived from the use of compound-specific isotopes studies. *Ecological Research*. 2013;28(5):683-95.
26. Nielsen J. Systems biology of metabolism. *Annu Rev Biochem*. 2017;86:245-75.
27. Heux S, Berges C, Millard P, Portais JC, Letisse F. Recent advances in high-throughput ¹³C-fluxomics. *Curr Opin Biotechnol*. 2017;43:104-9.
28. Bacher A, Chen F, Eisenreich W. Decoding biosynthetic pathways in plants by pulse-chase strategies using ¹³CO₂ as a universal tracer. *Metabolites*. 2016;6.
29. Römisch-Margl W, Schramek N, Radykewicz T, Ettenhuber C, Eylert E, Huber C, et al. ¹³CO₂ as a universal metabolic tracer in isotopologue perturbation experiments. *Phytochemistry*. 2007;68:2273-89.
30. Schramek N, Wang H, Römisch-Margl W, Keil B, Radykewicz T, Winzenhörlein B, et al. Artemisinin biosynthesis in growing plants of *Artemisia annua*. A ¹³CO₂ study. *Phytochemistry*. 2010;71:179-87.
31. Ishihara H, Obata T, Sulpice R, Fernie AR, Stitt M. Quantifying protein synthesis and degradation in *Arabidopsis* by dynamic ¹³CO₂ labeling and analysis of enrichment in individual amino acids in their free pools and in protein. *Plant Physiol*. 2015;168:74-93.
32. Cegelski L, Schaefer J. NMR determination of photorespiration in intact leaves using *in vivo* ¹³CO₂ labeling. *J Magn Reson*. 2006;178:1-10.

33. Alonso-Saez L, Galand PE, Casamayor EO, Pedros-Alio C, Bertilsson S. High bicarbonate assimilation in the dark by Arctic bacteria. *ISME J.* 2010;4(12):1581-90.
34. Hesselsoe M, Nielsen JL, Roslev P, Nielsen PH. Isotope labeling and microautoradiography of active heterotrophic bacteria on the basis of assimilation of $^{14}\text{CO}_2$. *Appl Environ Microbiol.* 2005;71(2):646-55.
35. Johnson BT, Romanenko VI. Xenobiotic perturbation of microbial growth as measured by CO_2 uptake in aquatic heterotrophic bacteria. *J Great Lakes Res.* 1984;10:245-50.
36. Miltner A, Kopinke F-D, Kindler R, Selesi D, Hartmann A, Kästner M. Non-phototrophic CO_2 fixation by soil microorganisms. *Plant and Soil.* 2005;269(1-2):193-203.
37. Wegener G, Bausch M, Holler T, Thang NM, Prieto Mollar X, Kellermann MY, et al. Assessing sub-seafloor microbial activity by combined stable isotope probing with deuterated water and ^{13}C -bicarbonate. *Environ Microbiol.* 2012;14(6):1517-27.
38. DeLorenzo S, Brauer SL, Edgmont CA, Herfort L, Tebo BM, Zuber P. Ubiquitous dissolved inorganic carbon assimilation by marine bacteria in the Pacific Northwest coastal ocean as determined by stable isotope probing. *PLoS One.* 2012;7(10):e46695.
39. Yakimov MM, La Cono V, Smedile F, Crisafi F, Arcadi E, Leonardi M, et al. Heterotrophic bicarbonate assimilation is the main process of *de novo* organic carbon synthesis in hadal zone of the Hellenic Trench, the deepest part of Mediterranean Sea. *Environ Microbiol Rep.* 2014;6(6):709-22.
40. Eylert E, Schär J, Mertins S, Stoll R, Bacher A, Goebel W, et al. Carbon metabolism of *Listeria monocytogenes* growing inside macrophages. *Mol Microbiol.* 2008;69(4):1008-17.
41. Lee WN, Byerley LO, Bergner EA, Edmond J. Mass isotopomer analysis: theoretical and practical considerations. *Biol Mass Spectrom.* 1991;20:451-8.
42. Meyer FM, Stülke J. Malate metabolism in *Bacillus subtilis*: distinct roles for three classes of malate-oxidizing enzymes. *FEMS Microbiol Lett.* 2013;339(1):17-22.
43. Glycine, serine and threonine metabolism - *Bacillus subtilis* subsp. *spizizenii* W23 [Internet]. 2019 [cited 26.10.2019]. Available from: https://www.kegg.jp/kegg-bin/highlight_pathway?scale=1.0&map=bss00260&keyword=glycine.
44. Schilling O, Frick O, Herzberg C, Ehrenreich A, Heinzle E, Wittmann C, et al. Transcriptional and metabolic responses of *Bacillus subtilis* to the availability of organic acids: transcription regulation is important but not sufficient to account for metabolic adaptation. *Appl Environ Microbiol.* 2007;73:499-507.
45. Mirouze N, Bidnenko E, Noirot P, Auger S. Genome-wide mapping of TnrA-binding sites provides new insights into the TnrA regulon in *Bacillus subtilis*. *Microbiologyopen.* 2015;4:423-35.
46. Cazzulo JJ, Stoppani AO. Effects of adenosine phosphates and nicotinamide nucleotides on pyruvate carboxylase from baker's yeast. *Biochem J.* 1969;112:755-62.
47. Zhu B, Stülke J. SubtiWiki in 2018: from genes and proteins to functional network annotation of the model organism *Bacillus subtilis*. *Nucleic Acids Res.* 2018;46:D743-D8.

48. Diesterhaft MD, Freese E. Role of pyruvate carboxylase, phosphoenolpyruvate carboxykinase, and malic enzyme during growth and sporulation of *Bacillus subtilis*. *J Biol Chem*. 1973;248:6062-70.
49. Chubukov V, Uhr M, Le Chat L, Kleijn RJ, Jules M, Link H, et al. Transcriptional regulation is insufficient to explain substrate-induced flux changes in *Bacillus subtilis*. *Mol Syst Biol*. 2013;9:709.
50. Kikuchi G, Motokawa Y, Yoshida T, Hiraga K. Glycine cleavage system: reaction mechanism, physiological significance, and hyperglycinemia. *Proc Jpn Acad Ser B Phys Biol Sci*. 2008;84:246-63.
51. Born Timothy L., S. BJ. Structure/function studies on enzymes in the diaminopimelate pathway of bacterial cell wall biosynthesis. *Current Opinion in Chemical Biology*. 1999;3:607-13.
52. Lysine biosynthesis - Reference pathway [Internet]. 2020 [cited 14.02.2020]. Available from: https://www.kegg.jp/kegg-bin/highlight_pathway?scale=1.0&map=map00300&keyword=Diaminopimelate.
53. Jitrapakdee S, Wallace JC. Structure, function and regulation of pyruvate carboxylase. *Biochem J*. 1999;340:1-16.

Uncorrected Proof




Fig. 1 Metabolic routes of ^{13}C -labelled bicarbonate after anaplerotic fixation during growth on different substrates. Scenarios A, B, and C show the expected labelling patterns from $H^{13}CO_3^-$ when using unlabelled glucose (Glc), lactate (Lac), or malate (Mal) as main organic carbon substrates. The bold arrows indicate main carbon fluxes. Red arrows show the respective fluxes from the supplied $H^{13}CO_3^-$ tracer through the metabolic network; blue arrows depict the carbon fluxes from the unlabelled organic substrates, respectively. Metabolites and products marked with a red circle are expected to receive ^{13}C -label originating from $H^{13}CO_3^-$.




Fig. 2 (A) Glucose consumption and biomass production by *B. subtilis* W23 growing in M9 medium containing 1 g/ L glucose. The circles represent the glucose concentration and

the triangles the biomass production over time. **(B) Incorporation of ^{13}C -carbon into microbial biomass by *B. subtilis* W23 growing in M9 medium containing 1 g/ L glucose and 1 g/ L $\text{NaH}^{13}\text{CO}_3^-$.** The diamonds represent the ^{13}C -incorporation into the biomass as determined by EA-IRMS measurements. The depicted values are mean values of three biological replicates. The squares represent the control experiment conducted with unlabelled bicarbonate which shows the natural abundance of ^{13}C -carbon of 1.1 % in the environment. **(C) Incorporation of ^{13}C -carbon into microbial biomass by *B. subtilis* W23 growing in M9 glucose medium containing 1 g/ L $\text{NaH}^{13}\text{CO}_3^-$ during the stationary phase.** The culture was supplied with the tracer 10 h after inoculation. The ^{13}C -abundance of the biomass (depicted as crosses) increased up to 2 %. In a control experiment, no $\text{H}^{13}\text{CO}_3^-$ was added. The ^{13}C -abundance of the biomass (depicted as squares) again mirrored the natural abundance of ^{13}C -carbon in the environment.




Fig. 3 Incorporation of ^{13}C -carbon from $\text{H}^{13}\text{CO}_3^-$ into the biomass of *B. subtilis* W 23 growing in M9 containing unlabelled glucose, lactate or malate as carbon sources, respectively. The control shows the natural abundance of ^{13}C -carbon. The depicted values are mean values of three biological replicates. The error bars represent the calculated standard deviation; in case of the control, the error bars are too small to be visible.




Fig. 4 ^{13}C -Excess (mol%) of the M+1 isotopomers of specific amino acids produced in $\text{H}^{13}\text{CO}_3^-$ -labelling experiments with glucose (A), lactate (B), and malate (C), respectively. Amino acids depicted as diamonds are derived from the TCA cycle. Amino acids depicted as circles are derived from pyruvate. Amino acids depicted as crosses are derived from the gluconeogenesis pathway. The ^{13}C -excess of the amino acids derived from pyruvate and the gluconeogenesis pathway are also displayed with a different scaling to improve visibility.




Fig. 5 Marker ratios of ^{13}C -excess between selected amino acids. **(A)** ^{13}C -excess (TCA cycle-derived amino acids) / ^{13}C -excess (pyruvate-derived amino acids). The coloured bars depict the ratios of the chosen amino acids in three independent labelling experiments with each of the three substrates, glucose (Glc), lactate (Lac) or malate (Mal). The grey bars show the mean values of ^{13}C -ratios for these fractions. The individual ratios as well as the mean values are significantly higher when an active CO_2 -fixation is required in the experiments with Glc and Lac compared to a background or random fixation of CO_2 in the experiment with Mal. p-Values were calculated by Student's t-test (unpaired): $p(\text{Glc/Lac}) = 0.05031$, $p(\text{Glc/Mal}) = 0.00003$, $p(\text{Lac/Mal}) = 0.00006$. **(B)** ^{13}C -excess (gluconeogenesis-derived

amino acids)/ ^{13}C -excess (pyruvate-derived amino acids). The coloured bars depict the ratios of the chosen amino acids when glucose or lactate were the growth substrates, respectively, and the grey bars show again the mean values of ^{13}C -ratios. p-Value was calculated by Student's t-test (unpaired): $p(\text{Glc/Lac}) = 0.00017$ (For detailed calculation see SI excel file F5 m+1 and ratio of label_SI Paper)

Uncorrected Proof




Fig. 6 Metabolic network observed for *B. subtilis* W23 growing in M9 medium containing $\text{H}^{13}\text{CO}_3^-$ as a tracer. The full red circles mark the location of the ^{13}C -carbon atom detected in amino acids. On this basis, the labelling profiles of their respective precursors were reconstructed. The equilibrium between the reactions of the reductive branch of the TCA cycle could transfer the ^{13}C -label from oxaloacetate to malate, fumarate, and succinate. The intrinsic symmetry of fumarate leads to the formation of a 0.5:0.5 mixture of [1- ^{13}C]- and [4- ^{13}C]malate, and [1- ^{13}C]- and [4- ^{13}C]oxaloacetate, respectively. This is indicated by the red half circles.




Fig. 7 Labelling experiment with a natural groundwater community and $\text{H}^{13}\text{CO}_3^-$ as a tracer. (A) Growth curves of the bacterial groundwater communities growing on different concentrations of organic fertiliser. The microorganisms that were supplemented with 30 mL of organic fertiliser are depicted as open circles. The microorganisms that were supplemented with 50 mL of organic fertiliser are depicted as full circles. (B) Growth curve of the combined bacterial groundwater community growing on organic fertiliser and soil extract. (C) ^{13}C -Excess of the M+1 isotopomers of specific amino acids produced by $\text{H}^{13}\text{CO}_3^-$ -labelling experiments. (D) Marker ratios of ^{13}C -excess between selected amino acids: ^{13}C -excess (TCA-cycle-derived amino acids) / ^{13}C -excess (pyruvate-derived amino acids). The coloured bars depict the ratios of the chosen amino acids in the labelling

experiments. The grey bars show the mean values of ^{13}C -ratios for these fractions. p-Values as calculated by student's t-test (unpaired): $p(\text{Glc/Lac}) = 0.05031$, $p(\text{Glc/Mal}) = 0.00003$, $p(\text{Glc/groundwater}) = 0.00002$, $p(\text{Lac/Mal}) = 0.00006$, $p(\text{Lac/groundwater}) = 0.00004$, $p(\text{Mal/groundwater}) = 0.00082$. **(E) Marker ratios of ^{13}C -excess between selected amino acids: ^{13}C -excess (gluconeogenesis-derived amino acids)/ ^{13}C -excess (pyruvate-derived amino acids).** The coloured bars depict the ratios of the chosen amino acids in the labelling experiments and the grey bars show again the mean values of ^{13}C -ratios. p-Values as calculated by student's t-test (unpaired): $p(\text{Glc/Lac}) = 0.00017$, $p(\text{Glc/Mal}) = 0.00002$, $p(\text{Glc/groundwater}) = 0.10712$, $p(\text{Lac/Mal}) = 0.00166$, $p(\text{Lac/groundwater}) = 0.00103$, $p(\text{Mal/groundwater}) = 0.06284$.

Uncorrected Proof