

# Author Query Form

Journal: ALL

Article: 14470

Dear Author,

During the copyediting of your manuscript, the following queries arose.

Please refer to the query reference callout numbers in the page proofs and respond to each by marking the necessary comments using the PDF annotation tools.

Please remember illegible or unclear comments and corrections may delay publication.

Many thanks for your assistance.

**AUTHOR:** Please note that missing content in references have been updated where we have been able to match the missing elements without ambiguity against a standard citation database, to meet the reference style requirements of the journal. It is your responsibility to check and ensure that all listed references are complete and accurate.

Query reference	Query	Remarks
1	AUTHOR: Please check and confirm the edit made in the article title.	
2	AUTHOR: Please confirm that given names (blue) and surnames/family names (vermillion) have been identified correctly.	
3	AUTHOR: Please verify that the linked ORCID identifiers are correct for each author. The ORCID ID for Buters seems to be invalid. Please check and supply the correct ORCID ID for this author.	
4	AUTHOR: Please check and confirm the edit made in the sentence 'The global warming-oriented...tree and grass pollen of breathed air in Europe, as documented in aerobiological studies (4).'	
5	AUTHOR: Please suggest whether the term 'minimalizing' can be changed as 'minimizing' in the sentence 'The optimal slope and...the curve and the data points.'	
6	AUTHOR: Figure 1 has been saved at a low resolution of 111 dpi. Please resupply at 600 dpi. Check required artwork specifications at <a href="https://authorservices.wiley.com/asset/photos/electronic_artwork_guidelines.pdf">https://authorservices.wiley.com/asset/photos/electronic_artwork_guidelines.pdf</a>	
7	AUTHOR: As per journal style, author contributions section has been removed. Please check.	
8	AUTHOR: As per journal style, maximum of 9 references are allowed in Letter to Editor article type. Please check and limit to 9 references in this article.	

# Funding Info Query Form

Please confirm that the funding sponsor list below was correctly extracted from your article: that it includes all funders and that the text has been matched to the correct FundRef Registry organization names. If a name was not found in the FundRef registry, it may not be the canonical name form, it may be a program name rather than an organization name, or it may be an organization not yet included in FundRef Registry. If you know of another name form or a parent organization name for a “not found” item on this list below, please share that information.


FundRef name	FundRef Organization Name
European Academy of Allergy and Clinical Immunology	European Academy of Allergy and Clinical Immunology

# 1 The development of birch pollen seasons over 30 years in Munich, Germany—An EAACI Task Force report\*

To the Editor

4 The global warming-oriented climate change is able to induce changes in the presentation of allergic disorders<sup>1</sup> due to (a) increase in the annual sum of some important airborne allergenic pollen,<sup>2</sup> (b)

increase in the pollen allergenic potency,<sup>3</sup> and (c) earlier shifts in airborne tree and grass pollen of breathed air in Europe, as documented in aerobiological studies.<sup>4</sup> Studies on the period of times with pollen flight do not necessarily correlate directly with symptom



**FIGURE 1** Overview on the start and end of birch pollen season, the number of days  $\geq 100$  birch pollen/m<sup>3</sup> and the total pollen (annual sum) measured at the pollen station Frauenlobstrasse 9-11 in 80337 Munich, Germany from 1988 till 2018, except of 1995. For every plot, the slope of the fitted regression line indicating a (linear) trend, and the statistical significance of the trend (in terms of *P*-value) is mentioned. The number of days  $\geq 100$  birch pollen/m<sup>3</sup> has a statistically significant trend of increase during the studied period. Dashed red line represents the linear regression line; blue line represents a least squares fit; thick light blue line represents the confidence interval for the aforementioned least squares fit

\*For the Members of EAACI Task Force "Definition of clinical-relevant thresholds of allergen-exposure": Oliver Pfaar, Kostas Karatzas, Katharina Bastl, Uwe Berger, Jeroen Buters, Ulf Darsow, Pascal Demoly, Stephen R. Durham, Carmen Galán, Regula Gehrig, Roy Gerth van Wijk, Lars Jacobsen, Nikos Katsifarakis, Ludger Klimek, Annika Saarto, Mikhail Sofiev, Michel Thibaudon, Barbora Werchan, Karl-Christian Bergmann.

burden of hay fever sufferers as long they have not included symptom data from subjects with allergic rhinitis or rhinoconjunctivitis. This shortcoming was taken into account when the new EAACI pollen season definition was developed and published to describe the period of time in which pollen allergy sufferers develop their allergic symptoms.<sup>5</sup>

The definition has been shown to be robust<sup>6</sup> and, even more importantly, reflects the onset and end of symptoms due to airborne birch and grass pollen.<sup>7,8</sup> Hence, using the EAACI definition of pollen season we are now able to better understand changes in the time periods during which hay fever subjects suffer.

In this communication, we report that the EAACI definitions are an appropriate tool for the demonstration of a change in periods of the birch pollen season in Munich, Germany.

The German Pollen Information Service Foundation ([www.pollenstiftung.de](http://www.pollenstiftung.de)) has quantified airborne pollen in Munich since 1988 using a Hirst-type volumetric spore trap.

The data from the pollen trap in Munich from 1988 until 2018, except for 1995, were used to calculate<sup>9</sup> the time of onset and end of birch pollen season over the 30-year period, including the peak pollen days and the total annual sum of birch pollen.

In order to identify trends of the studied pollen season (PS) characteristics, we fitted a linear regression model for each of them and evaluated the statistical significance using a *t* test for the hypothesis that there is no association between the identified trend and our PS indicators.<sup>10</sup>

5 The optimal slope and intersection (coefficients) were calculated by minimizing the sum of squared errors (SSE) between the curve and the data points. We calculated the *P*-value for each plot and used *P* < .05 as the cutoff for the significance level. Furthermore, 95% confidence intervals of the sample distribution were calculated for each of the pollen season.

The analysis using the EAACI definition showed a variable but suggestive trend toward an earlier start and end of birch pollen season, although statistically not significant. We estimated that the mean overall change in the PS start is equal to 6 days, while the change in the PS end is approx. 8 days, over the 30 years of study. The mean length of the PS was equal to 30 days, with a standard deviation equal to approx. 8 days, demonstrating a slight decreasing trend, reflecting a duration of the PS shorter by approx. 2 days in the 30 years of study (Figure 1).

Furthermore, the number of peak pollen days per year (number of days  $\geq 100$  birch pollen/m<sup>3</sup>) was variable but increasingly more frequent (slope = 0.2, *P* < .05) in later years. The annual birch pollen count shows a slight increase over this period.

The knowledge of the earlier beginning of the birch pollen season and the onset of allergic symptoms can serve to adapt drug therapy and the start of allergen immunotherapy against birch pollen allergy.

In conclusion, using the harmonized and recently validated EAACI criteria for birch pollen season definition we demonstrated a tendency to an earlier start of the season over the last 30 years in

Munich, Germany, and a statistically significant increase in the number of days with high pollen counts ( $>100$  birch pollen/m<sup>3</sup>).

## ACKNOWLEDGMENT

The authors thank the team of pollen analysts of the German Pollen Information Service Foundation in Munich.

## CONFLICT OF INTEREST

No conflict of interest declared by KCB, KK, BW, MW, and TT Dr Pfaar reports grants and personal fees from ALK-Abelló, grants and personal fees from Allergopharma, grants and personal fees from Stallergenes Greer, grants and personal fees from HAL Allergy Holding BV/HAL Allergie GmbH, grants and personal fees from Bencard Allergie GmbH/Allergy Therapeutics, grants and personal fees from Lofarma, grants from Biomay, grants from Circassia, grants and personal fees from ASIT Biotech Tools SA, grants and personal fees from Laboratorios LETI/LETIPharma, personal fees from MEDA Pharma/MYLAN, grants and personal fees from Anergis SA, personal fees from Mobile Chamber Experts (a GA2LEN Partner), personal fees from Indoor Biotechnologies, grants from GlaxoSmithKline, personal fees from Astellas Pharma Global, personal fees from EUFOREA, personal fees from ROXALL, personal fees from Novartis, personal fees from Sanofi-Aventis, personal fees from Med Update Europe GmbH, personal fees from streamedup! GmbH, outside the submitted work. Dr Buters reports grants from Government of Bavaria, during the conduct of the study, and personal fees from speaker's fees also on climate change, outside the submitted work.

## FUNDING INFORMATION

The measurements of birch pollen were done by the German Pollen Information Service Foundation. The work of the EAACI TF has been supported by the European Academy of Allergy and Clinical Immunology (EAACI, Zurich, Switzerland).

Karl-Christian Bergmann<sup>1</sup>

Jeroen Buters<sup>2</sup> 

Kostas Karatzas<sup>3</sup> 

Thomas Tasioulis<sup>3</sup>

Barbora Werchan<sup>1</sup>

Matthias Werchan<sup>1</sup>

Oliver Pfaar<sup>4</sup> 

2 3

<sup>1</sup>German Pollen Information Service Foundation, Berlin, Germany

<sup>2</sup>ZAUM-Center of Allergy & Environment, Technical University Munich, Munich, Germany

<sup>3</sup>Environmental Informatics Research Group, Department of Mechanical Engineering, Aristotle University of Thessaloniki, Thessaloniki, Greece

<sup>4</sup>Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Marburg, Germany

## Correspondence

Oliver Pfaar, Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Marburg, Germany.  
Email: oliver@pfaar.org

## ORCID

Jeroen Buters  <https://orcid.org/0000-0003-3581-5472>

Kostas Karatzas  <https://orcid.org/0000-0002-1033-5985>

Oliver Pfaar  <https://orcid.org/0000-0003-4374-9639>

## REFERENCES

1. D'Amato G, Holgate ST, Pawankar R, et al. Meteorological conditions, climate change, new emerging factors, and asthma and related allergic disorders. A statement of the World Allergy Organization. *World Allergy Organ J.* 2015;8(1):25.
2. Ziello C, Sparks TH, Estrella N, et al. Changes to airborne pollen counts across Europe. *PLoS ONE.* 2012;7(4):e34076.
3. Cecchi L, D'Amato G, Annesi-Maesano I. External exposome and allergic respiratory and skin diseases. *J Allergy Clin Immunol.* 2018;141:846-857.
4. Cecchi L, D'Amato G, Ayres JG, et al. Projections of the effects of climate change on allergic asthma: the contribution of aerobiology. *Allergy* 2010;65:1073-1081.
5. Pfaar O, Bastl K, Berger U, et al. Defining pollen exposure times for clinical trials of allergen immunotherapy for pollen-induced rhinoconjunctivitis – an EAACI position paper. *Allergy* 2017;72:713-722.
6. Karatzas K, Riga M, Berger U, Werchan M, Pfaar O, Bergmann KC. Computational validation of the recently proposed pollen season definition criteria. *Allergy* 2018;73:5-7.
7. Karatzas K, Katsifarakis N, Riga M, et al. New European Academy of Allergy and Clinical Immunology definition on pollen season mirrors symptom load for grass and birch pollen-induced allergic rhinitis. *Allergy* 2018;73:1851-1859.
8. Pfaar O, Karatzas K, Bastl K, et al. Pollen season is reflected on symptom load for grass and birch pollen-induced allergic rhinitis in different geographic areas – an EAACI Task Force Report. *Allergy* 2020;75(5):1099-1106.
9. Rojo J, Picornell A, Oteros J. AeRobiology: the computational tool for biological data in the air. *Methods Ecol Evol.* 2019;10:1371-1376.
10. Indrayan A, Malhotra RK. *Medical biostatistics.* New York: Chapman and Hall/CRC; 2018.