

Raaschou-Nielsen Ole (Orcid ID: 0000-0002-1223-0909)  
 Boutron-Ruault Marie-Christine (Orcid ID: 0000-0002-5956-5693)  
 Nagel Gabriele (Orcid ID: 0000-0001-6185-8535)  
 Severi Gianluca (Orcid ID: 0000-0001-7157-419X)  
 So Rina (Orcid ID: 0000-0002-2113-3887)

## Long-term exposure to air pollution and liver cancer incidence in six

### European cohorts

Rina So<sup>1,\*</sup>, Jie Chen<sup>2</sup>, Amar J. Mehta<sup>3,4</sup>, Shuo Liu<sup>1</sup>, Maciej Strak<sup>2,6</sup>, Kathrin Wolf<sup>7</sup>, Ulla A. Hvidtfeldt<sup>8</sup>, Sophia Rodopoulou<sup>9</sup>, Massimo Stafoggia<sup>10,11</sup>, Jochem O. Klompmaker<sup>2,6</sup>, Evangelia Samoli<sup>9</sup>, Ole Raaschou-Nielsen<sup>8,12</sup>, Richard Atkinson<sup>13</sup>, Mariska Bauwelinck<sup>14</sup>, Tom Bellander<sup>10,15</sup>, Marie-Christine Boutron-Ruault<sup>16</sup>, Jørgen Brandt<sup>12,17</sup>, Bert Brunekreef<sup>2</sup>, Giulia Cesaroni<sup>11</sup>, Hans Concin<sup>18</sup>, Francesco Forastiere<sup>19,20</sup>, Carla H. van Gils<sup>21</sup>, John Gulliver<sup>22</sup>, Ole Hertel<sup>23</sup>, Barbara Hoffmann<sup>24</sup>, Kees de Hoogh<sup>25,26</sup>, Nicole Janssen<sup>6</sup>, Youn-hee Lim<sup>1</sup>, Rudi Westendorp<sup>4, 27</sup>, Jeanette T. Jørgensen<sup>1</sup>, Klea Katsouyanni<sup>9,28</sup>, Matthias Ketzel<sup>12, 29</sup>, Anton Lager<sup>30</sup>, Alois Lang<sup>18</sup>, Petter L. Ljungman<sup>10,31</sup>, Patrik K.E. Magnusson<sup>32</sup>, Gabriele Nagel<sup>18,33</sup>, Mette K. Simonsen<sup>34</sup>, Göran Pershagen<sup>10,15</sup>, Raphael S. Peter<sup>33</sup>, Annette Peters<sup>7,35</sup>, Matteo Renzi<sup>11</sup>, Debora Rizzuto<sup>36</sup>, Torben Sigsgaard<sup>37</sup>, Danielle Vienneau<sup>25,26</sup>, Gudrun Weinmayr<sup>33</sup>, Gianluca Severi<sup>16, 38</sup>, Daniela Fecht<sup>39</sup>, Anne Tjønneland<sup>1,8</sup>, Karin Leander<sup>10,31</sup>, Gerard Hoek<sup>2</sup>, Zorana J. Andersen<sup>1</sup>

<sup>1</sup> Section of Environmental Health, Department of Public Health, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark

<sup>2</sup> Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands

<sup>3</sup> Statistics Denmark, Copenhagen, Denmark

<sup>4</sup> Section of Epidemiology, Department of Public Health, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

<sup>6</sup> National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

<sup>7</sup> Institute of Epidemiology, Helmholtz Zentrum München, Neuherberg, Germany

<sup>8</sup> Danish Cancer Society Research Centre, Copenhagen, Denmark

<sup>9</sup> Department of Hygiene, Epidemiology and Medical Statistics, National and Kapodistrian University of Athens, Medical School, Athens, Greece

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/ijc.33743](https://doi.org/10.1002/ijc.33743)

<sup>10</sup> Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

<sup>11</sup> Department of Epidemiology, Lazio Region Health Service / ASL Roma 1, Rome, Italy

<sup>12</sup> Department of Environmental Science, Aarhus University, Roskilde, Denmark

<sup>13</sup> Population Health Research Institute and MRC-PHE Centre for Environment and Health, St George's, University of London, London, United Kingdom

<sup>14</sup> Interface Demography – Department of Sociology, Vrije Universiteit Brussel, Brussels, Belgium

<sup>15</sup> Centre for Occupational and Environmental Medicine, Region Stockholm, Stockholm, Sweden

<sup>16</sup> CESP, UMR 1018, Université Paris-Saclay, Inserm, Gustave Roussy, Villejuif, France

<sup>17</sup> iClimate – Aarhus University interdisciplinary Centre for Climate Change, Roskilde, Denmark

<sup>18</sup> Agency for Preventive and Social Medicine, Bregenz, Austria

<sup>19</sup> Environmental Research Group, School of Public Health, Imperial College, London, United Kingdom

<sup>20</sup> Institute for Biomedical Research and Innovation (IRIB), National Research Council, Palermo, Italy

<sup>21</sup> Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, the Netherlands

<sup>22</sup> Centre for Environmental Health and Sustainability & School of Geography, Geology and the Environment, University of Leicester, Leicester, United Kingdom

<sup>23</sup> Department of Bioscience, Aarhus University, Roskilde, Denmark

<sup>24</sup> Institute of Occupational, Social and Environmental Medicine, Centre for Health and Society, Medical Faculty, Heinrich-Heine-University, Dusseldorf, Germany

<sup>25</sup> Swiss Tropical and Public Health Institute, Basel, Switzerland

<sup>26</sup> University of Basel, Basel, Switzerland

<sup>27</sup> Center for Healthy Aging, University of Copenhagen, Copenhagen, Denmark

<sup>28</sup> Environmental Research Group, School of Public Health, Imperial College London, London, United Kingdom

<sup>29</sup> Global Centre for Clean Air Research (GCARE), University of Surrey, Guildford, United Kingdom

<sup>30</sup> Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden

<sup>31</sup> Department of Cardiology, Danderyd University Hospital, Stockholm, Sweden

<sup>32</sup> Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

<sup>33</sup> Institute of Epidemiology and Medical Biometry, Ulm University, Ulm, Germany

<sup>34</sup> Department of Neurology and Parker Institute, Bispebjerg and Frederiksberg Hospital, Denmark

<sup>35</sup> Ludwig-Maximilians University, Munich, Germany

<sup>36</sup> Department of Neurobiology, Care Sciences, and Society, Karolinska Institutet and Stockholm University and The Stockholm Gerontology Research Center, Stockholm, Sweden

<sup>37</sup> Department of Public Health, Environment Occupation and Health, Danish Ramazzini Centre, Aarhus University, Aarhus, Denmark

<sup>38</sup> Department of Statistics, Computer Science and Applications “G. Parenti” (DISIA), University of Florence, Italy

<sup>39</sup> UK Small Area Health Statistics Unit, MRC Centre for Environment and Health, School of Public Health, Imperial College London, London, UK

**\*Corresponding author:** Rina So, Section of Environmental Health, Department of Public Health, Faculty of Health and Medical Sciences, University of Copenhagen, Øster Farimagsgade 5, Copenhagen, Denmark. E-mail: [rina.so@sund.ku.dk](mailto:rina.so@sund.ku.dk); [rinani616@gmail.com](mailto:rinani616@gmail.com)

Keywords: liver cancer incidence, air pollution, particulate matter; cohort study.

Abbreviations: BC, black carbon; CEANS, Cardiovascular Effects of Air Pollution and Noise in Stockholm (cohort); CI, confidence interval; CPS-II, Cancer Prevention Study II; DCH, Diet, Cancer, and Health (cohort); DNC, Danish Nurse Cohort; E3N, Etude Epidémiologique auprès de femmes de la Mutuelle Générale de l'Education Nationale (cohort); ELAPSE, Effects of Low-Level Air Pollution (project); EPIC-MORGEN, EPIC-Monitoring Project on Risk Factor (cohort); EPIC-NL, Dutch European Investigation into Cancer and Nutrition (cohort); EPIC-PROSPECT, EPIC-Chronic Diseases in the Netherlands; ESCAPE, European Study of Cohorts for Air Pollution Effects (project); HR, hazard ratio; ICD-9, International Classification of diseases 9<sup>th</sup> version; ICD-10, International Classification of diseases 10<sup>th</sup> version; MAPLE, Mortality–Air Pollution Associations in Low-Exposure

Environments (project); NO<sub>2</sub>, nitrogen dioxide; NO<sub>x</sub>, nitrogen oxides; O<sub>3</sub>, ozone; PM, particulate matter; PM<sub>2.5</sub>, particulate matter with diameter < 2.5 μm; SALT, Stockholm Screening Across the Lifespan Twin study (cohort); SDPP, Stockholm Diabetes Prevention Program (cohort); SES, socioeconomic status; Sixty, Stockholm cohort of 60-year-olds (cohort); SNAC-K, Swedish National Study on Aging and Care in Kungsholmen (cohort); VHM&PP, Vorarlberg Health Monitoring and Prevention Programme (cohort).

The article category: Cancer Epidemiology

**Novelty and Impact:** While air pollution is classified as carcinogenic and risk factor for lung cancer, the evidence for cancers in other organs is limited. In the pooled analysis of six European cohorts with 330,064 subjects, the authors detected associations between air pollution and liver cancer incidence, even at low levels, below current EU standards. Corroborating results from several earlier substantially smaller studies, this work brings solid evidence that air pollution is a risk factor for liver cancer.

## Abstract

Particulate matter air pollution and diesel engine exhaust have been classified as carcinogenic for lung cancer, yet few studies have explored associations with liver cancer. We used six European adult cohorts which were recruited between 1985 and 2005, pooled within the 'Effects of low-level air pollution: A study in Europe' (ELAPSE) project, and followed for the incidence of liver cancer until 2011 to 2015. The annual average exposure to nitrogen dioxide (NO<sub>2</sub>), particulate matter with diameter < 2.5 µm (PM<sub>2.5</sub>), black carbon (BC), warm-season ozone (O<sub>3</sub>), and eight elemental components of PM<sub>2.5</sub> (copper, iron, zinc, sulfur, nickel, vanadium, silicon, potassium) were estimated by European-wide hybrid land-use regression models at participants' residential addresses. We analyzed the association between air pollution and liver cancer incidence by Cox proportional hazards models adjusting for potential confounders. Of 330,064 cancer-free adults at baseline, 512 developed liver cancer during a mean follow-up of 18.1 years. We observed positive linear associations between NO<sub>2</sub> (hazard ratio, 95% confidence interval: 1.17, 1.02-1.35 per 10 µg/m<sup>3</sup>), PM<sub>2.5</sub> (1.12, 0.92-1.36 per 5 µg/m<sup>3</sup>), and BC (1.15, 1.00-1.33 per 0.5 10<sup>-5</sup>/m) and liver cancer incidence. Associations with NO<sub>2</sub> and BC persisted in two-pollutant models with PM<sub>2.5</sub>. Most components of PM<sub>2.5</sub> were associated with the risk of liver cancer, with the strongest associations for sulfur and vanadium, which were robust to adjustment for PM<sub>2.5</sub> or NO<sub>2</sub>. Our study suggests that ambient air pollution may increase the risk of liver cancer, even at concentrations below current EU standards.

## Introduction

Ambient air pollution is a major environmental stressor, posing a huge health burden related to increased risk of cardiometabolic, respiratory disease, and lung cancer<sup>1</sup>. A number of components presented in air pollution are carcinogenic, including polycyclic aromatic hydrocarbons, volatile organic compounds, and other heavy metals<sup>2</sup>. Particulate matter (PM)<sup>2</sup> and diesel engine exhaust<sup>3</sup> are classified as carcinogenic to humans, largely based on literature related to lung cancer<sup>2,4</sup>. However, the epidemiological evidence on air pollution and cancers other than lung cancer remains limited and inconclusive<sup>5</sup>.

Primary liver or hepatic cancer is the second leading cause of cancer death for men and the sixth for women, accounting for nearly 782,000 deaths (8.2% of all cancer deaths) globally in 2018<sup>6</sup>. Alcohol use, cigarette smoking, and Hepatitis B and C virus infections are the main risk factors<sup>7</sup>. Several plausible biological mechanisms support a link between ambient air pollution and liver cancer. Exposure to PM with diameter < 2.5  $\mu\text{m}$  (PM<sub>2.5</sub>) in mice led to liver fibrosis as well as non-alcoholic steatohepatitis-like phenotype<sup>8,9</sup>, an increasingly important etiology of liver cancer<sup>10</sup>. Exposure to diesel exhaust in rats caused oxidative stress with DNA damage, apoptosis, and upregulation of DNA repair in the liver<sup>11,12</sup>. Inhalation of particles can result in gastrointestinal exposure through the mucociliary clearance from the airways<sup>13</sup> or cross the alveolar-capillary barrier and reach the liver via the circulatory system<sup>14,15</sup>. In several human studies<sup>16-19</sup>, air pollution has been associated with increased serum levels of hepatic enzymes such as  $\gamma$ -glutamyltranspeptidase, aspartate aminotransferase, and alanine transaminase, markers of liver damage usually caused by inflammation, the main mechanism by which air pollution induces adverse health effects<sup>20</sup>.

There are only five epidemiological studies on long-term exposure (i.e., mean air pollution exposures of one year or more) to air pollution and liver cancer with somewhat mixed results<sup>19,21-24</sup>. A cohort study from Taiwan, with 22,062 subjects and 464 liver cancer cases, detected an association with PM<sub>2.5</sub> and found that elevated serum alanine transaminase levels mediated this association<sup>19</sup>. A study in the Danish Diet, Cancer, and Health (DCH) cohort (54,160 adults, 57 cases) reported an association with traffic density within 200m of the residence, but not with nitrogen oxides (NO<sub>x</sub>)<sup>22</sup>. A study in four

European cohorts (174,770 adults, 279 cases) which took part in ‘The European Study of Cohorts for Air Pollution Effects’ (ESCAPE) project, found a positive but statistically non-significant association with PM<sub>2.5</sub> and nitrogen dioxide (NO<sub>2</sub>)<sup>23</sup>. The American Cancer Prevention Study II (CPS-II) cohort, with 623,048 subjects and 1,003 liver cancer deaths, found no association with PM<sub>2.5</sub>, NO<sub>2</sub>, or ozone (O<sub>3</sub>)<sup>24</sup>. A US study (ecological study at a county level) with 56,245 cases of hepatocellular carcinoma, the most common histological type of liver cancer, detected a strong positive association with PM<sub>2.5</sub><sup>21</sup>. Only ESCAPE study had data on PM elemental components and reported associations with sulfur, silicon, nickel, and iron components of PM<sub>2.5</sub><sup>23</sup>.

Within the “Effects of Low-level Air Pollution: a Study in Europe” (ELAPSE) collaboration, which built on ESCAPE cohorts, we aimed to examine the association between long-term exposure to air pollution and liver cancer incidence and identify relevant sources by analyzing eight specific elements of PM<sub>2.5</sub>. In contrast to the ESCAPE project, which analyzed the individual cohort separately in a standardized way and applied meta-analysis, we performed a pooled data analysis, applied a European-wide harmonized air pollution exposure assessment, and had additional years of follow-up, providing enhanced statistical power to examine the association between air pollution and liver cancer.

## Materials and methods

### Study population

We used the framework of the ELAPSE project, under which nine European cohorts were pooled to study health effects related to low-level air pollution<sup>25</sup>, stored on a secure server in Utrecht, and made available for remote analyses.

Of the nine pooled cohorts from the ELAPSE project, we used six from five European countries, which had information on follow-up for liver cancer incidence available: 1) ‘Cardiovascular Effects of Air Pollution and Noise in Stockholm’ (CEANS) from Stockholm county of Sweden, which included four sub-cohorts: ‘Swedish National Study on Aging and Care in Kungsholmen’ (SNAC-K)<sup>26</sup>, ‘Stockholm Screening Across the Lifespan Twin study’ (SALT)<sup>27</sup>, ‘Stockholm cohort of 60-year-olds’

(Sixty)<sup>28</sup>, and ‘Stockholm Diabetes Prevention Program’ (SDPP)<sup>29</sup>; 2) DCH<sup>30</sup> from Copenhagen and Aarhus of Denmark; 3) ‘Danish Nurse Cohort’ (DNC)<sup>31</sup> from entire Denmark, which included two sub-cohorts from recruitment rounds in 1993 and 1999; 4) ‘Dutch European Investigation into Cancer and Nutrition’ (EPIC-NL)<sup>32</sup> from four cities in the Netherland, consisting of ‘EPIC-Monitoring Project on Risk Factors’ (EPIC-MORGEN) and ‘EPIC-Chronic Diseases in the Netherlands’ (EPIC-PROSPECT); 5) ‘Etude Epidémiologique auprès de femmes de la Mutuelle Générale de l’Education Nationale’ (E3N)<sup>33</sup> from entire France; and 6) ‘Vorarlberg Health Monitoring and Prevention Programme’ (VHM&PP)<sup>34</sup> from Vorarlberg, Austria. Cohorts were recruited between 1985 and 2005 and were followed until 2011 to 2015. Individual-level information on smoking, employment status, alcohol use, education level, and area-level socioeconomic status (SES) have been harmonized across the cohorts. A more detailed description of the cohorts included in this analysis can be found in the Supplementary material page 1-7.

#### Liver cancer definition

We obtained the cancer diagnosis data from national and state cancer registries, except for E3N, where cancer was defined from biannual questionnaires self-reports confirmed by oncologist review of pathological reports or from death certificates. We defined liver cancer incidence as the first diagnosis of primary cancer in liver during the follow-up, according to the International Classification of diseases 9<sup>th</sup> version (ICD-9) or 10<sup>th</sup> version (ICD-10) code (155 for ICD-9 and C22 for ICD-10)<sup>23</sup>, and excluded persons with any cancer diagnosis before cohort baseline.

#### Air pollution exposure assessment

As our main exposure assessment, we used Europe-wide hybrid land-use regression (LUR) models at a fine spatial scale (100 m × 100 m grids) to estimate annual mean exposure to air pollutants (NO<sub>2</sub>, PM<sub>2.5</sub>, black carbon [BC], and O<sub>3</sub> [warm-season]) and eight elemental components of PM<sub>2.5</sub> for the year 2010 at the participants’ residential addresses of the baseline, described in detail elsewhere<sup>35,36</sup>.



The models for NO<sub>2</sub>, PM<sub>2.5</sub>, black carbon [BC], and O<sub>3</sub> [warm-season])<sup>35</sup> were developed by supervised linear regression (SLR), based on the European Environment Agency (EEA) AirBase daily concentration data for 2010 for PM<sub>2.5</sub>, NO<sub>2</sub>, and O<sub>3</sub>, and ESCAPE monitoring data for BC, which was not available from EEA. For annual estimates of BC, PM<sub>2.5</sub> absorbance data based on reflectance measurement of the filters during 2009 and 2010 were used and treated as 2010 annual mean concentrations. The annual warm-season average concentrations of O<sub>3</sub> were calculated based on the maximum running 8-hour means for each day. The input data for the LUR models included land use and traffic data, satellite observations, and dispersion model estimates. Ordinary kriging was used to additionally explain the residuals of spatial variation from the LUR model. The models explained spatial variation in the measured concentration well; the R<sup>2</sup> for NO<sub>2</sub>, PM<sub>2.5</sub>, BC, and O<sub>3</sub> was 0.59, 0.72, 0.54, and 0.69, respectively.

The models for PM<sub>2.5</sub> components were developed based on measurement data from the ESCAPE monitoring campaigns from 2008 to 2011 by two methods: supervised linear regression and random forest<sup>36</sup>. While model performance for explaining within-area variability in measured concentration was similar in two, the random forest method was better performed in explaining the overall variability of pollutant concentration levels across Europe than the supervised linear regression method. Eight elements were a priori selected based on their toxicity and representation of major pollution sources: copper, iron, and zinc mainly from non-tailpipe traffic emissions (i.e., brake and tire wear), sulfur from long-range transport of secondary inorganic aerosols from sulfur-containing fossil fuels combustion, nickel and vanadium from coal or mixed oil burning in buildings/ships, silicon from crustal dust, and potassium from biomass burning<sup>36</sup>.

As a sensitivity analysis, we also assessed the annual mean levels of NO<sub>2</sub>, PM<sub>2.5</sub>, BC, and O<sub>3</sub> for each year from recruitment to end of follow-up by a back-extrapolation method incorporating residential history (available only for DCH, VHM&PP, CEANS, and EPIC-NL), as described elsewhere in detail<sup>25</sup>. Briefly, we back-extrapolated the exposure estimates for the year 2010 from the LUR model, using both a difference and a ratio method with the Danish Eulerian Hemispheric Model (DEHM)<sup>37</sup> because the European Environment Agency (EEA) AirBase data did not provide continuous measurement of

Accepted Article

monitoring data during the study period. DEHM data was complete with giving annual averages for all four pollutants at 26 x 26 km spatial resolution across Europe at least back to 1990 and covered all of the study area (downscaled from the original 50 x 50 km resolution using bilinear interpolation). The differences or the ratios of exposure levels between each year and 2010, estimated from DEHM models, were calculated larger spatial scale of NUTS-1 (Nomenclature of territorial units for statistics), allowing different spatial trends within Europe (for DCH, VHM&PP, CEANS, each cohort considered as one region, and EPIC-NL has four regions), and were added or multiplied the exposure estimates for the year 2010 from the LUR model at each residential. When residential history was incorporated, if someone moved within the same NUTS-1 region, then the ratio or difference values for each year after moving are the same as before.

Additionally, as sensitivity analyses, we used the NO<sub>2</sub>, PM<sub>2.5</sub>, and BC estimates from models developed within the ESCAPE, which were developed for each study area<sup>38,39</sup>, and the PM<sub>2.5</sub> from the Canadian 'Mortality–Air Pollution Associations in Low-Exposure Environments' (MAPLE) project<sup>40</sup>.

#### Statistical analysis

We used stratified Cox proportional hazards models with age as the underlying time scale to examine the association between air pollution and liver cancer incidence. The start of follow-up was the participants' age at cohort entry, and the end of follow-up was the age at the time of the diagnosis of liver cancer (event), the first occurrence of any other cancer, date of death, date of emigration, loss to follow-up, or the end of follow-up, whichever came first. We included the sub-cohort indicator as strata to account for baseline hazard heterogeneity across the cohorts.

We included one air pollutant at a time as a continuous variable and evaluated the association with liver cancer incidence with increasing adjustment for variables chosen a priori: Model 1 included age (time axis), sex (strata), sub-cohort (strata), and the cohort baseline year; Model 2 additionally included smoking (never, former, current) and employment status (employed, other); and Model 3 (main model) additionally included neighborhood SES (mean income in 2001). Estimates for main pollutants were expressed as hazard ratio (HR) with 95% confidence interval (95% CI) for increments

of 10  $\mu\text{g}/\text{m}^3$  for  $\text{NO}_2$ , 5  $\mu\text{g}/\text{m}^3$  for  $\text{PM}_{2.5}$ , 0.5  $10^{-5}/\text{m}$  for BC, 10  $\mu\text{g}/\text{m}^3$  for  $\text{O}_3$ , and interquartile range increase for  $\text{PM}_{2.5}$  components.

To evaluate the shape of the exposure-response function between air pollutants and liver cancer incidence, we applied natural cubic splines with two degrees of freedom, which was selected based on the lowest Akaike Information Criteria (AIC) among various degrees of freedom (between 2 and 4) (AIC results not shown). To investigate the associations below the current air quality standards, we additionally applied the main model (model 3) to subsets where we only included participants with concentrations below a certain value. We evaluated cut-points of 40 (the WHO guideline and the EU standard), 30, and 20  $\mu\text{g}/\text{m}^3$  for  $\text{NO}_2$ , 25 (the EU standard), 20, 15, 12 (the US-EPA NAAQS), and 10 (the WHO guideline)  $\mu\text{g}/\text{m}^3$  for  $\text{PM}_{2.5}$ , 3, 2.5, 2, 1.5, and 1.0  $10^{-5}/\text{m}$  for BC, and 120, 100, and 80  $\mu\text{g}/\text{m}^3$  for  $\text{O}_3$ .

We fitted two-pollutant models for  $\text{NO}_2$ ,  $\text{PM}_{2.5}$ , BC, and  $\text{O}_3$  to examine the effects of pollutants independently from each other. For  $\text{PM}_{2.5}$  components, we fitted two-pollutant models with  $\text{PM}_{2.5}$  and  $\text{NO}_2$  as the second pollutant to assess whether associations with the component remained after adjusting for generic  $\text{PM}_{2.5}$  and  $\text{NO}_2$ , which is a marker for traffic tailpipe emissions and other fossil fuel combustion sources. The latter is especially important for the non-tailpipe components of copper, iron, and zinc.

We investigated effect modification of the associations between air pollutants and liver cancer by age (< 65 years,  $\geq$  65 years), alcohol intake (low: <4g/day, medium: 4-15 g/day, high: >15 g/day), and smoking status (never, former, current), by including an interaction term in the model and testing with the likelihood ratio test.

We performed several sensitivity analyses: 1) In order to account for temporal variation and spatial trend in air pollution and residential mobility, we applied the time-varying analysis with back-extrapolated time-varying exposure for  $\text{NO}_2$ ,  $\text{PM}_{2.5}$ , BC, and  $\text{O}_3$  (excluding DNC and E3N) with controlling of time trend (strata per a year or 5-year of follow-up time). 2) To examine the robustness of results to using different exposure metrics, we incorporated the main model (model 3) for the  $\text{NO}_2$ ,  $\text{PM}_{2.5}$ , and BC from the ESCAPE model (excluding DNC and E3N) and for the  $\text{PM}_{2.5}$  from MAPLE. 3)

To investigate the impact of further adjustment of potential confounders, which were not available in VHM&PP, Sixty, and SNAC-K, we applied the main model with and without the further adjustment to the subsets of the pooled cohort with the available information on potential confounders, including educational level (low: primary school or less, medium: up to secondary school or equivalent, or high: university degree or more) and alcohol intake (Low: <4g/day, Medium:4-15 g/day, or High: ≥15 g/day).

4) To evaluate the impact of an individual cohort on the association, we applied the main model to the subsets of data, excluding one cohort at a time.

All statistical tests were two-sided, and p-values of < 0.05 were considered statistically significant. We performed all analyses and graphical presentations in R, version 3.4.0, with common R scripts developed within the ELAPSE project.

## Results

### Description of the study population and exposure

Of the total of 367,404 participants from six pooled cohorts, we excluded 3,348 with missing information on the date of start or end of follow-up, 10,446 with cancer before enrollment, 136 with missing information on the prevalent cancer status, 1,830 with missing air pollution exposure data, and 21,580 with missing data on the individual- and area-level risk factors, leaving 330,064 participants for the final analysis. The number of excluded subjects in each (sub) cohort is presented in the Supplementary material page 1-7.

Over a mean follow-up time of 18.1 years (5,971,185 person-years), 512 participants developed liver cancer. Compare to those free of liver cancer at the end of follow-up, those who developed liver cancer were older and more likely to be men, current or former smokers, unemployed, moderate or high alcohol drinkers, highly educated, and live in the higher-income neighborhood at baseline (Table 1). The mean concentrations of NO<sub>2</sub> and PM<sub>2.5</sub> for the year 2010 at the residential address of baseline ranged from 19.8 (in CEANS) to 35.1 µg/m<sup>3</sup> (in EPIC-NL) for NO<sub>2</sub>, and 8.1 (in CEANS) to 17.5 µg/m<sup>3</sup> (in EPIC-NL) for PM<sub>2.5</sub>, respectively, which were well below the current EU standards of 40 µg/m<sup>3</sup> for NO<sub>2</sub>

and 25  $\mu\text{g}/\text{m}^3$  for  $\text{PM}_{2.5}$  (detailed descriptive statistics on air pollution levels for each cohort are provided in the supplementary material page 1-7). Varying levels of exposure were observed between the individual cohorts with generally lower  $\text{PM}_{2.5}$  and BC in northern countries (Figure S1 in Supplementary material). For  $\text{PM}_{2.5}$ , exposure contrast within cohorts was smaller than for BC and  $\text{NO}_2$ . BC and  $\text{NO}_2$  were highly correlated in all cohorts (The mean of cohort-specific Pearson correlations is 0.83), whereas  $\text{PM}_{2.5}$  was moderately to highly correlated with BC and  $\text{NO}_2$  (The mean of cohort-specific Pearson correlations is 0.57 with BC, 0.62 with  $\text{NO}_2$ . For the correlation per each cohort, see Table S1 in Supplementary material).  $\text{O}_3$  was negatively correlated with  $\text{PM}_{2.5}$  and especially with  $\text{NO}_2$  and BC (the mean of correlations is -0.38, -0.64, and -0.58 for  $\text{PM}_{2.5}$ ,  $\text{NO}_2$ , and BC, respectively).

#### Association between liver cancer incidence and air pollution exposure

We found positive associations of all four main pollutants with the risk of liver cancer in Model1, which attenuated after adjustment for smoking status and employment status in Model2 (Table 2). In a fully adjusted model, we detected an association between  $\text{NO}_2$  and liver cancer incidence with a HR of 1.17 (95% CI 1.02-1.35) per 10  $\mu\text{g}/\text{m}^3$ , while associations with  $\text{PM}_{2.5}$  (HR: 1.12 [95% CI 0.92-1.36] per 5  $\mu\text{g}/\text{m}^3$ ) and BC (HR: 1.15 [95% CI 1.00-1.33] per 0.5  $10^{-5}/\text{m}$ ) were statistically non-significant. The modest attenuation in HRs for  $\text{NO}_2$ ,  $\text{PM}_{2.5}$ , and BC from Model 1 to Model 2 was mainly due to adjustment for smoking, while the inclusion of neighborhood SES in the model modestly increased HRs. However, differences in HRs between models 1, 2, and 3 were small, suggesting limited confounding. A statistically significant negative association was observed with  $\text{O}_3$ .

In the two pollutant models, the associations for  $\text{NO}_2$  remained positive after adjusting for  $\text{PM}_{2.5}$  or BC (HR: 1.19 [95% CI 1.00 to 1.43] and 1.22 [95% CI] 0.87 to 1.70, respectively), while those for  $\text{PM}_{2.5}$  attenuated to unity after adjusting for  $\text{NO}_2$  or BC. The HRs for BC attenuated to unity after adjusting for  $\text{NO}_2$  and remained unchanged after adjusting for  $\text{PM}_{2.5}$  (HR: 1.16 [95% CI: 0.96 to 1.41]). Furthermore, the associations with  $\text{NO}_2$ ,  $\text{PM}_{2.5}$ , and BC attenuated to unity after adjustment  $\text{O}_3$  but remained unchanged for  $\text{O}_3$ , possibly reflecting negative correlations between  $\text{O}_3$  and other urban or

traffic-related pollutants than NO<sub>2</sub> and BC. Due to the high correlation between BC and NO<sub>2</sub>, as well as NO<sub>2</sub> and O<sub>3</sub>, these two pollutant models should be interpreted with caution.

We observed no deviations from linearity for the associations with PM<sub>2.5</sub>, NO<sub>2</sub>, and BC (Figure 1). Associations with NO<sub>2</sub> and PM<sub>2.5</sub> persisted below current EU standards of 40 and 25 µg/m<sup>3</sup> for NO<sub>2</sub> and PM<sub>2.5</sub>, respectively, but leveled off at below 30 µg/m<sup>3</sup> for NO<sub>2</sub> and 12 µg/m<sup>3</sup> for PM<sub>2.5</sub> (Table 3). The associations with BC were persisted in the subset with the concentration below 1.5 10<sup>-5</sup>/m.

The associations between NO<sub>2</sub> and liver cancer were statistically significantly stronger in older participants, while no interaction was detected with alcohol intake and smoking status (Table S2 in Supplementary material).

Observed associations with NO<sub>2</sub>, PM<sub>2.5</sub>, and BC were robust to including time-varying air pollution concentrations and control for time trends (Table S3 in Supplementary material). Analyses using alternative air pollution exposure estimates from ESCAPE and MAPLE project showed stronger associations with liver cancer, but presented the overlapped confidence intervals (Table S4 and Table S5 in Supplementary material) compared to those with ELAPSE exposure model.

Associations were also robust to additional adjustments for education level or alcohol intake (Table S6 in Supplementary material) and to the exclusion of one cohort at a time except for association with PM<sub>2.5</sub>, which attenuated to unity after exclusion of VHM&PP (Table S7 in Supplementary material).

#### Association between liver cancer incidence and PM<sub>2.5</sub> components exposure

Single pollutant models for PM<sub>2.5</sub> components estimated by supervised linear regression showed statistically significant HRs for almost all components, with the strongest associations with sulfur and vanadium (Figure 2; Table S8 in Supplementary material). For PM<sub>2.5</sub> components estimated with the random forest method, weaker associations were observed than those with the supervised linear regression method for all components, statistically significant only for sulfur and vanadium (Table S8

in Supplementary material). Associations were mostly robust (slightly attenuated or enhanced) to adjustment for PM<sub>2.5</sub> and NO<sub>2</sub> (Figure 2).

## Discussion

In this pooled analysis of six cohorts from five European countries, we detected associations between long-term exposure to NO<sub>2</sub>, PM<sub>2.5</sub>, and BC and liver cancer incidence. The exposure-response curves were linear for all three pollutants, and the associations persisted below the current EU standards for NO<sub>2</sub> and PM<sub>2.5</sub>. We found associations with sulfur and vanadium components of PM<sub>2.5</sub>.

Previous studies on air pollution and liver cancer generally reported associations, though most not statistically significant<sup>19,21-24</sup>. In our study, HR per 10 µg/m<sup>3</sup> increase in NO<sub>2</sub> was 1.17 (95% CI 1.02-1.35), stronger but comparable to that of 1.10 (95% CI 0.93-1.30) from the ESCAPE<sup>23</sup>. The DCH cohort study<sup>22</sup> reported HR of 1.66 (95% CI 0.70-3.94) per 100 µg/m<sup>3</sup> increase in NO<sub>x</sub>, and the American CPS-II study<sup>24</sup> found no association between NO<sub>2</sub> and liver cancer mortality (HR: 1.03 [95% CI 0.93-1.14] per 6.5 ppb [~12 µg/m<sup>3</sup>] increase in NO<sub>2</sub>). In the American CPS-II study<sup>24</sup>, only primary liver cancer death was included, and since liver cancer is a highly fatal cancer type, results from this study are comparable with those from ours on liver cancer incidence. With re-calculated HRs of each study per the same unit as ours (per 5 µg/m<sup>3</sup> for PM<sub>2.5</sub>) for the comparison, our findings of the association between PM<sub>2.5</sub> and liver cancer with a HR of 1.12 (95% CI 0.92-1.36) was weaker than one reported in ESCAPE study (HR: 1.34 [95% CI 0.76-2.35]), comparable to those reported in the US study<sup>21</sup> with a HR of 1.12 (95% CI 1.03-1.21), and stronger than Pan et al.<sup>19</sup> study with a HR of 1.08 (95% CI 0.98-1.17) and the American CPS-II study<sup>24</sup> with a mortality rate ratio of 1.06 (95% CI 0.93-1.18). Our finding of an association between BC and liver cancer incidence with a HR of 1.15 (95% CI 1.00-1.33) per 0.5 10<sup>-5</sup>/m showed a more precise estimate compared to the ESCAPE finding with PM<sub>2.5</sub> absorbance (BC equivalent) with a HR of 1.21 (95% CI 0.68-2.15) per 10<sup>-5</sup>/m increase. Two pollutant models showed robust associations with NO<sub>2</sub> and BC after adjustment of PM<sub>2.5</sub>, indicating the relevance of traffic emission-related pollution for liver cancer. However, we cannot determine whether NO<sub>2</sub> per se or the associated gaseous pollutants and particles from local combustion sources are responsible for the



observed association. Finally, we found an inverse association between O<sub>3</sub> and liver cancer, whereas the only other study on O<sub>3</sub> and liver cancer, the American CPS-II study, found no association<sup>24</sup>. Despite the varying size and statistical power to detect associations with this rare cancer type, all studies to date on NO<sub>2</sub>, PM<sub>2.5</sub>, and BC, report HRs above 1.

We present novel findings on the association between specific elemental components of PM<sub>2.5</sub> and liver cancer. A single previous study<sup>23</sup> on elemental components in PM and liver cancer incidence, detected associations with sulfur, silicon, nickel, and iron components of PM<sub>2.5</sub>. Our association with the sulfur component of PM<sub>2.5</sub> is consistent with this studies<sup>23</sup> and indicates the possible relevance of long-range transported secondary inorganic aerosols from sulfur-containing fossil fuels combustion. Furthermore, the strong association with the vanadium component of PM<sub>2.5</sub> is in line with the large population-based cohort study in Rome<sup>41</sup> showing a strong association between the vanadium component of PM<sub>10</sub> and liver cirrhosis, as well as experimental studies<sup>42,43</sup> showing the association of inhaled vanadium with oxidative stress and cell alteration suggestive for liver regeneration. Still, overall evidence on which components and sources of air pollution are most relevant for liver cancer development is premature and demands attention in future research.

Our main strength is the large sample size obtained by pooling six European cohorts with a long follow-up over 18 years and sufficient statistical power to examine the association between air pollution and this rare cancer type. Furthermore, we benefited from the European-wide air pollution exposure model which provided comparable air pollution data over the six European cohorts, detailed data on relevant confounders, and liver cancer incidence information from cancer registries with high validity. Limitations include lack of information on occupational exposures such as benzene (one of the hepatotoxic chemicals), alcohol intake, and hepatitis B or C infection status. Additional adjustment for alcohol intake (Table S6 in Supplementary material) did not affect the main results. Furthermore, we adjusted for SES, a strong determinant of hepatitis B or C infection<sup>44,45</sup>. Furthermore, the differences in the two modeling approaches used in ESCAPE and ELAPSE projects may explain the difference in risk estimates in the sensitivity analysis (Table S4). Briefly, unlike our main assessment method of Europe-wide hybrid LUR model, the ESCAPE model was the LUR model developed for each study area, using



local predictor data. The discrepancy in distributions of air pollution exposure concentrations between two exposure assessment methods can be checked elsewhere<sup>25</sup>.

In conclusion, our study suggests that long-term exposure to air pollution may increase the risk of liver cancer, even at concentrations below current EU standards, and adds evidence of detrimental health effect of air pollution on cancers other than lung cancer

### **Funding**

This work was supported by funding from Health Effects Institute Research Agreement (#4954-RFA14-3/16-5-3) and by the the Novo Nordisk Foundation Challenge Programme (#NNF17OC0027812). The funders had no role in the design of the study; the collection, analysis, and interpretation of the data; the writing of the manuscript; and the decision to submit the manuscript for publication.

### **Conflict of Interest**

Marie-Christine Boutron-Ruault has a potential personal conflict of interest regarding two conferences sponsored by pharmaceutical companies as follows: 1) 03/07/2020 - 30/07/2020 MAYOLI-SPINDLER: Symposium: Pancreatology in practice in 2020 e-JFHOD 2020 Conference « Why do I see more and more pancreatic cancers? »; 2) 04/12/2020 - 04/12/2020- GILEAD e-conference Weight gain and HIV infection in 2020.

The other authors have no conflicts of interest to disclose.

### **Disclaimers**

The research described in this article was conducted under contract to the Health Effects Institute (HEI), an organization jointly funded by the United States Environmental Protection Agency (EPA) (Assistance Award No. R-82811201) and certain motor vehicle and engine manufacturers. The contents of this article do not necessarily reflect the views of HEI, or its sponsors, nor do they necessarily reflect the

views and policies of the EPA or motor vehicle and engine manufacturers.

### **Data Availability Statement**

The exposure maps are available on request from Dr Kees de Hoogh (c.dehoogh@swisstph.ch). The ELAPSE study protocol is available at <http://www.elapseproject.eu/>. Further information and a detailed statistical analysis plan is available on reasonable request from the corresponding author (rina.so@sund.ku.dk).

### **Ethics statement**

All original cohort studies were approved by the local institutional medical review board and ethics committees in accordance with the national regulations.

## Reference

1. Cohen AJ, Brauer M, Burnett R, Anderson HR, Frostad J, Estep K, Balakrishnan K, Brunekreef B, Dandona L, Dandona R, Feigin V, Freedman G, Hubbell B, Jobling A, Kan H, Knibbs L, Liu Y, Martin R, Morawska L, Pope CA, Shin H, Straif K, Shaddick G, Thomas M, van Dingenen R, van Donkelaar A, Vos T, Murray CJL, Forouzanfar MH. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet*. 2017;389:1907-1918.
2. International Agency for Research on Cancer (IARC) Working Group on the Evaluation of Carcinogenic Risks to Humans. Outdoor Air Pollution. *IARC Monogr Eval Carcinog Risks Hum*. 2016;109.
3. International Agency for Research on Cancer (IARC) Working Group on the Evaluation of Carcinogenic Risks to Humans. Diesel and Gasoline Engine Exhausts and Some Nitroarenes. *IARC Monogr Eval Carcinog Risks Hum*. 2014;105.
4. Raaschou-Nielsen O, Andersen ZJ, Beelen R, Samoli E, Stafoggia M, Weinmayr G, Hoffmann B, Fischer P, Nieuwenhuijsen MJ, Brunekreef B, Xun WW, Katsouyanni K, Dimakopoulou K, Sommar J, Forsberg B, Modig L, Oudin A, Oftedal B, Schwarze PE, Nafstad P, De Faire U, Pedersen NL, Östenson CG, Fratiglioni L, Penell J, Korek M, Pershagen G, Eriksen KT, Sørensen M, Tjønneland A, Ellermann T, Eeftens M, Peeters PH, Meliefste K, Wang M, Bueno-de-Mesquita B, Key TJ, de Hoogh K, Concin H, Nagel G, Vilier A, Grioni S, Krogh V, Tsai MY, Ricceri F, Sacerdote C, Galassi C, Migliore E, Ranzi A, Cesaroni G, Badaloni C, Forastiere F, Tamayo I, Amiano P, Dorronsoro M, Trichopoulou A, Bamia C, Vineis P, Hoek G. Air pollution and lung cancer incidence in 17 European cohorts: Prospective analyses from the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Lancet Oncol*. 2013;14:813-822.
5. Turner MC, Andersen ZJ, Baccarelli A, Diver WR, Gapstur SM, Pope CA, Prada D, Samet J, Thurston G, Cohen A. Outdoor air pollution and cancer: An overview of the current evidence and public health recommendations. *CA Cancer J Clin*. 2020;0:1-20.

6. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018;68:394-424.
7. Global Burden of Disease Liver Cancer Collaboration. The Burden of Primary Liver Cancer and Underlying Etiologies From 1990 to 2015 at the Global, Regional, and National Level: Results From the Global Burden of Disease Study 2015. *JAMA Oncol.* 2017;3:1683-1691.
8. Laing S, Wang G, Briazova T, Zhang C, Wang A, Zheng Z, Gow A, Chen AF, Rajagopalan S, Chen LC, Sun Q, Zhang K. Airborne particulate matter selectively activates endoplasmic reticulum stress response in the lung and liver tissues. *Am J Physiol Cell Physiol.* 2010;299:C736-49.
9. Zheng Z, Xu X, Zhang X, Wang A, Zhang C, Hüttemann M, Grossman LI, Chen LC, Rajagopalan S, Sun Q, Zhang K. Exposure to ambient particulate matter induces a NASH-like phenotype and impairs hepatic glucose metabolism in an animal model. *J Hepatol.* 2013;58:148-154.
10. Wong CR, Nguyen MH, Lim JK. Hepatocellular carcinoma in patients with non-alcoholic fatty liver disease. *World J Gastroenterol.* 2016;22:8294-8303.
11. Danielsen PH, Risom L, Wallin H, Autrup H, Vogel U, Loft S, Møller P. DNA damage in rats after a single oral exposure to diesel exhaust particles. *Mutat Res.* 2008;637:49-55.
12. Dybdahl M, Risom L, Møller P, Autrup H, Wallin H, Vogel U, Bornholdt J, Daneshvar B, Dragsted LO, Weimann A, Poulsen HE, Loft S. DNA adduct formation and oxidative stress in colon and liver of Big Blue rats after dietary exposure to diesel particles. *Carcinogenesis.* 2003;24:1759-1766.
13. Conklin DJ. From lung to liver: How does airborne particulate matter trigger NASH and systemic insulin resistance? *J Hepatol.* 2013;58:8-10.
14. Li D, Li Y, Li G, Zhang Y, Li J, Chen H. Fluorescent reconstitution on deposition of PM2.5 in lung and extrapulmonary organs. *Proc Natl Acad Sci U S A.* 2019;116:2488-2493.

15. Kim JW, Park S, Lim CW, Lee K, Kim B. The role of air pollutants in initiating liver disease. *Toxicol Res.* 2014;30:65-70.
16. Markevych I, Wolf K, Hampel R, Breitner S, Schneider A, von Klot S, Cyrys J, Heinrich J, Döring A, Beelen R, Koenig W, Peters A. Air Pollution and Liver Enzymes. *Epidemiology.* 2013;24:934-935.
17. Kim K-N, Lee H, Kim JH, Jung K, Lim Y, Hong Y. Physical Activity- and Alcohol-dependent Association Between Air Pollution Exposure and Elevated Liver Enzyme Levels : An Elderly Panel Study. *J Prev Med Public Heal.* 2015;48:151-169.
18. Kim HJ, Min JY, Seo YS, Min KB. Association of ambient air pollution with increased liver enzymes in Korean adults. *Int J Environ Res Public Health.* 2019;16.
19. Pan WC, Wu C Da, Chen MJ, Huang YT, Chen CJ, Su HJ, Yang HI. Fine Particle Pollution, Alanine Transaminase, and Liver Cancer: A Taiwanese Prospective Cohort Study (REVEAL-HBV). *J Natl Cancer Inst.* 2016;108:1-7.
20. Brook RD, Brook JR, Rajagopalan S. Air Pollution : The “Heart” of the Problem. *Curr Hypertens Rep.* 2003;5:32-39.
21. VoPham T, Bertrand KA, Tamimi RM, Laden F, Hart JE. Ambient PM<sub>2.5</sub> air pollution exposure and hepatocellular carcinoma incidence in the United States. *Cancer Causes Control.* 2018;29:563-572.
22. Raaschou-Nielsen O, Andersen ZJ, Hvidberg M, Jensen SS, Ketzel M, Sørensen M, Hansen J, Loft S, Overvad K, Tjønneland A. Air pollution from traffic and cancer incidence: A Danish cohort study. *Environ Health.* 2011;10:67.
23. Pedersen M, Andersen ZJ, Stafoggia M, Weinmayr G, Galassi C, Sørensen M, Eriksen KT, Tjønneland A, Loft S, Jaensch A, Nagel G, Concin H, Tsai MY, Grioni S, Marcon A, Krogh V, Ricceri F, Sacerdote C, Ranzi A, Sokhi R, Vermeulen R, de Hoogh K, Wang M, Beelen R, Vineis P, Brunekreef B, Hoek G, Raaschou-Nielsen O. Ambient air pollution and primary liver cancer incidence in four European cohorts within the ESCAPE project. *Environ Res.* 2017;154:226-233.

24. Turner MC, Krewski D, Diver WR, Pope CA 3rd, Burnett RT, Jerrett M, Marshall JD, Gapstur SM. Ambient air pollution and cancer mortality in the cancer prevention study II. *Environ Health Perspect.* 2017;125:087013.
25. Hvidtfeldt UA, Severi G, Andersen ZJ, Atkinson R, Bauwelinck M, Bellander T, Boutron-Ruault M-C, Brandt J, Brunekreef B, Cesaroni G, Chen J, Concin H, Forastiere F, van Gils CH, Gulliver J, Hertel O, Hoek G, Hoffmann B, de Hoogh K, Janssen N, Jöckel K-H, Jørgensen JT, Katsouyanni K, Ketzel M, Klompmaker JO, Krog NH, Lang A, Leander K, Liu S, Ljungman PLS, Magnusson PKE, Mehta AJ, Nagel G, Oftedal B, Pershagen G, Peter RS, Peters A, Renzi M, Rizzuto D, Rodopoulou S, Samoli E, Schwarze PE, Sigsgaard T, Simonsen MK, Stafoggia M, Strak M, Vienneau D, Weinmayr G, Wolf K, Raaschou-Nielsen O, Fecht D. Long-term low-level ambient air pollution exposure and risk of lung cancer – A pooled analysis of 7 European cohorts. *Environ Int.* 2021;146:106249.
26. Lagergren M, Fratiglioni L, Hallberg IR, Berglund J, Elmståhl S, Hagberg B, Holst G, Rennemark M, Sjölund BM, Thorslund M, Wiberg I, Winblad B, Wimo A. A longitudinal study integrating population, care and social services data. The Swedish National study on Aging and Care (SNAC). *Aging Clin Exp Res.* 2004;16:158-168.
27. Magnusson PKE, Almqvist C, Rahman I, Ganna A, Viktorin A, Walum H, Halldner L, Lundström S, Ullén F, Långström N, Larsson H, Nyman A, Gumpert CH, Råstam M, Anckarsäter H, Cnattingius S, Johannesson M, Ingelsson E, Klareskog L, de Faire U, Pedersen NL, Lichtenstein P. The Swedish Twin Registry: Establishment of a Biobank and Other Recent Developments. *Twin Res Hum Genet.* 2013;16:317-329.
28. Wändell PE, Wajngot A, de Faire U, Hellénus ML. Increased prevalence of diabetes among immigrants from non-European countries in 60-year-old men and women in Sweden. *Diabetes Metab.* 2007;33:30-36.
29. Eriksson AK, Ekblom A, Granath F, Hilding A, Efendic S, Östenson CG. Psychological distress and risk of pre-diabetes and type 2 diabetes in a prospective study of swedish middle-aged men and women. *Diabet Med.* 2008;25:834-842.

- Accepted Article
30. Tjønneland A, Olsen A, Boll K, Stripp C, Christensen J, Engholm G, Overvad K. Study design, exposure variables, and socioeconomic determinants of participation in Diet, Cancer and Health: a population-based prospective cohort study of 57,053 men and women in Denmark. *Scand J Public Health*. 2007;35:432-441.
  31. Hundrup YA, Simonsen MK, Jørgensen T, Obel EB. Cohort profile: The danish nurse cohort. *Int J Epidemiol*. 2012;41:1241-1247.
  32. Beulens JWJ, Monninkhof EM, Verschuren WMM, van der Schouw YT, Smit J, Ocke MC, Jansen EHJM, van Dieren S, Grobbee DE, Peeters PHM, Bueno-de-Mesquita HB. Cohort profile: the EPIC-NL study. *Int J Epidemiol*. 2010;39:1170-1178.
  33. Clavel-Chapelon F. Cohort Profile: The French E3N Cohort Study. *Int J Epidemiol*. 2015;44:801-809.
  34. Ulmer H, Kelleher CC, Fitz-Simon N, Diem G, Concin H. Secular trends in cardiovascular risk factors: an age-period cohort analysis of 698,954 health examinations in 181,350 Austrian men and women. *J Intern Med*. 2007;261:566-576.
  35. de Hoogh K, Chen J, Gulliver J, Hoffmann B, Hertel O, Ketzel M, Bauwelinck M, van Donkelaar A, Hvidtfeldt UA, Katsouyanni K, Klompmaker J, Martin R V., Samoli E, Schwartz PE, Stafoggia M, Bellander T, Strak M, Wolf K, Vienneau D, Brunekreef B, Hoek G. Spatial PM2.5, NO2, O3 and BC models for Western Europe – Evaluation of spatiotemporal stability. *Environ Int*. 2018;120:81-92.
  36. Chen J, de Hoogh K, Gulliver J, Hoffmann B, Hertel O, Ketzel M, Weinmayr G, Bauwelinck M, van Donkelaar A, Hvidtfeldt UA, Atkinson R, Janssen NAH, Martin R V., Samoli E, Andersen ZJ, Oftedal BM, Stafoggia M, Bellander T, Strak M, Wolf K, Vienneau D, Brunekreef B, Hoek G. Development of Europe-wide models for particle elemental composition using supervised linear regression and random forest. *Environ Sci Technol*. 2020;54:15698-15709.
  37. Brandt J, Silver JD, Frohn LM, Geels C, Gross A, Hansen AB, Hansen KM, Hedegaard GB, Skjøth CA, Villadsen H, Zare A, Christensen JH. An integrated model study for Europe and

North America using the Danish Eulerian Hemispheric Model with focus on intercontinental transport of air pollution. *Atmos Environ.* 2012;53:156-176.

38. Beelen R, Hoek G, Vienneau D, Eeftens M, Dimakopoulou K, Pedeli X, Tsai M-Y, Künzli N, Schikowski T, Marcon A, Eriksen KT, Raaschou-Nielsen O, Stephanou E, Patelarou E, Lanki T, Yli-Tuomi T, Declercq C, Falq G, Stempfelet M, Birk M, Cyrus J, von Klot S, Nádor G, Varró MJ, Dèdelè A, Gražulevičienė R, Mölter A, Lindley S, Madsen C, Cesaroni G, Ranzi A, Badaloni C, Hoffmann B, Nonnemacher M, Krämer U, Kuhlbusch T, Cirach M, de Nazelle A, Nieuwenhuijsen M, Bellander T, Korek M, Olsson D, Strömgren M, Dons E, Jerrett M, Fischer P, Wang M, Brunekreef B, de Hoogh K. Development of NO<sub>2</sub> and NO<sub>x</sub> land use regression models for estimating air pollution exposure in 36 study areas in Europe – The ESCAPE project. *Atmos Environ.* 2013;72:10-23.
39. Eeftens M, Beelen R, de Hoogh K, Bellander T, Cesaroni G, Cirach M, Declercq C, Dèdelè A, Dons E, de Nazelle A, Dimakopoulou K, Eriksen K, Falq G, Fischer P, Galassi C, Gražulevičienė R, Heinrich J, Hoffmann B, Jerrett M, Keidel D, Korek M, Lanki T, Lindley S, Madsen C, Mölter A, Nádor G, Nieuwenhuijsen M, Nonnemacher M, Pedeli X, Raaschou-Nielsen O, Patelarou E, Quass U, Ranzi A, Schindler C, Stempfelet M, Stephanou E, Sugiri D, Tsai M-Y, Yli-Tuomi T, Varró MJ, Vienneau D, Klot S von, Wolf K, Brunekreef B, Hoek G. Development of Land Use Regression Models for PM<sub>2.5</sub>, PM<sub>2.5</sub> Absorbance, PM<sub>10</sub> and PM<sub>coarse</sub> in 20 European Study Areas; Results of the ESCAPE Project. *Environ Sci Technol.* 2012;46:11195-11205.
40. Brauer M, Brook JR, Christidis T, Chu Y, Crouse DL, Erickson A, Hystad P, Li C, Martin R V., Meng J, Pappin AJ, Pinault LL, Tjepkema M, Donkelaar A Van, Weichenthal S, Burnett RT, van Donkelaar A, Weichenthal S, Burnett RT. Mortality-Air Pollution Associations in Low-Exposure Environments (MAPLE): Phase 1. *Research Report 203 Boston, MA:Health Effects Institute.* 2019.
41. Orioli R, Solimini AG, Michelozzi P, Forastiere F, Davoli M, Cesaroni G. A cohort study on long-term exposure to air pollution and incidence of liver cirrhosis. *Environ Epidemiol.* 2020;4:e109.



- Accepted Article
42. Fortoul TI, Rodriguez-Lara V, Gonzalez-Villalva A, Rojas-Lemus M, Cano-Gutierrez G, Ustarroz-Cano M, Colin-Barenque L, Montaña LF, García-Pelez I, Bizarro-Nevaras P, Lopez-Valdez N, Falcon-Rodriguez CI, Jimenez-Martínez RS, Ruiz-Guerrero ML, Lpez-Zepeda LS, Morales-Rivero A, Muñoz-Rivera-Cambas A. Vanadium inhalation in a mouse model for the understanding of air-suspended particle systemic repercussion. *J Biomed Biotechnol.* 2011;2011.
  43. Cano-Gutiérrez G, Acevedo-Nava S, Santamaría A, Altamirano-Lozano M, Cano-Rodríguez MC, Fortoul TI. Hepatic megalocytosis due to vanadium inhalation: Participation of oxidative stress. *Toxicol Ind Health.* 2012;28:353-360.
  44. Omland LH, Osler M, Jepsen P, Krarup H, Weis N, Christensen PB, Roed C, Sørensen HT, Obel N. Socioeconomic status in HCV infected patients - risk and prognosis. *Clin Epidemiol.* 2013;5:163-172.
  45. Meffre C, Le Strat Y, Delarocque-Astagneau E, Dubois F, Antona D, Lemasson J-M, Warszawski J, Steinmetz J, Coste D, Meyer J-F, Leiser S, Giordanella J-P, Gueguen R, Desenclos J-C. Prevalence of hepatitis B and hepatitis C virus infections in France in 2004: social factors are important predictors after adjusting for known risk factors. *J Med Virol.* 2010;82:546-555.

## Figure legends

### **Figure 1. Estimated exposure-response curves for associations between (A) NO<sub>2</sub>, (B) PM<sub>2.5</sub>, (C) BC, and (D) O<sub>3</sub> concentration at the residential addresses of baseline and liver cancer incidence.**

Abbreviation: NO<sub>2</sub>, nitrogen dioxide; PM<sub>2.5</sub>, particulate matter with aerodynamic diameters of less than 2.5 μm; BC, black carbon; O<sub>3</sub>, ozone.

Natural splines with 2 degrees of freedom. Black solid lines indicate hazard ratio values, and dashed lines indicate their 95% confidence intervals. Grey vertical dotted lines mean the values used for the subset analyses.

Models were adjusted for age (time scale), sex (strata), sub-cohort (strata), calendar year of baseline, smoking status, employment status, and mean income at the neighborhood level in 2001.

### **Figure 2. Associations between PM<sub>2.5</sub> components and liver cancer incidence.**

Abbreviation: PM<sub>2.5</sub>, particulate matter with aerodynamic diameters of less than 2.5 μm; NO<sub>2</sub>, nitrogen dioxide; Cu, copper; Fe, iron; Zn, zinc; S, sulfur; Ni, nickel; V, vanadium; Si, Silicon; K, potassium.

Models were adjusted for age (time scale), sex (strata), sub-cohort (strata), calendar year of baseline, smoking status, employment status, and mean income at the neighborhood level in 2001.

Associations were expressed as hazard ratios with 95% confidence intervals per interquartile range increase for each of PM<sub>2.5</sub> components: 3.7 ng/m<sup>3</sup> for Cu, 55.8 ng/m<sup>3</sup> for Fe, 10.7 ng/m<sup>3</sup> for Zn, 212.2 ng/m<sup>3</sup> for S, 0.8 ng/m<sup>3</sup> for Ni, 1.7 ng/m<sup>3</sup> for V, 24.1 ng/m<sup>3</sup> for Si, 82.3 ng/m<sup>3</sup> for K.

PM<sub>2.5</sub> components were estimated by the supervised linear regression method

**Table 1. Descriptive statistics for 330,064 participants at baseline by liver cancer incidence status at the end of follow-up.**

Characteristic	Total (N=330,064)	No liver cancer (N=329,552)	Liver cancer (N=512)	P-value <sup>c</sup>
Age, years (Mean ± SD)	48.2 ± 13.4	48.2 ± 13.4	55.8 ± 8.9	<.001
Age category, N (%)				0.003
< 65 years old	305,744 (92.6)	305,288 (92.6)	456 (89.1)	
≥ 65 years old	24,320 (7.4)	24,264 (7.4)	56 (10.9)	
Women, N (%)	220,292 (66.7)	220,104 (66.8)	188 (36.7)	<.001
Smoking status, N (%)				<.001
Never	180,703 (54.7)	180,461 (54.8)	242 (47.3)	
Former smoker	65,465 (19.8)	65,361 (19.8)	104 (20.3)	
Current Smoker	83,896 (25.4)	83,730 (25.4)	166 (32.4)	
Unemployed, N (%)	94,225 (28.5)	93,973 (28.5)	252 (49.2)	<.001
Intake of alcohol <sup>a</sup> , N (%)				<.001
Low (<4 g/day)	35,413 (23.1)	35,380 (23.1)	33 (17.9)	
Medium (4-15 g/day)	58,047 (37.9)	57,996 (37.9)	51 (27.7)	
High (15> g/day)	59,593 (38.9)	59,493 (38.9)	100 (54.3)	

<b>Education level<sup>b</sup>, N (%)</b>				<.001
<b>Low</b>	21,148 (11.8)	21,105 (11.8)	43 (19.6)	
<b>Medium</b>	70,980 (39.7)	70,886 (39.7)	94 (42.9)	
<b>High</b>	86,504 (48.4)	86,422 (48.4)	82 (37.4)	
<b>Mean income at neighborhood level in 2001, € (Mean ± SD)</b>	19,496.2 ± 5426.5	19,494.2 ± 5,428.0	20,791.4 ± 4,161.3	<.001
<b>Air pollutants for the year 2010 (Mean ± SD)</b>				
<b>NO<sub>2</sub>, µg/m<sup>3</sup></b>	24.9 ± 8.0	24.9 ± 8.0	24.8 ± 7.4	0.71
<b>PM<sub>2.5</sub>, µg/m<sup>3</sup></b>	15.0 ± 3.2	15.0 ± 3.2	14.9 ± 2.9	0.45
<b>BC, 10<sup>-5</sup>/m</b>	1.5 ± 0.4	1.5 ± 0.4	1.5 ± 0.4	0.35
<b>O<sub>3</sub>, µg/m<sup>3</sup></b>	85.6 ± 9.0	85.6 ± 9.0	85.9 ± 9.2	0.38
<b>PM<sub>2.5</sub> components (Mean ± SD)<sup>d</sup></b>				
<b>Copper, ng/m<sup>3</sup></b>	3.5 ± 2.6	3.5 ± 2.4	3.5 ± 2.6	0.82
<b>Iron, ng/m<sup>3</sup></b>	87.6 ± 46.8	87.9 ± 40.9	87.6 ± 46.8	0.85
<b>Zinc, ng/m<sup>3</sup></b>	16.9 ± 11.3	16.1 ± 10.9	17.0 ± 11.3	0.10
<b>Sulfur, ng/m<sup>3</sup></b>	659.1 ± 142.1	638.8 ± 126.2	659.2 ± 142.1	0.001
<b>Nickel, ng/m<sup>3</sup></b>	0.8 ± 0.7	0.7 ± 0.7	0.8 ± 0.7	0.10

<b>Vanadium, ng/m<sup>3</sup></b>	1.4 ± 1.4	1.3 ± 1.7	1.4 ± 1.4	0.58
<b>Silicon, ng/m<sup>3</sup></b>	96.2 ± 21.1	97.4 ± 19.1	96.2 ± 21.1	0.20
<b>Potassium, ng/m<sup>3</sup></b>	167.2 ± 52.4	168.7 ± 53.8	167.2 ± 52.4	0.52

Abbreviation: N, number; SD, standard deviation; NO<sub>2</sub>, nitrogen dioxide; PM<sub>2.5</sub>, particulate matter with aerodynamic diameters of less than 2.5 µm; BC, black carbon; O<sub>3</sub>, ozone.

<sup>a</sup> n=153,043

<sup>b</sup> n=178,632, low: primary school or less, medium: up to secondary school or equivalent, or high: university degree or more

<sup>c</sup> t-test for a continuous variable and chi-square test for a discrete variable to test the difference of a participant characteristic variable between cases and non-cases.

<sup>d</sup> Based on the supervised linear regression exposure model

**Table 2. Associations between long-term exposure to air pollution and the risk for liver cancer incidence.**

Air pollutant	Model 1 <sup>a</sup>	Model 2 <sup>b</sup>	Model 3 <sup>c</sup>	Two-pollutant model:			
				Model 3+NO <sub>2</sub>	Model 3 +PM <sub>2.5</sub>	Model 3+BC	Model 3+O <sub>3</sub>
NO <sub>2</sub>	1.14 (1.00 to 1.31)	1.12 (0.98 to 1.29)	1.17 (1.02 to 1.35)	-	1.19 (1.00 to 1.43)	1.22 (0.87 to 1.70)*	0.95 (0.78 to 1.16)
PM <sub>2.5</sub>	1.10 (0.91 to 1.33)	1.09 (0.90 to 1.32)	1.12 (0.92 to 1.36)	0.96 (0.75 to 1.23)	-	0.98 (0.76 to 1.27)	0.88 (0.70 to 1.11)
BC	1.13 (0.98 to 1.30)	1.11 (0.97 to 1.28)	1.15 (1.00 to 1.33)	0.96 (0.68 to 1.36)*	1.16 (0.96 to 1.41)	-	0.94 (0.77 to 1.14)
O <sub>3</sub>	0.69 (0.58 to 0.84)	0.71 (0.59 to 0.86)	0.70 (0.58 to 0.85)	0.67 (0.51 to 0.88)	0.66 (0.52 to 0.82)	0.66 (0.51 to 0.86)	-

Abbreviation: HR, Hazard ratio; CI, confidence interval; NO<sub>2</sub>, nitrogen dioxide; PM<sub>2.5</sub>, particulate matters with aerodynamic diameters of less than 2.5 µm; BC, black carbon; O<sub>3</sub>, ozone.

Results are presented as hazard ratio and 95% confidence interval [HR (95% CI)] for the following increments: 5 µg/m<sup>3</sup> for PM<sub>2.5</sub>, 10 µg/m<sup>3</sup> for NO<sub>2</sub>, 0.5 10<sup>-5</sup> /m for BC and 10 µg/m<sup>3</sup> for O<sub>3</sub>.

\*: The results from the model with NO<sub>2</sub> and BC are difficult to interpret because of their high correlation, which reached 0.83. <sup>a</sup> Model1 was adjusted for age (time scale), sex (strata), sub-cohort (strata), and calendar year of baseline.

<sup>b</sup> Model2 was adjusted for age (time scale), sex (strata), sub-cohort (strata), and calendar year of baseline, smoking status, and employment status.

<sup>c</sup> Model3 was adjusted for age (time scale), sex (strata), sub-cohort (strata), calendar year of baseline, smoking status, employment status, and mean income at the neighborhood level in 2001.

**Table 3. Associations between long-term exposure to air pollution and incident liver cancer, below various cut-points.**

<b>Air pollutants</b>	<b>Subset<sup>a</sup></b>	<b>Participants, N</b>	<b>Cases, N</b>	<b>Hazard ratio (95% CI)<sup>b</sup></b>
<b>NO<sub>2</sub></b>	Full dataset	330,064	512	1.17 (1.02 to 1.35)
	<40 µg/m <sup>3</sup> (EU standard)	315,023	498	1.20 (1.03 to 1.40)
	<30 µg/m <sup>3</sup>	252,154	392	1.03 (0.83 to 1.28)
	<20 µg/m <sup>3</sup>	90,300	135	1.01 (0.57 to 1.77)
<b>PM<sub>2.5</sub></b>	Full dataset	330,064	512	1.12 (0.92 to 1.36)
	<25 µg/m <sup>3</sup> (EU standard)	330,024	512	1.12 (0.92 to 1.36)
	<20 µg/m <sup>3</sup>	320,759	505	1.12 (0.92 to 1.36)
	<15 µg/m <sup>3</sup>	153,720	264	1.56 (0.96 to 2.55)
	<12 µg/m <sup>3</sup>	52,349	72	1.02 (0.32 to 3.26)
	<10 µg/m <sup>3</sup>	24,495	24	0.39 (0.06 to 2.73)
<b>BC</b>	Full dataset	330,064	512	1.15 (1.00 to 1.33)
	< 3.0 10 <sup>-5</sup> /m	329,305	512	1.17 (1.01 to 1.35)
	< 2.5 10 <sup>-5</sup> /m	324,258	506	1.16 (0.99 to 1.35)
	< 2.0 10 <sup>-5</sup> /m	299,519	471	1.16 (0.97 to 1.38)

	< 1.5 10 <sup>-5</sup> /m	142,778	206	1.29 (0.91 to 1.84)
	< 1.0 10 <sup>-5</sup> /m	34,477	37	0.57 (0.20 to 1.61)
<b>O<sub>3</sub></b>	Full dataset	330,064	512	0.70 (0.58 to 0.85)
	< 120 µg/m <sup>3</sup>	330,064	512	0.70 (0.58 to 0.85)
	< 100 µg/m <sup>3</sup>	324,120	507	0.67 (0.55 to 0.81)
	< 80 µg/m <sup>3</sup>	97,767	142	0.65 (0.45 to 0.95)

Abbreviation: N, number; CI, confidence interval; NO<sub>2</sub>, nitrogen dioxide; PM<sub>2.5</sub>, particulate matters with aerodynamic diameters of less than 2.5 µm; BC, black carbon; O<sub>3</sub>, ozone.

<sup>a</sup>Participants with concentrations above a cut-point were excluded.

<sup>b</sup>From models adjusting for age (time scale), sex (strata), sub-cohort (strata), calendar year of baseline, smoking status, employment status, and mean income at the neighborhood level in 2001.



**Novelty and Impact:**

Air pollution contains a number of known carcinogens. While air pollution is classified as carcinogenic and is a known risk factor for lung cancer, the evidence for cancers in other organs is limited. In this large European study, the authors detected associations between air pollution components and liver cancer incidence, even at levels that are below current EU standards. These results corroborate findings from several earlier, substantially smaller studies, and suggest that ambient air pollution may increase the risk of liver cancer.



