

# Association of eating motives with anthropometry, body composition, and dietary intake in healthy German adults

Matthias Staudigel<sup>a,\*1</sup>, Beate Brandl<sup>b,1</sup>, Jutta Roosen<sup>a</sup>, Eva Kiesswetter<sup>c</sup>, Hanna M. Siebentritt<sup>c</sup>, Nina Wawro<sup>d,e</sup>, Jakob Linseisen<sup>d,e,f</sup>, Thomas Skurk<sup>b,g</sup>, Dorothee Volkert<sup>c</sup>, Hans Hauner<sup>g,h</sup>

<sup>a</sup> *Technical University of Munich, TUM School of Management, Chair of Marketing and Consumer Research, Alte Akademie 16, 85354, Freising, Germany*

<sup>b</sup> *Technical University of Munich, ZIEL – Institute for Food & Health, Core Facility Human Studies, Gregor-Mendel-Str. 2, 85354, Freising, Germany*

<sup>c</sup> *Friedrich-Alexander-University Erlangen-Nürnberg, Institute for Biomedicine of Aging, Kobergerstr. 60, 90408, Nürnberg, Germany*

<sup>d</sup> *Helmholtz Zentrum München, Independent Research Group Clinical Epidemiology, Ingolstädter Landstr. 1, 85764 Neuherberg, Germany*

<sup>e</sup> *Ludwig-Maximilians-Universität München, Medical Faculty, Institute for Medical Information Processing, Biometry, and Epidemiology (IBE), Marchioninistr. 15, 81377 München*

<sup>f</sup> *Chair of Epidemiology, University of Augsburg, University Hospital Augsburg, Stenglinstr. 2, 86156 Augsburg, Germany*

<sup>g</sup> *Else Kröner-Fresenius-Center for Nutritional Medicine, Technical University of Munich, Gregor-Mendel-Str. 2, 85354, Freising, Germany*

<sup>h</sup> *Technical University of Munich, Klinikum rechts der Isar, Institute for Nutritional Medicine, Georg-Brauchle-Ring 62, 80992, Munich, Germany*

## \* Corresponding author:

Dr. Matthias Staudigel

Technical University of Munich  
TUM School of Management  
Chair of Marketing and Consumer Research

Alte Akademie 16  
85354 Freising, Germany

Phone: +49 8161 71 3869

E-mail: [matthias.staudigel@tum.de](mailto:matthias.staudigel@tum.de)

<sup>1</sup> Matthias Staudigel and Beate Brandl contributed equally to this work

# 1 Association of eating motives with anthropometry, body composition, 2 and dietary intake in healthy German adults

3

## 4 Abstract

5 Effective policies to address poor food choices and dietary patterns need to consider the complex set  
6 of motives affecting eating behavior. This study examined how different eating motives are associated  
7 with anthropometry, body composition, and dietary intake. Our analysis is based on a cross-sectional  
8 sample with 429 healthy adults in three different age groups collected in Germany from 2016 to 2018.  
9 Dietary intake, Body Mass Index (BMI), waist circumference (WC), and fat-free mass (FFM) were  
10 measured by standardized methods. Eating motives were measured using The Eating Motivation  
11 Scale (TEMS). Regressing dietary intakes and anthropometric indicators on TEMS motives, we  
12 identify the main sources of variation in diet and nutritional status separately for men and women.  
13 Results indicated the *Health* motive to be positively associated with FFM ( $B \pm SE = 1.72 \pm 0.44$ ) and  
14 negatively with WC ( $B \pm SE = -3.23 \pm 0.81$ ) for men. For women, the *Need & Hunger* motive was  
15 positively associated with FFM ( $B \pm SE = 1.63 \pm 0.44$ ) and negatively with WC ( $B \pm SE = -2.46 \pm$   
16  $0.81$ ). While *Liking* and *Habits* were the most frequently stated eating motives, we did not find them  
17 to be significantly related to the nutritional status. Other motives were associated with dietary intake  
18 but not anthropometry or body composition. The *Price* motive was positively and the *Convenience*  
19 motive was negatively associated with energy ( $B \pm SE = 63.77 \pm 19.98$ ;  $B \pm SE = -46.96 \pm 17.12$ )  
20 and carbohydrate intake ( $B \pm SE = 7.15 \pm 2.65$ ;  $B \pm SE = -5.98 \pm 2.27$ ) for men. The results  
21 highlight the need for more differentiated analyses of eating motives, beyond comparing the relative  
22 importance of motives based on mean values, towards the association of motives with dietary intake  
23 and nutritional status.

24 **Keywords:** Food choice; Motivation; Dietary intake; Nutritional status; Policy implications

## 25 1. Introduction

26 Unhealthy diets and poor nutrition are among the top risk factors worldwide for obesity,  
27 cardiovascular diseases, and diabetes. The World Health Organization (WHO) estimates that more  
28 than 1.9 billion adults were overweight and 650 million of these were obese in 2016 (World Health  
29 Organization, 2020a) and that the prevalence of diabetes among adults increased from 4.7% in 1980  
30 to 8.5% in 2014 (World Health Organization, 2020b). The scientific and public debate about  
31 appropriate and effective policies to counter these developments increasingly acknowledges that  
32 poor food choices and dietary patterns are the results of multiple motives and determinants  
33 (Naughton et al., 2015; Pollard et al., 1998; Pollard et al., 2002; Sproesser, Moraes et al., 2019; Steptoe  
34 et al., 1995). Hence, there is no ‘one-size-fits-all’ solution (Hawkes et al., 2015; Just & Gabrielyan,  
35 2016; Lusk, 2017) as the factors affecting food choice and eating behavior are complex (Köster, 2009).

36 Literature has discussed a wide range of different motives that affect food choice (Renner et al., 2012).  
37 Compiling, extending, and consolidating the empirical evidence on motives for eating behavior since  
38 Steptoe et al.’s (1995) seminal study, Renner et al. (2012) developed The Eating Motivation Scale  
39 (TEMS). Their measure includes 15 factors, which are *Liking, Habits, Need & Hunger, Health,*  
40 *Convenience, Pleasure, Traditional Eating, Natural Concerns, Sociability, Price, Visual Appeal, Weight*  
41 *Control, Affect Regulation, Social Norms, and Social Image* (Renner et al., 2012).

42 A growing number of studies has investigated the relative importance of food choice motives in  
43 general, across socio-demographics, by body weight, and across different countries and cultures.  
44 *Liking* or *Sensory Appeal, Habits, Need & Hunger, Health, Convenience* and *Price* are consistently  
45 reported as the most prominent drivers of food choice (Chambers et al., 2016; Phan & Chambers,  
46 2016a; Renner et al., 2012; Sproesser, Moraes et al., 2019; Sproesser, Ruby et al., 2018; Steptoe et al.,  
47 1995). Results for different age groups suggest that factors driving younger consumers’ eating  
48 behavior are more “short-term oriented” (Renner et al., 2012) including *Liking, Need & Hunger,*  
49 *Pleasure, Convenience, and Visual Appeal*. For older people, in contrast, long-term motives such as  
50 *Health* and *Natural Concerns* have been found to be more relevant based on TEMS (Renner et al.,  
51 2012). Another recurring finding is that women rated most motives significantly higher on average  
52 than men (Renner et al., 2012; Steptoe et al., 1995).

53 Food choice motives have also been shown to vary with nutritional status indicated mainly by the  
54 Body Mass Index (BMI). Renner et al. (2012) found that participants in the normal-weight range  
55 reported their eating behavior to be driven more frequently by *Liking, Health, and Need & Hunger*.  
56 The motives *Weight Control, Affect Regulation, and Social Norms* were more prevalent among

57 participants who had overweight or obesity. Similar findings are reported by Sproesser, Moraes et al.  
58 (2019) and Rempe et al. (2019).

59 While it is important to quantify the self-reported motives in their own right, targeted public policy  
60 measures need information on how these different motives drive actual eating behavior, and intake  
61 of critical nutrients, as well as how they affect both the nutrition and health status. Studies have  
62 shown that motives are associated with choice and consumption of individual food groups (Hebden  
63 et al., 2015; Phan & Chambers, 2016b; Souza et al., 2020) and vary over eating occasions (Chambers  
64 et al., 2016; Phan & Chambers, 2016a, 2018). Motives have further been shown to be associated with  
65 preferences for specific food product attributes such as 'organic' or 'local' (Hasselbach & Roosen,  
66 2015; Honkanen et al., 2006) or functional properties in food products (Ares & Gámbaro, 2007).

67 However, empirical evidence on how motives drive nutritional and health outcomes such as dietary  
68 intakes and indicators of nutritional status other than BMI is scarce. The present study contributes to  
69 the literature by providing empirical evidence on how different eating motives are associated with  
70 variation in anthropometric parameters, body composition, and dietary intake. Collecting evidence  
71 on the relative importance of motives and their association with consumption of specific food groups,  
72 diets, or meal occasions, is important for understanding drivers of food choice. At the same time,  
73 investigating which motives drive health- and nutrition-relevant behavior and outcomes, may yield  
74 specific insights on the levers that health and nutrition policies should be addressing.

75  
76 Our analysis is based on a unique dataset from a study with 429 healthy adults at three defined age  
77 groups conducted in Germany from 2016 to 2018 (Brandl et al. 2020). The collected data include  
78 information on usual dietary intake (Mitry et al., 2019) for a wide range of energy and macronutrients,  
79 anthropometric measures (BMI and waist circumference), and body composition. Additionally,  
80 participants reported socio-demographic variables and their eating motivations by completing The  
81 Eating Motivation Scale (TEMS) (Renner et al., 2012). We regressed anthropometric indicators, body  
82 composition, and energy and nutrient intakes on motives from TEMS controlling for socio-  
83 demographic characteristics to identify associations that are responsible for variation in dietary  
84 intake and nutritional status separately for men and women. This approach goes one step further  
85 than existing literature, which has mostly analysed relationships of motives with nutritional status or  
86 dietary intakes based on simple correlations or for specific subsets of motives. Since motives have  
87 been found to be correlated among each other as well as with socio-demographic characteristics,  
88 parameters of such analyses are likely to be biased when confounders are not accounted for.

## 89 **2. Methods**

### 90 **2.1. Study design and participants**

91 Data for this study come from the phenotyping program of the *enable* cluster of nutrition research  
92 performed between February 2016 and February 2018. The cluster's major objective was to identify  
93 determinants of a healthy nutrition and lifestyle for defined stages of life (Brandl et al., 2020). For this  
94 cross-sectional study, healthy volunteers were recruited from three different age groups, each of  
95 which is characterized by specific nutritional requirements and transitions in dietary habits and food  
96 selection: young adults aged 18-25 years (n=94), adults aged 40-65 years ("middle agers", n=205),  
97 and older adults aged 75-85 years (n=160). While young adults and middle agers were recruited at  
98 one study site (Freising), participants in the older adults group came from two sites (Freising and  
99 Nuremberg, two cities in Bavaria, Germany). Participants were recruited by advertising in  
100 kindergarten, newspapers, senior homes, and other media in the region of Freising and Nuremberg,  
101 Germany. In total 459 healthy subjects were comprehensively phenotyped. Apart from  
102 anthropometry and body composition analysis, participants completed standardized questionnaires  
103 regarding their dietary intake and eating motives. The study was approved by the ethics committee  
104 of the Faculty of Medicine of Technical University of Munich (#492/17S) and by the ethics committee  
105 of Friedrich-Alexander-Universität Erlangen-Nürnberg (#291\_15 B). It was registered on German  
106 Clinical Trial register (DRKS-ID: DRKS DRKS00009797). Written informed consent was obtained from  
107 all participants prior to the assessments. For detailed information on selection and eligibility criteria  
108 and study procedure see Brandl et al. (2020).

### 109 **2.2. Dietary intake, anthropometry, and body composition**

110 In the different age groups dietary habits were recorded by using a food frequency questionnaire  
111 (FFQ2) referring to a twelve-months period (Nöthlings et al., 2007) and two 24h-food lists (Freese et  
112 al., 2014). Based on these questionnaires, the "usual intake" of energy, carbohydrates, fat, protein,  
113 fiber, and alcohol for every participant was estimated by a multi-step procedure. First, the probability  
114 of consuming a certain food item was estimated using logistic regression models on the repeated 24h-  
115 food-list data, controlling for age, sex, BMI, smoking, physical activity level, education level, and the  
116 consumption frequency for each food item derived from the FFQ. Second, the consumption amount  
117 for each food item on a consumption day was estimated on the basis of the Second Bavarian Food  
118 Consumption Survey (BVS II) (Himmerich et al., 2003). The usual food intake was then derived by  
119 multiplying consumption probability and quantity of each food item. All usual food intakes were then

120 linked to the “Bundeslebensmittelschlüssel” (BLS, German Nutrient Data Base) to derive the usual  
121 intake of energy and nutrients. Details on the procedure are described in detail in Mitry et al. (2019).

122 All anthropometric parameters (body height, body weight, waist and hip circumference) were  
123 measured in the morning following an overnight fast using established standard operation procedure  
124 (SOP). Body composition including fat free mass (FFM) and fat mass (FM) was measured by  
125 bioelectrical impedance using the Seca mBCA 515 device (Seca GmbH & Co KG, Hamburg, Germany).

### 126 2.3. Eating motives

127 All participants completed the German short version of the eating motivation survey (TEMS)  
128 consisting of 15 motivational factors, each measured by three items (Renner et al., 2012). For  
129 example, the items for the factor *Affect Regulation* started with “I eat what I eat, ...” followed by 1) “...  
130 because I am sad”, 2) “... because I am frustrated”, and 3) “... because I feel lonely”. Each item was  
131 measured on a 7-point Likert scale whose extremes were labeled as 1 “never” to 7 “always”.

132 In order to compare the quality of our TEMS data with those of previous studies (Rempe et al., 2019;  
133 Renner et al., 2012; Sproesser, Ruby et al., 2018), we tested the original model with a confirmatory  
134 factor analysis (CFA) allowing for correlated factors (‘sem’ and ‘alpha’ commands in Stata 14). Results  
135 of the CFA showed a very good fit as indicated by a comparative fit index (CFI) of .94, a root-mean-  
136 square error of approximation (RMSEA) of .043, a standardized root mean square residual (SRMR) of  
137 .045 and a  $\chi^2/df$ -value of 1.8 (Kline, 2011). These values compare very well to or even exceed  
138 previous assessments of TEMS (Rempe et al., 2019; Renner et al., 2012; Sproesser, Ruby et al., 2018).  
139 The factor loadings were significant ( $p < .001$ ), with values larger than .4. Factor correlations ranged  
140 from -.18 (*Price* and *Sociability*) to .53 (*Need & Hunger* and *Habits*). The Cronbach’s Alpha values for  
141 the 15 factors indicated very good reliabilities, with the lowest value being .62 for *Need & Hunger*. For  
142 more details, see Table A1 in the appendix.

### 143 2.4. Data analysis

144 We analyzed the relation between eating motives and anthropometry, body composition, and dietary  
145 intake, respectively, via multiple regression models based on eq. (1):

$$146 \quad (1) \quad Y_i = (\alpha + \sum_{j=1}^{15} \beta_j \cdot Motive_{ji} + \sum_{k=2}^4 \delta_k \cdot Age\ Group_{ki} + \delta_5 \cdot Middle\ Income_i +$$

$$147 \quad \delta_6 \cdot High\ Income_i + \sum_{l=2}^6 \eta_l \cdot Education_{li}) \cdot (D_{Male,i} + D_{Female,i}) + \varepsilon_i$$

148 Dependent variables  $Y_i$  were either anthropometric measures (BMI and waist circumference), body  
149 composition (FFM), or energy and nutrient intakes (energy, protein, carbohydrates, fat, fiber, alcohol)  
150 of participant  $i$ . The key explanatory variables were the 15 motivation factors ( $Motive_{ji}$ ). We  
151 controlled for the effects of age groups, income, and education level, as these affect most of the  
152 dependent variables and have been shown to also correlate with the eating motives in previous  
153 studies (Renner et al., 2012). For these sets of categorical variables,  $Age\ Group_1$ ,  $Low\ Income$ , and  
154  $Education_1$  were omitted as reference categories to avoid singularity. All independent variables  
155 including the constant were multiplied by the term  $(D_{Male,i} + D_{Female,i})$ , where  $D_{Male}$  and  $D_{Female}$   
156 are binary indicators for male and female participants, respectively. This way we obtain a full set of  
157 motive parameters separately for men and women, which has two advantages. First, we have a larger  
158 sample size yielding more precise coefficient estimates. Second, we can immediately test for equality  
159 of male and female coefficients for each motive.

160 The set of equations for all dependent variables was estimated via seemingly unrelated regression  
161 (SUR) (Zellner, 1962), to account for correlated error terms within individuals across equations. For  
162 each equation, we test for equality of each parameter of all 30 motive-sex combinations with zero  
163 based on the standard errors and corresponding p-values provided by the regression. We consider  
164 multiplicity by controlling for the false discovery rate (FDR) using Benjamini and Hochberg's (1995)  
165 procedure. The p-values we report in the text represent adjusted p-values, i.e. so-called "q-values"  
166 derived by Stata's *qqvalue* command. We test for equality of male and female parameters for each  
167 motive using simple Wald tests after the regression (using Stata's *test* command).

168

### 169 **3. Results**

#### 170 *3.1. Sample characteristics*

171 From the original sample, 29 observations were dropped because of incomplete dietary data and one  
172 person was dropped due to a missing BMI value, yielding a final sample of 429 participants. Table 1  
173 shows definitions and summary statistics for the sociodemographic variables as well as national  
174 statistics for Germany for comparison. Men and women had equal shares in the sample, which is  
175 representative for sex/gender. 22% of respondents belong to the young adults' group, 47% to the  
176 middle-age group, and 31% were older adults. Given this study's focus on specific phases of life, all  
177 age groups included had a higher share than in the national statistics. The three binary variables for  
178 income represent lower, middle, and upper tertiles of the income distribution in our data. The

179 proportion of participants with a monthly household income < 2000 € was comparable to official  
 180 statistics, while the group with monthly incomes between 2000 € and 3000 € had a higher and the  
 181 group with monthly incomes above 3,000 € had a lower share. The average education level was high,  
 182 with 54% of participants graduating from schools that qualify for university admission compared to  
 183 32.5% in the German population.

184 **Table 1**

185 Summary statistics for sociodemographic variables for the total sample (n = 429)

Variable	Definition	Mean ( $\pm$ SD) / Percentage	Germany
Male	= 1, if respondent is male	49.2%	49.3%
Age	Respondent's age in years	53.8 ( $\pm$ 20.7)	
Age Cohort 1 (young adults)	= 1 if Age is 18-25 years	21.7%	8.8 %
Age Cohort 2 (middle agers)	= 1 if Age is 40-65 years	47.3%	36.8%
Age Cohort 3 (older adults, FS)	= 1 if Age is 75-85 years (Freising sample)	8.4%	9.5%
Age Cohort 4 (older adults, N)	= 1 if Age is 75-85 years (Nuremberg sample)	22.6%	
Low Income	= 1, if monthly household income < 2000 €	39.9%	36.6%
Middle Income	= 1, if monthly household income $\geq$ 2000 € and < 3000 €	28.9%	21.9%
High Income	= 1, if monthly household income $\geq$ 3000 €	31.2%	41.4%
General secondary	= 1, if highest education is "General secondary school (Hauptschule)"	16.8%	29.6%
Intermediate secondary	= 1, if highest ed. is "Intermediate secondary school (Mittlere Reife)"	24.7%	23.3%
Polytechnical secondary	= 1, if highest ed. is "Polytechnical secondary school (Polytechnische Oberschule)"	1.9%	6.6%
Technical college qualification	= 1, if highest ed. is "Advanced technical college entrance qualification (Fachhochschulreife)"	10.5%	32.5%
International baccalaureate	= 1, if highest ed. is "International baccalaureate (Abitur)"	44.1%	
Other	= 1, if degree from abroad or other degree	2.1%	0.2%

186 *Note:* Official statistics for *Male* and *Education Groups* from the German Statistical Yearbook  
 187 (Destatis, 2019), for *Age Cohorts* from Destatis (2021), *Income tertiles* from Destatis (2018).

188



189 Table 2 shows summary statistics for anthropometric indicators, body composition, and usual dietary  
 190 intakes for the overall sample as well as for the male and female subsamples.

191 **Table 2**

192 Summary statistics for anthropometric indicators and usual daily nutrient intakes for the total  
 193 sample and for male and female subsamples

Variable name [unit]		Overall (n = 429)		Male (n = 211)		Female (n = 218)		<i>t</i>	Effect size (Cohen's <i>d</i> ) <sup>a)</sup>	
		M	SD	M	SD	M	SD			
<i>Anthropometric measures and body composition</i>										
BMI	SECA Body Mass Index [kg/m <sup>2</sup> ]	26.0	4.5	26.7	4.2	25.3	4.7	3.28	***	0.32
FFM	SECA Fat free mass [%]	68.7	10.0	73.8	8.2	63.7	9.1	12.06	***	1.16
Waist	Waist circumference [cm]	91.3	14.9	97.3	13.9	85.5	13.5	8.98	***	0.87
<i>Energy and nutrient intake</i>										
Energy	Energy [kcal/d]	1936	388	2168	357	1711	264	15.01	***	1.46
Protein	Protein [g/d]	75	16	84	14	67	12	10.45	***	1.01
Carbs	Carbohydrates, absorbable [g/d]	204	47	225	47	183	36	13.02	***	1.26
Fat	Fat [g/d]	83	17	93	16	75	13	12.84	***	1.24
Fibre	Fibre [g/d]	21	6	21	7	20	6	1.62		0.16
Alcohol	Alcohol (Ethanol) [g/d]	8	8	12	10	3	2	12.48	***	1.22

194 *Note:* M Mean, SD Standard deviation. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.1$ , based on t-tests. <sup>a)</sup> Effect size for  
 195 differences in variables between sexes.

196

197 Table 3 presents the mean values of the 15 motivation factors for the overall sample, as well as for  
 198 the male and female subsamples. Additionally, we included the values reported by Renner et al.  
 199 (2012) for comparison purposes. Top-rated factors of eating motivation overall were *Liking* with a  
 200 mean score of 5.6, *Health* (4.7), *Need & Hunger* (4.5), *Natural Concerns* (4.5), and *Habits* (4.2). Factors  
 201 with the lowest average scores were *Price* (3.3), *Social Norms* (2.7), *Visual Appeal* (2.6), *Affect*  
 202 *Regulation* (2.0), and *Social Image* (1.5).

203 Compared to Renner et al. (2012), the present sample had lower values for most motivation factors,  
 204 especially for *Habits, Visual Appearance, Traditional Eating, and Affect Regulation*. Exceptions with  
 205 higher values in the present sample were *Natural Concerns, Weight Control, and Social Norms*.

206 We found significantly lower factor values for men compared to women for 11 out of 15 motives. The  
 207 male and female subgroups differed most prominently for *Natural Concerns, Traditional Eating, Affect*  
 208 *Regulation, Health, Sociability, and Social Norms*. No differences occurred for *Price, Liking, Habits, and*  
 209 *Visual Appeal*.

210

211 **Table 3**

212 Mean (M) and standard deviation (SD) of the motivation factors from this study (total sample, male  
 213 and female subsamples) and by the study of Renner et al. (2012)

	Overall (n = 429)		Male (n = 211)		Female (n = 218)		<i>t</i>	Effect size (Cohen's <i>d</i> ) <sup>a)</sup>	Renner et al. (2012) (n = 1040)	
	M	SD	M	SD	M	SD			M	SD
<i>Liking</i>	5.6	1.1	5.6	1.0	5.7	1.1	1.49	0.14	5.9	0.7
<i>Health</i>	4.7	1.2	4.4	1.3	4.9	1.1	4.13	***	4.7	1.0
<i>Need &amp; Hunger</i>	4.5	1.2	4.4	1.2	4.6	1.1	2.06	**	4.8	0.8
<i>Natural Concerns</i>	4.5	1.5	4.1	1.6	4.9	1.4	5.56	***	4.0	1.4
<i>Habits</i>	4.2	1.3	4.1	1.3	4.2	1.4	1.05		4.8	0.9
<i>Convenience</i>	4.1	1.4	4	1.5	4.3	1.4	2.31	**	4.5	1.0
<i>Pleasure</i>	4.1	1.3	3.9	1.3	4.3	1.3	3.12	***	4.3	0.9
<i>Sociability</i>	3.9	1.5	3.6	1.4	4.2	1.5	3.94	***	3.7	1.1
<i>Weight Control</i>	3.7	1.5	3.6	1.6	3.9	1.5	1.92	*	3.3	1.3
<i>Traditional Eating</i>	3.6	1.4	3.3	1.3	3.9	1.4	4.41	***	4.1	0.9
<i>Price</i>	3.3	1.4	3.3	1.4	3.2	1.4	-0.04		3.7	1.0
<i>Social Norms</i>	2.7	1.5	2.5	1.4	3	1.5	3.70	***	2.4	0.7
<i>Visual Appeal</i>	2.6	1.1	2.5	1.1	2.6	1.2	1.27		3.5	0.9
<i>Affect Regulation</i>	2.0	1.2	1.7	1.0	2.3	1.4	5.26	***	2.7	1.2
<i>Social Image</i>	1.5	0.7	1.4	0.6	1.6	0.8	2.36	**	2.0	0.7

214 *Note:* M Mean, SD Standard deviation. \*\*\*  $p < 0.01$ , \*\*  $p < 0.05$ , \*  $p < 0.1$ , based on t-tests. <sup>a)</sup> Effect size for  
 215 differences in motive scores between sexes.

216

217

218

219 3.2. Regressions for anthropometric indicators

220 Table 4 presents the results of multiple regression models for anthropometry and body composition  
221 separate for men and women based on eq. (1). Results without differentiating coefficients by sex are  
222 depicted Table A2 in the appendix. Regarding goodness of fit,  $R^2$ -values range from 34% for BMI to  
223 69% for FFM. These values are relatively high for models explaining the variation in anthropometric  
224 measures influenced by a complex set of biological, personal, social, or environmental factors (Chou  
225 et al., 2004; Schmeiser, 2009; Schroeter & Lusk, 2008). Estimated coefficients show similar effects for  
226 men and women for some eating motives, but pronounced differences for others as indicated by  
227 magnitude, sign, and/or tests for equality (significance levels from these tests are depicted in the  $\Delta$ -  
228 column). We report corrected p-values (i.e., “q-values”) based on Benjamini and Hochberg (1995) in  
229 the following.

230  
231 For males, the *Health* motive showed particularly pronounced effects. The estimated parameters  
232 suggest that a 1-point increase in the valuation of the health motive (on a scale from 1 to 7) was  
233 associated with a decrease of 1.72%-points increase in FFM ( $p < .001$ ), and a decrease of 3.23 cm in  
234 waist circumference ( $p < .001$ ). The *Health* coefficient in the BMI model showed a considerable  
235 magnitude (-.74), but the estimate was less precise and not significantly different from zero. Higher  
236 scores for *Affect Regulation* as an eating motive was linked to a lower share of FFM (-1.26;  $p = .068$ )  
237 and a higher waist circumference (2.38;  $p = .070$ ). Also here, the coefficient for BMI was substantial  
238 in size but not significant. We were unable to reject hypotheses of equality between the parameters  
239 for these variables between women and men, except for the case of *Affect Regulation* and waist  
240 circumference.

241  
242 Results for the female sample indicate that a strong *Need & Hunger* motive was negatively associated  
243 with waist circumference (-2.46;  $p = .030$ ) and positively with FFM (1.63;  $p < .001$ ). These coefficients  
244 were also statistically different from those estimated for men. High values in the *Traditional Eating*  
245 motive were associated with higher BMI (0.77;  $p = .060$ ) and lower FFM (-1.04;  $p = .060$ ). These  
246 coefficients were not different from those obtained for men.

247  
248 While the estimated parameters for *Weight Control* and *Sociability* were not significantly different  
249 from zero, the tests for equality between men and women indicate differences for FFM (in the case of  
250 *Weight Control*) and waist circumference. While *Weight Control* seems to be negatively associated  
251 with FFM and positively with waist circumference for men, the opposite seems to hold for women.

252 **Table 4**

253 Regression results for anthropometric measures and body composition

	BMI (kg/m <sup>2</sup> )					FFM (%)					Waist (cm)					
	Male		Female		Δ	Male		Female		Δ	Male		Female		Δ	
	B	SE	B	SE		B	SE	B	SE		B	SE	B	SE		
<i>Liking</i>	0.05	0.28	0.05	0.27		0.06	0.43	0.31	0.42		0.20	0.78	0.43	0.77		
<i>Habits</i>	-0.31	0.25	0.24	0.22	†	0.08	0.39	-0.61	0.34		-0.39	0.72	-0.36	0.63		
<i>Need &amp; Hunger</i>	0.18	0.27	-0.69	0.28	††	0.05	0.42	<b>1.63</b>	0.44	*** †††	0.56	0.77	<b>-2.46</b>	0.81	** †††	
<i>Health</i>	-0.74	0.28	-0.27	0.30		<b>1.72</b>	0.44	***	0.78	0.46	<b>-3.23</b>	0.81	***	-1.48	0.84	
<i>Convenience</i>	0.39	0.22	0.08	0.20		0.01	0.33	-0.29	0.31		0.59	0.62	0.46	0.58		
<i>Pleasure</i>	-0.04	0.27	-0.20	0.25		-0.28	0.41	-0.18	0.39		0.01	0.76	0.32	0.72		
<i>Traditional Eating</i>	0.35	0.29	<b>0.77</b>	0.25	*	-0.26	0.44	<b>-1.04</b>	0.38	*	1.52	0.82	1.36	0.70		
<i>Natural Concerns</i>	0.24	0.21	0.21	0.25		-0.53	0.33	-0.41	0.38		1.02	0.61	-0.45	0.71		
<i>Sociability</i>	-0.43	0.24	-0.03	0.20		0.33	0.37	0.52	0.31		-1.66	0.69	0.37	0.57	††	
<i>Price</i>	-0.04	0.25	-0.06	0.22		-0.22	0.39	0.34	0.34		0.26	0.72	0.65	0.63		
<i>Visual Appeal</i>	0.27	0.32	0.07	0.27		-0.57	0.49	-0.27	0.41		-0.01	0.90	1.32	0.76		
<i>Weight Control</i>	0.31	0.20	-0.04	0.18		-0.60	0.31	0.17	0.28	†	1.00	0.56	-0.47	0.52	†	
<i>Affect Regulation</i>	0.36	0.31	0.18	0.24		<b>-1.26</b>	0.48	*	-0.44	0.37	<b>2.38</b>	0.89	*	0.52	0.68	†
<i>Social Norms</i>	0.04	0.26	-0.03	0.21		0.06	0.40	0.08	0.33		-0.31	0.74	0.00	0.60		
<i>Social Image</i>	-0.01	0.54	-0.68	0.38		1.56	0.83	0.71	0.58		0.15	1.53	-2.08	1.07		
<i>Age group 2 (middle agers)<sup>a</sup></i>	5.04	0.87	5.24	0.92		-10.55	1.44	-10.24	1.54		17.69	2.38	14.20	2.57		
<i>Age group 3 (older adults, FS)<sup>a</sup></i>	4.61	1.20	3.62	1.16		-12.44	2.05	-12.99	1.92		20.20	3.59	15.98	3.03		
<i>Age group 4 (older adults, N)<sup>a</sup></i>	4.46	1.00	4.99	0.94		-15.46	1.49	-16.91	1.44		23.13	2.67	16.75	2.89		
<i>Constant</i>	21.78	2.10	21.11	2.50		81.38	3.42	68.12	3.84		83.18	6.47	80.22	6.68		
<i>R<sup>2</sup></i>	0.34					0.69					0.52					
<i>N</i>	429					429					429					

254 Note: \*\*\*, \*\*, and \* indicate significance at the 0.01-, 0.05-, and 0.1-level of regression coefficients based on adjusted p-values (Benjamini &  
 255 Hochberg, 1995); Δ-column: †††  $p < 0.01$ , ††  $p < 0.05$ , and †  $p < 0.1$ , based on post-regression Wald tests of equality between coefficients for males  
 256 and females. Standard errors in parentheses. BMI, Body Mass Index; FFM, fat free mass; <sup>a</sup>Reference: *Young adults*.

257 3.3. Regressions for energy and macronutrient intake

258 Tables 5a and 5b show the results of regression models for energy and nutrient intake. Results  
259 without differentiating coefficients by sex are depicted Table A3 in the appendix. The values of  $R^2$   
260 range from 0.30 to 0.46 and indicate a reasonable share of explained variation in dependent variables.  
261 Results suggest that the motives playing a relevant role for intakes differ to a certain degree from  
262 those relevant for anthropometry and body composition.

263  
264 Among men, the *Health* motive is linked to intakes of those nutrients that can be clearly identified as  
265 “healthy” or “unhealthy”. Specifically, the *Health* motive is positively associated with fiber intake  
266 (1.58;  $p < .001$ ) and negatively associated with alcohol intake (-1.67;  $p = .030$ ). The *Price* motive is  
267 significantly and positively associated with intakes of energy (63.77;  $p = .030$ ), carbohydrates (7.15;  
268  $p = .090$ ), and alcohol (1.23;  $p = .050$ ). Additionally, high scores on the *Habit* motive are positively  
269 related to higher intakes of carbohydrates (7.75;  $p = .090$ ) and high scores on the *Convenience* motive  
270 are linked to lower intake of energy (-46.96;  $p = .090$ ) and carbohydrates (-5.98;  $p = .090$ ). A final  
271 noteworthy result is the significant negative relation of *Social Norm* with alcohol intakes (-1.33;  $p =$   
272 .050).

273  
274 For women, a positive association of the *Health* motive was only found for fibre intake (1.41;  $p = .015$ ).  
275 We were not able to reject further null hypotheses of coefficients being significantly different from  
276 zero for women. Often, the estimates point to the same direction for men and a look at the relative  
277 size of the coefficients, e.g., for *Visual Appeal*, *Health*, and *Need & Hunger* suggests non-negligible  
278 effects. However, from a statistical perspective, these estimated parameters show too large standard  
279 errors for rejecting the null hypothesis.

280  
281 We find significantly higher coefficients for men compared to women for the motives *Habits* (energy,  
282 carbohydrates, fat, and fibre), *Need & Hunger* (fibre), and *Price* (energy, protein), and higher  
283 coefficients for women for *Pleasure* (protein), *Social Norms* (alcohol), and *Health* (alcohol).

284  
285

286 **Table 5a**

287 Regression results for energy, carbohydrate and fat intake

	Energy (kcal/day)					Carbohydrates (g/day)					Fat (g/day)					
	Male		Female		Δ	Male		Female		Δ	Male		Female		Δ	
	B	SE	B	SE		B	SE	B	SE		B	SE	B	SE		
<i>Liking</i>	18.80	21.78	-28.70	21.47		1.89	2.89	-4.06	2.85		0.19	1.00	-1.35	0.99		
<i>Habits</i>	41.52	19.88	-24.53	17.36	††	<b>7.75</b>	2.64	*	-2.77	2.30	†††	1.22	0.91	-1.04	0.80	†
<i>Need &amp; Hunger</i>	16.84	21.51	27.00	22.37		-0.23	2.85	5.04	2.97		1.15	0.99	1.08	1.03		
<i>Health</i>	25.45	22.44	22.86	23.46		5.99	2.98	2.07	3.11		0.77	1.03	0.87	1.08		
<i>Convenience</i>	<b>-46.96</b>	17.12	*	-17.52	16.09	<b>-5.98</b>	2.27	*	-2.25	2.14		-1.50	0.79	-0.41	0.74	
<i>Pleasure</i>	-31.97	21.04	14.92	20.01		-4.10	2.79	0.60	2.66		-1.38	0.97	0.78	0.92		
<i>Traditional Eating</i>	9.45	22.64	1.21	19.31		-2.29	3.00	1.87	2.56		1.09	1.04	-0.35	0.89		
<i>Natural Concerns</i>	13.00	16.85	17.04	19.62		0.43	2.24	3.87	2.60		0.88	0.77	0.13	0.90		
<i>Sociability</i>	3.61	19.11	-13.74	15.72		0.93	2.54	-4.33	2.09		-0.50	0.88	0.41	0.72		
<i>Price</i>	<b>63.77</b>	19.98	**	5.59	17.53	††	<b>7.15</b>	2.65	*	0.69	2.33	†	2.29	0.92	0.40	0.81
<i>Visual Appeal</i>	-1.05	24.92	45.34	21.15		-1.46	3.31	5.65	2.81		0.05	1.15	1.72	0.97		
<i>Weight Control</i>	-23.08	15.68	-13.40	14.51		-2.67	2.08	-2.44	1.93		-1.45	0.72	-1.05	0.67		
<i>Affect Regulation</i>	-19.01	24.75	-25.16	18.84		-1.78	3.28	-2.12	2.50		-0.81	1.14	-1.19	0.87		
<i>Social Norms</i>	-28.47	20.56	10.83	16.74		-1.82	2.73	0.68	2.22		-0.78	0.95	0.61	0.77		
<i>Social Image</i>	67.00	42.51	7.51	29.77		7.64	5.64	1.01	3.95		2.37	1.95	0.16	1.37		
<i>Age group 2 (middle agers)<sup>a</sup></i>	15.48	83.27	14.70	58.88		-11.89	11.15	-7.10	7.44		5.02	3.97	4.53	3.10		
<i>Age group 3 (older adults, FS)<sup>a</sup></i>	154.91	121.93	162.07	83.48		0.79	16.80	18.05	10.32		12.66	5.71	8.54	4.68		
<i>Age group 4 (older adults, N)<sup>a</sup></i>	56.86	95.34	114.02	67.39		-1.05	12.50	14.36	8.84		5.67	4.21	6.29	3.37		
<i>Constant</i>	1552.88	211.66	1493.37	173.35		167.35	28.62	164.82	23.15		67.47	9.42	63.94	7.99		
<i>R<sup>2</sup></i>		0.46					0.35					0.39				
<i>N</i>		429					429					429				

288 Note: \*\*\*, \*\*, and \* indicate significance at the 0.01-, 0.05-, and 0.1-level of regression coefficients based on adjusted p-values (Benjamini &  
289 Hochberg, 1995); Δ-column: †††  $p < 0.01$ , ††  $p < 0.05$ , and †  $p < 0.1$ , based on post-regression Wald tests of equality between coefficients for males  
290 and females. Standard errors in parentheses. BMI, Body Mass Index; FFM, fat free mass; <sup>a</sup>Reference: *Young adults*.

291 **Table 5b**

292 Regression results for protein, fibre, and alcohol intakes

	Protein (g/day)					Fibre (g/day)					Alcohol (g/day)						
	Male		Female		Δ	Male		Female		Δ	Male		Female		Δ		
	B	SE	B	SE		B	SE	B	SE		B	SE	B	SE			
<i>Liking</i>	0.82	0.93	-0.20	0.91		-0.14	0.40	-0.41	0.39		0.86	0.48	0.07	0.48			
<i>Habits</i>	0.82	0.85	-0.95	0.74		0.86	0.36	-0.49	0.32	†††	-0.61	0.44	0.03	0.38			
<i>Need &amp; Hunger</i>	1.23	0.91	0.22	0.95		-0.26	0.39	0.98	0.41	††	0.38	0.48	-0.59	0.50			
<i>Health</i>	1.34	0.95	1.38	1.00		<b>1.58</b>	0.41	***	<b>1.41</b>	0.43	**	<b>-1.67</b>	0.50	**	0.12	0.52	††
<i>Convenience</i>	-1.13	0.73	-0.89	0.68		-0.68	0.31	-0.21	0.29		-0.70	0.38	-0.15	0.36			
<i>Pleasure</i>	-1.95	0.89	1.17	0.85	††	-0.43	0.39	-0.22	0.37		0.71	0.47	0.10	0.44			
<i>Traditional Eating</i>	0.77	0.96	-0.80	0.82		-0.57	0.41	-0.51	0.35		0.87	0.50	0.05	0.43			
<i>Natural Concerns</i>	0.56	0.72	-0.45	0.83		0.58	0.31	0.40	0.36		0.22	0.37	0.26	0.43			
<i>Sociability</i>	-0.38	0.81	-0.01	0.67		0.20	0.35	-0.20	0.29		0.82	0.42	0.06	0.35			
<i>Price</i>	1.57	0.85	-0.59	0.75	†	0.10	0.37	-0.39	0.32		<b>1.23</b>	0.44	**	0.26	0.39		
<i>Visual Appeal</i>	-0.10	1.06	2.18	0.90		-0.30	0.46	0.63	0.39		0.66	0.55	-0.27	0.47			
<i>Weight Control</i>	0.16	0.67	1.03	0.62		-0.21	0.29	-0.21	0.27		-0.06	0.35	0.19	0.32			
<i>Affect Regulation</i>	-0.18	1.05	-1.14	0.80		-0.34	0.45	-0.16	0.34		-0.54	0.55	-0.20	0.42			
<i>Social Norms</i>	-1.15	0.87	0.57	0.71		-0.24	0.38	-0.10	0.31		<b>-1.33</b>	0.46	**	0.06	0.37	††	
<i>Social Image</i>	3.44	1.81	-0.02	1.27		1.35	0.78	0.60	0.54		0.09	0.94	0.26	0.66			
<i>Age group 2 (middle agers)<sup>a</sup></i>	0.42	3.46	0.32	2.94		-2.95	1.51	-2.05	1.25		2.69	2.28	0.51	0.47			
<i>Age group 3 (older adults, FS)<sup>a</sup></i>	1.95	5.02	2.17	4.13		0.58	2.35	2.13	1.82		4.47	3.28	0.47	0.76			
<i>Age group 4 (older adults, N)<sup>a</sup></i>	-2.31	3.99	-0.55	3.25		0.02	1.59	0.37	1.43		3.05	2.38	0.37	0.61			
<i>Constant</i>	56.31	7.98	57.29	7.65		14.76	3.52	14.94	3.52		5.67	6.31	2.12	1.36			
<i>R<sup>2</sup></i>			0.39					0.30					0.39				
<i>N</i>			429					429					429				

293 Note: \*\*\*, \*\*, and \* indicate significance at the 0.01-, 0.05-, and 0.1-level of regression coefficients based on adjusted p-values (Benjamini &  
294 Hochberg, 1995); Δ-column: †††  $p < 0.01$ , ††  $p < 0.05$ , and †  $p < 0.1$ , based on post-regression Wald tests of equality between coefficients for males  
295 and females. Standard errors in parentheses. BMI, Body Mass Index; FFM, fat free mass; <sup>a</sup>Reference: *Young adults*.

296 **4. Discussion**

297 The present study aimed to investigate associations of eating motives with anthropometric  
298 parameters, body composition, and dietary intake. The analysis was based on comprehensive data  
299 from the *enable* study (Brandl et al., 2020) and used the 15 TEMS factors (Renner et al., 2012) to  
300 measure eating motivation.

301 *4.1. Motives and their association with nutritional status and dietary intake*

302 A major result of this study was that there was a difference between whether a motive is ranked highly  
303 on average and whether that motive is associated with variation in anthropometric parameters. Only  
304 two out of the top five motives (*Health, Need & Hunger*) showed significant associations with  
305 anthropometric parameters, while effects for *Liking* and *Habits*, in particular, were absent. *Liking*  
306 appears to be the most frequent eating motive throughout for people with healthy and unhealthy diets  
307 and has low discriminatory power (Glanz et al., 1998; Pollard et al., 1998). Likewise, *Habits* as a major  
308 determinant of eating behavior (Köster, 2009; Pollard et al., 2002) can be “good” or “bad” and may  
309 thus lead to more or less desirable nutritional outcomes.

310  
311 *Health* emerged as a central motive with strong positive effects on desirable values of both  
312 anthropometric parameters and nutrient intakes. While we find significant associations for men only,  
313 most estimated parameters for women show the same direction and substantial magnitudes  
314 compared to coefficients of other motives. We were not able to reject hypotheses of equal parameters  
315 for the *Health* motive for men and women. Hence, we cannot interpret our findings in the sense that  
316 women’s anthropometry or dietary intakes are not affected by *Health* or that the effects differ from  
317 those for men. The restricted sample size as well as the high scores of the *Health* motive paired with  
318 a low variability in women (Mean = 4.9; SD = 1.1, Table 3) apparently impede a more precise  
319 parameter estimation.

320  
321 Our findings are in line with previous studies that showed a positive association of stronger health  
322 attitude or motivation to eat healthily with various desirable eating behaviors (Eertmans et al., 2005;  
323 Hearty et al., 2007; Naughton et al., 2015; Pollard et al., 1998; Rempe et al., 2020). At the same time,  
324 this result underlines that individual health motivation matters. While an obesogenic environment  
325 has been increasingly blamed as the main source of increasing waistlines (Swinburn & Egger, 2002),  
326 our results suggest that personal motivation to eat healthily still needs to be considered as an  
327 important driver of health outcomes and a relevant factor in policy design. The results on energy



328 nutrient intakes suggest that the *Health* motive is especially relevant for unambiguously healthy or  
329 unhealthy nutrients. It is positively associated with fiber intake and negatively associated with  
330 alcohol intake. The positive but insignificant coefficients for energy, carbohydrates, protein, and fat  
331 suggest that the *Health* motive is only weakly related to specific dietary intakes.

332  
333 The association of *Need & Hunger* with anthropometric parameters confirms previous studies on  
334 intuitive eating (Tylka, 2006). We found a strongly positive association of this factor with FFM and a  
335 strongly negative one with waist circumference for women. Characterized as “eating based on  
336 perceived internal states”, intuitive or adaptive eating has been found to be negatively related to BMI,  
337 supportive of weight maintenance, and positively associated with physical health indicators, dietary  
338 intake and behavior and psychological well-being (Augustus-Horvath & Tylka, 2011; Tylka, 2006; van  
339 Dyke & Drinkwater, 2014). The pronounced differences between men and women may be related to  
340 “pressure for thinness” and “internalization of a thin ideals” (Tylka, 2006) imposed by media and  
341 society, with potentially adverse effects as women then respond more to external cues. Hence, this  
342 factor may have more discriminatory power for women. While increased pressure can also be exerted  
343 on men regarding muscle mass and strength, most studies were so far conducted in female samples  
344 (van Dyke & Drinkwater, 2014).

345  
346 Results suggest that *Traditional Eating* may be an unfavorable factor for anthropometric parameters  
347 in women. Literature has only recently begun to address the question of what constitutes traditional  
348 eating and food products (Sproesser, Imada et al., 2018; Sproesser, Ruby et al., 2019), emphasizes  
349 that it depends on the context, but also suggests that traditional is more related to unprocessed,  
350 natural, and healthier food. The single items in the TEMS factor (“...because it belongs to certain  
351 situations”; “...out of traditions (e.g. family traditions, special occasions)”; “...because I grew up with  
352 it”), however, suggest that traditional food in our study sample may refer to old-fashioned Bavarian-  
353 style foods, recipes, and cooking practices heavy in meat and fat, as well as large portion sizes and  
354 rigid meal times. In this particular case, high scores on *Traditional Eating* may be inappropriate for  
355 modern, more sedentary lifestyles.

356  
357 Our results indicate that higher scores of *Affect Regulation*, i.e., eating in response to sadness,  
358 frustration, and loneliness as measured by TEMS, are associated with lower FFM and higher Waist  
359 Circumference. This finding is in line with literature on eating as a strategy for regulating negative  
360 affective states (Canetti et al., 2002; Macht, 2008; Macht & Simons, 2011). The strong coefficients for  
361 FFM and Waist circumference and the observation that the results hold for the male sample only

362 suggest that emotional eating is particularly related to increased fat accumulation in the abdominal  
363 region. In contrast, *Affect Regulation* did not show any association with nutrient intakes. One potential  
364 reason could be inflated self-reports on the TEMS items as persons with overweight or obesity may  
365 be sensitized for the drivers of their eating behavior (Macht & Simons, 2011). A second explanation  
366 may be underreporting of hedonic or guilt-related food intake during phases of negative affective  
367 states (Haftenberger et al., 2010).

368  
369 *Convenience*, the motivation to choose food because it is most convenient, quick, and easy to prepare,  
370 is consistently negatively associated with intakes of most nutrients. However, this finding was more  
371 pronounced among men and significant for energy and carbohydrate intake. Previous literature has  
372 pointed to the emergence and easy accessibility of “convenience” foods (such as fast food, take-away,  
373 snacks, frozen meals, etc.) as a potentially important driver of soaring obesity rates (Dixon et al.,  
374 2006). Our results suggest that people with a high *Convenience* motive may have a rather low  
375 preoccupation with food and the aim to keep food preparation simple leads to lower overall intake.  
376 Hence, it may not be the convenience part in “convenience” foods that is detrimental to nutritional  
377 status, but rather other characteristics such as being high-fat, high-energy foods.

378  
379 The positive association of the *Price* motive with the intake of energy, carbohydrates, and alcohol – in  
380 particular among men – is in line with literature that connects energy density and energy costs with  
381 obesity. In this framework poorer households are assumed to economize on energy costs leading to  
382 higher intakes of palatable, energy-dense foods (Drewnowski & Specter, 2004). While our results  
383 support these arguments, the *Price* motives was unrelated to anthropometric parameters, hence  
384 higher intakes do not seem to translate into increased body mass or body fat in the long term. A  
385 potential explanation could be that higher physical activity at work or during leisure may compensate  
386 for higher intakes.

387  
388 A final motivational factor with interesting results regarding dietary intake parameters is *Social*  
389 *Norms*, where higher values are related to lower alcohol intake for men. Social norms have been  
390 identified as one of the best predictors of alcohol consumption (Neighbors et al., 2007). The literature  
391 distinguishes descriptive norms (i.e., the perceived prevalence of drinking by a typical member of the  
392 peer group) and injunctive norms (i.e., the perception of how much others such as friends or parents  
393 approve of someone’s drinking behavior) (Neighbors et al., 2007). Both a higher perception of the  
394 usual alcohol quantity consumed and higher perceived approval of drinking predict higher quantities  
395 consumed. Against this background, our result is surprising at first sight given the formulation of this

396 factor's items ('...because it would be impolite not to eat it'; '... to avoid disappointing someone who  
397 is trying to make me happy'; ... because I am supposed to eat it'). We would expect someone scoring  
398 higher on these to be more susceptible towards invitations of others to have a drink. However, the  
399 negative correlation we find suggests that this motive rather measures a general tendency to adhere  
400 to what others or society deem appropriate. Apparently, people who pay much attention to the  
401 judgement of others (report to) drink less.

402

#### 403 4.2. Relation between TEMS and other approaches to identify motives

404 The endeavor of this study was to investigate the association of a comprehensive set of eating motives  
405 with nutritional status and dietary intake. We used the TEMS as a "concise questionnaire that allows  
406 for a comprehensive, systematic, and psychometric sound measurement and investigation of motives  
407 for normal eating behavior" (Renner et al., 2012). As each TEMS factor is based on just three items, a  
408 discussion is in order whether the motives participants endorse on this concise and simple  
409 instrument and their relationships to anthropomorphic data align well with those found for more  
410 specialized, extensive instruments such as the Three-factor Eating Questionnaire (TFEQ) (Stunkard  
411 & Messick, 1985), the Dutch Eating Behavior Questionnaire (DEBQ) (van Strien et al., 1986), and  
412 others. We would briefly like to highlight how "core, higher order motives of food choice" (Renner et  
413 al., 2012) such as health, body weight control, affect regulation, or hunger relate to anthropometry  
414 when measured by TEMS versus measured by other instruments.

415

416 Our result of a higher *Health* motive being related to healthier anthropometry and body composition  
417 (for males) aligns well with findings from studies using different instruments to measure eating for  
418 health reasons. These include the FCQ (Eertmans et al., 2005; Pollard et al., 1998), self-administered  
419 questionnaires (Hearty et al., 2007), a Healthy Eating Motivation Score based on the FCQ and the  
420 Health and Taste Attitude Scale (HTAS) (Naughton et al., 2015), as well as TEMS itself (Rempe et al.,  
421 2020). Likewise, the finding that higher *Health* values are positively correlated with healthier intake  
422 patterns for specific nutrients (higher fiber, lower alcohol) is in line with studies using the Health and  
423 Taste Attitude Scale (Roininen et al., 1999) to study preference for reformulated or functional food  
424 products.

425

426 The same holds for the *Need & Hunger* motive, where we find a negative association with fat-free mass  
427 and a positive one with waist circumference (for females). This is coherent with research reporting  
428 high scores on different versions of the Intuitive Eating Scale (Tylka, 2006) to be associated with

429 lower BMI (Augustus-Horvath & Tylka, 2011; Tylka, 2006; van Dyke & Drinkwater, 2014). The result  
430 of a higher score for *Affect Regulation* being negatively related to a healthy nutritional status is in line  
431 with studies that measure emotional eating using the “Ways of Coping Checklist” (Laitinen et al.,  
432 2002), the TFEQ (Konttinen, Männistö et al., 2010; Konttinen, Silventoinen et al., 2010; Péneau et al.,  
433 2013) as well as the DEBQ (Pothos et al., 2009; Sung et al., 2009; van Strien et al., 2009).

434  
435 Finally, *Weight Control* is a major TEMS factor that relates to measures of cognitive restraint (TFEQ)  
436 or dietary restraint (DEBQ), respectively. Our results for this motive were rather inconclusive and  
437 insignificant. Coefficients indicate that higher values of *Weight Control* for men relate positively to  
438 BMI and WC, and negatively to FFM. Coefficients for women significantly differed from these and  
439 pointed to opposite directions. Results suggest that higher scores on this factor were associated with  
440 lower intakes of energy, carbohydrates, and fat. Literature provides a mixed picture as well. Many  
441 studies find positive correlations with BMI or overweight based on the TFEQ (Anglé et al., 2009;  
442 Cappelleri et al., 2009; Lauzon-Guillain et al., 2006) and the DEBQ (Lluch et al., 2000; Olea López &  
443 Johnson, 2016; Snoek et al., 2007) . However, Johnson et al. (2012) report that these are mostly seen  
444 in normal-weight samples, while for samples of persons with overweight or obesity, the relation is  
445 often negative. The authors interpret positive restraint-BMI associations such that restraint may be  
446 “acting as a marker for overeating tendencies”. Price et al. (2015) show that restraint measures from  
447 different instruments (TFEQ, DEBQ) load on the same factor, which is positively associated with BMI.  
448 The authors interpret this relation to reflect “unsuccessful attempts at dietary control”.

449  
450 In summary, these examples suggest that the relations of TEMS motives with anthropometry are  
451 congruent to findings for other measures from more specialized and extensive instruments. TEMS  
452 motives seem able to capture relevant constructs accurately and to provide a large set of potential  
453 predictors of anthropometric measures and their confounders at the same time. The simple  
454 comparison of different instruments and their relations to anthropometry over distinct samples and  
455 studies as done here needs to be confirmed by either comparing relations for different measures  
456 based on the same sample or more systematic meta-analyses.

457  
458 A related question concerns the comparability of TEMS motives and behavioral measures of related  
459 constructs such as the “eating in the absence of hunger” test and their relation to anthropomorphic  
460 data. While such tests have been mostly used in research on children (French et al., 2012; Goldschmidt  
461 et al., 2017), Carnell and Wardle (2007) compared parental reports on the Child Eating Behaviour  
462 Questionnaire (CEBQ) to results from several behavioral tests. They found that higher values for the

463 subscale “satiety responsiveness” were associated with lower intakes in the “eating in the absence of  
464 hunger” test and higher values in the “enjoyment of food” subscale with higher intakes in the test.  
465 They concluded that “the CEBQ is capable of capturing important facets of children’s eating behavior  
466 that have previously been assessed only with behavioral tests”. These findings should inspire future  
467 research comparing behavioral and self-reported measures for adults as well.

468  
469 Another example is the social facilitation of eating where a recent review and meta-analysis (Ruddock  
470 et al., 2019) has shown, which effort is required to obtain empirical evidence by experiments and  
471 naturalistic observations. While the evidence strongly suggests that eating with friends and family  
472 increases food and energy intake, there is no evidence yet, whether the phenomenon of social  
473 facilitation of eating affects the long-term energy balance positively (Ruddock et al., 2021).  
474 Accordingly, there are no insights on the relation to anthropometric measures. Our findings indicate  
475 higher *Sociability* to be related to lower BMI, higher FFM, and inconclusive evidence on waistlines  
476 (lower for men, higher for females). Future research is needed on how self-reported measures  
477 compare to high-effort behavioral tests and procedures.

478

#### 479 *4.3. Implications for public health interventions*

480 The results of the present study suggest a series of implications for design and strategic levers of  
481 public health measures. The identification of the *Health* motive as a major factor in relation to  
482 anthropometric parameters may be used in two ways. The nearest thought is attempting to increase  
483 those persons’ *Health* motivation, where it is low. However, changing the underlying attitudes, values,  
484 and social norms may be complex and difficult, promising only little success. Alternatively, one could  
485 approach the group with low health motivation through factors on which they have a higher score.  
486 Motives for which this study found a low or negative correlation with the health motive are *Price*,  
487 *Visual Appeal*, *Affect Regulation*, *Social Norms*, and *Social Image*. A more promising strategy could be  
488 to make healthy foods more appealing regarding these factors, in particular those that are  
489 significantly related to the nutritional status. This could be done, for example, through social  
490 marketing campaigns, but certainly also through product development, innovation, and marketing  
491 strategies in the food industry, the food service sector, and retailing.

492

493 A similar case could be made for *Traditional Eating*, where eating is guided by specific meals, meal  
494 occasions, or sensory components. A lever would be to let people make new experiences, opening up  
495 alternatives to rigid structures and standard dishes. Such a strategy might be particularly fruitful, if

496 people were provided with healthier alternatives that match their taste, needs, and routines. For  
497 *Affect Regulation*, with its prominent role in the male sample, a strategy would be to provide men with  
498 alternative means to cope with their emotions other than eating and drinking. From a broader  
499 perspective it would also be a societal and institutional challenge to reduce potential stressors and  
500 sources for negative emotions at work and in private lives in the first place. Results for *Need & Hunger*  
501 as a beneficial factor for nutritional status suggest increasing awareness on feelings of hunger and  
502 satiety as well as equipping people with techniques and methods to listen more carefully to internal  
503 cues.

#### 504 4.4. *Strengths, limitations and avenues for future research*

505 The major strength of the study is that it could rely on comprehensive data for eating motivations,  
506 nutrient intakes, and anthropometric parameters from a large sample covering different age groups.  
507 The results highlight that it is important to investigate the associations of motives with eating  
508 behavior to derive meaningful and promising strategies for public health measures. The results also  
509 suggest that it is important to look to look separately at nutrition behavior, i.e. the observed dietary  
510 intake, and nutritional status, i.e. anthropometry and body composition. While some motives are  
511 consistently related to both stages, some are exclusively related to nutritional status and others  
512 exclusively with dietary intake. This has a major effect on the implications that follow from the study  
513 of motives. If we would only measure the effect of the price motivation on dietary intake but not on  
514 nutritional status, we would agree with previous literature that price-sensitive consumers would  
515 consume more energy and have a higher chance of developing obesity. However, as we have  
516 additional data that indicate no direct relation to nutritional status, we are more reluctant to draw  
517 such a conclusion.

518  
519 Of course, this raises questions about the reasons for the discrepancy between dietary intake and  
520 nutritional status. One missing piece in the equation is the level of physical activity that may be higher  
521 in people with a high score on the price motive. However, additional checks with several indicators  
522 for physical activity did not confirm this hypothesis. Another limitation is that the cross sectional  
523 nature of our data set only allows us to study correlations but not direct causation. This could affect  
524 especially coefficients for *Weight Control* motive, where persons with overweight or obesity may  
525 report a higher motivation to control their weight leading to reverse causation. Also, as Macht and  
526 Simons (2011) have argued, persons with overweight or obesity may be more sensitized regarding  
527 the underlying reasons of their eating behavior so that their self-reports on *Affect Regulation* may be  
528 more accurate than for participants with normal weight.

529 A last point touches issues of intra- and inter-individual variability in dietary intake or motives. The  
530 implications of day-to-day variability on measurements of usual food and nutrient intakes have been  
531 already discussed by literature (Palaniappan et al., 2003). We are confident that we safeguarded our  
532 measures of dietary intake from high variability in 24 h-recalls by the careful procedure we apply to  
533 obtain usual dietary intake as described above. What may be more relevant is that both motives and  
534 intakes present in the short- and medium-term period surrounding data collection may be different  
535 from those that are responsible for observed anthropometry and body composition in the long-term.  
536 Wahl et al. (2020) recently showed that motives from TEMS measured in single-time-point  
537 questionnaires, interpreted as “situation-stable dispositions (traits)” correlated well with “in-the-  
538 moment (state) assessments”, i.e. experienced eating motives in the moment of consumption.  
539 However, trait motives had higher average values than state motives and intraindividual motive  
540 profiles differed substantially between participants. The authors argue that representativeness  
541 heuristics or response biases may have contributed to these observations. While we exclusively  
542 measure trait motives, we cannot rule that these, too, vary over time. Hence, it could be that a range  
543 of motives is rather stable over time and correlates to long-term nutritional status, while other  
544 motives may be more medium to short term (or more salient during the study period) and correlate  
545 more closely with the nutrition behavior observed during the study. To resolve these questions, it  
546 would be helpful to have longitudinal data with repeated measurement of motives to assess whether  
547 they are stable as traits or whether some of them vary in a more pronounced way than others.

## 548 **5. Conclusion**

549  
550 This study identified specific eating motives based on TEMS as potential drivers of energy and  
551 macronutrient intake and nutritional status. Controlling for sex, age, income, and education, we found  
552 motives related to *Health, Need & Hunger, Traditional Eating, and Affect Regulation* to be strongly  
553 associated with BMI, FFM, and waist circumference. While motives related to *Liking* and *Habits* were  
554 stated as most frequent drivers of eating behavior, we did not find a significant relation to the  
555 nutritional status. Other motives such as *Price* and *Convenience* were associated with intake, but not  
556 anthropometric measures. The results highlight the need for more differentiated analyses regarding  
557 eating motives, beyond simple comparisons of relative importance, towards the association of  
558 motives to consumption patterns, dietary intake, and nutritional status. Such analyses would yield  
559 even more specific and targeted implications for nutrition interventions and counselling. Future  
560 research should seek to collect more longitudinal data to trace changes in motives and nutrition in  
561 the long term and control for unobservable third-factor effects.

562

563 **Author contributions**

564 Data base establishment: J.R. sociodemographics and TEMS; B.B., T.S., D.V., E.K., H.S., H.H. dietary  
565 intakes, anthropometric parameters, and body composition. J.L. and N.W. computation of usual  
566 intakes of participants. M.S. and B.B. conceived and designed the structure of the statistical analysis,  
567 M.S. analyzed and interpreted results, M.S. and B.B. conceptualized the manuscript and wrote the  
568 paper. M.S., B.B., E.K., J.R., H.S., T.S., D.V., J.L., N.W., H.H. revised it critically. All authors have read and  
569 approved the final version of the article.

570

571 **Conflict of interest**

572 The authors declare no conflict of interest. This work is funded by the German Ministry of Education  
573 and Research (BMBF, grant no. 01EA1409A). The funding sponsor had no role in the design of the  
574 study; in the collection, analyses or interpretation of data; in the writing of the manuscript, and in the  
575 decision to publish the results.

576

577 **Data availability**

578 The data used for this study are part of *Project Z* within the *enable* cluster of nutrition research.  
579 Researchers who are interested in working with these data can request access at the study center  
580 (please contact Dr. Beate Brandl, beate.brand@tum.de). Applicants will have to sign a data use  
581 agreement including a confirmation that they will comply with the General Data Protection  
582 Regulation (GDPR).

583

584 **Acknowledgments**

585 We thank Laura Lepiarczyk for her outstanding support.



## References

- Anglé, S., Engblom, J., Eriksson, T., Kautiainen, S., Saha, M.-T., Lindfors, P., Lehtinen, M., & Rimpelä, A. (2009). Three factor eating questionnaire-R18 as a measure of cognitive restraint, uncontrolled eating and emotional eating in a sample of young Finnish females. *International Journal of Behavioral Nutrition and Physical Activity*, 6(1), 41. <https://doi.org/10.1186/1479-5868-6-41>
- Ares, G., & Gábaro, A. (2007). Influence of gender, age and motives underlying food choice on perceived healthiness and willingness to try functional foods. *Appetite*, 49(1), 148–158. <https://doi.org/10.1016/j.appet.2007.01.006>
- Augustus-Horvath, C. L., & Tylka, T. L. (2011). The acceptance model of intuitive eating: A comparison of women in emerging adulthood, early adulthood, and middle adulthood. *Journal of Counseling Psychology*, 58(1), 110–125. <https://doi.org/10.1037/a0022129>
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(1), 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
- Brandl, B., Skurk, T., Rennekamp, R., Hannink, A., Kiesswetter, E., Freiherr, J., Ihsen, S., Roosen, J., Klingenspor, M., Haller, D., Krautwurst, D., Hofmann, T., Linseisen, J., Volkert, D., & Hauner, H. (2020). A phenotyping platform to characterize healthy individuals across four stages of life - The enable study. *Frontiers in Nutrition*, 7, 582387. <https://doi.org/10.3389/fnut.2020.582387>
- Canetti, L., Bachar, E., & Berry, E. M. (2002). Food and emotion. *Behavioural Processes*, 60(2), 157–164. [https://doi.org/10.1016/S0376-6357\(02\)00082-7](https://doi.org/10.1016/S0376-6357(02)00082-7)
- Cappelleri, J. C., Bushmakin, A. G., Gerber, R. A., Leidy, N. K., Sexton, C. C., Lowe, M. R., & Karlsson, J. (2009). Psychometric analysis of the Three-Factor Eating Questionnaire-R21: Results from a large diverse sample of obese and non-obese participants. *International Journal of Obesity*, 33(6), 611–620. <https://doi.org/10.1038/ijo.2009.74>
- Carnell, S., & Wardle, J. (2007). Measuring behavioural susceptibility to obesity: Validation of the child eating behaviour questionnaire. *Appetite*, 48(1), 104–113. <https://doi.org/10.1016/j.appet.2006.07.075>
- Chambers, D., Phan, U. T. X., Chanadang, S., Maughan, C., Sanchez, K., Di Donfrancesco, B., Gomez, D., Higa, F., Li, H., Chambers, E., & Esen, E. (2016). Motivations for food consumption during specific eating occasions in Turkey. *Foods (Basel, Switzerland)*, 5(2). <https://doi.org/10.3390/foods5020039>
- Chou, S.-Y., Grossman, M., & Saffer, H. (2004). An economic analysis of adult obesity: Results from the Behavioral Risk Factor Surveillance System. *Journal of Health Economics*, 23(3), 565–587. <https://doi.org/10.1016/j.jhealeco.2003.10.003>
- Destatis. (2018). *Wirtschaftsrechnungen 2013: Einkommens- und Verbrauchsstichprobe - Einkommensverteilung in Deutschland. Fachserie 15: Heft 6.*
- Destatis. (2019). *Statistisches Jahrbuch Deutschland 2019.* Statistisches Bundesamt. [https://www.statistischebibliothek.de/mir/receive/DEAusgabe\\_mods\\_00004527](https://www.statistischebibliothek.de/mir/receive/DEAusgabe_mods_00004527)
- Destatis. (2021). *Bevölkerung: Deutschland, Stichtag, Altersjahre 2017.* <https://www-genesis.destatis.de/genesis/online>
- Dixon, J. M., Hinde, S. J., & Banwell, C. L. (2006). Obesity, convenience and “phood”. *British Food Journal*, 108(8), 634–645. <https://doi.org/10.1108/00070700610682328>

- Drewnowski, A., & Specter, S. E. (2004). Poverty and obesity: The role of energy density and energy costs. *The American Journal of Clinical Nutrition*, 79(1), 6–16.  
<https://doi.org/10.1093/ajcn/79.1.6>
- Eertmans, A., an Victoir, Vansant, G., & van den Bergh, O. (2005). Food-related personality traits, food choice motives and food intake: Mediator and moderator relationships. *Food Quality and Preference*, 16(8), 714–726. <https://doi.org/10.1016/j.foodqual.2005.04.007>
- Freese, J., Feller, S., Harttig, U., Kleiser, C., Linseisen, J., Fischer, B., Leitzmann, M. F., Six-Merker, J., Michels, K. B., Nimptsch, K., Steinbrecher, A., Pischon, T., Heuer, T., Hoffmann, I., Jacobs, G., Boeing, H., & Nöthlings, U. (2014). Development and evaluation of a short 24-h food list as part of a blended dietary assessment strategy in large-scale cohort studies. *European Journal of Clinical Nutrition*, 68(3), 324–329. <https://doi.org/10.1038/ejcn.2013.274>
- French, S. A., Epstein, L. H., Jeffery, R. W., Blundell, J. E., & Wardle, J. (2012). Eating behavior dimensions. Associations with energy intake and body weight. A review. *Appetite*, 59(2), 541–549. <https://doi.org/10.1016/j.appet.2012.07.001>
- Glanz, K., Basil, M., Maibach, E., Goldberg, J., & Snyder, D. (1998). Why Americans eat what they do. *Journal of the American Dietetic Association*, 98(10), 1118–1126.  
[https://doi.org/10.1016/S0002-8223\(98\)00260-0](https://doi.org/10.1016/S0002-8223(98)00260-0)
- Goldschmidt, A. B., Crosby, R. D., Cao, L., Pearson, C. M., Utzinger, L. M., Pacanowski, C. R., Mason, T. B., Berner, L. A., Engel, S. G., Wonderlich, S. A., & Peterson, C. B. (2017). Contextual factors associated with eating in the absence of hunger among adults with obesity. *Eating Behaviors*, 26, 33–39. <https://doi.org/10.1016/j.eatbeh.2017.01.005>
- Haftenberger, M., Heuer, T., Heidemann, C., Kube, F., Krems, C., & Mensink, G. B. M. (2010). Relative validation of a food frequency questionnaire for national health and nutrition monitoring. *Nutrition Journal*, 9, 36. <https://doi.org/10.1186/1475-2891-9-36>
- Hasselbach, J. L., & Roosen, J. (2015). Motivations behind preferences for local or organic food. *Journal of International Consumer Marketing*, 27(4), 295–306.  
<https://doi.org/10.1080/08961530.2015.1022921>
- Hawkes, C., Smith, T. G., Jewell, J., Wardle, J., Hammond, R. A., Friel, S., Thow, A. M., & Kain, J. (2015). Smart food policies for obesity prevention. *The Lancet*, 385(9985), 2410–2421.  
[https://doi.org/10.1016/S0140-6736\(14\)61745-1](https://doi.org/10.1016/S0140-6736(14)61745-1)
- Hearty, A. P., McCarthy, S. N., Kearney, J. M., & Gibney, M. J. (2007). Relationship between attitudes towards healthy eating and dietary behaviour, lifestyle and demographic factors in a representative sample of Irish adults. *Appetite*, 48(1), 1–11.  
<https://doi.org/10.1016/j.appet.2006.03.329>
- Hebden, L., Chan, H. N., Louie, J. C., Rangan, A., & Allman-Farinelli, M. (2015). You are what you choose to eat: Factors influencing young adults' food selection behaviour. *Journal of Human Nutrition and Dietetics : The Official Journal of the British Dietetic Association*, 28(4), 401–408. <https://doi.org/10.1111/jhn.12312>
- Himmerich, S., Gedrich, K., & Karg G. (2003). *Bayerische Verzehrsstudie (BVS) II - Abschlussbericht (in German) - Im Auftrag des Bayerischen Staatsministeriums fuer Umwelt, Gesundheit und Verbraucherschutz.*
- Honkanen, P., Verplanken, B., & Olsen, S. O. (2006). Ethical values and motives driving organic food choice. *Journal of Consumer Behaviour*, 5(5), 420–430. <https://doi.org/10.1002/cb.190>
- Johnson, F., Pratt, M., & Wardle, J. (2012). Dietary restraint and self-regulation in eating behavior. *International Journal of Obesity*, 36(5), 665–674. <https://doi.org/10.1038/ijo.2011.156>
- Just, D. R., & Gabrielyan, G. (2016). Food and consumer behavior: why the details matter. *Agricultural Economics*, 47(S1), 73–83. <https://doi.org/10.1111/agec.12302>

- Kline, R. B. (2011). *Principles and practice of structural equation modeling* (3. ed.). *Methodology in the social sciences*. Guilford Press.  
<http://site.ebrary.com/lib/academiccompletetitles/home.action>
- Konttinen, H., Männistö, S., Sarlio-Lähteenkorva, S., Silventoinen, K., & Haukkala, A. (2010). Emotional eating, depressive symptoms and self-reported food consumption. A population-based study. *Appetite*, *54*(3), 473–479. <https://doi.org/10.1016/j.appet.2010.01.014>
- Konttinen, H., Silventoinen, K., Sarlio-Lähteenkorva, S., Männistö, S., & Haukkala, A. (2010). Emotional eating and physical activity self-efficacy as pathways in the association between depressive symptoms and adiposity indicators. *The American Journal of Clinical Nutrition*, *92*(5), 1031–1039. <https://doi.org/10.3945/ajcn.2010.29732>
- Köster, E. P. (2009). Diversity in the determinants of food choice: A psychological perspective. *Food Quality and Preference*, *20*(2), 70–82. <https://doi.org/10.1016/j.foodqual.2007.11.002>
- Laitinen, J., Ek, E., & Sovio, U. (2002). Stress-related eating and drinking behavior and body mass index and predictors of this behavior. *Preventive Medicine*, *34*(1), 29–39.  
<https://doi.org/10.1006/pmed.2001.0948>
- Lauzon-Guillain, B. de, Basdevant, A., Romon, M., Karlsson, J., Borys, J.-M., & Charles, M. A. (2006). Is restrained eating a risk factor for weight gain in a general population? *The American Journal of Clinical Nutrition*, *83*(1), 132–138. <https://doi.org/10.1093/ajcn/83.1.132>
- Lusk, J. L. (2017). Economics and obesity policy. *International Journal of Obesity (2005)*, *41*(6), 831–834. <https://doi.org/10.1038/ijo.2017.5>
- Macht, M. (2008). How emotions affect eating: A five-way model. *Appetite*, *50*(1), 1–11.  
<https://doi.org/10.1016/j.appet.2007.07.002>
- Macht, M., & Simons, G. (2011). Emotional Eating. In I. Nyklíček, A. Vingerhoets, & M. Zeelenberg (Eds.), *Emotion Regulation and Well-Being* (1st ed., pp. 281–295). Springer Science+Business Media LLC. [https://doi.org/10.1007/978-1-4419-6953-8\\_17](https://doi.org/10.1007/978-1-4419-6953-8_17)
- Mitry, P., Wawro, N., Six-Merker, J., Zoller, D., Jourdan, C., Meisinger, C., Thierry, S., Nöthlings, U., Knüppel, S., Boeing, H., & Linseisen, J. (2019). Usual dietary intake estimation based on a combination of repeated 24-h food lists and a food frequency questionnaire in the KORA FF4 cross-sectional study. *Frontiers in Nutrition*, *6*, 145.  
<https://doi.org/10.3389/fnut.2019.00145>
- Naughton, P., McCarthy, S. N., & McCarthy, M. B. (2015). The creation of a healthy eating motivation score and its association with food choice and physical activity in a cross sectional sample of Irish adults. *The International Journal of Behavioral Nutrition and Physical Activity*, *12*, 74.  
<https://doi.org/10.1186/s12966-015-0234-0>
- Neighbors, C., Lee, C. M., Lewis, M. A., Fossos, N., & Larimer, M. E. (2007). Are social norms the best predictor of outcomes among heavy-drinking college students? *Journal of Studies on Alcohol and Drugs*, *68*(4), 556–565. <https://doi.org/10.15288/jsad.2007.68.556>
- Nöthlings, U., Hoffmann, K., Bergmann, M. M., & Boeing, H. (2007). Fitting portion sizes in a self-administered food frequency questionnaire. *The Journal of Nutrition*, *137*(12), 2781–2786.  
<https://doi.org/10.1093/jn/137.12.2781>
- Palaniappan, U., Cue, R. I., Payette, H., & Gray-Donald, K. (2003). Implications of day-to-day variability on measurements of usual food and nutrient intakes. *The Journal of Nutrition*, *133*(1), 232–235. <https://doi.org/10.1093/jn/133.1.232>
- Péneau, S., Ménard, E., Méjean, C., Bellisle, F., & Hercberg, S. (2013). Sex and dieting modify the association between emotional eating and weight status. *The American Journal of Clinical Nutrition*, *97*(6), 1307–1313. <https://doi.org/10.3945/ajcn.112.054916>


- Phan, U. T.X., & Chambers, E. (2016a). Application of an eating motivation survey to study eating occasions. *Journal of Sensory Studies*, 31(2), 114–123. <https://doi.org/10.1111/joss.12197>
- Phan, U. T.X., & Chambers, E. (2016b). Motivations for choosing various food groups based on individual foods. *Appetite*, 105, 204–211. <https://doi.org/10.1016/j.appet.2016.05.031>
- Phan, U. T.X., & Chambers, E. (2018). Motivations for meal and snack times: Three approaches reveal similar constructs. *Food Quality and Preference*, 68, 267–275. <https://doi.org/10.1016/j.foodqual.2018.03.018>
- Pollard, J., Kirk, S. F. L., & Cade, J. E. (2002). Factors affecting food choice in relation to fruit and vegetable intake: A review. *Nutrition Research Reviews*, 15(2), 373–387. <https://doi.org/10.1079/NRR200244>
- Pollard, T. M., Steptoe, A., & Wardle, J. (1998). Motives underlying healthy eating: Using the Food Choice Questionnaire to explain variation in dietary intake. *Journal of Biosocial Science*, 30(2), 165–179. <https://doi.org/10.1017/s0021932098001655>
- Pothos, E. M., Tapper, K., & Calitri, R. (2009). Cognitive and behavioral correlates of BMI among male and female undergraduate students. *Appetite*, 52(3), 797–800. <https://doi.org/10.1016/j.appet.2009.03.002>
- Price, M., Higgs, S., & Lee, M. (2015). Self-reported eating traits: Underlying components of food responsiveness and dietary restriction are positively related to BMI. *Appetite*, 95, 203–210. <https://doi.org/10.1016/j.appet.2015.07.006>
- Rempe, H. M., Sproesser, G., Gingrich, A., Skurk, T., Brandl, B., Hauner, H., Renner, B., Volkert, D., Sieber, C. C., Freiberger, E., & Kiesswetter, E. (2020). The relationship between healthy eating motivation and protein intake in community-dwelling older adults with varying functional status. *Nutrients*, 12(3). <https://doi.org/10.3390/nu12030662>
- Rempe, H. M., Sproesser, G., Gingrich, A., Spiegel, A., Skurk, T., Brandl, B., Hauner, H., Renner, B., Volkert, D., Sieber, C. C., Freiberger, E., & Kiesswetter, E. (2019). Measuring eating motives in older adults with and without functional impairments with The Eating Motivation Survey (TEMS). *Appetite*, 137, 1–20. <https://doi.org/10.1016/j.appet.2019.01.024>
- Renner, B., Sproesser, G., Strohbach, S., & Schupp, H. T. (2012). Why we eat what we eat. The Eating Motivation Survey (TEMS). *Appetite*, 59(1), 117–128. <https://doi.org/10.1016/j.appet.2012.04.004>
- Roininen, K., Lähteenmäki, L., & Tuorila, H. (1999). Quantification of consumer attitudes to health and hedonic characteristics of foods. *Appetite*, 33(1), 71–88. <https://doi.org/10.1006/appe.1999.0232>
- Ruddock, H. K., Brunstrom, J. M., & Higgs, S. (2021). The social facilitation of eating: Why does the mere presence of others cause an increase in energy intake? *Physiology & Behavior*, 240, 113539. <https://doi.org/10.1016/j.physbeh.2021.113539>
- Ruddock, H. K., Brunstrom, J. M., Vartanian, L. R., & Higgs, S. (2019). A systematic review and meta-analysis of the social facilitation of eating. *The American Journal of Clinical Nutrition*, 110(4), 842–861. <https://doi.org/10.1093/ajcn/nqz155>
- Schmeiser, M. D. (2009). Expanding wallets and waistlines: The impact of family income on the BMI of women and men eligible for the Earned Income Tax Credit. *Health Economics*, 18(11), 1277–1294. <https://doi.org/10.1002/hec.1430>
- Schroeter, C., & Lusk, J. L. (2008). Economic factors and body weight: An empirical analysis. *Journal of Agricultural and Applied Economics*, 40(2), 523–538. <https://doi.org/10.22004/AG.ECON.46991>
- Souza, A. M., Bezerra, I. W. L., Pereira, G. S., Torres, K. G., Costa, R. M., & Oliveira, A. G. (2020). Relationships between motivations for food choices and consumption of food groups: A

- prospective cross-sectional survey in manufacturing workers in Brazil. *Nutrients*, 12(5). <https://doi.org/10.3390/nu12051490>
- Sproesser, G., Imada, S., Furumitsu, I., Rozin, P., Ruby, M. B., Arbit, N., Fischler, C., Schupp, H. T., & Renner, B. (2018). What Constitutes Traditional and Modern Eating? The Case of Japan. *Nutrients*, 10(2). <https://doi.org/10.3390/nu10020118>
- Sproesser, G., Moraes, J. M. M., Renner, B., & Alvarenga, M. D. S. (2019). The Eating Motivation Survey in Brazil: Results From a Sample of the General Adult Population. *Frontiers in Psychology*, 10, 2334. <https://doi.org/10.3389/fpsyg.2019.02334>
- Sproesser, G., Ruby, M. B., Arbit, N., Akotia, C. S., Alvarenga, M. D. S., Bhangaokar, R., Furumitsu, I., Hu, X., Imada, S., Kaptan, G., Kaufer-Horwitz, M., Menon, U., Fischler, C., Rozin, P., Schupp, H. T., & Renner, B. (2019). Understanding traditional and modern eating: The TEP10 framework. *BMC Public Health*, 19(1), 1606. <https://doi.org/10.1186/s12889-019-7844-4>
- Sproesser, G., Ruby, M. B., Arbit, N., Rozin, P., Schupp, H. T., & Renner, B. (2018). The Eating Motivation Survey: Results from the USA, India and Germany. *Public Health Nutrition*, 21(3), 515–525. <https://doi.org/10.1017/S1368980017002798>
- Step toe, A., Pollard, T. M., & Wardle, J. (1995). Development of a measure of the motives underlying the selection of food: the food choice questionnaire. *Appetite*, 25(3), 267–284.
- Sung, J., Lee, K., & Song, Y.-M. (2009). Relationship of eating behavior to long-term weight change and body mass index: The Healthy Twin study. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 14(2-3), e98-105. <https://doi.org/10.1007/BF03327806>
- Swinburn, B., & Egger, G. (2002). Preventive strategies against weight gain and obesity. *Obesity Reviews : An Official Journal of the International Association for the Study of Obesity*, 3(4), 289–301. <https://doi.org/10.1046/j.1467-789x.2002.00082.x>
- Tylka, T. L. (2006). Development and psychometric evaluation of a measure of intuitive eating. *Journal of Counseling Psychology*, 53(2), 226–240. <https://doi.org/10.1037/0022-0167.53.2.226>
- van Dyke, N., & Drinkwater, E. J. (2014). Relationships between intuitive eating and health indicators: Literature review. *Public Health Nutrition*, 17(8), 1757–1766. <https://doi.org/10.1017/S1368980013002139>
- van Strien, T., Herman, C. P., & Verheijden, M. W. (2009). Eating style, overeating, and overweight in a representative Dutch sample. Does external eating play a role? *Appetite*, 52(2), 380–387. <https://doi.org/10.1016/j.appet.2008.11.010>
- Wahl, D. R., Villinger, K., Blumenschein, M., König, L. M., Ziesemer, K., Sproesser, G., Schupp, H. T., & Renner, B. (2020). Why We Eat What We Eat: Assessing Dispositional and In-the-Moment Eating Motives by Using Ecological Momentary Assessment. *JMIR MHealth and UHealth*, 8(1), e13191. <https://doi.org/10.2196/13191>
- World Health Organization. (2020a, April 1). *Fact Sheet - Obesity and overweight*. World Health Organization. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
- World Health Organization. (2020b, June 8). *Fact Sheet - Diabetes*. World Health Organization. <https://www.who.int/news-room/fact-sheets/detail/diabetes>
- Zellner, A. (1962). An efficient method of estimating Seemingly Unrelated Regressions and tests for aggregation bias. *Journal of the American Statistical Association*, 57(298), 348–368. <https://doi.org/10.1080/01621459.1962.10480664>

## Appendix

**Figure A1**

Mean values of the motivation factors for the full sample (n = 429; blue) versus the study by Renner et al. (2012; n = 1,040; blue) and for male (n = 211; orange) and female (n = 218; grey) subsamples.



**Table A1**

Correlations and internal consistencies of 15 TEMS factors (n = 429).

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1 <i>Liking</i>															
2 <i>Habits</i>	.19 *														
3 <i>Need &amp; Hunger</i>	.29 *	.59 *													
4 <i>Health</i>	.30 *	.30 *	.56 *												
5 <i>Convenience</i>	.12	.31 *	.36 *	.23 *											
6 <i>Pleasure</i>	.27 *	.33 *	.45 *	.40 *	.28 *										
7 <i>Traditional Eating</i>	.06	.42 *	.46 *	.33 *	.25 *	.59 *									
8 <i>Natural Concerns</i>	.15 *	.06	.24 *	.58 *	.02	.32 *	.28 *								
9 <i>Sociability</i>	.16 *	.13	.25 *	.29 *	.20 *	.48 *	.58 *	.33 *							
10 <i>Price</i>	.04	.36 *	.30 *	.08	.33 *	.27 *	.35 *	-.18 *	.21 *						
11 <i>Visual Appeal</i>	.05	.25 *	.21 *	-.01	.19 *	.39 *	.53 *	.01	.39 *	.40 *					
12 <i>Weight Control</i>	.04	.16 *	.14	.31 *	.25 *	.21 *	.22 *	.20 *	.20 *	.22 *	.22 *				
13 <i>Affect Regulation</i>	-.10	.12	.02	-.08	.15 *	.30 *	.35 *	.04	.17 *	.21 *	.45 *	.20 *			
14 <i>Social Norms</i>	.06	.27 *	.24 *	.10	.21 *	.35 *	.56 *	.13	.41 *	.27 *	.51 *	.26 *	.40 *		
15 <i>Social Image</i>	-.07	.19 *	.12	.03	.18 *	.25 *	.39 *	.09	.23 *	.28 *	.43 *	.15 *	.55 *	.54 *	
<b>Cronbach's Alpha</b>	<b>.87</b>	<b>.84</b>	<b>.62</b>	<b>.86</b>	<b>.92</b>	<b>.77</b>	<b>.78</b>	<b>.87</b>	<b>.89</b>	<b>.84</b>	<b>.74</b>	<b>.84</b>	<b>.91</b>	<b>.89</b>	<b>.83</b>

Note: \*  $p < 0.01$ .

**Table A2**

Regression results for anthropometric measures – no separation of coefficients by sex

	BMI (kg/m <sup>2</sup> )		FFM (%)		Waist (cm)			
	B	SE	B	SE	B	SE		
<i>Liking</i>	0.04	0.19	0.25	0.30	0.47	0.56		
<i>Habits</i>	-0.04	0.16	-0.19	0.25	-0.60	0.48		
<i>Need &amp; Hunger</i>	-0.22	0.20	<b>0.73</b>	0.30	* -1.12	0.57		
<i>Health</i>	<b>-0.51</b>	0.20	*	<b>1.30</b>	0.32	*** -2.26	0.59	***
<i>Convenience</i>	0.25	0.14	-0.28	0.22	0.63	0.41		
<i>Pleasure</i>	-0.12	0.18	-0.26	0.29	0.13	0.54		
<i>Traditional Eating</i>	<b>0.53</b>	0.19	*	-0.62	0.29	1.24	0.54	
<i>Natural Concerns</i>	0.21	0.16	-0.47	0.25	0.27	0.46		
<i>Sociability</i>	-0.17	0.15	0.39	0.24	-0.42	0.44		
<i>Price</i>	-0.02	0.17	0.06	0.26	0.48	0.48		
<i>Visual Appeal</i>	0.13	0.20	-0.30	0.31	0.75	0.59		
<i>Weight Control</i>	0.14	0.13	-0.14	0.21	0.40	0.38		
<i>Affect Regulation</i>	0.25	0.19	<b>-0.74</b>	0.29	*	1.32	0.55	
<i>Social Norms</i>	-0.06	0.16	0.20	0.25	-0.46	0.47		
<i>Social Image</i>	-0.39	0.31	0.86	0.48	-1.10	0.90		
<i>Male</i>	1.53	0.41	10.13	0.63	11.81	1.18		
<i>Age group 2 (middle agers)<sup>a</sup></i>	5.01	0.62	-10.33	0.96	15.24	1.80		
<i>Age group 3 (older adults, FS)<sup>a</sup></i>	3.71	0.85	-12.75	1.32	17.39	2.48		
<i>Age group 4 (older adults, N)<sup>a</sup></i>	4.56	0.68	-15.89	1.05	18.89	1.97		
<i>Constant</i>	20.98	1.60	69.03	2.48	77.14	4.65		
<i>R<sup>2</sup></i>	0.31		0.67		0.48			
<i>N</i>	429		429		429			

Note: \*\*\*, \*\*, and \* indicate significance at the 0.01-, 0.05-, and 0.1-level based on adjusted p-values (Benjamini & Hochberg, 1995); Standard errors in parentheses. BMI, Body Mass Index; FFM, fat free mass; <sup>a</sup>Reference: *Young adults*.



**Table A3**

Regression results for energy and nutrient intakes – no separation of coefficients by sex

	Energy (kcal/day)		Carbohydrates (g/day)		Fat (g/day)		Protein (g/day)		Fibre (g/day)		Alcohol (g/day)		
	B	SE	B	SE	B	SE	B	SE	B	SE	B	SE	
<i>Liking</i>	-7.63	15.37	-1.45	2.05	-0.66	0.70	0.12	0.66	-0.27	0.28	0.51	0.34	
<i>Habits</i>	5.32	12.96	1.92	1.73	0.00	0.59	-0.18	0.55	0.11	0.24	-0.24	0.29	
<i>Need &amp; Hunger</i>	22.40	15.52	2.47	2.07	0.95	0.71	0.64	0.66	0.28	0.28	0.18	0.35	
<i>Health</i>	24.86	16.22	4.41	2.16	0.83	0.74	1.39	0.69	1.63	0.29	***	-0.92	0.36
<i>Convenience</i>	-22.41	11.22	-2.72	1.50	-0.68	0.51	-0.54	0.48	-0.29	0.20	-0.45	0.25	
<i>Pleasure</i>	-9.75	14.63	-2.06	1.95	-0.35	0.67	-0.42	0.62	-0.41	0.27	0.50	0.33	
<i>Traditional Eating</i>	6.71	14.69	0.35	1.96	0.36	0.67	-0.13	0.63	-0.47	0.27	0.41	0.33	
<i>Natural Concerns</i>	13.28	12.58	1.50	1.68	0.57	0.57	0.15	0.54	0.46	0.23	0.23	0.28	
<i>Sociability</i>	-5.14	12.01	-1.98	1.60	0.10	0.55	-0.19	0.51	-0.07	0.22	0.40	0.27	
<i>Price</i>	31.82	13.16	3.28	1.76	1.31	0.60	0.53	0.56	-0.23	0.24	0.72	0.29	
<i>Visual Appeal</i>	25.29	16.08	2.46	2.15	0.99	0.73	1.25	0.69	0.26	0.29	0.18	0.36	
<i>Weight Control</i>	-22.37	10.48	-3.17	1.40	-1.28	0.48	0.47	0.45	-0.24	0.19	-0.05	0.23	
<i>Affect Regulation</i>	-18.67	15.03	-1.36	2.01	-0.89	0.69	-0.65	0.64	-0.16	0.27	-0.36	0.34	
<i>Social Norms</i>	-6.40	12.91	-0.12	1.72	-0.10	0.59	-0.27	0.55	-0.11	0.23	-0.55	0.29	
<i>Social Image</i>	22.86	24.51	2.33	3.27	0.84	1.12	1.07	1.05	0.77	0.45	0.20	0.55	
<i>Male</i>	441.02	32.26	40.62	4.30	17.37	1.47	16.05	1.38	1.47	0.59	8.37	0.72	
<i>Age group 2 (middle agers)<sup>a</sup></i>	14.85	48.96	-8.35	6.53	4.38	2.24	-0.29	2.09	-2.61	0.89	1.75	1.09	
<i>Age group 3 (older adults, FS)<sup>a</sup></i>	186.12	67.60	15.80	9.02	10.66	3.09	1.60	2.88	1.73	1.23	2.83	1.51	
<i>Age group 4 (older adults, N)<sup>a</sup></i>	84.40	53.83	8.69	7.18	5.43	2.46	-2.09	2.30	0.10	0.98	1.42	1.20	
<i>Constant</i>	1340.92	126.74	151.92	16.91	58.16	5.79	50.14	5.41	14.42	2.30	-0.63	2.83	
<i>R<sup>2</sup></i>	0.42		0.30		0.36		0.35		0.26		0.34		
<i>N</i>	429		429		429		429		429		429		

Note: \*\*\*, \*\*, and \* indicate significance at the 0.01-, 0.05-, and 0.1-level based on adjusted p-values (Benjamini & Hochberg, 1995); Standard errors in parentheses. BMI, Body Mass Index; FFM, fat free mass; <sup>a</sup>Reference: *Young adults*.