

OPEN

Neurology[®]

The most widely read and highly cited peer-reviewed neurology journal
The Official Journal of the American Academy of Neurology



Neurology Publish Ahead of Print
DOI: 10.1212/WNL.0000000000200699

Genome-wide Association and Meta-analysis of Age-at-Onset in Parkinson Disease: Evidence From
COURAGE-PD Consortium

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Neurology[®] Published Ahead of Print articles have been peer reviewed and accepted for publication. This manuscript will be published in its final form after copyediting, page composition, and review of proofs. Errors that could affect the content may be corrected during these processes.

Author(s):

Sandeep Grover, PhD¹; Ashwin Ashok Kumar Sreelatha, MTech¹; Lasse Pihlstrom, MD²; Cloé Domenighetti, MSc³; Claudia Schulte, MSc^{4,5}; Pierre-Emmanuel Sugier, PhD³; Milena Radivojkov-Blagojevic, MSc⁶; Peter Lichtner, PhD⁶; Océane Mohamed, MSc³; Berta Portugal, PhD³; Zied Landoulsi, PhD⁷; Patrick May, PhD⁷; Dheeraj BOBBILI, PhD⁷; Connor Edsall, PhD⁸; Felix Bartusch, MSc^{9,10}; Maximilian Hanussek, MSc^{9,10}; Jens Krüger, PhD⁹; Dena G Hernandez, PhD⁸; Cornelis Blauwendraat, PhD⁸; George D Mellick, PhD¹¹; Alexander Zimprich, MD¹²; Walter Pirker, MD¹³; Manuela Tan, MSc¹⁴; Ekaterina Rogaeva, PhD¹⁵; Anthony Lang, MD^{15,16,17,18}; Sulev Koks, MD, PHD^{19,20}; Pille Taba, MD, PHD^{21,22}; Suzanne Lesage, PHD²³; Alexis Brice, PhD²³; Jean-Christophe Corvol, MD, PHD^{23,24}; Marie-Christine Chartier-Harlin, PHD²⁵; Eugénie Mutez, MD, PhD²⁵; Kathrin Brockmann, MD^{4,5}; Angela B Deuschländer, MD^{26,27,28}; Georges M Hadjigeorgiou, MD^{29,30}; Efthimos Dardiotis, MD³⁰; Leonidas Stefanis, MD, PHD^{31,32}; Athina Maria Simitsi, MD, PHD³²; Enza Maria Valente, MD, PHD^{33,34}; Simona Petrucci, MD, PHD^{35,36}; Letizia Straniero, PHD^{37,38}; Anna Zecchinelli, MD³⁹; Gianni Pezzoli, MD³⁹; Laura Brighina, MD, PHD^{40,41}; Carlo Ferrarese, MD, PHD^{40,41}; Grazia Annesi, PHD⁴²; Andrea Quattrone, MD⁴³; Monica Gagliardi, PHD⁴²; Lena F Burbulla, PHD^{5,44,45,46}; Hirotaka Matsuo, MD, PHD⁴⁷; Yusuke Kawamura, MD⁴⁷; Nobutaka Hattori, MD, PHD⁴⁸; Kenya Nishioka, MD, PHD⁴⁸; Sun Ju Chung, MD, PHD⁴⁹; Yun Joong Kim, MD, PHD⁵⁰; Lukas Pavelka, MD⁵¹; Bart PC van de Warrenburg, MD, PHD⁵²; Bastiaan R Bloem, MD, PHD⁵²; Andrew B. Singleton, PHD⁸; Jan Aasly, MD⁵³; Mathias Toft, MD, PHD⁵⁴; Leonor Correia Guedes, MD, PHD^{55,56}; Joaquim J Ferreira, MD, PHD^{55,57}; Soraya Bardien, PHD⁵⁸; Jonathan Carr, PHD⁵⁹; Eduardo Tolosa, MD, PHD^{60,61}; Mario Ezquerro, PHD⁶²; Pau Pastor, MD, PHD^{63,64}; Monica Diez-Fairen, MSc^{63,64}; Karin Wirdefeldt, MD, PHD^{65,66}; Nancy L Pedersen, PHD⁶⁶; Caroline Ran, PHD⁶⁷; Andrea C Belin, PHD⁶⁷; Andreas Puschmann, MD, PHD⁶⁸; Clara Hellberg, MD⁶⁸; Carl E Clarke, MD⁶⁹; Karen E Morrison, MD⁷⁰; Dimitri Krainc, MD, PHD⁴⁴; Matt J Farrer, PHD⁷¹; Rejko Kruger, MD^{7,72,73,74}; Alexis Elbaz, PhD³; Thomas Gasser, MD^{4,5}; Manu Sharma, PHD^{1,4} on behalf of and the Comprehensive Unbiased Risk Factor Assessment for Genetics and Environment in Parkinson's Disease (COURAGE-PD) consortium

Corresponding Author:

Manu Sharma, manu.sharma@uni-tuebingen.de

Affiliation Information for All Authors: 1. Centre for Genetic Epidemiology, Institute for Clinical Epidemiology and Applied Biometry, University of Tübingen, Germany. 2. Department of Neurology, Oslo University Hospital, Oslo, Norway. 3. Université Paris-Saclay, UVSQ, Univ. Paris-Sud, Inserm, Team "Exposome, heredity, cancer and health", CESP, 94807, Villejuif, France. 4. Department for Neurodegenerative Diseases, Hertie Institute for Clinical Brain Research, University of Tübingen, Germany. 5. German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany. 6. Institute of Human Genetics, Helmholtz Zentrum München, Neuherberg, Germany. 7. Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Esch-sur-Alzette, Luxembourg. 8. Molecular Genetics Section, Laboratory of Neurogenetics, NIA, NIH, Bethesda, MD 20892, USA. 9. Group of Applied Bioinformatics, University of Tübingen, Tübingen, Germany. 10. High Performance and Cloud Computing Group ZDV, University of Tübingen, Tübingen, Germany. 11. Griffith Institute for Drug Discovery, Griffith University, Don Young Road, Nathan, Queensland, Australia. 12. Department of Neurology, Medical University of Vienna, Austria. 13. Department of Neurology, Wilhelminenspital, Austria. 14. Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology, University College London, London, UK. 15. Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Toronto, Ontario, Canada. 16. Edmond J. Safra Program in Parkinson's Disease, Morton and Gloria Shulman Movement Disorders Clinic, Toronto Western Hospital, UHN, Toronto, Ontario, Canada. 17. Division of Neurology, University of Toronto, Toronto, Ontario, Canada. 18. Krembil Brain Institute, Toronto, Ontario, Canada. 19. Centre for Molecular Medicine and Innovative Therapeutics, Murdoch University, Murdoch, Australia. 20. Perron Institute for Neurological and Translational Science, Nedlands, Western Australia, Australia. 21. Department of Neurology and Neurosurgery, University of Tartu, Estonia. 22. Neurology Clinic, Tartu University Hospital, Tartu, Estonia. 23. Sorbonne Université (SU) Unité Mixte de Recherche (UMR) 1127, INSERM U1127, CNRS UMR 7225, Institut du Cerveau et de la Moelle épinière, ICM, Paris, France. 24. Assistance Publique Hôpitaux de Paris, Department of Neurology, CIC Neurosciences, Paris, France. 25. Univ. Lille, Inserm, CHU Lille, UMR-S 1172 - JPArc - Centre de Recherche Lille Neurosciences & Cognition, F-59000 Lille, France. 26. Department of Neurology, Ludwig Maximilians University of Munich, Germany. 27. Department of Neurology, Max Planck Institute of Psychiatry, Munich, Germany. 28. Department of Neurology and Department of Clinical Genomics, Mayo Clinic Florida, Jacksonville, FL, USA. 29. Department of Neurology, Medical School, University of Cyprus, Nicosia, Cyprus. 30. Department of Neurology, Laboratory of Neurogenetics, University of Thessaly, University Hospital of Larissa, Larissa, Greece. 31. Center of Clinical Research, Experimental Surgery and Translational Research, Biomedical Research Foundation of the Academy of Athens, Athens, Greece. 32. 1st Department of Neurology, Eginition Hospital, Medical School, National and Kapodistrian University of Athens, Athens, Greece. 33. Department of Molecular Medicine, University of Pavia, Pavia, Italy. 34. Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) Mondino Foundation, Pavia,

Italy. 35. UOC Medical Genetics and Advanced Cell Diagnostics, S. Andrea University Hospital, Rome, Italy. 36. Department of Clinical and Molecular Medicine, University of Rome, Rome, Italy. 37. Department of Biomedical Sciences - Humanitas University, Milan, Italy. 38. Humanitas Clinical and Research Center, IRCCS, Via Manzoni 56, 20089 Rozzano, Milan, Italy. 39. Parkinson Institute, Azienda Socio Sanitaria Territoriale (ASST) Gaetano Pini/CTO, Milano, Italia. 40. Department of Neurology, San Gerardo Hospital, Milan, Italy. 41. Center for Neuroscience, University of Milano Bicocca, Monza, Italy. 42. Institute for Biomedical Research and Innovation, National Research Council, Mangone, Cosenza, Italy. 43. Institute of Neurology, Department of Medical and Surgical Sciences, Magna Graecia University, Catanzaro, Italy. 44. Department of Neurology, Northwestern University Feinberg School of Medicine, Chicago, Illinois 60611, United States. 45. Metabolic Biochemistry, Biomedical Center (BMC), Faculty of Medicine, Ludwig-Maximilians University, Munich, Germany. 46. Munich Cluster for Systems Neurology (SyNergy), Munich, Germany. 47. Department of Integrative Physiology and Bio-Nano Medicine, National Defense Medical College, Saitama 359-8513, Japan. 48. Department of Neurology, Juntendo University School of Medicine, Bunkyo-ku, Tokyo 113-8421, Japan. 49. Department of Neurology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, South Korea. 50. Department of Neurology, Yonsei University College of Medicine, Seoul, South Korea. 51. Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Belval, Luxembourg. 52. Radboud University Medical Centre, Donders Institute for Brain, Cognition and Behaviour, Department of Neurology, Nijmegen, The Netherlands. 53. Department of Neurology, St Olav's Hospital and Norwegian University of Science and Technology, Trondheim, Norway. 54. Department of Neurology, Oslo University Hospital, Oslo, Norway. 55. Instituto de Medicina Molecular João Lobo Antunes, Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal. 56. Department of Neurosciences and Mental Health, Neurology, Hospital de Santa Maria, Centro Hospitalar Universitário Lisboa Norte (CHULN), Lisbon, Portugal. 57. Laboratory of Clinical Pharmacology and Therapeutics, Faculdade de Medicina, Universidade de Lisboa, Lisbon, Portugal. 58. Division of Molecular Biology and Human Genetics, Department of Biomedical Sciences, Faculty of Medicine and Health Sciences, Stellenbosch University, South Africa. 59. Division of Neurology, Department of Medicine, Faculty of Medicine and Health Sciences, Stellenbosch University, South Africa. 60. Parkinson's disease & Movement Disorders Unit, Neurology Service, Hospital Clínic de Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), University of Barcelona, Barcelona, Spain. 61. Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas (CIBERNED: CB06/05/0018-ISCI) Barcelona, Spain. 62. Lab of Parkinson Disease and Other Neurodegenerative Movement Disorders, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Institut de Neurociències, Universitat de Barcelona, ES-08036 Barcelona, Catalonia. 63. Fundació per la Recerca Biomèdica i Social Mútua Terrassa, Terrassa, Barcelona, Spain. 64. Movement Disorders Unit, Department of Neurology, Hospital Universitari Mutua de Terrassa, Terrassa, Barcelona, Spain. 65. Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden. 66. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden. 67. Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden. 68. Lund University, Skåne University Hospital, Department of Clinical Sciences Lund, Neurology, Getingevägen 4, 221 85, Lund, Sweden. 69. University of Birmingham and Sandwell and West Birmingham Hospitals NHS Trust, United Kingdom. 70. Faculty of Medicine, Health and Life Sciences, Queens University, Belfast, United Kingdom. 71. Department of Neurology, McKnight Brain Institute, University of Florida, Gainesville, FL, USA. 72. Parkinson Research Clinic, Centre Hospitalier de Luxembourg, Luxembourg, Luxembourg. 73. Transversal Translational Medicine, Luxembourg Institute of Health (LIH), Strassen, Luxembourg. 74. Neurology, Centre Hospitalier de Luxembourg, Luxembourg, Luxembourg

Equal Author Contribution:

Contributions:

Sandeep Grover: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Study concept or design; Analysis or interpretation of data

Ashwin Ashok Kumar Sreelatha: Major role in the acquisition of data; Analysis or interpretation of data

Lasse Pihlstrom: Drafting/revision of the manuscript for content, including medical writing for content; Analysis or interpretation of data

Cloé Domenighetti: Drafting/revision of the manuscript for content, including medical writing for content

Claudia Schulte: Drafting/revision of the manuscript for content, including medical writing for content

Pierre-Emmanuel Sugier: Drafting/revision of the manuscript for content, including medical writing for content

Milena Radivojkov-Blagojevic: Drafting/revision of the manuscript for content, including medical writing for content

Peter Lichtner: Drafting/revision of the manuscript for content, including medical writing for content

Océane Mohamed: Drafting/revision of the manuscript for content, including medical writing for content

Berta Portugal: Drafting/revision of the manuscript for content, including medical writing for content

Zied Landoulsi: Drafting/revision of the manuscript for content, including medical writing for content
Patrick May: Drafting/revision of the manuscript for content, including medical writing for content
Dheeraj BOBBILI: Drafting/revision of the manuscript for content, including medical writing for content
Connor Edsall: Drafting/revision of the manuscript for content, including medical writing for content
Felix Bartusch: Drafting/revision of the manuscript for content, including medical writing for content
Maximilian Hanussek: Drafting/revision of the manuscript for content, including medical writing for content
Jens Krüger: Drafting/revision of the manuscript for content, including medical writing for content
Dena G Hernandez: Drafting/revision of the manuscript for content, including medical writing for content
Cornelis Blauwendraat: Drafting/revision of the manuscript for content, including medical writing for content
George D Mellick: Drafting/revision of the manuscript for content, including medical writing for content
Alexander Zimprich: Drafting/revision of the manuscript for content, including medical writing for content
Walter Pirker: Drafting/revision of the manuscript for content, including medical writing for content
Manuela Tan: Drafting/revision of the manuscript for content, including medical writing for content
Ekaterina Rogaeva: Drafting/revision of the manuscript for content, including medical writing for content
Anthony Lang: Drafting/revision of the manuscript for content, including medical writing for content
Sulev Koks: Drafting/revision of the manuscript for content, including medical writing for content
Pille Taba: Drafting/revision of the manuscript for content, including medical writing for content
Suzanne Lesage: Drafting/revision of the manuscript for content, including medical writing for content
Alexis Brice: Drafting/revision of the manuscript for content, including medical writing for content
Jean-Christophe Corvol: Drafting/revision of the manuscript for content, including medical writing for content
Marie-Christine Chartier-Harlin: Drafting/revision of the manuscript for content, including medical writing for content
Eugenie Mutez: Drafting/revision of the manuscript for content, including medical writing for content
Kathrin Brockmann: Drafting/revision of the manuscript for content, including medical writing for content
Angela B Deuschländer: Drafting/revision of the manuscript for content, including medical writing for content
Georges M Hadjigeorgiou: Drafting/revision of the manuscript for content, including medical writing for content
Efthimos Dardiotis: Drafting/revision of the manuscript for content, including medical writing for content
Leonidas Stefanis: Drafting/revision of the manuscript for content, including medical writing for content
Athina Maria Simitsi: Drafting/revision of the manuscript for content, including medical writing for content
Enza Maria Valente: Drafting/revision of the manuscript for content, including medical writing for content
Simona Petrucci: Drafting/revision of the manuscript for content, including medical writing for content
Letizia Straniero: Drafting/revision of the manuscript for content, including medical writing for content
Anna Zecchinelli: Drafting/revision of the manuscript for content, including medical writing for content
Gianni Pezzoli: Drafting/revision of the manuscript for content, including medical writing for content
Laura Brighina: Drafting/revision of the manuscript for content, including medical writing for content
Carlo Ferrarese: Drafting/revision of the manuscript for content, including medical writing for content
Grazia Annesi: Drafting/revision of the manuscript for content, including medical writing for content
Andrea Quattrone: Drafting/revision of the manuscript for content, including medical writing for content
Monica Gagliardi: Drafting/revision of the manuscript for content, including medical writing for content
Lena F Burbulla: Drafting/revision of the manuscript for content, including medical writing for content
Hirotaka Matsuo: Drafting/revision of the manuscript for content, including medical writing for content
Yusuke Kawamura: Drafting/revision of the manuscript for content, including medical writing for content
Nobutaka Hattori: Drafting/revision of the manuscript for content, including medical writing for content
Kenya Nishioka: Drafting/revision of the manuscript for content, including medical writing for content
Sun Ju Chung: Drafting/revision of the manuscript for content, including medical writing for content
Yun Joong Kim: Drafting/revision of the manuscript for content, including medical writing for content
Lukas Pavelka: Drafting/revision of the manuscript for content, including medical writing for content
Bart PC van de Warrenburg: Drafting/revision of the manuscript for content, including medical writing for content
Bastiaan R Bloem: Drafting/revision of the manuscript for content, including medical writing for content
Andrew B. Singleton: Drafting/revision of the manuscript for content, including medical writing for content
Jan Aasly: Drafting/revision of the manuscript for content, including medical writing for content
Mathias Toft: Drafting/revision of the manuscript for content, including medical writing for content
Leonor Correia Guedes: Drafting/revision of the manuscript for content, including medical writing for content
Joaquim J Ferreira: Drafting/revision of the manuscript for content, including medical writing for content
Soraya Bardiën: Drafting/revision of the manuscript for content, including medical writing for content
Jonathan Carr: Drafting/revision of the manuscript for content, including medical writing for content
Eduardo Tolosa: Drafting/revision of the manuscript for content, including medical writing for content
Mario Ezquerra: Drafting/revision of the manuscript for content, including medical writing for content
Pau Pastor: Drafting/revision of the manuscript for content, including medical writing for content
Monica Diez-Fairen: Drafting/revision of the manuscript for content, including medical writing for content

Karin Wirdefeldt: Drafting/revision of the manuscript for content, including medical writing for content
Nancy L Pedersen: Drafting/revision of the manuscript for content, including medical writing for content
Caroline Ran: Drafting/revision of the manuscript for content, including medical writing for content
Andrea C Belin: Drafting/revision of the manuscript for content, including medical writing for content
Andreas Puschmann: Drafting/revision of the manuscript for content, including medical writing for content
Clara Hellberg: Drafting/revision of the manuscript for content, including medical writing for content
Carl E Clarke: Drafting/revision of the manuscript for content, including medical writing for content
Karen E Morrison: Drafting/revision of the manuscript for content, including medical writing for content
Dimitri Krainc: Drafting/revision of the manuscript for content, including medical writing for content
Matt J Farrer: Drafting/revision of the manuscript for content, including medical writing for content
Rejko Kruger: Drafting/revision of the manuscript for content, including medical writing for content
Alexis Elbaz: Drafting/revision of the manuscript for content, including medical writing for content
Thomas Gasser: Drafting/revision of the manuscript for content, including medical writing for content
Manu Sharma: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Study concept or design; Analysis or interpretation of data

Figure Count:

2

Table Count:

1

Search Terms:

[53] Case control studies, [95] Association studies in genetics, [337] Burden of disease, [313] Basal ganglia, [165] Parkinson's disease/Parkinsonism

Acknowledgment:

Study Funding:

The COURAGE-PD consortium is conducted under a partnership agreement between 35 studies. The COURAGE-PD consortium is supported by the EU Joint Program for Neurodegenerative Disease research (JPND); <https://www.neurodegenerationresearch.eu/initiatives/annual-calls-for-proposals/closed-calls/risk-factors-2012/risk-factor-call-results/courage-pd/>, Grant ID: 01ED1406).

Disclosures:

M. Sharma is supported by the grants from the German Research Council (DFG/SH 599/6-1), MSA Coalition, and Michael J Fox Foundation. This work was also supported by grants from the Luxembourg National Research Fund (FNR) within the PEARL programme (FNR/P13/6682797 to R. Krüger) and the National Centre for Excellence in Research in Parkinson's disease (NCER-PD PD to R. Krüger, P. May, Z. Landoulsi). Z. Landoulsi and P. May were supported by the DFG Research Unit FOR2715 (INTER/DFG/17/11583046) and FOR2488 (INTER/DFG/19/14429377). C. Domenighetti is the recipient of a doctoral grant from Université Paris-Saclay, France. T. Gasser reports personal fees from UCB Pharma, personal fees from Novartis, personal fees from Teva, personal fees from MedUpdate, grants from The Michael J Fox Foundation for Parkinson's Research, grants from Bundesministerium für Bildung und Forschung (BMBF), grants from Deutsche Forschungsgemeinschaft (DFG), other from "Joint Programming for Neurodegenerative Diseases"(JPND) program, funded by the European

Commission, outside the submitted work; In addition, T. Gasser has a patent Patent Number: EP1802749 (A2) KASPP (LRRK2) gene, its production and use for the detection and treatment of neurodegenerative disorders issued. D. Krainc is the Founder of Lysosomal Therapeutics Inc. and Vanqua Bio, and serves on the Scientific Advisory Boards of The Silverstein Foundation, Vanqua Bio, Intellia Therapeutics, Prevail Therapeutics and is a Venture Partner at OrbiMed. BvdW received research support from Radboud university medical center, ZonMW, Gossweiler Foundation, and Hersenstichting, and has served on a scientific advisory board of uniQure. The remaining authors report no disclosures relevant to the manuscript.

Preprint DOI:

Received Date:

2021-11-17

Accepted Date:

2022-03-23

Handling Editor Statement:

Submitted and externally peer reviewed. The handling editor was Peter Hedera, MD, PhD.

Abstract

Background and Objectives

Considerable heterogeneity exists in the literature concerning genetic determinants of the age of onset (AAO) of Parkinson's disease (PD), which could be attributed to lack of well-powered replication cohorts. The previous largest GWAS identified *SNCA* and *TMEM175* loci on chromosome (Chr) 4 with a significant influence on AAO of PD, these have not been independently replicated. The present study aims to conduct a meta-analysis of GWAS of PD AAO and validate previously observed findings in worldwide populations.

Methods

A meta-analysis was performed on PD AAO GWAS of 30 populations of predominantly European ancestry from the Comprehensive Unbiased Risk Factor Assessment for Genetics and Environment in Parkinson's Disease (COURAGE-PD) consortium. This was followed up by combining our study with the largest publicly available European ancestry dataset compiled by the International Parkinson disease Genomics Consortium (IPDGC).

Results

The COURAGE-PD included a cohort of 8,535 patients with PD (91.9%: Europeans, 9.1%: East-Asians). The average AAO in the COURAGE-PD dataset was 58.9 years (SD=11.6), with an under-representation of females (40.2%). The heritability estimate for AAO in COURAGE-PD was 0.083 (SE=0.057). None of the loci reached genome-wide significance ($P < 5 \times 10^{-8}$). Nevertheless, the COURAGE-PD dataset confirmed the role of the previously published *TMEM175* variant as genetic determinant of AAO of PD with Bonferroni-corrected nominal levels of significance ($P < 0.025$): (rs34311866; $\beta(\text{SE})_{\text{COURAGE}} = 0.477(0.203)$, $P_{\text{COURAGE}} = 0.0185$). The subsequent meta-analysis of COURAGE-PD and IPDGC datasets ($N_{\text{total}} = 25,950$) led to the identification of two genome-wide significant association signals on Chr 4, including the previously reported *SNCA* locus (rs983361; $\beta(\text{SE})_{\text{COURAGE+IPDGC}} = 0.720(0.122)$, $P_{\text{COURAGE+IPDGC}} = 3.13 \times 10^{-9}$) and a novel *BST1* locus (rs4698412; $\beta(\text{SE})_{\text{COURAGE+IPDGC}} = -0.526(0.096)$, $P_{\text{COURAGE+IPDGC}} = 4.41 \times 10^{-8}$).

Discussion

Our study further refines the genetic architecture of Chr 4 underlying the AAO of the PD phenotype through the identification of *BST1* as a novel AAO PD locus. These findings open a new direction for the development of treatments to delay the onset of PD.

Keywords: Burden of disease, Age at onset, Duration of disease, Parkinson's disease, Genetic heritability

Introduction

In 2019, over 8.51 million individuals (95% uncertainty interval [UI] 7.3–9.8) had PD globally¹. This disease is one of the fastest-growing neurodegenerative diseases globally, with an estimated 30.9% increase in the number of patients with PD in 2019 compared to 2010. However, the prevalence of a disease depends on both the incidence and duration of disease, making an earlier age at onset of PD an essential contributor to the overall burden of

the disease. While less than 5% of patients with PD harbor pathogenic mutations in known monogenic PD genes, the majority are sporadic with predominantly late age of onset (AAO)^{2,3}. A better understanding of genetic factors influencing variability in AAO in sporadic patients could lead to a better understanding of PD pathophysiology.

The emergence of genome-wide association studies (GWAS) has resulted in a rapidly expanding list of loci harboring disease-susceptibility variants for the sporadic form of the disorder⁴⁻⁶. To date, genetic variants at 78 loci have been identified for sporadic PD⁶. Despite advances in understanding the genetic basis of PD incidence, the heritability underlying PD AAO remains largely unexplained. A recent global effort involving 28,568 sporadic PD patients of European ancestry led to the identification of two loci, *SNCA* and *TMEM175*, as risk factors for an earlier AAO, both of which are also known to play a role in α -synuclein-linked mechanisms underlying PD pathology^{1,7,8}. More recently, a meta-analysis including 5,166 Chinese patients with PD led to the identification of another locus *NDN/PWRN4*⁹. Despite the large disparity in sample size and the genetic loci identified by the two studies, both works estimated similar total heritability of AAO of 10-14%^{7,9}. They also showed an inverse correlation between a polygenic risk score (PRS) and AAO based on risk loci for PD on individuals of similar ancestry, suggesting overlap between the pathways underlying disease susceptibility and AAO in PD.

Recent studies have underscored the relevance of inclusion of ethnic diversity in genomic research^{9,10}. The COURAGE-PD (COMprehensive Unbiased Risk Factor Assessment for Genetics and Environment in Parkinson's Disease) is a worldwide collaboration consortium comprising 35 PD study cohorts which aims to address this disparity to some extent in PD research¹¹. The present study aims to perform an AAO GWAS in COURAGE-PD and to investigate the validity of previously observed loci by conducting one of the largest meta-analysis of PD AAO GWAS to date by combining previous International Parkinson's Disease Genomics Consortium (IPDGC) AAO GWAS (n=17, 415) with newly generated

COURAGE-PD AAO GWAS (n=8,535), resulting in a combined dataset of 25,950 patients with PD. Lastly, we investigate the influence of a PD PRS on PD AAO to dissect the potential overlapping etiology.

Methods

Study cohorts and participants

The COURAGE-PD consortium comprises data from 15,849 patients with PD and 11,444 controls of predominantly European ancestry from 35 cohorts with a major contribution from the Genetic Epidemiology of Parkinson's disease (GEOPD) Consortium (www.geopd.net). Quality control (QC) of genome-wide data was performed in each COURAGE-PD study cohort. See **eMethods** for more details, including collected phenotypic data. AAO was defined based on the initial manifestation of motor symptoms associated with PD, as described elsewhere⁶. Post-imputation, only sporadic PD patients with data available on AAO and not overlapping with previous IPDGC AAO GWAS were included in the present study, leaving 8,535 samples from 30 cohorts. These comprised 26 European and four East-Asian ancestry cohorts.

Genotype-phenotype analysis

Regression analysis and meta-analysis of study-specific estimates

Linear regression analysis of imputed dosages with AAO was performed in each study cohort using an additive model, implemented in *rvtests*, correcting for gender and the first five principal components¹². The selection of five principal components was based on study cohort specific scree plots. The scree plot flattened out after the third factor for majority of study cohorts, with few exceptions, where five factors explained the highest proportion of the total variance. This was followed by combining study-specific results through an inverse variance weighted (IVW) fixed-effect meta-analyses conducted using METAL^{13, 14}. Additionally, only those variants which were successfully genotyped in at least 2/3rd of study cohorts were included for further interpretation. Similarly, the variants with I^2 statistic $\geq 50\%$

were considered to have substantial heterogeneity and were excluded from further interpretation. We also employed additive random-effect meta-analyses using the DerSimonian-Laird estimator to check the influence of heterogeneity on our findings¹⁵. The quantile-quantile (QQ) plot was generated using R to judge the potential influence of population stratification on the overall significance of the effect estimates. We considered a $P < 5 \times 10^{-8}$ as genome-wide significant and a $P < 1 \times 10^{-6}$ as suggestive evidence for a potential association⁹. We also considered a Bonferroni-corrected $P < 0.025$ for reporting replication signals originating from two SNPs (rs356203 (*SNCA*), rs34311866 (*TMEM175*)) that reached a genome-wide significance in the previous largest meta-analysis of AAO of PD⁷. The results were visualized using R generated Manhattan and LocusZoom generated regional association plots¹⁶. We conducted LD score regression with LDSC (using summary-level data) to estimate heritability explained by the PD AAO GWAS¹⁷. We also performed a meta-analysis of COURAGE-PD AAO (n=8535) with the previous largest AAO meta-analysis comprising IPDGC dataset (n=17,415) to discover potentially new loci and improve heritability estimates⁷.

Correlation between case-control GWAS and AAO GWAS

We used two approaches to assess the correlation between PD case-control GWAS meta-analysis and COURAGE-PD AAO GWAS meta-analysis. Firstly, we computed the genome-wide genetic correlation between PD status and PD AAO in COURAGE-PD dataset using the cross-trait LD score regression method¹⁷. Secondly, we used effect estimates of significant genetic variants ($P < 5 \times 10^{-8}$) identified by combining of COURAGE-PD case-control GWAS meta-analysis dataset with the IPDGC-PD case-control GWAS meta-analysis dataset to generate individual-specific polygenic risk scores (PRS) in the COURAGE-PD AAO population, using PRSice2¹⁸. Linear regression analysis of PRS with AAO was performed, correcting for gender and the first five principal components.

Subgroup analysis and power computation

A subgroup analysis was performed to explore the influence of ethnicity and gender on the AAO GWAS as well as the correlation between case-control and AAO GWAS meta-analyses. The power was estimated using QUANTO 1.2.4¹⁹.

Expression quantitative trait loci analysis

We further explored the potential influence of novel variants identified in the present study on the expression traits using the gene expression data from the Genotype-Tissue Expression Project using the GTEx portal (gtexportal.org, Data Source: GTEx Analysis Release V8 (dbGaP Accession phs000424.v8.p2) and UK Brain Expression Consortium (UKBEC) using the Braineac portal (braineac.org).^{20, 21}

Standard protocol approvals, registrations, and patient consents

The study was conducted at the University of Tübingen, and the ethical approval was obtained by the local institutional review board (IRB) of respective study sites. All the study participants provided signed informed consent.

Data availability

Summary statistics of COURAGE-PD AAO GWAS used in the meta-analysis are available from the corresponding author upon reasonable request. In addition, IPDGC summary statistics for AAO GWAS was downloaded from the IPDGC website (<https://pdgenetics.org/resources>). Significant SNPs of risk of PD based on meta-analysis of COURAGE-PD and IPDGC datasets, used in the PRS calculation can be found in the original publication (Grover et al. in preparation). Relevant programming scripts used for the present work are available at the github website of Center of Genetic Epidemiology (CGE) at Tübingen (https://github.com/CGEatTuebingen/Ageatonsset_GWAS_Courage-PD).

Results

Main study outcome variable

The final cohort after QC included a total of 8,535 patients with PD, 7,847 of European ancestry (91.9%) and 688 of East-Asian ancestry (9.1%). The average AAO in the COURAGE-PD dataset was 58.9 years (SD=11.6), with an under-representation of females (40.2%) (see eTable 1). We did not observe any major influence of gender or ethnicity on AAO. Furthermore, the average AAO was slightly lower than that reported in the IPDGC dataset (62.1 years; SD=12.1), a difference that was statistically significant ($P<0.05$).

Genetic heritability of the study outcome

Using summary-level data, the total estimated heritability (h^2) in the COURAGE-PD dataset was 0.083 (SE=0.057). Similar heritability estimates were observed in the European sub-cohort ($h^2=0.079$, SE=0.061). However, the heritability estimates in the Asian sub-cohort could not be reliably computed due to an insufficient number of patients. Additionally, we failed to achieve any improvement in heritability estimates, although with improved accuracy by combining COURAGE-PD with IPDGC dataset ($h^2=0.078$, SE=0.018).

Genome-wide meta-analysis

COURAGE-PD

GWAS meta-analysis

The genomic inflation factor λ was 1.016 (See eFigure 1 for the QQ plot). None of the loci reached genome-wide significance (Figure 1). We observed one locus reaching suggestive genome-wide significance level, *PDZPHIP* (chr5) ($\beta(\text{SE})_{\text{COURAGE}}=-1.456(0.293)$, $P_{\text{COURAGE}}=6.91 \times 10^{-7}$). However, stratifying the analyses by ethnicities, we did not observe any suggestive involvement of *PDZPHIP* locus in the European sub-cohort. (see eTable 2). Interestingly, despite being a smaller sub-cohort, *SUGCT* locus on chromosome (Chr) 7 was detected as a suggestive locus in the East-Asian sub-cohort ($\beta(\text{SE})_{\text{COURAGE-EASIAN}}=13.681(2.769)$, $P_{\text{COURAGE-EASIAN}}=7.80 \times 10^{-7}$). Furthermore, the stratified analysis

provided suggestive evidence of three loci, *RHEB* (chr8) in males ($\beta(\text{SE})_{\text{COURAGE-M}} = -1.112(0.222)$, $P_{\text{COURAGE-M}} = 5.15 \times 10^{-7}$), and *MTHFD1L* (chr6) ($\beta(\text{SE})_{\text{COURAGE-F}} = -1.995(0.402)$, $P_{\text{COURAGE-F}} = 6.78 \times 10^{-7}$) and *KNH3* (chr12) in females ($\beta(\text{SE})_{\text{COURAGE-F}} = 2.176(0.432)$, $P_{\text{COURAGE-F}} = 4.59 \times 10^{-7}$) (see eTable 2).

In the replication of previously reported variants, only the *TMEM175* variant (rs34311866: $\beta(\text{SE})_{\text{COURAGE}} = 0.477(0.203)$, $P_{\text{COURAGE}} = 0.018$) reached Bonferroni-corrected nominal levels of significance in the COURAGE-PD dataset. Nevertheless, the *SNCA* variant also showed a trend towards association (rs356203: $\beta(\text{SE})_{\text{COURAGE}} = 0.362(0.172)$, $P_{\text{COURAGE}} = 0.035$).

Meta-analysis of COURAGE-PD and IPDGC datasets

The meta-analysis of COURAGE-PD and IPDGC datasets led to the identification of two loci that reached genome-wide significance (see eTable 2; Figure 2). The *SNCA* variant, rs983361, was the most strongly associated SNP, with the presence of allele T (frequency=0.204) leading to an average delay in AAO by 0.72 years ($\beta(\text{SE})_{\text{COURAGE+IPDGC}} = 0.720(0.122)$, $P_{\text{COURAGE+IPDGC}} = 3.13 \times 10^{-9}$). This association, however, appeared to be driven by the strong association reported by IPDGC dataset, with negligible effect detected in the COURAGE-PD dataset ($P_{\text{COURAGE/COURAGE-EUR (rs983361)}} = 0.022$; Not detected in East-Asian sub-population) (see eFigure 2A), which was also reflected in the loss of genome-wide significance, when using an additive random effect model ($P = 2.98 \times 10^{-6}$). On the other hand, another independent locus on the same chromosome, *BST1* (rs4698412) showed similar effects in COURAGE-PD and IPDGC datasets ($\beta(\text{SE})_{\text{COURAGE}} = -0.633(0.175)$, $P_{\text{COURAGE}} = 2.95 \times 10^{-4}$; $\beta(\text{SE})_{\text{IPDGC}} = -0.480(0.115)$, $P_{\text{IPDGC}} = 3.04 \times 10^{-5}$), and the combination of both estimates resulted in the identification of a novel genome-wide significant *BST1* locus for AAO ($\beta(\text{SE})_{\text{COURAGE+IPDGC}} = -0.526(0.096)$, $P_{\text{COURAGE+IPDGC}} = 4.41 \times 10^{-8}$) (see eFigure 2B). The rs4698412 allele A (frequency=0.562) at the locus led to an average earlier AAO of 0.526 years in PD patients. No genetic heterogeneity was detected in the observed association

($I^2=0$; Heterogeneity $P=0.465$). Furthermore, we did not observe any change in the effect estimates, when using the additive random effect model ($P=4.41 \times 10^{-8}$).

The previously reported *TMEM175* (rs34311866) showed suggestive association in the combined analysis ($\beta(\text{SE})_{\text{COURAGE+IPDGC}}=0.589(0.114)$, $P_{\text{COURAGE+IPDGC}}=2.64 \times 10^{-7}$) that appeared to be driven by previously reported findings in the IPDGC dataset ($\beta(\text{SE})_{\text{IPDGC}}=0.642(0.139)$, $P_{\text{IPDGC}}=3.72 \times 10^{-6}$) (see eFigure 2C). Another locus *AL391867.1/RP11-342F21.1* (rs62582905), a locus of unknown biological significance also crossed the threshold of suggestive association in the same analysis ($\beta(\text{SE})_{\text{COURAGE+IPDGC}}=-1.456(0.293)$, $P_{\text{COURAGE}}=6.62 \times 10^{-7}$) (see eTable 2). However, unlike the *TMEM175* association, the association with *AL391867.1/RP11-342F21.1* was observed to be stronger in the COURAGE-PD dataset ($\beta(\text{SE})_{\text{COURAGE}}=-1.925(0.447)$, $P_{\text{COURAGE}}=1.64 \times 10^{-5}$).

We carried out a sensitivity analysis by excluding the Asian sub-cohort from the COURAGE dataset, followed by combining with the IPDGC dataset. Similar findings were observed for the two genome-wide significant loci (*SNCA* rs983361: $P_{\text{COURAGE(EUR)+IPDGC}}=3.13 \times 10^{-9}$, *BST1* rs4698412: $P_{\text{COURAGE(EUR)+IPDGC}}=6.27 \times 10^{-8}$) (see eTable 2). A similar sensitivity analysis for the previously reported APOE $\epsilon 4$ locus also showed suggestive association with PD AAO (*APOE* rs429358: $\beta(\text{SE})_{\text{COURAGE(EUR)+IPDGC}}=0.711(0.145)$, $P_{\text{COURAGE(EUR)+IPDGC}}=9.33 \times 10^{-7}$). However, the association was primarily driven by highly significant findings in the IPDGC dataset ($\beta(\text{SE})_{\text{IPDGC}}=0.754(0.171)$, $P_{\text{IPDGC}}=9.86 \times 10^{-6}$; $\beta(\text{SE})_{\text{COURAGE(EUR)}}=0.599(0.275)$, $P_{\text{COURAGE(EUR)}}=0.029$).

Correlation between genetic risk for PD and PD AAO

Using complete GWAS summary datasets for COURAGE-PD case-control and COURAGE-PD AAO, we observed a non-significant negative genetic correlation between PD and PD AAO ($r_g=-0.291$, $SE=0.224$; $P=0.186$). Furthermore, a slightly stronger genetic correlation was observed when restricting our correlation analysis to European sub-cohorts only ($r_g=-0.315$; $SE=0.252$; $P=0.211$).

When using PRS based on the significant loci detected in the meta-analysis of COURAGE-PD and IPDGC European datasets, as reported elsewhere, we observed that each unit increase in SD in PRS lead to a significant decrease in AAO in COURAGE-PD by 0.58 years ($\beta(\text{SE})_{\text{COURAGE}}=-0.581(0.149)$, $P_{\text{COURAGE}}=9.35 \times 10^{-5}$). Despite the significant findings, the PRS explained only 0.59% of the genetic proportion of PD heritability.

Expression quantitative trait analysis of novel *BST1* locus

The mining of the Genotype-Tissue Expression (GTEx) portal showed that rs4698412 representing the *BST1* locus is a highly significant expression quantitative trait locus (eQTL) for *CD38* in the basal ganglia (caudate, nucleus accumbens and putamen) and cortex (NES=-0.32 – -0.44; $P < 1 \times 10^{-10}$) (**Table 1**). The expression analysis also showed a strong dosage effect with a consistent lower expression in the presence of AA genotype compared to GG genotype with a higher expression, irrespective of brain tissue type. In addition, we also found that SNP modulates the expression of *BST1* in whole blood. However, the effect was considerably lower in comparison to that observed on *CD38* expression levels in brain tissues (NES=-0.071; $P=1.7 \times 10^{-6}$). The follow-up of association of rs4698412 with expression in brain tissues in the UKBEC database further confirmed the role of basal ganglia with *CD38* as the most significantly associated expressed gene in the putamen ($P=7.1 \times 10^{-6}$) (**Table 1**)

Discussion

The identification of genetic determinants that modify the disease progression will not only help to increase our understanding of PD etiopathogenesis, but also enables the development of strategies that could be used for therapeutic intervention for at-risk carriers. Our study not only validates previously reported AAO PD loci in the COURAGE-PD dataset, but our meta-analysis with IPDGC data also provides the first genome-wide significant evidence that the known *BST1* PD risk locus affects AAO. Interestingly, the variant, rs4698412, representing the *BST1* locus, showed a similar large effect in COURAGE-PD and

IPDGC providing strong evidence that this is a bona-fide genetic locus for PD AAO. Lastly, using significant SNPs from the meta-analysis of COURAGE-PD and IPDGC case-control datasets, we demonstrate an inverse association between a PD PRS and AAO of PD.

Numerous genetic loci for familial and sporadic PD have been well characterized. The existence of overlapping loci between familial and sporadic PD suggests a complex but interconnected relationship between PD and age. Several meta-analyses of candidate genes and GWAS have previously recognized the *BST1* locus as a locus that could influence the development of sporadic late-onset PD^{6, 22-24}. Notably, the *BST1* locus has been demonstrated to play a role in both Asian and European PD populations.^{6, 22-24} The genome-wide significant *BST1* variant, rs4698412 observed in our AAO meta-analysis, is also identical to the top *BST1* variant reported in the latest PD GWAS meta-analysis⁶. Interestingly, regional plots showed that the genome-wide significant variant, rs4694812 was neither the top genetic variant in the *BST1* locus in IPDGC nor COURAGE AAO PD datasets. While, rs4694819 (r^2 with rs4694812<0.6) was the most significant variant in the COURAGE AAO dataset, rs11724635 (r^2 with rs4698412=1.0) was the most significant variant in the IPDGC AAO dataset (e**Figure 2B**).

BST1 was first identified as a gene encoding a cell surface receptor on bone marrow stromal cells (bone marrow stromal cell antigen 1) with a role in promoting the growth of hematopoietic stromal cells²⁵. In addition to its role as a receptor, it also exhibits ADP-ribosyl cyclase activity, leading to the generation of cyclic ADP-ribose (cADPR), with a role in intrinsic Ca^{2+} regulation²⁶. The dual functional protein, a highly conserved glycosylphosphatidylinositol (GPI)-anchored glycoprotein (also known as CD157), is now known to be expressed in a wide variety of tissues, including vascular endothelium and follicular dendritic cells, with an ability to perform a wide variety of immune system and inflammation-related cellular functions²⁷. The initial identification of *BST1/CD157* as a potential risk locus for sporadic late-onset PD in a GWAS in the Japanese population by

Satake *et al.* 2009, led to several functional studies aimed at deciphering its potential neuronal role in influencing PD phenotype²². Several knockout mouse model studies have shown that *BST1* can influence social behaviour. However, the studies failed to demonstrate any influence on motor functioning, the cardinal feature which is impaired in patients with PD^{28, 29}. The eQTL analysis demonstrated a highly significant effect of the *BST1* locus, rs4694812, on gene expression, with the A allele resulting in a decreased expression of *CD38*, a paralog of *CD157*, in a dose-dependent manner. *CD38* and *CD157* are contiguous gene duplicates, which belong to same gene family with a similar role of dual functional protein and an ability to modulate social behavior^{30, 31}. Interestingly, unlike *CD157*, *CD38* knock out mice have been shown to have higher locomotor activity³². Furthermore, the highly significant increased expression of *CD38* was mainly observed in the striatum, a region directly implicated in motor dysfunction in PD. Interestingly, a statistically underpowered brain imaging study in humans suggested that allele A of *BST1* SNP rs4698412 leads to deficits in the right lingual gyrus region in the brain during the progression of PD³³. This brain region is known to play a role in spatial orientation and visuospatial information processing. However, specific molecular and neuronal pathways influenced by altered *CD38* expression in basal ganglia, with a potential role in triggering earlier AAO in sporadic PD, remain unclear.

SNCA is one of the most consistently observed significant loci in both early and late-onset PD and has been suggested to play a critical role in the age-related hierarchy of disease onset. While monogenic PD, often with relatively early onset, is attributed to rare point mutations and multiple copies of the *SNCA* gene, susceptibility to late-onset PD is attributed to common variants^{6, 10, 34-36}. In addition to being a leading locus in the largest GWAS of sporadic PD to date, the locus was also recently reported to be a top locus in influencing AAO in Europeans in a meta-analysis comprising IPDGC and 23andMe datasets (n=28,568)⁷. An SNP present towards the 3' end (rs356203) of the *SNCA* gene was observed as the strongest genome-wide significant variant originating from the region ($P=1.9 \times 10^{-12}$). Based on the

conditional analysis, the study also identified an independent signal at the 5' end of the gene, rs983361 ($P=6.8 \times 10^{-6}$). A recent GWAS of AAO in 5166 East Asian (Chinese) PD patients further reported a slightly weaker signal originating from another independent *SNCA* variant, rs3775458 ($P=9.92 \times 10^{-7}$)⁹. Using the 1000 genome phase 3 dataset, we failed to detect any LD among the three variants in both European and East Asian populations (data not shown here). Upon screening of the *SNCA* locus in the COURAGE-PD dataset, we observed nominal significance of all the three variants ($P_{\text{COURAGE (rs356203)}}=0.035$, $P_{\text{COURAGE (rs3775458)}}=0.005$, $P_{\text{COURAGE (rs983361)}}=0.022$), possibly suggesting a consistence influence of different loci around the *SNCA* region in determining AAO in different worldwide PD populations. The combining of our dataset with IPDGC further showed an independent genome-wide significant signal originating from the 3' end of the *SNCA* gene (rs983361), as shown in the results section above. Notably, we also observed an independent signal at the 5' end (rs356203). However, the variant was excluded for further interpretation due to high heterogeneity observed when combining IPDGC and COURAGE datasets ($\beta(\text{SE})_{\text{COURAGE+IPDGC}}=-0.591(0.097)$, $P_{\text{COURAGE+IPDGC}}=9.28 \times 10^{-10}$; $I^2=61.9\%$).

Another PD locus, *TMEM175*, was previously shown to reach genome-wide significance in an AAO study³⁷. Similar to *SNCA*, our study also demonstrated replication of the *TMEM175* locus in the COURAGE-PD AAO dataset with a nominal level of significance ($P=0.018$). The subsequent combining of the non-synonymous coding variant, rs3431186 (p.M393T), representing the genome-wide significant locus, in the IPDGC dataset with the COURAGE-PD, resulted in suggestive level of association without any underlying heterogeneity ($P_{\text{COURAGE+IPDGC}}=2.64 \times 10^{-7}$; $I^2=0.0$). On the contrary, a recent East-Asian GWAS failed to observe any signal originating from the locus, possibly suggesting contribution of the locus mainly in the European populations.⁹ A previous study also reported a borderline significant association of the variant rs429358 representing *APOE ε4* locus with PD-AAO ($P=5.69 \times 10^{-8}$) in a combined dataset ($n=28568$) comprising IPDGC and 23andMe

datasets⁷. The study, however, suggested that the association at the locus could be an age-related effect with a highly significant association with the age of controls ($P=1.49 \times 10^{-5}$). The variant also resulted in a suggestive association upon merging of COURAGE-PD European dataset only with the IPDGC dataset ($P=9.3 \times 10^{-7}$). These findings are consistent with the failure to detect association of *APOE* $\epsilon 4$ locus with PD-AAO in the recently reported East-Asian GWAS⁹. However, being a longevity marker, the suggestive finding of *APOE* $\epsilon 4$ locus in Europeans must be interpreted with caution.

Our study has several strengths and limitations. Our study provides the largest independent dataset for testing the reliability of previously discovered AAO loci in a highly diverse and predominantly European population. Another strength of our study was the availability of data on AAO on all the study participants as opposed to age of diagnosis, often used as a proxy for AAO. One of the significant limitations of our findings was the lack of ready access to the recently published East-Asian AAO GWAS dataset that prevented us from drawing any conclusion on the validity of the novel *BST1* locus in the East-Asian population. Likewise, the unavailability of the 23andMe dataset to us has precluded us from making an unequivocal claim on our *BST1* findings. Hopefully, the inclusion of other datasets, such as 23andMe and East-Asian GWAS datasets, will help further to refine the signals originating from the *BST1* locus. We also suggest that loci identified through meta-analysis in the COURAGE-PD dataset (*PDZPH1P*) and subsequent stratification by gender (*RHEB*, *MTHFD1L*, *KNH3*) and ethnicity (*MOAP1/TMEM251 1*, *SUGCT*) be meta-analysed with these unavailable datasets. Another limitation was our inability to conduct gene-gene interaction due to the limited sample size in the present study. The possibility of complex interactions among various loci on Chr 4 in modulating AAO cannot be ruled out. A recent study showed the association of several genome-wide significant loci on the X Chr with PD³⁸. It is also possible that some of these variants may also modulate AAO. However, due to potential analytic challenges from calling and imputation of X Chr genotypes, to model

uncertainty associated with random X Chr inactivation, we excluded the X Chr variants from the present analysis³⁹. And lastly, it is hoped that in the future, the availability of a larger dataset would enable us to integrate additional layers of genetic data, including rare and copy-number variants⁴⁰.

Our findings clearly highlight the importance of combining GWAS from diverse populations representative of worldwide populations to refine the genetic architecture underlying a complex trait like AAO. Our COURAGE-PD dataset suggests a role for additional pathways in addition to α -synuclein mechanisms of modulating PD pathogenesis and influencing AAO in worldwide PD populations.

WNL-2022-200706_efig-<http://links.lww.com/WNL/C86>

WNL-2022-200706_emth-<http://links.lww.com/WNL/C87>

WNL-2022-200706_etab-<http://links.lww.com/WNL/C88>

References

1. Soldner F, Stelzer Y, Shivalila CS, et al. Parkinson-associated risk variant in distal enhancer of α -synuclein modulates target gene expression. *Nature* 2016;533:95-99.
2. Lill CM. Genetics of Parkinson's disease. *Mol Cell Probes* 2016;30:386-396.
3. Bloem BR, Okun MS, Klein C. Parkinson's disease. *Lancet* 2021;397:2284-2303.
4. Nalls MA, Pankratz N, Lill CM, et al. Large-scale meta-analysis of genome-wide association data identifies six new risk loci for Parkinson's disease. *Nat Genet* 2014;46:989-993.
5. Chang D, Nalls MA, Hallgrímsdóttir IB, et al. A meta-analysis of genome-wide association studies identifies 17 new Parkinson's disease risk loci. *Nat Genet* 2017;49:1511-1516.
6. Nalls MA, Blauwendraat C, Vallerga CL, et al. Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet Neurol* 2019;18:1091-1102.
7. Blauwendraat C, Heilbron K, Vallerga CL, et al. Parkinson's disease age at onset genome-wide association study: Defining heritability, genetic loci, and α -synuclein mechanisms. *Mov Disord* 2019;34:866-875.
8. Jinn S, Drolet RE, Cramer PE, et al. TMEM175 deficiency impairs lysosomal and mitochondrial function and increases α -synuclein aggregation. *Proc Natl Acad Sci U S A* 2017;114:2389-2394.
9. Li C, Ou R, Chen Y, et al. Genetic Modifiers of Age at Onset for Parkinson's Disease in Asians: A Genome-Wide Association Study. *Mov Disord* 2021.

10. Foo JN, Chew EGY, Chung SJ, et al. Identification of Risk Loci for Parkinson Disease in Asians and Comparison of Risk Between Asians and Europeans: A Genome-Wide Association Study. *JAMA Neurol* 2020;77:746-754.
11. Grover S, Kumar-Sreelatha AA, Bobbili DR, et al. Replication of a Novel Parkinson's Locus in a European Ancestry Population. *Mov Disord* 2021.
12. Zhan X, Hu Y, Li B, Abecasis GR, Liu DJ. RVTESTS: an efficient and comprehensive tool for rare variant association analysis using sequence data. *Bioinformatics* 2016;32:1423-1426.
13. Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 2010;26:2190-2191.
14. Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst* 1959;22:719-748.
15. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-188.
16. Pruim RJ, Welch RP, Sanna S, et al. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* 2010;26:2336-2337.
17. Bulik-Sullivan BK, Loh PR, Finucane HK, et al. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 2015;47:291-295.
18. Choi SW, Mak TS, O'Reilly PF. Tutorial: a guide to performing polygenic risk score analyses. *Nat Protoc* 2020;15:2759-2772.
19. Gauderman WJ. Sample size requirements for matched case-control studies of gene-environment interaction. *Stat Med* 2002;21:35-50.
20. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* 2015;348:648-660.
21. Ramasamy A, Trabzuni D, Guelfi S, et al. Genetic variability in the regulation of gene expression in ten regions of the human brain. *Nat Neurosci* 2014;17:1418-1428.
22. Satake W, Nakabayashi Y, Mizuta I, et al. Genome-wide association study identifies common variants at four loci as genetic risk factors for Parkinson's disease. *Nat Genet* 2009;41:1303-1307.
23. Li J, Luo J, Liu L, Fu H, Tang L. The association between CD157/BST1 polymorphisms and the susceptibility of Parkinson's disease: a meta-analysis. *Neuropsychiatr Dis Treat* 2019;15:1089-1102.
24. Sharma M, Ioannidis JP, Aasly JO, et al. Large-scale replication and heterogeneity in Parkinson disease genetic loci. *Neurology* 2012;79:659-667.
25. Kaisho T, Ishikawa J, Oritani K, et al. BST-1, a surface molecule of bone marrow stromal cell lines that facilitates pre-B-cell growth. *Proc Natl Acad Sci U S A* 1994;91:5325-5329.
26. Fliegert R, Gasser A, Guse AH. Regulation of calcium signalling by adenine-based second messengers. *Biochem Soc Trans* 2007;35:109-114.
27. Ortolan E, Augeri S, Fissolo G, Musso I, Funaro A. CD157: From immunoregulatory protein to potential therapeutic target. *Immunol Lett* 2019;205:59-64.
28. Kasai S, Yoshihara T, Lopatina O, Ishihara K, Higashida H. Selegiline Ameliorates Depression-Like Behavior in Mice Lacking the CD157/BST1 Gene, a Risk Factor for Parkinson's Disease. *Front Behav Neurosci* 2017;11:75.
29. Lopatina O, Yoshihara T, Nishimura T, et al. Anxiety- and depression-like behavior in mice lacking the CD157/BST1 gene, a risk factor for Parkinson's disease. *Front Behav Neurosci* 2014;8:133.
30. Quarona V, Zaccarello G, Chillemi A, et al. CD38 and CD157: a long journey from activation markers to multifunctional molecules. *Cytometry B Clin Cytom* 2013;84:207-217.

31. Higashida H, Liang M, Yoshihara T, et al. An immunohistochemical, enzymatic, and behavioral study of CD157/BST-1 as a neuroregulator. *BMC Neurosci* 2017;18:35-35.
32. Jin D, Liu HX, Hirai H, et al. CD38 is critical for social behaviour by regulating oxytocin secretion. *Nature* 2007;446:41-45.
33. Shen YT, Wang JW, Wang M, et al. BST1 rs4698412 allelic variant increases the risk of gait or balance deficits in patients with Parkinson's disease. *CNS Neurosci Ther* 2019;25:422-429.
34. Polymeropoulos MH, Lavedan C, Leroy E, et al. Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 1997;276:2045-2047.
35. Pihlstrøm L, Blauwendraat C, Cappelletti C, et al. A comprehensive analysis of SNCA-related genetic risk in sporadic parkinson disease. *Ann Neurol* 2018;84:117-129.
36. Du B, Xue Q, Liang C, et al. Association between alpha-synuclein (SNCA) rs11931074 variability and susceptibility to Parkinson's disease: an updated meta-analysis of 41,811 patients. *Neurol Sci* 2020;41:271-280.
37. Blauwendraat C, Reed X, Krohn L, et al. Genetic modifiers of risk and age at onset in GBA associated Parkinson's disease and Lewy body dementia. *Brain* 2020;143:234-248.
38. Le Guen Y, Napolioni V, Belloy ME, et al. Common X-Chromosome Variants Are Associated with Parkinson Disease Risk. *Ann Neurol* 2021;90:22-34.
39. Deng WQ, Mao S, Kalnapekns A, et al. Analytical strategies to include the X-chromosome in variance heterogeneity analyses: Evidence for trait-specific polygenic variance structure. *Genet Epidemiol* 2019;43:815-830.
40. Rudakou U, Yu E, Krohn L, et al. Targeted sequencing of Parkinson's disease loci genes highlights SYT11, FGF20 and other associations. *Brain* 2021;144:462-472.

Figure 1. Manhattan plot of COURAGE-PD age at onset GWAS .




Figure 2. Manhattan plot of meta-analysis of COURAGE-PD and IPDGC age at onset GWAS.




Table 1. eQTL lookup of BST1 SNP rs4698412 from GTEx in brain tissue

Table 1. eQTL lookup of *BST1* SNP rs4698412 from GTEx and UKBEC in brain tissue.

Database	Gene	Symbol	P-Value	NES	Tissue
<i>GTEx</i>	ENSG00000004468	<i>CD38</i>	3.3e-16	-0.44	Caudate (basal ganglia)
	ENSG00000004468	<i>CD38</i>	5.5e-15	-0.39	Cortex
	ENSG00000004468	<i>CD38</i>	1.4e-13	-0.39	Nucleus accumbens (basal ganglia)
	ENSG00000004468	<i>CD38</i>	1.4e-11	-0.32	Putamen (basal ganglia)
	ENSG00000004468	<i>CD38</i>	6.6e-6	-0.21	Frontal Cortex (BA9)
	ENSG00000237765	<i>FAM200B</i>	1.0e-5	0.23	Cerebellar Hemisphere
	ENSG00000004468	<i>CD38</i>	1.2e-5	-0.26	Anterior cingulate cortex (BA24)
	ENSG00000237765	<i>FAM200B</i>	1.4e-5	0.23	Cortex
	ENSG00000004468	<i>CD38</i>	1.4e-5	-0.22	Hypothalamus
	<i>UKBEC</i>	ENSG00000118564	<i>FBXL5</i>	5.1e-7	NA
ENSG00000004468		<i>CD38</i>	7.1e-6	NA	Putamen (basal ganglia)
ENSG00000004468		<i>CD38</i>	2.1e-5	NA	Hippocampus
ENSG00000137449		<i>CPEB2</i>	2.4e-5	NA	Medulla

GTEx: Genotype-Tissue Expression Project, NA: Not available, UKBEC: UK Brain Expression Consortium

Neurology®

Genome-wide Association and Meta-analysis of Age-at-Onset in Parkinson Disease: Evidence From COURAGE-PD Consortium

Sandeep Grover, Ashok Kumar Sreelatha Ashwin, Lasse Pihlstrom, et al.

Neurology published online May 26, 2022

DOI 10.1212/WNL.0000000000200699

This information is current as of May 26, 2022

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/early/2022/05/26/WNL.0000000000200699.full
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Association studies in genetics http://n.neurology.org/cgi/collection/association_studies_in_genetics Basal ganglia http://n.neurology.org/cgi/collection/basal_ganglia Burden of disease http://n.neurology.org/cgi/collection/burden_of_disease Case control studies http://n.neurology.org/cgi/collection/case_control_studies Parkinson's disease/Parkinsonism http://n.neurology.org/cgi/collection/parkinsons_disease_parkinsonism
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

