

Musiol Stephanie (Orcid ID: 0000-0001-8356-4343)
 Standl Marie (Orcid ID: 0000-0002-5345-2049)
 Schmidt-Weber Carsten (Orcid ID: 0000-0002-3203-8084)
 Ussar Siegfried (Orcid ID: 0000-0001-7575-0920)

Original Article

Dietary digestible carbohydrates are associated with higher prevalence of asthma in humans and with aggravated lung allergic inflammation in mice

Stephanie Musiol^{1,*,#}, Carla P. Harris^{2,3,*}, Ruth Karlina^{4,5,#}, Johanna M. Gostner⁶, Birgit Rathkolb^{5,7,8}, Benjamin Schnautz¹, Evelyn Schneider¹, Lisa Mair⁶, Ernesto Elorduy Vergara^{9,10}, Claudia Flexeder², Sibylle Koletzko^{3,11}, Carl-Peter Bauer¹², Tamara Schikowski¹³, Dietrich Berdel¹⁴, Andrea von Berg¹⁴, Gunda Herberth¹⁵, Jan Rozman^{5,7,16}, Martin Hrabe de Angelis^{5,7,17}, Marie Standl^{2,10}, Carsten B. Schmidt-Weber^{1,10}, Siegfried Ussar^{4,5,18}, Francesca Alessandrini¹

¹ Center of Allergy & Environment (ZAUM), Technical University of Munich (TUM) and Helmholtz Zentrum München, German Research Center for Environmental Health, Munich, Germany

² Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

³ Dr. von Hauner Children's Hospital, University Hospital, LMU of Munich, Munich, Germany

⁴ RG Adipocytes & Metabolism, Institute for Diabetes & Obesity, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

⁵ German Center for Diabetes Research (DZD), Neuherberg, Germany

⁶ Institute of Medical Biochemistry, Biocenter, Medical University of Innsbruck, Innsbruck, Austria.

⁷ Institute of Experimental Genetics, German Mouse Clinic, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

⁸ Institute of Molecular Animal Breeding and Biotechnology, Gene Center, Ludwig-Maximilians-University Munich, Munich, Germany

⁹ Institute of Lung Biology and Disease, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/all.15589](https://doi.org/10.1111/all.15589)

This article is protected by copyright. All rights reserved.

¹⁰ German Center of Lung Research (DZL), Munich, Germany

¹¹ Department of Pediatrics, Gastroenterology and Nutrition, School of Medicine Collegium Medicum University of Warmia and Mazury, Olsztyn, Poland

¹² Department of Pediatrics, Technical University of Munich, Munich, Germany

¹³ IUF-Leibniz Research Institute for Environmental Medicine, Düsseldorf, Germany

¹⁴ Research Institute, Department of Pediatrics, Marien-Hospital Wesel, Wesel, Germany

¹⁵ Department of Environmental Immunology, Helmholtz Centre for Environmental Research-UFZ, Leipzig, Germany

¹⁶ Czech Centre for Phenogenomics, Institute of Molecular Genetics of the Czech Academy of Sciences, Průmyslová 595, 252 50 Vestec, Czech Republic

¹⁷ Chair of Experimental Genetics, TUM School of Life Sciences (SoLS), Technische Universität München, Freising, Germany

¹⁸ Department of Medicine, Technical University of Munich (TUM), Munich, Germany

Current address: Eurofins BioPharma Product Testing Munich GmbH, Planegg, Germany

*equal contribution

Corresponding author

Francesca Alessandrini, PhD

ZAUM Center of Allergy & Environment

Technical University of Munich and Helmholtz Zentrum München

Ingolstädter Landstr. 1

85764 Neuherberg, Germany

Tel. +49 (0)89 3187 2524

Email: franci@helmholtz-munich.de

Short running title: Carbohydrate intake enhance lung allergic inflammation

Word count: 4241

List of abbreviations

AAI: Allergic airway inflammation

ALA: alpha-linolenic acid

APE: Aqueous pollen extract

BAL: Bronchoalveolar lavage

FACS: Fluorescence activated cell sorting

GAE: Gallic acid equivalents

GINIplus: German Infant study on the Influence of Nutrition Intervention plus environmental and genetic influences on allergy development

GTT: Glucose tolerance test

HDM: House dust mite

HFD: High-fat diet

LA: linoleic acid

LF-HA: Low fat, high starch (*Amylum*)

LF-HS: Low fat, high sucrose

LISA: Life-style related factors on the development of the Immune System and Allergies in East and West Germany

MUFA: monounsaturated fatty acids

NEFA: non-esterified fatty acids

PAS: Periodic acid–Schiff

PBS: Dulbecco's phosphate-buffered saline, no calcium, no magnesium

SFA: saturated fatty acids, linoleic acid=LA, and alpha-linolenic acid=ALA

T_H: T helper

tIgE: total immunoglobulin E

Treg: regulatory T cell

VAT: Visceral adipose tissue

Acknowledgments: *GINplus Cohort:* The authors thank all the families for their participation in the GINplus study. Furthermore, we thank all members of the GINplus Study Group for their excellent work. The GINplus Study group consists of the following: Institute of Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg (Heinrich J, Brüske I, Schulz H, Flexeder C, Zeller C, Standl M, Schnappinger M, Ferland M, Thiering E, Tiesler C); Department of Pediatrics, Marien-Hospital, Wesel (Berdel D, von Berg A, Filipiak-Pittroff B); Ludwig-Maximilians-University of Munich, von Hauner Children's Hospital (Koletzko S, Werkstetter K); Child and Adolescent Medicine, University Hospital rechts der Isar of the Technical University Munich (Bauer CP, Hoffmann U); IUF-Environmental Health Research Institute, Düsseldorf (Schikowski T, Link E, Klümper C, Krämer U, Sugiri D). *LISA Cohort:* The authors thank all the families for their participation in the LISA study. Furthermore, we thank all members of the LISA Study Group for their excellent work. The LISA Study group consists of the following: Helmholtz Zentrum München, German Research Center for Environmental Health, Institute of Epidemiology, Munich (Heinrich J, Schnappinger M, Brüske I, Ferland M, Schulz H, Zeller C, Standl M, Thiering E, Tiesler C, Flexeder C); Department of Pediatrics, Municipal Hospital "St. Georg", Leipzig (Borte M, Diez U); Marien Hospital Wesel, Department of Pediatrics, Wesel (von Berg A, Berdel D, Stiers G, Maas B); Pediatric Practice, Bad Honnef (Schaaf B); Helmholtz Centre of Environmental Research-UFZ, Department of Environmental Immunology, Leipzig (Herberth G, Schilde M, Bauer M, Röder S); Technical University Munich, Department of Pediatrics, Munich (Hoffmann U, Paschke M,

Marra S); Clinical Research Group Molecular Dermatology, Department of Dermatology and Allergy, Technische Universität München (TUM), Munich (Ollert M, J. Grosch). For the experimental study, the authors wish to thank the animal caretakers of the Helmholtz Center Munich, Johanna Grosch and the technicians of the GMC clinical chemistry laboratory for technical assistance. We are very grateful to Dr. Annette Schuhmacher, Ssniff Spezialdiäten GmbH and to Mr. Steven Yeung, Research Diets, Inc. for excellent assistance regarding the diets used in the study.

Abstract

Background: Dietary carbohydrates and fats are intrinsically correlated within the habitual diet. We aimed to disentangle the associations of starch and sucrose from those of fat, in relation to allergic sensitization, asthma and rhinoconjunctivitis prevalence in humans, and to investigate underlying mechanisms using murine models.

Methods: Epidemiological data from participants of two German birth cohorts (age 15) were used in logistic regression analyses testing cross-sectional associations of starch and sucrose (and their main dietary sources) with aeroallergen sensitization, asthma and rhinoconjunctivitis, adjusting for correlated fats (saturated, monounsaturated, omega-6 and omega-3 polyunsaturated) and other covariates. For mechanistic insights, murine models of aeroallergen-induced allergic airway inflammation (AAI) fed with a low-fat-high-sucrose or -high-starch versus a high-fat diet were used to characterize and quantify disease development. Metabolic and physiologic parameters were used to track outcomes of dietary interventions and cellular and molecular responses to monitor the development of AAI. Oxidative stress biomarkers were measured in murine sera or lung homogenates.

Results: We demonstrate a direct association of dietary sucrose with asthma prevalence in males, while starch was associated with higher asthma prevalence in females. In mice, high-carbohydrate feeding, despite scant metabolic effects, aggravated AAI compared to high-fat in both sexes, as displayed by humoral response, mucus hypersecretion, lung inflammatory cell infiltration and T_H2 - T_H17 profiles. Compared to high-fat, high-carbohydrate intake was associated with increased pulmonary oxidative stress, signals of metabolic switch to glycolysis and decreased systemic anti-oxidative capacity.

Conclusion: High consumption of digestible carbohydrates is associated with increased prevalence of asthma in humans and aggravated lung allergic inflammation in mice, involving oxidative stress-related mechanisms.

Keywords: carbohydrates, nutrition, asthma, allergic airway inflammation, oxidative stress

Ethical approval

The GINIplus and LISA birth cohort studies were conducted in accordance with the ethical standards laid down in the Declaration of Helsinki and approved by the local ethics committees (Bavarian General Medical Council, Institutional Review Board of the University of Leipzig, Saxonian State Chamber of Physicians and Medical Council of North Rhine-Westphalia). All participants and their families gave written informed consent before study onset. Mouse experiments were conducted according to the European Convention for Animal Care and Use of Laboratory Animals and were approved by local ethics committee and government authorities (approval numbers: 55.2-1-54-2532-156-12, ROB-55.2-2532.Vet_02-14-172 and ROB-55.2-2532.Vet_02-18-94).

Conflict of interest

The authors declare no conflict of interest.

Author Contributions: Epidemiological data collection: SK, CPB, TS, DB, AvB, GH. Epidemiological study design: CPH, CF, MS and FA. Epidemiological data analysis: CPH, MS. Mouse experimental design: SM, RK, BR, JR, CBS-W, SU and FA. Conduction of experiments: SM, RK, JMG, BR, BS, ES, LM, EEV and FA. Experimental data analysis: SM, RK, JMG, BR, EEV, FA. Supervision: MS, CBS-W, SU, FA. Writing original draft: SM, CPH and FA. Review&Editing: All Authors. Funding acquisition: MHdA, MS, CBS-W, SU and FA.

Funding: *GINIplus Funding:* The GINIplus study was mainly supported for the first three years of the Federal Ministry for Education, Science, Research and Technology (interventional arm) and Helmholtz Zentrum Munich (former GSF) (observational arm). The four-year, six-year, 10-year, and 15-year follow-up examinations of the GINIplus study were covered from the respective budgets of the five study centers (Helmholtz Zentrum Munich (former GSF), Research Institute at Marien-Hospital Wesel, LMU Munich, TU Munich, and from six years onwards also from IUF – Leibniz Research-Institute for Environmental Medicine at the University of Düsseldorf), and a grant from the Federal Ministry for Environment (IUF Düsseldorf, FKZ 20462296). Further, the 15-year follow-up examination of the GINIplus study was supported by the Commission of the European Communities, the seventh Framework Program: MeDALL project, and

as well by the Mead Johnson and Nestlé companies. *LISA Funding*: The LISA study was mainly supported by grants from the Federal Ministry for Education, Science, Research, and Technology, and in addition from Helmholtz Zentrum Munich (former GSF), Helmholtz Centre for Environmental Research—UFZ, Leipzig, Research Institute at Marien-Hospital Wesel, Pediatric Practice, Bad Honnef for the first two years. The four-year, six-year, 10-year and 15-year follow-up examinations of the LISA study were covered from the respective budgets of the involved partners (Helmholtz Zentrum Munich (former GSF), Helmholtz Centre for Environmental Research—UFZ, Leipzig, Research Institute at Marien-Hospital Wesel, Pediatric Practice, Bad Honnef, IUF—Leibniz-Research Institute for Environmental Medicine at the University of Düsseldorf) and in addition by a grant from the Federal Ministry for Environment (IUF Düsseldorf, FKZ 20462296). Further, the 15-year follow-up examination of the LISA study was supported by the Commission of the European Communities, the seventh Framework Program: MeDALL project. *Mouse studies*: Mouse studies were supported by the HMGU Allergy Projects, Project #8 - G-509000-018, from the Helmholtz Zentrum München. German Mouse Clinic was supported by the German Federal Ministry of Education and Research (Infrafrontier grant 01KX1012); German Center for Diabetes Research (DZD).

Data availability statement

Due to data protection reasons, the datasets generated and/or analyzed during the current study cannot be made publicly available. The datasets are available on reasonable request, provided the release is consistent with the consent given by the GINIplus and LISA study participants and approved by the studies' steering committees. Ethical approval might be obtained for the release and a data transfer agreement from the legal department of the Helmholtz Zentrum München must be accepted. Requests should be addressed to Marie Standl (marie.standl@helmholtz-muenchen.de).

Introduction

The incidence of allergic diseases, such as rhinoconjunctivitis and asthma, has been steadily increasing in the last decades, especially in children ^{1,2}. Currently, an interaction of genetic, host, and environmental factors is discussed to be involved in asthma development ³. Westernized countries have experienced a substantial increase in asthma incidence, mainly attributed to changes in lifestyle, including diets rich in fats, added sugars, and highly-processed foods ^{4,5}. Whilst the effects of dietary fats have been widely investigated, pointing towards a functional link between obesity and asthma ⁶⁻⁸, the role of digestible carbohydrates (sugars and starches) in allergic diseases remains unclear. Addressing this evidence gap is crucial, as the two macronutrients are tightly interrelated, limiting our understanding of their isolated effects.

Indeed, prolonged high-sugar intake has been shown to be associated with obesity, especially if combined with a sedentary lifestyle ⁹. Additionally, high-sugar intake has been linked to the development of cardiovascular diseases, pointing to oxidative stress as one key mechanism in this process ¹⁰. With respect to allergic airway inflammation (AAI), recent studies have shown associations between sugar-sweetened beverage intake and asthma in both children ¹¹⁻¹⁴ and adults ^{15, 16}. However, these effects are believed to result from the excess free fructose present in such beverages ^{16,17}. The role of sucrose, a common source of added sugar in the human diet ¹⁸, calls for further research, especially considering that the already elevated consumption of sugar-rich foods has increased even further during the COVID-19 pandemic ¹⁹. Additionally, starchy foods with a high glycemic load, such as potatoes, or refined grains (processed to remove the protein- and fat-rich germ and fibre-rich bran), have also been associated with increased risk of cardiometabolic diseases, and contribute substantially more calories to the typical Western diet than do added sugars ²⁰. In addition, while reducing total dietary fat tends to be accompanied by an increased intake of carbohydrates ²¹, many high-fat products are also rich in added sugars, leading to expected correlations between starch and sucrose with fat intake. Therefore, the aims of the present study were to disentangle the effects of these two types of digestible carbohydrate from that of dietary fat on the development of aeroallergen-induced allergic disease. Therefore, we carried out epidemiological analyses among participants of two German birth cohort studies, testing the association of sucrose and starch, as well as their major dietary sources, with allergic sensitization, asthma and rhinoconjunctivitis, adjusting for dietary fat intake. Furthermore, to investigate possible underlying mechanisms, we used mouse models of aeroallergen-induced AAI and compared lung allergic response following low-fat,

high-sucrose (or high-starch) with high-fat diet feeding. Here we demonstrate that consumption of a diet rich in digestible carbohydrates is associated with higher asthma prevalence in humans and increased risk of AAI in mice. Moreover, mice on a low-fat, high-carbohydrate diet display increased pulmonary oxidative stress, signals of metabolic switch favoring glycolysis and decreased systemic anti-oxidative capacity, which may support the increased allergic response. These findings call for further dietary intervention studies investigating whether reducing the consumption of digestible carbohydrates serves to support prevention and therapy of asthma.

Methods

Epidemiological data

Data from participants of two German birth cohorts, GINIplus²² and LISA²³, were obtained through questionnaires or medical examinations carried out at the 15-year follow-up assessment. Detailed descriptions of the food frequency questionnaire used for dietary assessment, and the assessment of sensitization status, asthma, rhinoconjunctivitis and covariates, are provided in the online supplement.

Murine experimental protocols

C57BL/6J mice were fed a low-fat diet containing high-sucrose (LF-HS), high-starch (LF-HA) or a high-fat diet (HFD) for 11 weeks. Balb/c mice were fed LF-HA, HFD or a regular chow diet for 20 weeks (Table S2). To induce AAI, *Dermatophagoides farinae* (HDM)²⁴ or ragweed aqueous pollen (APE)²⁵ extracts were instilled in C57BL/6J and Balb/c mice, respectively, in the last two weeks of experiments. At sacrifice, serum was tested for total immunoglobulin E (tIgE) and metabolites, and lungs were subjected to bronchoalveolar lavage (BAL) analysis²⁵. Lung and visceral adipose tissue (VAT) were processed for flow cytometric analysis and lung tissue for additional RNA/protein isolation and histology. For methodological details see online supplement.

Statistical analysis

Epidemiological analyses in humans were performed stratified by sex using the statistical software R, version 4.4.1²⁶. Descriptive characteristics of the study population were presented as medians (25th; 75th percentile) for continuous variables, and counts (%) for categorical variables. Dietary variables were presented as their relative contribution to total energy intake (%EI). Sex-differences in population characteristics were tested by Wilcoxon rank-sum test or Pearson's chi-squared test (for continuous and categorical variables, respectively). Correlations of each carbohydrate variable with total fat, and different fat subtypes (saturated=SFA, monounsaturated=MUFA, linoleic acid=LA, and alpha-linolenic acid=ALA) were tested by Spearman's rank correlation. The association of

dietary carbohydrates, namely sucrose, sugary foods, sugary drinks, starch, starchy vegetables, and refined grains (independent variables) with allergic sensitization, asthma, and rhinoconjunctivitis (dependent variables), were assessed by logistic regression. In order to determine whether associations with asthma were mainly driven by participants with an allergic phenotype (allergic asthma), sensitivity analyses were performed in a subset of the study population with positive aeroallergen sensitization (N=715). Unadjusted analyses, then analyses adjusting for pre-selected covariates were performed. For independent variables presenting a significant correlation with dietary fats (at p-values<0.001), the respective regression model was further adjusted for the relevant fat subtypes. Effects are presented as odds ratios and 95% confidence intervals (OR (95% CI)) for an interquartile range increase in the respective independent variable. For murine studies, statistical analyses were performed by one-way or two-way analysis of variance (ANOVA) with Bonferroni's post-hoc test or by Student's unpaired two-tailed t-test. Data were analyzed using Prism software version 6 (GraphPad software Inc., La Jolla, CA, USA). Threshold of significance: P<0.05.

Results

Digestible carbohydrates are associated with increased odds of allergic sensitization and asthma

Descriptive characteristics of the study population are presented in Table 1A. Males presented significantly higher levels of allergic sensitization (50.7%) and asthma (6.9%) than females (40.9% and 4.1%, respectively). There was no sex difference in the prevalence of rhinoconjunctivitis (21.0% in females and 20.2% in males). Total sucrose intake represented around 10% of total energy intake (10.6% and 9.98% in females and males, respectively), and total starch intake around 27% (27.8% and 26.2% in females and males, respectively). Correlations between carbohydrates and fat variables are presented in Table 1B. Sugary drinks were negatively correlated with fat intake in males only, whereas all other correlations were mostly similar in both sexes. While sucrose presented significant negative correlations with MUFA and LA, sugary foods were positively correlated with SFA. Starch was negatively correlated with SFA, MUFA and ALA, and the same was observed for refined grains (except for ALA in females). Starchy vegetables were positively correlated with LA. The various associations with different fat types, each type having unique effects on health, show that simply adjusting analyses for total fat intake would not be sufficient to account for the complex interrelations between different dietary carbohydrates and fats. For example, while both starch and sucrose were inversely correlated with total fat, LA accounts largely for this association in the case of sucrose, whereas in the case of starch, the correlation is mostly due to SFA.

Associations of dietary intake variables (sucrose, sugary foods, sugary drinks, starch, starchy vegetables, and refined grains) with allergic sensitization, asthma, allergic asthma, and rhinoconjunctivitis, are displayed in Fig.1 and Table S1. A significant positive association was observed between sucrose intake and asthma in males (OR=1.68, 95% CI= (1.12;2.51), $p=0.011$), whereas no significant associations were observed with sugary foods or sugary drinks. In females, total starch was positively associated with asthma, (1.80 (1.02;3.15), $p=0.041$), and starchy vegetables were positively associated with allergic sensitization (1.25 (1.05;1.49), $p=0.013$). There were no significant associations with rhinoconjunctivitis in either sex. Sensitivity analyses assessing allergic asthma in the subset of aeroallergen sensitized participants indicated similar trends for sucrose in males (1.67 (1.02;2.75), $p=0.044$), however the OR of starch in females was reduced and not significant (1.36 (0.64;2.89), $p=0.420$). In contrast, sugary foods were significantly associated with allergic asthma in females (2.36 (1.01;5.51),

$p=0.047$). Nevertheless, given the small sample size available for sensitivity analyses and large confidence intervals, these associations must be interpreted with caution.

High-sucrose diet exacerbates murine HDM-induced AAI

To mechanistically explore the results obtained in our epidemiological analyses, we employed a model of HDM-induced AAI in male and female mice fed either with a sucrose-rich LF-HS or a HFD as a control (Fig.2A). As expected, HFD-fed males showed a stronger increase in body weight, body fat mass and weight of fat pads, and a lower tolerance to injected glucose compared to both LF-HS-fed males and to HFD-fed females (Fig.2B, Fig.S4A-C) ^{27, 28}. Accordingly, liver triglycerides, serum levels of insulin and glucose were increased in HFD-fed compared to LF-HS-fed male mice (Fig.S4D). Notably, both LF-HS and AAI had no impact on metabolic parameters compared to respective diet- and PBS-controls. The increased body weight (and fat mass) in female HFD HDM should be disregarded as it was present from begin of sensitization (Fig.2B, right). FACS analysis of VAT (Fig.S4E) revealed the expected increase of M1 macrophages in HFD-fed males ²⁹, their slight decrease following AAI in both sexes, and no diet- or treatment-induced variations in M2 macrophages, CD4⁺T or CD8⁺T cells.

In lungs AAI significantly increased BAL inflammatory cells numbers in LF-HS and HFD. Interestingly, total BAL cells and relative percentage of eosinophils were significantly higher in LF-HS compared to HFD in both sexes (Fig.2C, Fig. S5A), whereby female mice showed higher infiltrating macrophages, lymphocytes and neutrophils compared to males. Analysis of BAL T_H2, T_H17, pro-inflammatory cytokines and chemokines revealed HDM-induced increased tendencies in both sexes, reaching significance only in few parameters in LF-HS-fed animals (Fig.S5C). BAL protein concentration, a reliable parameter of increased lung barrier permeability³⁰ as expected ³¹ increased significantly following HDM challenge in both sexes (Fig. S5B). Serum tIgE, which in LF-HS-fed females was augmented compared to HFD-fed even before AAI induction, after HDM-challenge increased in both sexes, reaching significantly higher levels in LF-HS compared to HFD groups (Fig.2D). Lung histopathological analysis revealed increased perivascular and peribronchiolar inflammatory cell infiltration and mucus hypersecretion following AAI and, most importantly, both scores were significantly enhanced in LF-HS compared to HFD in both sexes (Fig.2E-I). To further characterize the type of lung immune response, we performed FACS analysis of lung tissue, which revealed a clear AAI-induced inflammatory type-2

phenotype, highlighted by a significant increase of CD4⁺T cell percentages in LF-HS HDM of both sexes, especially in females (Fig. 2J). Contrarily, CD8⁺T cells varied minimally in males, but decreased significantly in LF-HS HDM females compared to their diet and PBS controls (Fig.S3B). Whilst the percentage of GATA3⁺ cells strongly increase in in both sexes following AAI, FoxP3⁺ and RORγt⁺ cells underwent only minor variations. Furthermore, analysis of M1- and M2-BAL macrophages markers indicated an AAI-induced upregulation of both markers in LF-HS, stronger in females compared to males, whereas in HFD only *Nos2* was upregulated in females (Fig.2K).

Thus, consumption of high-sucrose diet leads to an aggravated type-2 phenotype compared to HFD in both sexes, but especially in females.

A high-starch diet exacerbates murine HDM-induced AAI

To clarify if the observed high-sucrose-induced AAI-enhancing effect applied also to starch, we compared AAI outcomes in mice fed a sucrose-rich LF-HS with mice fed a low-sucrose, high-starch (*Amylum*) (LF-HA) diet, using HFD-fed animals as reference (Fig.3A). LF-HA-fed mice gained more weight compared to LF-HS mice (Fig.3B), but no difference was observed in their glucose tolerance (Fig.3C). Serum clinical chemistry parameters did not differ between the two diets or treatment, only triglycerides and non-esterified fatty acids (NEFA) were slightly higher in LF-HA compared to LF-HS (Fig.S6A). Similarly, no difference in macrophage or T cell populations was detected between LF-HA and LF-HS in fat tissue (Fig.S6B). As for the lung response to AAI, no difference in BAL cell infiltration (Fig.3D), histopathological alterations and scores (Fig.3E-I), serum tIgE (Fig.3K), BAL protein and cytokines (Fig.S6C) was detected between LF-HA and LF-HS, whereby most of these parameters were strongly increased in high-carbohydrate groups compared to HFD. Lung FACS analysis revealed an AAI-induced T_H2-driven response similar in LF-HA and LF-HS whereas the T_H17 response was higher in LF-HA compared to HFD and to LF-HS, reflected also by higher IL-17A levels in BAL supernatant in this group (Fig.3L, S6C).

Herewith, we could demonstrate that a diet rich in starch potentiates a HDM-induced allergic response similarly to sucrose, but favoring mixed T_H2-T_H17 profiles.

Effects of a high-starch diet are not restricted to a specific allergen

To evaluate the allergen specificity of our findings, we employed an aero-allergen other than HDM in an AAI model previously established with ragweed-APE in female BALB/c mice²⁵ at the end of a 20-week feeding period with LF-HA or HFD as a control (Fig.4A). Whereas at week 7 LF-HA-fed mice showed, additionally to a lower body weight, higher glucose tolerance compared to HFD (Fig.S7A,C), at week 13 also mice fed LF-HA developed glucose intolerance, maintaining a lower body weight compared to HFD (Fig.S7B-C). Accordingly, perigonadal fat pads were significantly increased in sham-sensitized HFD-compared to LF-HA-fed mice (Fig.S7D-E). Additionally, serum cholesterol, triglyceride, NEFA and glucose were higher in HFD compared to LF-HA (Fig.S7F). AAI induction had no impact on the metabolic parameters evaluated.

APE-challenge increased lung inflammatory cell infiltration of both LF-HA- and HFD-fed mice, dominated by eosinophils and lymphocytes. Total BAL cells and eosinophils showed only a higher tendency in LF-HA compared to HFD (Fig.4B, above), BAL total protein was similar in the two challenged groups whereas the Th2 cytokines IL-4 and IL-13 were significantly higher in LF-HA- compared to HFD-fed animals, together with no IFN- γ alterations (Fig.4B, below). Also histopathological scoring of lungs revealed increased perivascular and peribronchiolar inflammatory cell infiltration and mucus hypersecretion in allergic LF-HA compared to HFD (Fig.4C-G). Additionally, BAL macrophages retrieved from LF-HA APE showed a significant upregulation of *Nos2*, and to a higher extent of *Arg1* and *Ccl17* (Fig.4H). The expression of both *Nos2* and *Ccl17* was higher in LF-HA compared to HFD, whereas for *Retn1a* no diet-related difference was shown. Lastly, LF-HA displayed a near-to-significantly stronger increase in serum tIgE compared to HFD (Fig.4I).

Thus, although the overall allergic response in the APE model was lower compared to HDM, we could confirm that high amounts of digestible carbohydrates exacerbates features of AAI in a non-allergen-specific manner.

High-carbohydrate diet increases pulmonary and systemic oxidative stress and induces a metabolic switch to glycolysis

To obtain insights into the mechanism of how carbohydrate-rich diets can aggravate AAI, and considering the important role of oxidative stress in AAI in both mice and humans^{32,33} we measured key indicators of oxidative stress in lung tissue and serum. Our results reveal an allergen-induced increased expression of

Gpx in LF-HA and of *Sod1* and *Sod2* in LF-HS compared to HFD (Fig.5A). On the same line, 8-isoprostane, a key marker of oxidative stress ³⁴, was increased in lung homogenates after challenge in both LF-HS and LF-HA and not in HFD. Noticeably, after challenge this marker was near-to-significantly increased in LF-HS and LF-HA compared to HFD (Fig.5B). Next, we investigated whether a glycolytic reprogramming may play a role in our model. The lung expression of *Sirtuin 1*, *HIF-1 α* and *Mct4*, important for sensing hypoxia and promoting glycolysis ^{35,36}, were higher in allergen-challenged LF-HS and LF-HA compared to HFD, although for *HIF-1 α* the difference was not significant. Contrarily, lung expression of *PGC1 α* , an indicator of mitochondrial oxidative metabolism ³⁷, was slightly downregulated in allergen-challenged LF-HS and LF-HA compared to PBS controls (Fig.5C). To investigate potential systemic effects of the increased pulmonary oxidative stress, we quantified serum anti-oxidative capacity in both C57BL/6J (LF-HS, Fig.5D) and Balb/c (LF-HA, Fig.5E) compared to respective HFD groups. Whilst peroxy radical scavenging capacity (trolox equivalents) was not altered by diet, nitrite concentrations were significantly lower in both LF-HS and LF-HA compared to HFD. Additionally, total phenolic antioxidant content (gallic acid equivalents, GAE) was significantly lower in LF-HA- compared to HFD-fed Balb/c animals, whereas no difference was reported between LF-HS- and HFD-fed C57BL/6J mice. Taken together, these results suggest that high levels of dietary carbohydrates induce augmented lung oxidative stress, signals of metabolic switch to glycolysis combined to systemic decreased anti-oxidative capacity compared to HFD.

Discussion

This study uses epidemiological data from two population-based cohorts of female and male adolescents to evaluate associations of different types of digestible carbohydrates with aeroallergen sensitization, asthma and rhinoconjunctivitis, and to investigate underlying mechanisms using murine models.

The observational study demonstrates associations between higher consumption of starch and sucrose and greater asthma prevalence in females and males, respectively. Given the inclusion of correlated dietary fats as covariates, the associations observed appear to be independent of inherent variations in fat intake. According to our findings in males, sucrose intake increases the risk for asthma. Except one study reporting increased atopic respiratory outcomes in the offspring of mothers consuming more sugar during pregnancy ¹¹, existing literature linking sugar intake to asthma focuses on consumption of sugary drinks ^{12, 15, 16, 38, 39}. In contrast, we observed no associations of asthma with sugary drinks. The low amount of sugary drink consumption (0.8 %EI in females and 2.8 %EI in males) and lower relative fructose content in drinks produced in Europe compared to the US ⁴⁰ may account for this discrepancy. Furthermore, the weak correlation between sugary drinks and total sucrose in our male population ($\rho=0.09$) shows that sugary drinks were not major contributors towards total sucrose intake. Additionally, we demonstrate that, among female adolescents, higher total starch intake is associated with asthma, and a similar non-significant trend is seen for refined grains (both are highly correlated, $\rho=0.83$). Sensitivity analysis among a subset of aeroallergen sensitized individuals indicated a smaller effect magnitude, suggesting that dietary starch increases the odds of having asthma irrespective of allergic sensitization. Nevertheless, given the reduced power due to the smaller sample size, this should be interpreted with caution. Our findings on total starch differ from those reported in an ecological study of asthma and allergies in 13-14 year-olds ⁴¹. However, the analyses in that study were not adjusted for positively-correlated protective foods such as fruit, vegetables, fibers, and whole grains, nor for inversely correlated fats. Despite accounting for numerous confounding factors in the present study, including known dietary, lifestyle, socioeconomic and anthropometric risk factors of allergic disease development, the possibility of residual confounding through other uncontrolled influences remains a limitation common to all epidemiological studies. Furthermore, the cross-sectional nature of our analysis reduces our ability to establish causal associations. Firstly, we cannot exclude the possibility of reverse causation, whereby symptomatic individuals might have purposefully altered the composition of their diet. Furthermore, dietary behaviors may change in adolescence due to physiological demands resulting from pubertal development, as well as increasing

autonomy. Nevertheless, when comparing average carbohydrate intakes among a subset of the study population who also had available dietary data at age 10 years (shown in Table S5), we observed only small changes over time ($\leq 1.5\%$ EI). In previous work addressing changes in diet from ages 10 to 15 years in the GINIplus cohort, we also found that although average intake changes occurred overall, intake levels relative to the rest of the study population often remained constant⁴². Interestingly, and in contrast to total starch, starchy vegetables were associated with allergic sensitization in females of our study. Indeed, prevalence of sensitization to potato has been reported at around 10%⁴³, and sensitization to cooked potatoes has been considered a risk factor for early development of pollen allergy⁴⁴. Overall, sex-specific relationships were observed in our epidemiological data, but not in the rodent models. These sex differences are difficult to explain yet not unexpected, given the age of the cohort participants, and the sex-specific trajectories of allergic diseases evident around the time of puberty, such as the known switch in asthma prevalence from male to female dominance⁴⁵. Different trajectories of different allergic diseases would also explain the lack of consistent associations with other outcomes, since for example, asthma prevalence increases around late childhood, while rhinoconjunctivitis prevalence increases steadily over time⁴⁶. Besides offering the above reflections, we cannot currently explain why in our cohorts total dietary starch is associated with asthma only in females. Given the lack of studies specifically assessing the role of starch in relation to asthma, there is little comparable evidence. Particularly, given the prominence of starch in the Western diet, this finding would have important public health implications. Nevertheless, further studies, including human intervention trials, are needed to confirm these findings before any definitive conclusions can be reached. The observation that sucrose was associated with asthma more clearly in males could be due to sex differences concerning glucose handling, as well as body fat storage and distribution, altering effects on systemic metabolism. In this context, males are more sensitive to sucrose and fructose compared to females, who store more lipids and have higher whole-body insulin sensitivity than males⁴⁷. On the other hand, it is also possible that females and males simply differ in the type of foods regularly consumed, the details of which may not be fully captured in the epidemiological data, and which obviously cannot be mimicked in mice experiments. Overall, we should also consider that the underrepresentation of participants from lower social classes constitutes a non-random loss-to-follow-up, which may limit the generalizability of our findings.

Rodent models of antigen-driven allergy are successfully used in mechanistic studies for preclinical research, mimicking specific aspects of human disease⁴⁸. To disentangle the effects of sucrose and starch from that of dietary fat on AAI development, we induced aeroallergen-induced AAI combined with LF-HS or LF-HA diets, and used HFD as a control, despite existing literature demonstrating controversial effects of HFD on allergies^{24, 49, 50}. Notably, by using HFD as a control we intended to avoid non-digestible carbohydrates contained in regular chow diets⁵¹. Nevertheless, we controlled for response in main allergic parameters following APE challenge and depicted no difference between HFD- and regular chow diet-fed animals. Our results show that high dietary sucrose and starch enhance typical T_H2 - or T_H2 - T_H17 -driven lung allergic inflammatory milieu, respectively^{48, 52, 53}, both comprising increased inflammatory infiltrate, mucus hypersecretion and serum tIgE levels compared to HFD. These data are in line with a recent study using an *Alternaria alternata* allergy model which demonstrates increased BAL and eosinophil cytokine content in high-sucrose- compared to high-fat-fed animals⁵⁴. High dietary starch, besides enhancing an HDM-driven T_H17 response compared to high sucrose, affected glucose tolerance, but only after longer-term feeding. Notably, starch possesses a high glycaemic index, which can probably explain both inflammatory and metabolic effects⁵⁵. Whereas male mice showed the anticipated higher susceptibility to high-fat, but not to high-carbohydrate feeding, female mice confirmed their higher susceptibility to allergy^{27, 28, 56, 57}. Consequently, in female mice, both M1 and M2 alveolar macrophages markers significantly increased following AAI in both sensitization models, indicating the presence of a mixed M1/M2 phenotype in the high-carbohydrate groups⁵⁸, consistently with the critical role of both phenotypes in asthma development⁵⁹. To shed light on the mechanism responsible for disease aggravation, we further investigated which source of excessive energy supply (i.e. sucrose, starch or fat) may lead to an increased oxidative activity that, along with insufficient antioxidant defense, causes local or systemic overproduction of reactive oxygen species (ROS)^{60, 61}. Increased expression of *Gpx*, *Sod1* and *Sod2* in mouse lungs following high-carbohydrate-feeding indicates increased activation of the antioxidant defense and thus, enhanced lung oxidative challenge compared to HFD-feeding. This was confirmed by the measurements of 8-isoprostane, which clearly demonstrate increased oxidative stress in lung tissue of high-carbohydrate- compared to HFD-fed animals. Notably, under hypoxic or inflammatory conditions, HIF-1 α activity can trigger a shift from oxidative phosphorylation to glycolysis, thus supporting immune cell activation and inflammation under high cost of glucose consumption (Warburg effect)^{62, 63}. The slight upregulation of HIF-1 α combined to the downregulation of PGC1 α , involved in Krebs-Cycle induction and mitochondrial ROS detoxification^{37, 64} point to

a glycolytic reprogramming in the lungs of LF-HS- and LF-HA-fed animals. Furthermore, the increased expression of both *Sirtuin1* and *Mct4* in LF-HS- compared to HFD further indicate the occurrence of a metabolic switch to glycolysis in lung tissue with consecutive lactate production, a process known to be important in allergic asthma ⁶⁵⁻⁶⁸. Furthermore, LF-HA-fed Balb/c mice sera revealed lower GAE, applied as a sum parameter of anti-oxidative capacity ⁶⁹. Contrarily, GAE in C57BL/6J mice was very high concentrated independently of diets. This likely represents a side-effect caused by artificial colorants present in the specific diet, some of which possess anti-oxidative capacities ⁷⁰. Conversely, nitrite concentrations were lower in sera of both LF-HS- and LF-HA-fed compared to HFD-fed mice, independently of the mouse strain. Noteworthy, food nitrate/nitrite-derived nitric oxide (NO) is a readily bioavailable supplementation of endogenous NO, which possesses protective properties exerted through the attenuation of oxidative stress in immune cells ^{71, 72} and may therefore play an important role in our model. Summing up, a systemic reduction of anti-oxidative capacity in high-carbohydrate-fed mice may increase glycolytic reprogramming and susceptibility to oxidative stress, promoting disease exacerbation.

Taken together, our study demonstrates for the first time a direct association of dietary sucrose and starch with asthma in humans, as well as the harmful effect of a diet rich in these carbohydrates on AAI development in mice. Following confirmation in human intervention trials, these findings could have important implications for dietary recommendations in the prevention and/or treatment of asthma.

References

1. Lundback B, Backman H, Lotvall J, Ronmark E. Is asthma prevalence still increasing? *Expert Rev Respir Med* 2016;10(1):39-51. doi:10.1586/17476348.2016.1114417
2. Asher MI, Montefort S, Bjorksten B, et al. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet* 2006;368(9537):733-43. doi:10.1016/S0140-6736(06)69283-0
3. Beasley R, Semprini A, Mitchell EA. Risk factors for asthma: is prevention possible? *Lancet* 2015;386(9998):1075-85. doi:10.1016/S0140-6736(15)00156-7
4. Barros R, Moreira A, Padrao P, et al. Dietary patterns and asthma prevalence, incidence and control. *Clin Exp Allergy* 2015;45(11):1673-80. doi:10.1111/cea.12544
5. Julia V, Macia L, Dombrowicz D. The impact of diet on asthma and allergic diseases. *Nat Rev Immunol* 2015;15(5):308-22. doi:10.1038/nri3830
6. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *Am J Respir Crit Care Med* 2007;175(7):661-6. doi:10.1164/rccm.200611-1717OC
7. Wood LG. Diet, Obesity, and Asthma. *Ann Am Thorac Soc* 2017;14(Supplement_5):S332-S338. doi:10.1513/AnnalsATS.201702-124AW
8. Ronmark E, Andersson C, Nystrom L, Forsberg B, Jarvholm B, Lundback B. Obesity increases the risk of incident asthma among adults. *Eur Respir J* 2005;25(2):282-8. doi:10.1183/09031936.05.00054304
9. Siervo M, Montagnese C, Mathers JC, Soroka KR, Stephan BC, Wells JC. Sugar consumption and global prevalence of obesity and hypertension: an ecological analysis. *Public Health Nutr* 2014;17(3):587-96. doi:10.1017/S1368980013000141
10. Prasad K, Dhar I. Oxidative stress as a mechanism of added sugar-induced cardiovascular disease. *Int J Angiol* 2014;23(4):217-26. doi:10.1055/s-0034-1387169
11. Bedard A, Northstone K, Henderson AJ, Shaheen SO. Maternal intake of sugar during pregnancy and childhood respiratory and atopic outcomes. *Eur Respir J* 2017;50:1700073. doi:10.1183/13993003.00073-2017

12. Padilha LL, Vianna EO, Vale ATM, Nascimento J, da Silva AAM, Ribeiro CCC. Pathways in the association between sugar sweetened beverages and child asthma traits in the 2nd year of life: Findings from the BRISA cohort. *Pediatr Allergy Immunol* 2020;31(5):480-488. doi:10.1111/pai.13243
13. Berentzen NE, van Stokkom VL, Gehring U, et al. Associations of sugar-containing beverages with asthma prevalence in 11-year-old children: the PIAMA birth cohort. *Eur J Clin Nutr* 2015;69(3):303-8. doi:10.1038/ejcn.2014.153
14. Yu R, Yang B, Cai L, Lu X, Wang X. Excess Free Fructose Beverages and Allergy in Children and Adolescents: Results From NHANES 2005-2006. *Ann Fam Med* 2018;16(5):408-418. doi:10.1370/afm.2292
15. Park S, Akinbami LJ, McGuire LC, Blanck HM. Association of sugar-sweetened beverage intake frequency and asthma among U.S. adults, 2013. *Prev Med* 2016;91:58-61. doi:10.1016/j.ypmed.2016.08.004
16. DeChristopher LR, Tucker KL. Excess free fructose, high-fructose corn syrup and adult asthma: the Framingham Offspring Cohort. *Br J Nutr* 2018;119(10):1157-1167. doi:10.1017/S0007114518000417
17. DeChristopher LR, Uribarri J, Tucker KL. The link between soda intake and asthma: science points to the high-fructose corn syrup, not the preservatives: a commentary. *Nutr Diabetes* 28 2016;6(11):e234. doi:10.1038/nutd.2016.46
18. Rippe JM, Sievenpiper JL, Le KA, White JS, Clemens R, Angelopoulos TJ. What is the appropriate upper limit for added sugars consumption? *Nutr Rev* 2017;75(1):18-36. doi:10.1093/nutrit/nuw046
19. Mediouni M, Madiouni R, Kaczor-Urbanowicz KE. COVID-19: How the quarantine could lead to the depreobesity. *Obes Med* 2020;19:100255. doi:10.1016/j.obmed.2020.100255
20. Ludwig DS, Hu FB, Tappy L, Brand-Miller J. Dietary carbohydrates: role of quality and quantity in chronic disease. *BMJ* 2018;361:k2340. doi:10.1136/bmj.k2340
21. Lieberman HR, Fulgoni VL, Agarwal S, Pasiakos SM, Berryman CE. Protein intake is more stable than carbohydrate or fat intake across various US demographic groups and international populations. *Am J Clin Nutr* 2020;112(1):180-186. doi:10.1093/ajcn/nqaa044
22. Berg A, Kramer U, Link E, et al. Impact of early feeding on childhood eczema: development after nutritional intervention compared with the natural course - the GINIplus study up to the age of 6 years. *Clin Exp Allergy* 2010;40(4):627-36. doi:10.1111/j.1365-2222.2009.03444.x

23. Heinrich J, Bolte G, Holscher B, et al. Allergens and endotoxin on mothers' mattresses and total immunoglobulin E in cord blood of neonates. *Eur Respir J* 2002;20(3):617-23. doi:10.1183/09031936.02.02322001
24. Everaere L, Ait-Yahia S, Molendi-Coste O, et al. Innate lymphoid cells contribute to allergic airway disease exacerbation by obesity. *J Allergy Clin Immunol* 2016;138(5):1309-1318. doi:10.1016/j.jaci.2016.03.019
25. Wimmer M, Alessandrini F, Gilles S, et al. Pollen-derived adenosine is a necessary cofactor for ragweed allergy. *Allergy* 2015;70(8):944-54. doi:10.1111/all.12642
26. R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
27. Hwang LL, Wang CH, Li TL, et al. Sex differences in high-fat diet-induced obesity, metabolic alterations and learning, and synaptic plasticity deficits in mice. *Obesity (Silver Spring)* 2010;18(3):463-9. doi:10.1038/oby.2009.273
28. Nishikawa S, Yasoshima A, Doi K, Nakayama H, Uetsuka K. Involvement of sex, strain and age factors in high fat diet-induced obesity in C57BL/6J and BALB/cA mice. *Exp Anim* 2007;56(4):263-72. doi:10.1538/expanim.56.263
29. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest* 2003;112(12):1796-808. doi:10.1172/JCI19246
30. Carlier FM, de Fays C, Pilette C. Epithelial Barrier Dysfunction in Chronic Respiratory Diseases. *Front Physiol* 2021;12:691227. doi:10.3389/fphys.2021.691227
31. Herbert CA, King CM, Ring PC, et al. Augmentation of permeability in the bronchial epithelium by the house dust mite allergen Der p1. *Am J Respir Cell Mol Biol* 1995;12(4):369-78. doi:10.1165/ajrcmb.12.4.7695916
32. Park CS, Kim TB, Lee KY, et al. Increased oxidative stress in the airway and development of allergic inflammation in a mouse model of asthma. *Ann Allergy Asthma Immunol* 2009;103(3):238-47. doi:10.1016/S1081-1206(10)60188-3
33. Andrianjafimasy M, Zerimech F, Akiki Z, et al. Oxidative stress biomarkers and asthma characteristics in adults of the EGEA study. *Eur Respir J* 2017;50(6) doi:10.1183/13993003.01193-2017
34. Alessandrini F, Beck-Speier I, Krappmann D, et al. Role of oxidative stress in ultrafine particle-induced exacerbation of allergic lung inflammation. *Am J Respir Crit Care Med* 2009;179(11):984-91. doi:10.1164/rccm.200807-1061OC

35. Lim JH, Lee YM, Chun YS, Chen J, Kim JE, Park JW. Sirtuin 1 modulates cellular responses to hypoxia by deacetylating hypoxia-inducible factor 1alpha. *Mol Cell* 2010;38(6):864-78. doi:10.1016/j.molcel.2010.05.023
36. Tan Z, Xie N, Banerjee S, et al. The monocarboxylate transporter 4 is required for glycolytic reprogramming and inflammatory response in macrophages. *J Biol Chem* 2015;290(1):46-55. doi:10.1074/jbc.M114.603589
37. Scarpulla RC. Metabolic control of mitochondrial biogenesis through the PGC-1 family regulatory network. *Biochim Biophys Acta* 2011;1813(7):1269-78. doi:10.1016/j.bbamcr.2010.09.019
38. Padilha LL, Vianna EO, Vale ATM, Nascimento J, da Silva AAM, Ribeiro CCC. Pathways in the association between sugar sweetened beverages and child asthma traits in the 2nd year of life: findings from the BRISA Cohort. *Pediatr Allergy Immuno* 2020; 31(5):480-488. doi:10.1111/pai.13243
39. Cisneros R, Gonzalez M, Brown P, Schweizer D. Soda consumption and hospital admissions among Californian adults with asthma. *J Asthma* 2017;54(4):371-375. doi:10.1080/02770903.2016.1218014
40. Varsamis P, Larsen RN, Dunstan DW, Jennings GL, Owen N, Kingwell BA. The sugar content of soft drinks in Australia, Europe and the United States. *Med J Aust* 2017;206(10):454-455. doi:10.5694/mja16.01316
41. Ellwood P, Asher MI, Bjorksten B, Burr M, Pearce N, Robertson CF. Diet and asthma, allergic rhinoconjunctivitis and atopic eczema symptom prevalence: an ecological analysis of the International Study of Asthma and Allergies in Childhood (ISAAC) data. ISAAC Phase One Study Group. *Eur Respir J* 2001;17(3):436-43. doi:10.1183/09031936.01.17304360
42. Harris C, Flexeder C, Thiering E, et al. Changes in dietary intake during puberty and their determinants: results from the GINIplus birth cohort study. *BMC Public Health* 2015;15:841. doi:10.1186/s12889-015-2189-0
43. Chiriac AM, Bourrain JL, Lopicard E, Molinari N, Demoly P. Prevalence of sensitization and allergy to potato in a large population. *J Allergy Clin Immunol Pract* 2017;5(2):507-509. doi:10.1016/j.jaip.2016.12.011
44. De Swert LF, Cadot P, Ceuppens JL. Diagnosis and natural course of allergy to cooked potatoes in children. *Allergy* 2007;62(7):750-7. doi:10.1111/j.1398-9995.2007.01368.x
45. Keller T, Hohmann C, Standl M, et al. The sex-shift in single disease and multimorbid asthma and rhinitis during puberty - a study by MeDALL. *Allergy* 2018;73(3):602-614. doi:10.1111/all.13312

46. Nissen SP, Kjaer HF, Host A, Nielsen J, Halken S. The natural course of sensitization and allergic diseases from childhood to adulthood. *Pediatr Allergy Immunol* 2013;24(6):549-55. doi:10.1111/pai.12108
47. Varlamov O, Bethea CL, Roberts CT, Jr. Sex-specific differences in lipid and glucose metabolism. *Front Endocrinol (Lausanne)* 2014;5:241. doi:10.3389/fendo.2014.00241
48. Alessandrini F, Musiol S, Schneider E, Blanco-Perez F, Albrecht M. Mimicking Antigen-Driven Asthma in Rodent Models-How Close Can We Get? *Front Immunol* 2020;11:575936. doi:10.3389/fimmu.2020.575936
49. Silva FMC, Oliveira EE, Gouveia ACC, et al. Obesity promotes prolonged ovalbumin-induced airway inflammation modulating T helper type 1 (Th1), Th2 and Th17 immune responses in BALB/c mice. *Clin Exp Immunol* 2017;189(1):47-59. doi:10.1111/cei.12958
50. Schroder T, Wiese AV, Ender F, et al. Short-term high-fat diet feeding protects from the development of experimental allergic asthma in mice. *Clin Exp Allergy* 2019;49(9):1245-1257. doi:10.1111/cea.13454
51. Pellizzon MA, Ricci MR. Correction to: The common use of improper control diets in diet-induced metabolic disease research confounds data interpretation: the fiber factor. *Nutr Metab (Lond)* 2018;15:23. doi:10.1186/s12986-018-0256-0
52. Yu QL, Chen Z. Establishment of different experimental asthma models in mice. *Exp Ther Med* 2018;15(3):2492-2498. doi:10.3892/etm.2018.5721
53. Chesne J, Braza F, Chadeuf G, et al. Prime role of IL-17A in neutrophilia and airway smooth muscle contraction in a house dust mite-induced allergic asthma model. *J Allergy Clin Immunol* 2015;135(6):1643-1645. doi:10.1016/j.jaci.2014.12.1872
54. Percopo CM, McCullough M, Limkar AR, Druey KM, Rosenberg HF. Impact of controlled high-sucrose and high-fat diets on eosinophil recruitment and cytokine content in allergen-challenged mice. *PLoS One* 2021;16(8):e0255997. doi:10.1371/journal.pone.0255997
55. Campbell GJ, Belobrajdic DP, Bell-Anderson KS. Determining the Glycaemic Index of Standard and High-Sugar Rodent Diets in C57BL/6 Mice. *Nutrients* 2018;10(7):856. doi:10.3390/nu10070856
56. Melgert BN, Postma DS, Kuipers I, et al. Female mice are more susceptible to the development of allergic airway inflammation than male mice. *Clin Exp Allergy* 2005;35(11):1496-503. doi:10.1111/j.1365-2222.2005.02362.x

57. Melgert BN, Oriss TB, Qi Z, et al. Macrophages: regulators of sex differences in asthma? *Am J Respir Cell Mol Biol* 2010;42(5):595-603. doi:10.1165/rcmb.2009-0016OC
58. Saradna A, Do DC, Kumar S, Fu QL, Gao P. Macrophage polarization and allergic asthma. *Transl Res* 2018;191:1-14. doi:10.1016/j.trsl.2017.09.002
59. Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000Prime Rep* 2014;6:13. doi:10.12703/P6-13
60. Bhatti JS, Bhatti GK, Reddy PH. Mitochondrial dysfunction and oxidative stress in metabolic disorders - A step towards mitochondria based therapeutic strategies. *Biochim Biophys Acta Mol Basis Dis* 2017;1863(5):1066-1077. doi:10.1016/j.bbadis.2016.11.010
61. Bonomini F, Rodella LF, Rezzani R. Metabolic syndrome, aging and involvement of oxidative stress. *Aging Dis* 2015;6(2):109-20. doi:10.14336/AD.2014.0305
62. Warburg O. On the origin of cancer cells. *Science* 1956;123(3191):309-14. doi:10.1126/science.123.3191.309
63. Kornberg MD. The immunologic Warburg effect: Evidence and therapeutic opportunities in autoimmunity. *Wiley Interdiscip Rev Syst Biol Med* 2020;12(5):e1486. doi:10.1002/wsbm.1486
64. Schonenberger MJ, Kovacs WJ. Hypoxia signaling pathways: modulators of oxygen-related organelles. *Front Cell Dev Biol* 2015;3:42. doi:10.3389/fcell.2015.00042
65. Rodgers JT, Lerin C, Haas W, Gygi SP, Spiegelman BM, Puigserver P. Nutrient control of glucose homeostasis through a complex of PGC-1alpha and SIRT1. *Nature* 2005;434(7029):113-8. doi:10.1038/nature03354
66. Ullah MS, Davies AJ, Halestrap AP. The plasma membrane lactate transporter MCT4, but not MCT1, is up-regulated by hypoxia through a HIF-1alpha-dependent mechanism. *J Biol Chem* 2006;281(14):9030-7. doi:10.1074/jbc.M511397200
67. Ostroukhova M, Goplen N, Karim MZ, et al. The role of low-level lactate production in airway inflammation in asthma. *Am J Physiol Lung Cell Mol Physiol* 2012;302(3):L300-7. doi:10.1152/ajplung.00221.2011
68. Kim SR, Lee KS, Park SJ, et al. Involvement of sirtuin 1 in airway inflammation and hyperresponsiveness of allergic airway disease. *J Allergy Clin Immunol* 2010;125(2):449-460. doi:10.1016/j.jaci.2009.08.009

69. Ainsworth EA, Gillespie KM. Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin-Ciocalteu reagent. *Nat Protoc* 2007;2(4):875-7. doi:10.1038/nprot.2007.102
70. Gostner JM, Becker K, Ueberall F, Fuchs D. The good and bad of antioxidant foods: An immunological perspective. *Food Chem Toxicol* 2015;80:72-79. doi:10.1016/j.fct.2015.02.012
71. Lundberg JO, Gladwin MT, Ahluwalia A, et al. Nitrate and nitrite in biology, nutrition and therapeutics. *Nat Chem Biol* 2009;5(12):865-9. doi:10.1038/nchembio.260
72. Yang T, Peleli M, Zollbrecht C, et al. Inorganic nitrite attenuates NADPH oxidase-derived superoxide generation in activated macrophages via a nitric oxide-dependent mechanism. *Free Radic Biol Med* 2015;83:159-66. doi:10.1016/j.freeradbiomed.2015.02.016

Figure legends

Fig. 1: Association of dietary digestible carbohydrates with allergic sensitization, asthma and rhinoconjunctivitis. Odds ratio (OR) and 95% confidence intervals (95%CI) presented for an interquartile range increase in contribution to total daily energy intake. Models adjusted for study (GINIplus intervention/observation arm/LISA), study region (Munich, Wesel, Bad Honnef, Leipzig), sex (female/male), age (years), body mass index (kg/m²), parental education (high >10 years / med-low ≤10 years), parental allergy (one or both parents reported ever having an atopic disease), moderate-to-vigorous physical activity (min/day), fruit and vegetable intake (%EI), whole grain intake (%EI), total energy intake (kcal) and fat intake (%EI, only fat subtypes presenting a significant correlation with the independent variable). * p<0.05.

Fig. 2: Impact of LF-HS versus HFD feeding in the development of AAI in mice. Male and female C57BL/6 mice were fed either a low fat-high sucrose (LF-HS) or a high-fat (HFD) diet for 11 weeks. After 9 weeks, mice were sensitized and challenged with HDM or PBS. **(A)** Treatment scheme. **(B)** Change of body weight. **(C)** Total and relative percentages of BAL cells (left, male; right, female). **(D)** Serum total IgE (tIgE). **(E-H)** Representative periodic acid–Schiff (PAS)-stained lung sections of female mice. Arrows: inflammatory infiltrate; arrowheads: mucus hypersecretion; scale bar: 50µm. **(I)** Histological scoring. **(J)** Flow cytometric analysis of lung tissue. **(K)** M1- and M2-markers of alveolar macrophages analyzed by real-time PCR. n=8/group/sex (Fig. 2B, n=16/group until week 9; Fig. 3I and 3J, n=5/group/sex); data expressed as boxplots indicating minimum, 25th percentile, median, 75th percentile, and maximum or as mean±SEM. (B, D) 2way- and (C,I-K) 1way-ANOVA with Bonferroni test. *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001 LF-HS PBS vs HFD PBS or as depicted; ##p<0.01; ###p<0.001 LF-HS HDM vs HFD HDM; ++p<0.01; +++p<0.0001 LF-HS HDM vs LF-HS PBS; °°°p<0.001 HFD PBS vs HFD HDM; ssss p<0.0001 HFD female vs HFD male; §§p<0.01; §§§p<0.001 LF-HS female vs LF-HS male.

Fig. 3: Impact of different carbohydrates on the development of AAI. Male C57BL/6 mice were fed either a high-sucrose/no-starch (LF-HS) or a low-sucrose/high-starch (LF-HA) low-fat diet or a HFD for 10 weeks. At week 8 mice were sensitized and challenged with HDM or PBS. Grey dotted lines: response

in HFD-HDM mice as reference. **(A)** Treatment scheme. **(B)** Change of body weight. **(C)** Glucose tolerance test (GTT) before sacrifice. **(D)** Total (left) and relative (right) percentages of BAL cells. **(E-H)** Representative PAS-stained lung sections. Arrows: inflammatory infiltrate; arrowheads: mucus hypersecretion; scale bar: 50µm. **(I)** Histological scoring. **(K)** Serum total IgE (tIgE). **(L)** Flow cytometric analysis of lung tissue. n=8/group (Fig. 3B, n=16/group until week 8; Fig. 3I, n=5/group; Fig. 3L, n=6/group); data are expressed as boxplots indicating minimum, 25th percentile, median, 75th percentile, and maximum or as mean±SEM. (B, C, K) 2way- and (D, I, L) 1way-ANOVA with Bonferroni test. *p<0.05;**p<0.01;***p<0.001;****p<0.0001 as indicated; ####p<0.0001 vs tIgE day0; +p<0.05; ++p<0.01; +++p<0.001; ++++p<0.0001 vs HFD HDM.

Fig. 4: Analysis of ragweed APE-induced AAI in mice fed with LF-HA versus HFD diet. Female Balb/c mice were fed with LF-HA or HFD for 20 weeks and sensitized to ragweed pollen extract (APE) or PBS last 2 weeks of experiment. **(A)** Treatment scheme. **(B)** Total BAL cells (upper panel); Th1/Th2 cytokine levels in BAL fluid (lower panel). **(C)** Histological scoring of PAS-stained lung sections as represented in **(D-G)**; arrows: inflammatory infiltrates; arrowheads: mucus hypersecretion; scale bar: 100µm. **(H)** M1- and M2-markers of alveolar macrophages analyzed by real-time PCR. **(I)** Serum total IgE (tIgE). n=5 mice/group; data represent one out of two independent experiments and are displayed as boxplots indicating minimum, 25th percentile, median, 75th percentile, and maximum or as mean±SEM. (I) 2way- and (B-H) 1way-ANOVA with Bonferroni test. *p<0.05;**p<0.01;***p<0.001;****p<0.0001.

Fig. 5: Analysis of markers of pulmonary oxidative stress, glycolysis & oxidative phosphorylation and serum antioxidant capacity. **(A)** Enzymes involved in antioxidant defense, glutathione peroxidase (Gpx), superoxide dismutase 1 and 2 (Sod 1 and 2) in lung tissue analyzed by real-time PCR. **(B)** 8-isoprostane concentrations in lung tissue. **(C)** Key enzymes involved in glycolytic reprogramming, Sirtuin 1, hypoxia-inducible factor-1α (*Hif-1α*) and monocarboxylate transporter (*Mct4*) and of an indicator of mitochondrial oxidative metabolism (*PGC1α*) in lung tissue analyzed by real-time PCR. **(D)** and **(E)** Measurement of gallic acid equivalents, trolox equivalents and nitrite in mouse serum of C57/BL6 (D) and Balb/c (E) mice after indicated treatment. Since no difference was


detected due to sensitization, for D-E data from allergic and non-allergic mice were pooled. A, C: n=6-13/group; B: n=7-8/group; D: n=16/group; E: n=10/group. Data are expressed as boxplots indicating minimum, 25th percentile, median, 75th percentile, and maximum or mean±SEM. (A-C) 1way-ANOVA with Bonferroni test, (D-E) Student's unpaired two-tailed t-test. *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001.

Table 1


A: Study population descriptive characteristics				B: Correlation of dietary carbohydrates with dietary fats									
	Females (N=830)	Males (N=743)	P-value	Females (N=830)					Males (N=743)				
Dietary carbohydrates				Total	SFA	MUFA	LA	ALA	Total	SFA	MUFA	LA	ALA
Total sucrose (%EI)	10.6 (8.20; 13.4)	9.98 (7.89; 12.8)	0.028	-0.19*	0.02	-0.26*	-0.33*	-0.23*	-0.19*	0.00	-0.27*	-0.30*	-0.23*
Sugary foods (%EI)	10.7 (6.60; 15.7)	9.73 (6.58; 14.4)	0.031	0.18*	0.33*	0.10	-0.10	-0.08	0.16*	0.24*	0.08	0.02	-0.08
Sugary drinks (%EI)	0.79 (0.17; 2.64)	2.24 (0.86; 6.11)	<0.01	-0.09	-0.06	-0.11	-0.06	-0.09	-0.19*	-0.19*	-0.17*	-0.11	-0.09
Total starch (%EI)	27.8 (22.9; 32.5)	26.2 (21.7; 31.5)	0.001	-0.36*	-0.37*	-0.33*	-0.09	-0.17*	-0.41*	-0.48*	-0.35*	-0.03	-0.17*
Starchy vegetables (%EI)	1.87 (1.17; 3.02)	1.70 (1.16; 2.75)	0.057	0.02	-0.03	0.02	0.15*	0.06	-0.02	-0.08	-0.01	0.14*	0.06
Refined grains (%EI)	28.1 (22.1; 34.1)	27.1 (20.9; 33.7)	0.019	-0.25*	-0.26*	-0.24*	-0.03	-0.06	-0.36*	-0.42*	-0.34*	-0.06	-0.06
Allergic outcomes (yes)				Correlations presented as Spearman's <i>Rho</i> , tested by Spearman's rank correlation. *Significant correlation (P value<0.001). Total=total fat. SFA=saturated fat. MUFA=monounsaturated fat. LA=linoleic acid (omega-6). ALA=alpha-linolenic acid (omega-3)									
Aeroallergen sensitization ^a	340 (41.1)	375 (50.5)	<0.01										
Asthma	34 (4.1)	52 (7.0)	0.016										
<i>Subset: allergic asthma^b</i>	23 (6.8)	44 (11.7)	0.032										
Rhinconjunctivitis	172 (20.7)	149 (20.1)	0.790										
Covariates													
Age (years)	15 (14.9; 15.1)	15 (14.9; 15.1)	0.198										
BMI (kg/m ²)	20.3 (18.8; 22.1)	20 (18.5; 22.1)	0.058										
Total energy intake (kcal)	1731(1342; 2154)	2348 (1883; 2804)	<0.01										
Fruit & vegetables (%EI)	5.87 (3.72; 9.28)	3.73 (2.14; 5.59)	<0.01										
Whole grains (%EI)	291 (92.2; 798)	273 (82.9; 716)	0.296										
Meat (%EI)	11.4 (7.1; 15.9)	13.8 (10.0; 19.0)	<0.01										
Fish (%EI)	1.01 (0.47; 1.77)	1.19 (0.63; 1.84)	0.001										
Dairy (%EI)	13.9 (9.0; 20.1)	14.8 (9.0; 21.3)	0.142										
MVPA (min/day)	7 (5.0; 10.5)	9 (6.5; 13.0)	<0.01										
Study arm													
<i>GINI observation arm</i>	275 (33.1)	247 (33.2)	0.127										
<i>GINI intervention arm</i>	264 (31.8)	205 (27.6)											
<i>LISA</i>	291 (35.1)	291 (39.2)											
Study centre													
<i>Munich</i>	453 (54.6)	425 (57.2)	0.270										
<i>Leipzig</i>	65 (7.8)	71 (9.6)											
<i>Bad Honnef</i>	37 (4.5)	29 (3.9)											
<i>Wesel</i>	275 (33.1)	218 (29.3)											
Parental education (high)	613 (73.9)	533 (71.7)	0.375										
Parental allergy (yes)	596 (71.8)	539 (72.5)	0.788										

Values are presented as counts (%) for categorical variables, or median (25th, 75th percentile) for continuous variables. Differences between sexes were tested by Chi-squared test or Wilcoxon rank-sum test, for categorical or continuous variables, respectively (*Significant difference, P-value<0.05). ^aIgE > 0.35 kU/L ^bSubset with positive aeroallergen sensitization (females=340; males=375). %EI=percentage of total daily energy intake. BMI=body mass index. MVPA=moderate to vigorous physical activity.


Sucrose




Sugary foods




Sugary drinks




Starch



Starchy vegetables



Refined grains






Figure 2_Musiol&Harris et al




Figure 3_Musiol&Harris et al




Figure 4_Musiol&Harris et al




Figure 5_Musiol&Harris et al