

EPA Public Access

Author manuscript

Eur Respir J. Author manuscript; available in PMC 2024 October 26.

About author manuscripts

Submit a manuscript

Published in final edited form as:

Eur Respir J. 2023 October ; 62(4): . doi:10.1183/13993003.01380-2023.

Clean Air in Europe for All: Taking Stock of the Proposed Revision to the Ambient Air Quality Directives. A Joint ERS, HEI, and ISEE Workshop Report

Michelle C Turner^{1,2,3}, Zorana Jovanovic Andersen⁴, Maria Neira⁵, Michal Krzyzanowski⁶, Ebba Malmqvist⁷, Alberto González Ortiz⁸, Gregor Kiesewetter⁹, Klea Katsouyanni⁶, Bert Brunekreef¹⁰, Erik Melén¹¹, Petter Ljungman¹¹, Margherita Tolotto¹², Francesco Forastiere⁶, Paul Dendale¹³, Richard Price¹⁴, Ole Bakke¹⁵, Sibylle Reichert¹⁶, Gerard Hoek¹⁰, Göran Pershagen¹¹, Annette Peters^{17,18,19}, Xavier Querol²⁰, Anna Gerometta²¹, Evangelia Samoli²², Iana Markevych^{23,24}, Romain Basthiste²⁵, Haneen Khreis²⁶, Pallavi Pant²⁷, Mark Nieuwenhuijsen^{1,2,3}, Jason D Sacks²⁸, Kjeld Hansen^{29,30}, Thomas Lymes³¹, Anne Stauffer³², Gary W Fuller³³, Hanna Boogaard²⁷, Barbara Hoffmann³⁴

¹Barcelona Institute for Global Health (ISGlobal), Barcelona, Spain

²Universitat Pompeu Fabra (UPF), Barcelona, Spain

³CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

⁴University of Copenhagen, Copenhagen, Denmark

⁵World Health Organisation (WHO), Geneva, Switzerland

⁶Imperial College London, London, UK

⁷Lund University, Lund, Sweden

⁸European Environment Agency (EEA), Copenhagen, Denmark

⁹International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

¹⁰Utrecht University, Utrecht, the Netherlands

- ¹¹Karolinska Institutet, Stockholm, Sweden
- ¹²European Environmental Bureau (EEB), Brussels, Belgium
- ¹³European Society of Cardiology (ESC), Sophia Antipolis, France
- ¹⁴European Cancer Organisation (ECO), Brussels, Belgium
- ¹⁵Standing Committee of European Doctors (CPME), Brussels, Belgium
- ¹⁶International Association of Mutual Benefit Societies (AIM), Brussels, Belgium

Corresponding author: Michelle C Turner, Barcelona Institute for Global Health (ISGlobal), Doctor Aiguader, 88, Barcelona, Spain, 08003. Tel. +34 932 147 397 michelle.turner@isglobal.org.

Conflicts of Interest:

The remaining authors have no conflict of interest to declare.

Disclaimer: The views expressed in this commentary are those of the authors and do not necessarily represent the views or policies of their parent organisation/body, nor of the funders.

¹⁸IBE, Medical Faculty, Ludwig Maximilians Universität, Munich, Germany

¹⁹Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, United States

²⁰Institute of Environmental Assessment and Water Research, IDAEACSIC, Barcelona, Spain

²¹Cittadini per l'aria onlus, Milan, Italy

²²Medical School, National and Kapodistrian University of Athens, Athens, Greece

²³Institute of Psychology, Jagiellonian University, Krakow, Poland

²⁴Health and Quality of Life in a Green and Sustainable Environment, SRIPD, Medical University of Plovdiv, Plovdiv, Bulgaria

²⁵City of Paris, Paris, France

²⁶University of Cambridge, Cambridge, UK

²⁷Health Effects Institute (HEI), Boston, United States

²⁸Center for Public Health and Environmental Assessment, Office of Research and Development, US Environmental Protection Agency (EPA), Research Triangle Park, NC, United States

²⁹European Lung Foundation, Sheffield, UK

³⁰Kristiania University College, Oslo, Norway

³¹Eurocities, Brussels, Belgium

³²Health and Environment Alliance, Brussels, Belgium

³³MRC Centre for Environment and Health, Imperial College London, London, UK

³⁴University of Düsseldorf, Düsseldorf, Germany

Abstract

Ambient air pollution is a major public health concern and comprehensive new legislation is currently being considered to improve air quality in Europe. The European Respiratory Society (ERS), Health Effects Institute (HEI), and International Society for Environmental Epidemiology (ISEE) organised a joint meeting on May 24, 2023 in Brussels, Belgium, to review and critically evaluate the latest evidence on the health effects of air pollution and discuss ongoing revisions of the European Ambient Air Quality Directives (AAQDs). A multi-disciplinary expert group of air pollution and health researchers, patient and medical societies, and policy representatives participated. This report summarises key discussions at the meeting.

Proposal for Revision of the European AAQDs and Potential to Improve Health

In 2021, the World Health Organization (WHO) released new Air Quality Guidelines (AQGs) based on a comprehensive synthesis of scientific evidence on the health effects

of air pollution [1]. WHO recommended that annual mean concentrations of particulate matter 2.5 microns in aerodynamic diameter ($PM_{2.5}$) and nitrogen dioxide (NO_2) should not exceed 5 µg/m³ and 10 µg/m³, respectively. The 2021 AQG levels are more stringent than the previous 2005 AQG levels of 10 µg/m³ for PM_{2.5} and 40 µg/m³ for NO₂ [2]. The current air quality legislation in Europe—the 2008 AAQDs—sets limit values for the annual mean of PM_{2.5} and NO₂ to 25 and 40 µg/m³, respectively [3].

Despite continued trends of decreasing air pollution concentrations in Europe [4], 95% and 75% of monitoring stations recorded values above 2021 WHO AQG levels for $PM_{2.5}$ and NO_2 , respectively, in 2021 [5]. $PM_{2.5}$ concentrations were highest in Central and Eastern Europe, and NO_2 in cities with high road traffic volume. Inequalities exist with regions with lower gross domestic product (GDP) per capita generally experiencing higher air pollution concentrations (Figure 1). The European Environment Agency (EEA) attributed 238,000 deaths to $PM_{2.5}$ concentrations above 5 μ g/m³ in the EU-27 in 2020 [6, 7]. These deaths are preventable and the estimate does not include millions of cases of non-fatal diseases, years lived with disability, attributable hospitalisations, or health effects from other pollutants [8].

Findings from the 3rd Clean Air Outlook, analysing existing and future policies in Europe, reveal substantial projected reductions in sulfur dioxide (SO₂), nitrogen oxides (NO_x), and primary PM_{2.5} emissions, and associated expected decreases in total PM_{2.5} concentrations for 2030 and 2050 [9], achieving the EU target of reducing premature deaths by 55% from 2005 to 2030 [10]. However, projected reductions of ammonia (NH₃) emissions (mainly from intensive livestock farming) are modest. Importantly, the currently foreseen reductions in emissions will still cause exceedances of the WHO AQG level for PM_{2.5} in large areas of Europe. Further mitigation potential exists and additional policy measures are needed including more stringent limit values, technical emission controls, accelerated energy transition, large-scale shifts towards safe and active mobility, among others (below).

To reduce the health and financial burden of air pollution, and achieve EU Green Deal goals, the European Commission (EC) published in October 2022 a proposal for revision of the EU AAQDs, which is currently being considered by the European Parliament and Council [11, 12]. The proposed new annual limit value for $PM_{2.5}$ by 2030 is 10 µg/m³ and for NO₂ 20 µg/m³. The proposal also includes non-binding target values and long-term objectives for O₃, as well as additional monitoring and modeling requirements, including of ultrafine particles (UFPs) and black carbon (BC), a regular review mechanism, and access to justice for non-compliance.

The proposed EC limit values would result in total gross benefits of 42 billion EUR/year that outweigh by 7 times mitigation costs of 5.6 billion EUR/year, and result in a positive net GDP impact of +0.38% [11, 12]. Although the EC proposal presents important steps toward cleaner air in Europe, it falls short of complete alignment with the 2021 WHO AQGs which would ensure additional health benefits and further reduce the health burden from air pollution [12–14].

Latest Science on Air Pollution and Health

The HEI-funded ELAPSE study is the most recent and comprehensive European study investigating health effects of low levels of air pollution [15–17]. ELAPSE documented adverse effects of long-term exposure to $PM_{2.5}$ and NO_2 at levels below current EU limit values, and to BC, with total and cardio-respiratory mortality in analysis of 28 million individuals.

Associations for $PM_{2.5}$ and NO_2 were stronger in ELAPSE than summary estimates from systematic reviews underpinning the 2021 WHO AQGs and European burden and impact assessments [6, 18]. The WHO AQG systematic reviews included studies published up to September 2018 world-wide and did not include ELAPSE findings, which were published later. To use the most relevant and recent evidence for Europe, the EC and EEA conducted additional analyses using the concentration-response functions of ELAPSE for $PM_{2.5}$ and NO_2 which resulted in higher attributable mortality estimates, indicating that the current health and financial burden of air pollution may be underestimated in Europe [12, 19, 20].

Recent research shows a range of impacts of air pollution on morbidity. Exposure to air pollution in pregnancy and early childhood can adversely impact lung function trajectories across the life course [21, 22], impair cognitive growth and development in children [23, 24], and accelerate cognitive decline in older adults [25, 26]. Findings from Stockholm, Sweden, indicated adverse effects of air pollution on lung function in infants and improvements in lung function in children with reductions of air pollution, even at levels below the EC proposal [27, 28]. There is robust evidence of adverse effects of PM_{2.5} on incidence of stroke, ischemic heart disease, atrial fibrillation, and heart failure [29]. ELAPSE and other studies in Europe showed a relationship between PM_{2.5} and lung cancer incidence, even at low air pollution concentrations [30] and of associations with cancers other than lung cancer [31–34].

For UFPs and BC/elemental carbon, which are currently not regulated, there is a need to further develop emission inventories, systematic monitoring, source apportionment, and research on health effects [1]. UFPs and BC are closely linked and likely responsible for systemic impacts of combustion-related particles on organs beyond the lung [35].

There has been an increase in pollutants from so called 'natural' sources, including desert dust and wildfires, closely linked with climate change and increase in extreme weather events such as droughts, heatwaves and storms [36–39]. Evidence on adverse health effects of pollution from wildfires and dust storms is growing [40, 41]. Pollution from natural sources often occurs simultaneously with heatwaves and during the high tropospheric ozone (O_3) season, raising concerns about additional adverse health effects of these synergistic exposures. In the current AAQDs, natural contributions are only considered regarding exceedances of PM limit values if they occur with anthropogenic (non-natural or resulting from human activities) emissions. However, in line with the 2021 WHO AQGs, measures to reduce exposure to natural source pollutants should be implemented; the need for reductions in anthropogenic emissions is also further emphasised [42].

There are increasing concerns about air pollution and climate change interactions on health, with initial research showing synergistic effects of short-term exposure to heat and $PM_{2.5}$ on respiratory and cardiovascular mortality [43]. Air pollution and climate mitigation and abatement policies have substantial overlaps, leading to important opportunities and cobenefits in exposure reduction and prevention of acute and chronic diseases.

Perspectives Towards Clean Air

Cities across Europe are taking actions to reduce air pollution. Implementing multimodal urban and transportation policies in policy packages is most effective at reducing emissions and ambient air pollution levels in cities, while at the same time targeting climate-friendly and health-promoting environments [42, 44]. In Paris, implementation of the first phases of low emission zones, decreasing speed limits, increasing bicycle paths, limiting traffic on streets near schools and closing of streets on specific days to cars were effective in decreasing $PM_{2.5}$ and NO_x . A new online catalogue of urban and transportation policy studies summarises the evidence-base for decision-makers [45]. Some of the most studied policies with emissions reductions included alternative fuel technologies, vehicle retrofitting, road pricing, low emission zones, and parking charges.

Focus on Eastern Europe

In Eastern and Southeastern Europe, there is a paucity of research on the health effects of air pollution. Over 70% of the Southeastern European population live in areas that exceed the current annual $PM_{2.5}$ EU limit value of 25 µg/m³ [46]. Main air pollution sources in the region include coal combustion for energy production and wood and coal for domestic heating and cooking. There is a need for air quality monitoring, investments in research, and sustained targeted actions [46, 47]. Previous initiatives included campaigns during the home heating season and individual behavioural nudging. Challenges linked to socio-economic conditions remain, including energy poverty, access to clean energy, and information for citizens.

Where Do We Go from Here?

The current policy debate regarding the EU AAQDs and alignment with the 2021 WHO AQGs [13] has important repercussions in Europe and also worldwide. Ambitious new European air quality legislation is of importance for motivating action. Although there are differences in legislative and policy-making processes in Europe compared to other regions (for example in consideration of achievability, cost-benefit, implementation and enforcement), the final version of the updated AAQDs in Europe will set an international benchmark.

It is imperative to support research to maintain a robust and up to date evidence base of emissions, exposure, and health effects of ambient air pollution, and to monitor the impacts of policies and practices to maximise public health benefits. There are opportunities to reinforce linkages of air pollution reduction with broader efforts to reduce environmental pollution, mitigation of and adaption to climate change, increase of sustainable agriculture

and food production, prevention of biodiversity loss, improved productivity, fiscal policies, and reduction of health care costs.

There is a need for further efforts to improve communication and information about air pollution to the general public, patients and patient organisations, health professionals, scientific societies, and decision-makers. Citizens, as well as citizen scientists, play an important role in ensuring clean air through improving awareness, exerting political pressure, supporting and performing research, and sharing and adapting knowledge on best practices. It is important to harmonize air pollution communications, indices, and alert systems across Europe, such as the EEA's mobile phone application on the European air quality index in 24 languages [48].

Further support to cities and networks sharing experiences in air quality interventions, local spatial planning and development, and policies are important public health measures showcasing successes and helping to avoid unwanted negative impacts. It is important to engage health professionals more systematically in policy discussions. Increased efforts to reinforce training of health and medical professionals on the environment, air pollution and health is needed. Emerging clinical guidance for healthcare professionals should be disseminated broadly [49].

The adverse health effects caused by air pollution are serious, debilitating diseases that result in a large burden to society. The ongoing revision of the EU AAQDs provide a unique opportunity to be bold in its ambitions, and to maximize public health benefits for Europe and beyond.

Acknowledgements:

The authors acknowledge the ERS, HEI, and ISEE for providing funding to hold the workshop. The event was also supported by Bloomberg Philanthropies. Detailed workshop materials are available on the HEI website : https://www.healtheffects.org/meeting/brussels-meeting-air-pollution-and-health-taking-stock-proposed-revision-ambient-air-quality.

The authors also acknowledge the following workshop participants who provided an opening address Virginijus Sinkevi ius, European Commissioner for Environment, Oceans and Fisheries and Javi Lopez, EU Parliament, Rapporteur for the Ambient Air Quality Directive (video address); and Alexander Simidchiev, Bulgarian National Assembly Parliament for participation as a discussant.

MCT is funded by a Ramón y Cajal fellowship (RYC-2017-01892) from the Spanish Ministry of Science, Innovation and Universities and co-funded by the European Social Fund. ISGlobal acknowledges support from the Spanish Ministry of Science and Innovation through the "Centro de Excelencia Severo Ochoa 2019-2023" Program (CEX2018-000806-S), and support from the Generalitat de Catalunya through the CERCA Program. IM is supported from the "NeuroSmog: Determining the impact of air pollution on the developing brain" (Nr. POIR.04.04.00-1763/18-00) which is implemented as part of the TEAM-NET programme of the Foundation for Polish Science, co-financed from EU resources, obtained from the European Regional Development Fund under the Smart Growth Operational Programme and by the "Strategic research and innovation program for the development of Medical University - Plovdiv" № BG-RRP-2.004-0007-C01, Establishment of a network of research higher schools, National plan for recovery and resilience, financed by the European Union - NextGenerationEU.

Hanna Boogaard and Pallavi Pant work at the Health Effects Institute, an organization jointly funded by the United States Environmental Protection Agency (EPA) and certain motor vehicle and engine manufacturers. The views expressed in this article are those of the authors and do not necessarily reflect the views of the Health Effects Institute or its sponsors.

References:

- 1. World Health Organization (WHO). WHO Global Air Quality Guidelines: Particulate Matter (PM2.5 and PM10), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide [Internet]. Geneva: World Health Organization; 2021.
- 2. World Health Organization (WHO). Air Quality Guidelines, Global Update 2005. Copenhagen, Denmark: WHO Regional Office for Europe; 2006.
- 3. EU (European Union). Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on Ambient Air Quality and Cleaner Air for Europe. 2008.
- European Environment Agency (EEA). Air Quality e-Reporting (AQ e-Reporting). https:// www.eea.europa.eu/en/datahub/datahubitem-view/3b390c9c-f321-490a-b25a-ae93b2ed80c1
- European Environment Agency (EEA). Europés Air Quality Status 2023. https:// www.eea.europa.eu/publications/europes-air-quality-status-2023
- Chen J, Hoek G. Long-term exposure to PM and all-cause and cause-specific mortality: A systematic review and meta-analysis. Environ Int 2020;143:105974.
- 7. European Environment Agency (EEA). Health impacts of air pollution in Europe, 2022. https:// www.eea.europa.eu/publications/air-quality-in-europe-2022/health-impacts-of-air-pollution
- Kienzler S, Soares J, González Ortiz A, Plass D. 2022. Estimating the Morbidity Related Environmental Burden of Disease due to Exposure to PM2.5, NO2 and O3 in Outdoor Ambient Air. (Eionet Report – ETC HE 2022/11). European Topic Centre on Human Health and the Environment.
- 9. European Commission (EC). The Third Clean Air Outlook. COM/2022/673 final/2.
- 10. European Commission (EC). COMMUNICATION FROM THE COMMISSION TO THE EUROPEAN PARLIAMENT, THE COUNCIL, THE EUROPEAN ECONOMIC AND SOCIAL COMMITTEE AND THE COMMITTEE OF THE REGIONS Pathway to a Healthy Planet for All EU Action Plan: 'Towards Zero Pollution for Air, Water and Soil'. COM/2021/400 final.
- 11. European Commission (EC). Proposal for a Directive of the European Parliament of the Council on Ambient Air Quality and Cleaner Air for Europe. COM/2022/542 final.
- 12. European Commission (EC) Staff Working Document. Impact Assessment Report. Accompanying the Document Proposal for a Directive of the European Parliament and of the Council on Ambient Air Quality and Cleaner Air for Europe (recast). SWD/2022/545 final.
- Hoffmann B, Boogaard H, de Nazelle A, Andersen ZJ, Abramson M, Brauer M, et al. WHO Air Quality Guidelines 2021–Aiming for healthier air for all: a joint statement by medical, public health, scientific societies and patient representative organisations. Int J Public Health 2021; 66:1604465.
- Boogaard H, Andersen ZJ, Brunekreef B, Forastiere F, Forsberg B, Hoek G, et al. Clean air in Europe for all: A call for more ambitious action. Environ Epidemiol 2023;7:e245. [PubMed: 37064425]
- 15. Brunekreef B, Strak M, Chen J, Andersen ZJ, Atkinson R, Bauwelinck M, et al. Mortality and Morbidity Effects of Long-Term Exposure to Low-Level PM2.5, BC, NO2, and O3: An Analysis of European Cohorts in the ELAPSE Project. Res Rep Health Eff Inst. 2021; 208.
- 16. Strak M, Weinmayr G, Rodopoulou S, Chen J, de Hoogh K, Andersen ZJ, et al. Long term exposure to low level air pollution and mortality in eight European cohorts within the ELAPSE project: pooled analysis. BMJ 2021;374:n1904. [PubMed: 34470785]
- 17. Stafoggia M, Oftedal B, Chen J, Rodopoulou S, Renzi M, Atkinson RW, et al. Long-term exposure to low ambient air pollution concentrations and mortality among 28 million people: results from seven large European cohorts within the ELAPSE project. Lancet Planet Health 2022;6:e9–e18 [PubMed: 34998464]
- Huangfu P, Atkinson R. Long-term exposure to NO2 and O3 and all cause and respiratory mortality: a systematic review and meta-analysis. Environ Int 2020;144:105998.
- Soares J, González Ortiz A, Gsella A, Horálek J, Plass D, Kienzler S. 2022. Health Risk Assessment of Air Pollution and the Impact of the New WHO Guidelines (Eionet Report – ETC HE 2022/10). European Topic Centre on Human Health and the Environment.

- Hoffmann B, Brunekreef B, Andersen ZJ, Forastiere F, Boogaard H. Benefits of future clean air policies in Europe. Proposed analyses of the mortality impacts of PM2.5 and NO2. Environ Epidemiol 2022;6:e221. [PubMed: 36249272]
- Melen E, Koppelman GH, Vicedo-Cabrera AM, Andersen ZJ, Bunyavanich S. Allergies to food and airborne allergens in children and adolescents: role of epigenetics in a changing environment. Lancet Child Adolesc Health 2022;6:P810–P189.
- 22. Agustí A, Melén E, DeMeo DL, Breyer-Kohansal R, Faner R. Pathogenesis of chronic obstructive pulmonary disease: understanding the contributions of gene-environment interactions across the lifespan. Lancet Respir Med 2022;10:512–524. [PubMed: 35427533]
- 23. Sunyer J, Esnaola M, Alvarez-Pedrerol M, Forns J, Rivas I, López-Vicente M, et al. Association between traffic-related air pollution in schools and cognitive development in primary school children: a prospective cohort study. PLoS Med 2015;12:e1001792.
- Guxens M, Lubczynska MJ, Perez-Crespo L, Muetzel RL, El Marroun H, Basagana X, et al. Associations of Air Pollution on the Brain in Children: A Brain Imaging Study. Res Rep Health Eff Inst 2022;209.
- 25. Thompson R, Smith RB, Bou Karim Y, Shen C, Drummond K, Teng C, et al. Air pollution and human cognition: A systematic review and meta-analysis. Sci Total Environ 2023;859:160234.
- 26. Wilker EH, Osman M, Weisskopf MG. Ambient air pollution and clinical dementia: systematic review and meta-analysis. BMJ 2023;381:e071620.
- 27. Yu Z, Merid SK, Bellander T, Bergström A, Eneroth K, Georgelis A, et al. Associations of improved air quality with lung function growth from childhood to adulthood: the BAMSE study. Eur Respir J 2023;61:2201783.
- Lundberg B, Gruzieva O, Eneroth K, Melén E, Persson Å, Hallberg J, et al. Air pollution exposure impairs lung function in infants. Acta Paediatr 2022;111:1788–1794. [PubMed: 35582781]
- de Bont J, Jaganathan S, Dahlquist M, Persson Å, Stafoggia M, Ljungman P. Ambient air pollution and cardiovascular diseases: An umbrella review of systematic reviews and meta-analyses. J Intern Med 2022;291:779–800. [PubMed: 35138681]
- Hvidtfeldt UA, Chen J, Andersen ZJ, Atkinson R, Bauwelinck M, Bellander T, et al. Long-term exposure to fine particle elemental components and lung cancer incidence in the ELAPSE pooled cohort. Environ Res 2021;193:110568.
- Turner MC, Andersen ZJ, Baccarelli A, Diver WR, Gapstur SM, Pope III CA, et al. Outdoor air pollution and cancer: an overview of the current evidence and public health recommendations. CA: Cancer J Clin 2020;70:460–479.
- 32. Hvidtfeldt UA, Chen J, Rodopoulou S, Strak M, de Hoogh K, Andersen ZJ, et al. Breast cancer incidence in relation to long-term low-level exposure to air pollution in the ELAPSE pooled cohort. Cancer Epidemiol Biomarkers Prev 2023;32:105–113. [PubMed: 36215200]
- 33. Chen J, Rodopoulou S, Strak M, de Hoogh K, Taj T, Poulsen AH, et al. Long-term exposure to ambient air pollution and bladder cancer incidence in a pooled European cohort: the ELAPSE project. Br J Cancer 2022;126:1499–1507. [PubMed: 35173304]
- 34. So R, Chen J, Mehta AJ, Liu S, Strak M, Wolf K, et al. Long-term exposure to air pollution and liver cancer incidence in six European cohorts. Int J Cancer 2021;149:1887–1897. [PubMed: 34278567]
- 35. Peters A. Ambient air pollution and Alzheimer's disease: the role of the composition of fine particles. Proc Natl Acad Sci U S A 2023;120:e2220028120.
- 36. Querol X, Tobias A, Perez N, Karanasiou A, Amato F, Stafoggia M, et al. Monitoring the impact of desert dust outbreaks for air quality for health studies. Environ Int 2019;130:104867.
- 37. Querol X, Perez N, Reche C, Ealo M, Ripoll A, Tur J, et al. African dust and air quality over Spain: Is it only dust that matters? Sci Total Environ 2019;686:737–752. [PubMed: 31195282]
- Salvador P, Pey J, Perez N, Querol X, Artinano B. Increasing atmospheric dust transport towards the western Mediterranean over 1948–2020. npj Climate and Atmospheric Science 2022;5:34.
- San-Miguel-Ayanz J, Durrant T, Boca R, Maianti P, Liberta G, Oom D, et al. Advance Report on Forest Fires in Europe, Middle East and North Africa 2022, EUR 31479 EN, Publications Office of the European Union, Luxembourg, 2023, ISBN 978–92-68–02143-9, doi:10.2760/091540, JRC133215.

- 40. Karanasiou A, Alastuey A, Amato F, Renzi M, Stafoggia M, Tobias A, et al. 2021. Short-term health effects from outdoor exposure to biomass burning emissions: A review. Sci Total Environ 2021;781:146739. [PubMed: 33798874]
- 41. Lwin KS, Tobias A, Chua PL, Yuan L, Thawonmas R, Ith S, et al. Effects of desert dust and sandstorms on human health: a scoping review. Geohealth 2023;7:e2022GH000728.
- 42. May AD, Khreis H, Mullen C. Option generation for policy measures and packages: An assessment of the KonSULT knowledgebase. Case Studies on Transport Policy 2018;6:311–318.
- 43. Rai M, Stafoggia M, de'Donato F, Scortichini M, Zafeiratou S, Vazquez Fernandez L, et al. Heat-related cardiorespiratory mortality: Effect modification by air pollution across 482 cities from 24 countries. Environ Int 2023;174:107825.
- 44. Glazener A, Sanchez K, Ramani T, Zietsman J, Nieuwenhiujsen MJ, Mindell JS, et al. Fourteen pathways between urban transportation and health: A conceptual model and literature review. Journal of Transport & Health 2021;21:101070.
- 45. Khreis H, Sanchez KA, Foster M, Burns J, Nieuwenhuijsen MJ, Jaikumar R, et al. Urban policy interventions to reduce traffic-related emissions and air pollution: A systematic evidence map. Environ Int 2023;172:107805.
- 46. Health Effects Institute (HEI). 2022. Trends in Air Quality and Health in Southeast Europe: A State of Global Air Special Report. Boston, MA: Health Effects Institute.
- 47. Spatial Bibliography Southeast Europe. Health Effects Institute 2023. https:// www.healtheffects.org/global/spatial-bibliography
- 48. European Environment Agency (EEA). European Air Quality Index App now available in all EU languages. https://www.eea.europa.eu/en/newsroom/news/european-air-quality-index-app
- 49. Jovanovic Andersen Z, Vicedo-Cabrera AM, Hoffmann B, Melén E. Climate change and respiratory disease: clinical guidance for healthcare professionals. Breathe (Sheff) 2023;19:220222.
- 50. European Environment Agency (EEA). 2023. European Environment and Health Atlas. European Environment Agency, Copenhagen, Denmark. Available online at https://discomap.eea.europa.eu/atlas




Figure 1. Combined risks and inequalities. PM_{2.5} vs GDP per capita [50]