

Loci for insulin processing and secretion provide insight into type 2 diabetes risk

Authors

K. Alaine Broadaway, Xianyong Yin,
Alice Williamson, ..., Claudia Langenberg,
Eleanor Wheeler, Karen L. Mohlke

Correspondence

eleanor.wheeler@mrc-epid.cam.ac.uk (E.W.),
mohlke@med.unc.edu (K.L.M.)

Broadaway et al. describe a genome-wide association meta-analysis in which they identify 36 proinsulin signals. Identification and integration of the proinsulin signals with glycemic traits, expression data in trait-relevant tissues, and functional follow-up provide hypotheses about potential mechanistic pathways for T2D loci.

Broadaway et al., 2023, *The American Journal of Human Genetics* 110, 284–299

February 2, 2023 © 2023 American Society of Human Genetics.
<https://doi.org/10.1016/j.ajhg.2023.01.002>



Loci for insulin processing and secretion provide insight into type 2 diabetes risk

K. Alaine Broadaway,¹ Xianyong Yin,^{2,3} Alice Williamson,^{4,5} Victoria A. Parsons,¹ Emma P. Wilson,¹ Anne H. Moxley,¹ Swarooparani Vadlamudi,¹ Arushi Varshney,⁶ Anne U. Jackson,³ Vasudha Ahuja,⁷ Stefan R. Bornstein,^{8,9,10} Laura J. Corbin,^{11,12} Graciela E. Delgado,¹³ Om P. Dwivedi,^{14,15} Lilian Fernandes Silva,¹⁶ Timothy M. Frayling,¹⁷ Harald Grallert,^{18,19,10} Stefan Gustafsson,²⁰ Liisa Hakaste,⁷ Ulf Hammar,²¹ Christian Herder,^{10,22,23} Sandra Herrmann,^{24,9} Kurt Højlund,²⁵ David A. Hughes,^{11,12} Marcus E. Kleber,^{13,26} Cecilia M. Lindgren,^{27,28,29,30} Ching-Ti Liu,³¹ Jian'an Luan,⁴ Anni Malmberg,³² Angela P. Moissl,^{33,34,13} Andrew P. Morris,³⁵

(Author list continued on next page)

Summary

Insulin secretion is critical for glucose homeostasis, and increased levels of the precursor proinsulin relative to insulin indicate pancreatic islet beta-cell stress and insufficient insulin secretory capacity in the setting of insulin resistance. We conducted meta-analyses of genome-wide association results for fasting proinsulin from 16 European-ancestry studies in 45,861 individuals. We found 36 independent signals at 30 loci (p value $< 5 \times 10^{-8}$), which validated 12 previously reported loci for proinsulin and ten additional loci previously identified for another glycemic trait. Half of the alleles associated with higher proinsulin showed higher rather than lower effects on glucose levels, corresponding to different mechanisms. Proinsulin loci included genes that affect prohormone convertases, beta-cell dysfunction, vesicle trafficking, beta-cell transcriptional regulation, and lysosomes/autophagy processes. We colocalized 11 proinsulin signals with islet expression quantitative trait locus (eQTL) data, suggesting candidate genes, including *ARSG*, *WIP11*, *SLC7A14*, and *SIX3*. The *NKX6-3/ANK1* proinsulin signal colocalized with a T2D signal and an adipose *ANK1* eQTL signal but not the islet *NKX6-3* eQTL. Signals were enriched for islet enhancers, and we showed a plausible islet regulatory mechanism for the lead signal in the *MADD* locus. These results show how detailed genetic studies of an intermediate phenotype can elucidate mechanisms that may predispose one to disease.

Introduction

Proinsulin is a precursor to insulin that is formed in pancreatic beta cells. Some proinsulin is secreted into the plasma during insulin biosynthesis and secretion, and

circulating levels of proinsulin relative to insulin are increased in individuals with type 2 diabetes (T2D) and pre-diabetes.^{1–3} Elevated proinsulin relative to insulin in individuals with pre-diabetes and T2D may be caused by increased demand on beta cells to release insulin, thereby

¹Department of Genetics, University of North Carolina, Chapel Hill, NC, USA; ²Department of Biostatistics, University of Michigan, Ann Arbor, MI, USA; ³Center for Statistical Genetics, University of Michigan, Ann Arbor, MI, USA; ⁴MRC Epidemiology Unit, Institute of Metabolic Science, University of Cambridge School of Clinical Medicine, Cambridge, UK; ⁵University of Cambridge Metabolic Research Laboratories, Wellcome Trust-MRC Institute of Metabolic Science, Department of Clinical Biochemistry, University of Cambridge, Cambridge, UK; ⁶Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA; ⁷Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland; ⁸Department of Internal Medicine, Metabolic and Vascular Medicine, MedicAl Faculty Carl Gustav Carus, Dresden, Germany; ⁹Helmholtz Zentrum München, Paul Langerhans Institute Dresden, University Hospital and Faculty of Medicine, TU Dresden, Dresden, Germany; ¹⁰German Center for Diabetes Research, Neuherberg, Germany; ¹¹Medical Research Council Integrative Epidemiology Unit at the University of Bristol, Bristol, UK; ¹²Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK; ¹³Medical Faculty Mannheim, Heidelberg University, Mannheim, BW, Germany; ¹⁴University of Helsinki, Helsinki, Finland; ¹⁵Folkhälsan Research Center, Helsinki, Finland; ¹⁶Institute of Clinical Medicine, University of Eastern Finland, Kuopio, Finland; ¹⁷College of Medicine and Health, Exeter University, Exeter, UK; ¹⁸Research Unit of Molecular Epidemiology, Helmholtz Zentrum München-German Research Center for Environmental Health, Neuherberg, Germany; ¹⁹Institute of Epidemiology, Helmholtz Zentrum München-German Research Center for Environmental Health, Neuherberg, Germany; ²⁰Department of Medical Sciences, Clinical Epidemiology, Uppsala University, Uppsala, Sweden; ²¹Department of Medical Sciences, Molecular Epidemiology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden; ²²Institute for Clinical Diabetology, German Diabetes Center, Leibniz Center for Diabetes Research at Heinrich Heine University Düsseldorf, Düsseldorf, Germany; ²³Department of Endocrinology and Diabetology, Medical Faculty and University Hospital Düsseldorf, Heinrich Heine University Düsseldorf, Düsseldorf, Germany; ²⁴Department of Internal Medicine, Prevention and Care of Diabetes, Medical Faculty Carl Gustav Carus, Dresden, Germany; ²⁵Steno Diabetes Center Odense, Odense, Denmark; ²⁶SYNLAB MVZ Humangenetik Mannheim, Mannheim, BW, Germany; ²⁷Oxford Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK; ²⁸Nuffield Department of Population Health, University of Oxford, Oxford, UK; ²⁹Wellcome Trust Centre Human Genetics, University of Oxford, Oxford, UK; ³⁰Broad Institute, Cambridge, MA, USA; ³¹Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA; ³²Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Helsinki, Finland; ³³Institute of Nutritional Sciences, Friedrich-Schiller-University, Jena, Germany; ³⁴Competence Cluster for Nutrition and Cardiovascular Health, Halle-Jena-Leipzig, Germany; ³⁵Centre for Genetics and Genomics Versus Arthritis, Centre for Musculoskeletal Research, The University of Manchester, Manchester, UK; ³⁶School of Health and Wellbeing, University of Glasgow, Glasgow, UK; ³⁷Chair of Food Chemistry and Molecular Sensory Science, Technische Universität München,

(Affiliations continued on next page)



Nikolaos Perakakis,^{8,9,10} Annette Peters,^{19,10} John R. Petrie,³⁶ Michael Roden,^{22,23,10} Peter E.H. Schwarz,^{24,9,10} Sapna Sharma,^{10,18,19,37} Angela Silveira,^{38,39} Rona J. Strawbridge,^{40,38} Tiinamaija Tuomi,^{7,15,41} Andrew R. Wood,⁴² Peitao Wu,³¹ Björn Zethelius,⁴³ Damiano Baldassarre,^{44,45} Johan G. Eriksson,^{46,47,48} Tove Fall,²¹ Jose C. Florez,^{49,50,51} Andreas Fritsche,^{52,53,10} Bruna Gigante,³⁸ Anders Hamsten,³⁸ Eero Kajantie,^{54,55,56,57} Markku Laakso,¹⁶ Jari Lahti,³² Deborah A. Lawlor,^{11,12} Lars Lind,²⁰ Winfried März,^{58,13} James B. Meigs,^{59,51,60} Johan Sundström,²⁰ Nicholas J. Timpson,^{11,12} Robert Wagner,^{52,53,10} Mark Walker,⁶¹ Nicholas J. Wareham,^{4,62} Hugh Watkins,⁶³ Inês Barroso,⁶⁴ Stephen O'Rahilly,⁶⁵ Niels Grarup,⁶⁶ Stephen C.J. Parker,^{6,67,2} Michael Boehnke,^{2,3} Claudia Langenberg,^{4,68,69} Eleanor Wheeler,^{4,70,*} and Karen L. Mohlke^{1,70,*}

encouraging the premature release of granules that contain a higher ratio of proinsulin to mature insulin.³ Conversely, reduced proinsulin-to-insulin levels could result from defects in proinsulin processing and folding prior to cleavage into insulin, early defects in vesicular processing, or altered proinsulin versus insulin degradation.⁴

Proinsulin can serve as a valuable intermediate phenotype to aid identification of genetic variations influencing hyperglycemia and T2D.⁵ Additionally, the allelic effect directions on glucose versus proinsulin can help differentiate known T2D loci into those involved in beta-cell stress versus defects in proinsulin processing and secretion.^{3,4,6-9} Previous proinsulin genome-wide association studies (GWASs) reported 16 signals at 13 genomic loci. These studies included a meta-analysis of 10,700 discovery participants that reported ten loci,⁵ a subsequent exome array study of Finnish individuals that identified two more loci with low-frequency (minor allele frequency [MAF] < 5%) variants,¹⁰ and a genetic study of participants with high risk for cardiovascular diseases (CVDs) that identified another locus.¹¹ To provide a comprehensive genetic analysis of proinsulin and gain insight into glycemic trait dysregulation, we performed a large meta-analysis of proinsulin GWASs. This study quadrupled the sample size of the largest previous meta-analysis and doubled the number of proinsulin association signals, implicating

candidate genes that regulate insulin processing and glucose regulation.

Subjects and methods

Cohort/study description

As part of the Meta-Analysis of Glycemic and Insulin traits Consortium (MAGIC), we conducted a meta-analysis of GWAS results for fasting proinsulin levels from 16 European-ancestry cohorts in up to 45,861 individuals (Table S1). Each of the 16 cohorts obtained institutional review board approval, collected trait and genotype data, assessed quality, and performed association analyses (Table S1). Each cohort performed imputation and reported all variants to Genome Reference Consortium Human Build 37/hg19.¹² Study participants who had diabetes, were on a diabetes treatment, or had fasting glucose ≥ 7 mmol/L, 2-h glucose ≥ 11.1 mmol/L, or hemoglobin A1c (HbA1c) $\geq 6.5\%$ (48 mmol/mol) were excluded. Fasting proinsulin values (pmol/L) were natural logarithm transformed and analyses adjusted for age, sex, population structure, and natural logarithm of fasting insulin (study-level details of fasting requirements, sample collection, and population structure adjustments are in Table S1). Study analysts ran models adjusted and unadjusted for body mass index (BMI). To control for type I error rate of low-frequency variants and to fully remove trait-covariate correlations, covariate adjustment was performed in two steps.¹³ Analysts first modeled natural logarithm of fasting proinsulin on all covariates and then inverse

Freising, Germany; ³⁸Department of Medicine Solna, Division of Cardiovascular Medicine, Karolinska Institutet, Stockholm, Sweden; ³⁹Oxford Biomedical Research Centre, Wellcome Centre for Human Genetics, University of Oxford, Oxford, UK; ⁴⁰Institute of Health and Wellbeing, Mental Health and Wellbeing, University of Glasgow, Glasgow, UK; ⁴¹Abdominal Center, Endocrinology, Helsinki University Hospital, Helsinki, Finland; ⁴²Genetics of Complex Traits, College of Medicine and Health, University of Exeter, Exeter, UK; ⁴³Department of Geriatrics, Uppsala University, Uppsala, Sweden; ⁴⁴Department of Medical Biotechnology and Translational Medicine, Università degli Studi di Milano, Milan, Italy; ⁴⁵Cardiovascular Prevention Area, Centro Cardiologico Monzino I.R.C.C.S., Milan, Italy; ⁴⁶Department of General Practice and Primary Health Care, Faculty of Medicine, University of Helsinki, Helsinki, Finland; ⁴⁷Folkhälsan Research Centre, Helsinki, Finland; ⁴⁸Department of Obstetrics and Gynecology, Yong Loo Lin School of Medicine, National University Singapore, Singapore, Singapore; ⁴⁹Diabetes Unit and Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA; ⁵⁰Programs in Metabolism and Medical & Population Genetics, Broad Institute, Cambridge, MA, USA; ⁵¹Department of Medicine, Harvard Medical School, Boston, MA, USA; ⁵²Department of Internal Medicine, Diabetology, Tübingen, Germany; ⁵³Institute for Diabetes Research and Metabolic Diseases, Helmholtz Center Munich, University of Tübingen, Tübingen, Germany; ⁵⁴Population Health Unit, Finnish Institute for Health and Welfare, Helsinki, Finland; ⁵⁵PEDEGO Research Unit, MRC Oulu, Oulu University Hospital and University of Oulu, Oulu, Finland; ⁵⁶Department of Clinical and Molecular Medicine, Norwegian University of Science and Technology, Trondheim, Norway; ⁵⁷Children's Hospital, Helsinki University Hospital and University of Helsinki, Helsinki, Finland; ⁵⁸Synlab Academy, SYNLAB Holding Deutschland GmbH, Mannheim, BW, Germany; ⁵⁹Department of Medicine, Division of General Internal Medicine, Massachusetts General Hospital, Boston, MA, USA; ⁶⁰Program in Medical and Population Genetics, Broad Institute, Cambridge, MA, USA; ⁶¹Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, UK; ⁶²Health Data Research UK, Gibbs Building, London, UK; ⁶³Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Oxford, UK; ⁶⁴Exeter Centre of Excellence for Diabetes Research, Genetics of Complex Traits, University of Exeter Medical School, University of Exeter, Exeter, UK; ⁶⁵MRC Metabolic Diseases Unit, Wellcome Trust-Medical Research Council Institute of Metabolic Science, University of Cambridge, Cambridge, UK; ⁶⁶Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark; ⁶⁷Department of Human Genetics, University of Michigan, Ann Arbor, MI, USA; ⁶⁸Computational Medicine, Berlin Institute of Health at Charité-Universitätsmedizin Berlin, Berlin, Germany; ⁶⁹Precision Healthcare University Research Institute, Queen Mary University of London, London, UK

⁷⁰These authors contributed equally

*Correspondence: eleanor.wheeler@mrc-epid.cam.ac.uk (E.W.), mohlke@med.unc.edu (K.L.M.)

<https://doi.org/10.1016/j.ajhg.2023.01.002>

normal transformed the residuals. Analysts then modeled the inverse normally transformed residuals on the covariates again and used these residuals in the final regression analysis. Analysts used an additive model in a linear/linear mixed-model framework with software including EPACTS, rvtests, and PLINK.^{14–16}

Study-level quality control (QC)

Central analysts assessed each cohort input file for QC by using EasyQC.¹⁷ We excluded variants with low minor allele count (<3) or low minor allele frequency (MAF < 0.005), low call rate (<95%), deviation from Hardy-Weinberg equilibrium (HWE) (p value < 0.00001), low imputation quality ($r^2 < 0.3$), or exceptionally large effect standard errors (SE > 10). We also examined quantile-quantile (QQ) plots by frequency bins, assessed trends in standard errors relative to sample size, and checked allele frequencies relative to their frequency in the Haplotype Reference Consortium (HRC). Systematic QC issues for a study were resolved prior to inclusion in the meta-analyses.

GWAS meta-analysis

We performed a fixed-effects inverse-variance-weighted meta-analysis by using METAL¹⁸ with effect size estimates and SE. We applied genomic control (GC) on summary statistics for each study and also following the meta-analysis. Post-meta-analysis inclusion criteria required that variants were represented by at least one-quarter of the maximum sample size, in at least two studies, and had an overall MAF > 0.005; we analyzed 9,533,557 variants. We defined a locus as a lead variant p value < 5×10^{-8} and all variants within 500 kb. We used SWISS (<https://github.com/statgen/swiss>) to identify the lead variant for each locus and combined adjacent loci whose lead variants exhibited linkage disequilibrium (LD) ($r^2 > 0.4$) to form an extended locus region. All LD calculations are based on 1000 Genomes Europeans unless otherwise noted. We estimated the proportion of variance explained by each variant as $2\beta^2f(1-f)$, where β is the effect size from METAL and f is the average effect allele frequency in the meta-analysis. We summed the variants' proportion of variance to estimate total fasting proinsulin variance explained.

Approximate conditional analysis

To identify conditionally distinct signals within a locus, we performed approximate conditional analysis by using GCTA.^{19,20} To reduce collinearity, we excluded any variant from designation as part of a distinct signal if its multiple regression r^2 on the other selected variants was greater than 0.8. Since no lead proinsulin variant was within 1 Mb of another, and we noted regions of extended LD surrounding at least one lead proinsulin variant, we analyzed all variants within 1 Mb of each lead variant or the extended locus region, whichever was larger. Given that GCTA depends on use of a large representative LD reference panel, we compared results from three genotype-level reference panels: METSIM ($n = 10,070$)²¹ and Fenland ($n = 8,925$)²² are the two largest studies in the meta-analysis that combined represent 38% of the total sample size and Electronic MEDical Records and GENomics (eMERGE, dbGaP: phs000888.v1.p1) ($n = 6,795$) is a European-only general research subset.²³ We defined a signal as conditionally distinct if a variant from GCTA representing the signal was identified with at least two of the three reference panels and the variants were proxies of each other ($r^2 > 0.8$). We additionally required variants to have consistent MAF across the summary data and the reference panels; the MAF of rs181143493 near

ARAPI was 0.12 in the proinsulin summary results and <0.01 in both the METSIM and eMERGE reference panels and therefore was excluded. Because of limitations in approximate conditional analysis with an external LD reference panel, we report at most three signals within a locus.

Colocalization with glycemic traits

We assessed signal overlap, or colocalization, between the 36 primary and secondary proinsulin signals and the conditionally distinct signals reported by three T2D studies: the European-ancestry component of DIABetes Meta-ANalysis of Trans-Ethnic association studies (DIAMANTE EUR),²⁴ the full multi-ancestry DIAMANTE analysis (DIAMANTE TA),²⁵ Asian Genetic Epidemiology Network (AGEN)/East Asian ancestry (EAS) DIAMANTE,²⁶ and four European-ancestry Meta-Analysis of Glucose and Insulin-related traits Consortium (MAGIC) glycemic traits: fasting glucose, fasting insulin, HbA1c, and glucose 2 h after a glucose challenge.²⁷ We tested for colocalization by using two strategies: colocalization based on pairwise LD ($r^2 > 0.8$) between the lead proinsulin variant and the lead variant for another trait and a Bayesian multi-trait colocalization approach, either HyPrColoc²⁸ or coloc.²⁹ Because of differences in ancestry across proinsulin versus AGEN and DIAMANTE TA, we ran HyPrColoc with proinsulin, DIAMANTE EUR, and the four MAGIC traits. We observed some issues with sensitivity when using HyPrColoc, including unstable trait clusters and deflated posterior probability for colocalization (PPFC) values when multiple signals in the cluster are marginally significant. While multi-trait HyPrColoc provided a beneficial first-pass assessment for colocalization, sensitivity analyses using pairwise colocalization helped fine-tune the specific studies that colocalized with our proinsulin data. Therefore, we compared HyPrColoc's multi-trait performance against a series of two-trait colocalization analyses (i.e., proinsulin and results for only one of the other five traits).

We performed HyPrColoc analyses by using predefined, approximately independent LD blocks and included all traits that had at least one variant with a p value < 10^{-4} within the LD block.³⁰ We selected the default HyPrColoc settings (prior.1 = 0.0001, prior.2 = 0.98). We then ran sensitivity analyses, varying the regional alignment thresholds from 0.6 to 0.9, the alignment thresholds from 0.6 to 0.9, and the prior.2 from 0.98 to 0.995. Since Bayesian colocalization methods may be sensitive to differences in ancestry across studies, we separately performed two-trait coloc analyses between proinsulin signals and genome-wide significant DIAMANTE TA signals and then proinsulin and AGEN T2D signals. We selected coloc's default prior probability of colocalization of 1×10^{-5} and ran sensitivity analyses varying the priors across 100 values. The cumulative sensitivity score for HyPrColoc and coloc was the proportion of scores that identified a colocalization and ranged from 0 (no sensitivity tests identify colocalization) to 1 (all sensitivity tests identify colocalization). Given limitations in colocalization approaches, we considered both Bayesian methods and LD; we considered the signals colocalized if the Bayesian posterior probability of colocalization was >0.6 and either the sensitivity score was >0.4 or LD $r^2 > 0.8$ between lead variants.

Characterization of proinsulin locus effect directions to other glycemic traits

To assess the direction of effect of proinsulin signals on T2D and common glycemic traits, we looked up associations for proinsulin lead variants in the summary results for T2D in the

forementioned three studies and the four glyce- mic traits in MAGIC studies.^{24–27} If a proinsulin lead signal was associated with T2D or fasting glucose (p value $< 10^{-4}$) or at least two out- comes in the same direction at a more lenient p value threshold (p value < 0.01), we reported the consensus direction of effect. To evaluate proinsulin variant association with additional glyce- mic traits, we performed similar look ups in the summary results for 34 glyce- mic traits analyzed in the METSIM study (Table S2);¹⁰ briefly, these traits included proinsulin, glucose, and insulin levels at fasting and after an oral glucose tolerance test (30–120 min) and calculated areas under the curve measures as well as C-peptide, HbA1c, insulinogenic index, Matsuda index, and T2D. We analyzed the 34 traits as a subset of a total of 1,076 baseline traits for association with variants imputed via a reference panel from a subset of METSIM with whole-genome sequencing.³¹ For glucose and insulin metabolic traits, we excluded individuals known to be diabetic at baseline. For each quantitative trait, we inverse normalized the trait, regressed on covariates (see Table S2 for covariates per trait), and inverse normalized the residuals. We carried out single-variant association tests by using a linear mixed model in SAIGE v.0.39 (<https://github.com/weizhouUMICH/SAIGE>) on the normalized residual trait values.

We additionally looked up proinsulin lead variants for loci not identified in T2D or glyce- mic trait association results. We used genetics.opentargets.org to find significant associations (p value $< 5 \times 10^{-4}$) with the lead variants at these loci.^{32,33} The online resource identifies associations from the GWAS Catalog,³⁴ Neale lab UK Biobank summary statistics (<http://www.nealelab.is/uk-biobank/>), SAIGE UK Biobank summary statistics,³⁵ and FinnGen Summary statistics.³⁶

Candidate genes

We obtained nearby genes' islet expression specificity index (iESI) deciles.³⁷ iESI deciles indicate the extent to which genes are both highly expressed in islets as well as the specificity for islet expres- sion versus ubiquitous expression across other tissues; values near zero represent genes that have low islet specificity or low expres- sion in islets and values near 10 represent genes whose expression is highly specific to islets. We define high iESI genes as those with a decile above 7. We consolidated gene labels across sources by using Entrez gene symbols.

Next, we performed colocalization of proinsulin signals with two eQTL datasets. First, a human islet RNA sequencing (RNA- seq)-based eQTL study from the InsPIRE consortium ($n = 420$),³⁸ which reported significant eQTLs for 4,312 genes (false discovery rate [FDR] $< 1\%$), and second, a subcutaneous adipose tissue RNA-seq study from 434 Finnish men in the METSIM study,³⁹ which reported at least one significant eQTL at 9,687 genes (FDR $< 1\%$). We used LD and HyPrColoc to test for colocalizations with genes within 1 Mb of each lead proinsulin variant; as described in the previous section, we used a multi-study frame- work with proinsulin, European-ancestry DIAMANTE,²⁴ MAGIC glyce- mic traits,²⁷ and one eQTL gene at a time, as well as testing with only proinsulin and each gene. We considered the signals col- ocalized if HyPrColoc PPFC scores were >0.6 and either the sensi- tivity score was >0.4 or LD $r^2 > 0.8$. We plotted signals by using LocusZoom.⁴⁰ Additionally, we performed summary Mendelian randomization (SMR)⁴¹ to begin assessing potential causal rela- tionships by using the genetic variants as an instrumental variable to test for the causative effect of gene expression on proinsulin. To account for multiple hypothesis testing, we used a Bonferroni-cor-

rected significance threshold. To evaluate evidence of pleiotropy from linkage between two distinct causal variants, we ran hetero- geneity in dependent instruments (HEIDI) as part of the SMR analysis.

Identification of extended credible set variants

We determined 99% credible sets by using regions ± 500 kb around each lead variant, using the following equation for Bayes factors:

$$\ln(BF) \propto 0.5 \frac{\beta^2}{SE^2}$$

where β and SE are the effect sizes and standard errors from the meta-analysis.⁴² For loci with multiple significant signals, we used the approximate conditional analysis option in GCTA, using eMERGE as the reference panel, to define credible sets. Variants with a low posterior probability are less likely to be causal; howev- er, variants that are not represented or poorly represented in the meta-analysis may erroneously be excluded from consideration as a putative causal variant. We therefore extended the credible set to include all variants in high LD ($r^2 > 0.8$ in 1000 Genomes European) with the lead variant. This approach recognizes variants that are not included in the meta-analysis as a result of analytic or technical factors (e.g., insertions or deletions [indels] are not imputed by HRC and variants with MAF $< 0.5\%$) as well as variants that are poorly represented in our meta-analysis as a result of fac- tors such as low sample size.

Coding and regulatory elements

To identify potential candidate genes for each signal, we consid- ered protein-coding genes within ~ 100 kb of the signal's lead variant,⁴³ with special attention to genes for which a coding variant is included in a signal's extended credible set and those that are highly and specifically expressed in islets. To identify genes through coding effects, we obtained annotation for all vari- ants in our extended credible set by using Variant Effect Predictor (VEP),⁴⁴ Sorting Intolerant from Tolerant (SIFT),⁴⁵ PolyPhen-2,⁴⁶ Combined Annotation-Dependent Depletion (CADD),^{47,48} and MutationAssessor.⁴⁹ For all functional predication tools, we selected default thresholds.

We tested proinsulin signals for regulatory element enrichment by using the following epigenomic annotations: chromatin states in islets, adipose, and skeletal muscle,⁵⁰ bulk assay for transposase- accessible chromatin with high-throughput sequencing (ATAC- seq) peaks;^{38,51} islet single-nucleus ATAC-seq (sn-ATAC) cluster peaks;⁵² and other islet chromatin annotations.⁵³ We used the genomic regulatory elements and GWAS overlap algorithm (GREGOR) to evaluate global enrichment of proinsulin-associated variants in epigenomic regulatory features.⁵⁴ GREGOR observes the signal overlap in annotated regulatory data among lead GWAS variants or their LD proxies ($r^2 > 0.8$) relative to expected overlap-based control variants matched to index variants for num- ber of variants in LD, minor allele frequency, and distance to near- est gene.

Transcriptional activity assays

Cell culture

We cultured INS1-derived rat insulinoma pancreatic beta-islet 832/ 13 cells (provided by C. Newgard, Duke University, Durham, NC) in RPMI 1640 medium (Corning, NY) supplemented with 10% FBS, 10 mM HEPES, 2 mM L-glutamine, 1 mM sodium pyruvate,

and 50 μ M 2-mercaptoethanol, and we cultured murine insulinoma MIN6 cells (provided by C. Rhodes, Joslin Diabetes Center, Boston, MA) in high-glucose DMEM (Sigma-Aldrich, St. Louis, MO) supplemented with 10% FBS, 1 mM sodium pyruvate, and 100 μ M 2-mercaptoethanol. All cells were maintained in a humidified incubator at 37°C with 5% CO₂, and prior to transfection, both cell lines tested negative for *Mycoplasma* contamination in accordance with the MycoAlert *Mycoplasma* Detection Kit (Lonza, Morristown, NJ).

Transcriptional reporter assays

To test for allelic differences in transcriptional activity, we performed dual-luciferase reporter assays as previously described.⁵⁵ We used genomic DNA of individuals homozygous for the reference or alternate alleles to amplify fragments surrounding rs10501320, cloned amplicons into the firefly luciferase reporter vector pGL4.23 (Promega, Madison, WI), and sequence-confirmed five purified clones for each allele, in each orientation (Azenta, Research Triangle Park, NC); alleles at additional variants within each amplicon were kept consistent (Table S3). 24 h prior to transfection, we seeded 832/13 and MIN6 cells in 24-well plates (200,000 cells per well). Upon reaching 90% confluence, we transfected 832/13 cells in duplicate with 500 ng of plasmid DNA and 1 μ L of Lipofectamine 3000 (Thermo Fisher Scientific, Waltham, MA) per well, and we transfected MIN6 cells in duplicate with 250 ng of plasmid DNA and 1 μ L Lipofectamine LTX (Thermo Fisher Scientific) per well; we co-transfected both 832/13 and MIN6 cells with 80 ng of pRL-TK *Renilla* (Promega) per well. We used two independent preparations of empty vector pGL4.23 as negative controls. After 48 h, we performed dual-luciferase reporter assays (Promega), normalized luciferase to *Renilla*, and calculated fold-change relative to empty vector controls by using two-sided t tests assuming equal variance ($\alpha = 0.05$). We independently repeated transfections on different days and observed consistent results. Results show ten biological replicates (separate transfections) and two averaged technical replicates (luciferase and *Renilla* readings).

Results

Identification of proinsulin association signals

We identified 28 loci associated at genome-wide significance (p value $< 5 \times 10^{-8}$) with proinsulin adjusted for BMI, including 16 loci >500 kb away from a previously reported proinsulin association (Tables 1 and S4, Figures S1 and S2). Combined, the 28 lead variants explained an estimated 8.9% of the total proinsulin variance in the meta-analysis, and the estimated percent of trait variance explained by each variant ranged from 2.1% (*STARD10*) to 0.07% (*JARID2*).

Association results for fasting proinsulin without BMI adjustment yielded results similar to those obtained in the BMI-adjusted analysis (Pearson correlation of effect estimates = 0.97; Figure S3 and Table S5). Variants at two additional loci, *SLC2A10* and *BCL11A*, which narrowly missed the significance threshold in the analysis with BMI adjustment (p value = 6×10^{-8} and 1.5×10^{-7} , respectively) attained genome-wide significance in the analysis without BMI adjustment (Table 1).

We performed subsequent approximate conditional analysis and identified six additional signals at genome-

wide significance located within 500 kb of the lead variant of five known proinsulin loci near *STARD10*, *MADD*, *PCSK1*, *SGSM2*, and *DDX31* (Tables 2 and S6, Figures S4 and S5). We identified three previously reported signals near *MADD*, including one signal that consists of a proinsulin-associated¹⁰ nonsense variant (rs35233100) that is now genome-wide significant after conditioning on the lead signal (rs10501320). Both the primary and secondary signals at the *SGSM2* locus have been previously reported.^{5,10,11} We also identify secondary signals located near *STARD10*, *PCSK1*, and *DDX31*. At *DDX31*, although both signals (rs368476 and rs7864386) were within 50 kb of the previously reported female-specific *DDX31* signal (rs306549),¹¹ neither was in high LD with the previously reported lead variant ($r^2 < 0.1$, Figure S5),⁵ validating the *DDX31* locus, but not the previously reported signal. For subsequent analyses, unless otherwise stated, we included the 28 primary signals and six conditionally distinct signals for proinsulin adjusted for BMI, as well as the two signals for proinsulin not adjusted for BMI, for a total of 36 signals at 30 loci.

This meta-analysis replicated four low-frequency (MAF < 0.05) proinsulin-associated signals originally identified in an exome array analysis of Finnish participants in the METSIM exome study¹⁰ (Table S7, Figures S6 and S7). We validated missense or nonsense lead variants in *TBC1D30*, *SGSM2*, and *MADD*, all of which were genome-wide significant in the meta-analysis even after excluding METSIM. The signal at the *KANK1* locus was only genome-wide significant in the full meta-analysis (lead variant rs146375546, p value = 4.3×10^{-11}), as the lead variant is rare in general European-ancestry populations but enriched in Finnish-ancestry populations (1000 Genomes MAF = 0.003 in 1000 Genomes European-ancestry populations versus 0.015 in the Finnish population). The replications of associations at the four low-frequency variants highlight the utility of exome arrays in finding low-frequency variants and the challenges in replicating variants that are not equally represented across populations.

Proinsulin signals and other glycemic traits

We compared all 36 proinsulin signals described above to up to 568 GWAS signals identified for T2D^{24–26} and up to 218 signals in four glycemic traits including fasting and 2-h glucose, HbA1c, and fasting insulin²⁷ (Tables S8–S10). We performed colocalization analysis and identified colocalizations for 15 proinsulin signals with signals for T2D ($N = 12$) or glycemic traits ($N = 9$): six previously known proinsulin signals near *STARD10*, *MADD*, *TCF7L2*, *SGSM2*, *SLC30A8*, and *C2CD4A/B* and nine additional proinsulin signals near *SIX3*, *TLE1*, *RNF6*, *PAM*, *NKX6-3*, *FAM185A*, *BCL11A*, *GIPR*, and *FAM46C*. We also identified colocalizations between an additional ten T2D or glycemic trait loci that were associated with proinsulin at a less stringent significance threshold ($5 \times 10^{-8} < p$ value $< 1 \times 10^{-4}$) (Table S8). Eight proinsulin loci (*STX16*, *DLC1*, *SLC7A14*,

Table 1. Thirty loci associated with plasma proinsulin levels

Locus	rs ID	Chr	Position	EA/NEA	EAF	Beta	Std Err	p value
<i>SIX3</i>	rs12712928	2	45,192,080	C/G	0.16	0.09	0.01	1.5×10^{-21}
<i>ELAPOR1</i>	rs74920406	1	109,704,525	C/T	0.96	0.15	0.02	3.7×10^{-16}
<i>TLE1</i>	rs2796441	9	84,308,948	G/A	0.59	0.05	0.01	9.6×10^{-14}
<i>TPD52</i>	rs1346146	8	81,047,278	T/C	0.45	0.05	0.01	2.0×10^{-13}
<i>GIPR</i>	rs10423928	19	46,182,304	A/T	0.22	0.06	0.01	7.6×10^{-12}
<i>STX16</i>	rs218473	20	57,235,980	C/T	0.32	0.05	0.01	1.5×10^{-10}
<i>DLC1</i>	rs2977105	8	12,794,444	C/T	0.82	0.06	0.01	1.0×10^{-9}
<i>FAM46C</i>	rs826415	1	118,153,977	T/G	0.67	0.04	0.01	1.3×10^{-9}
<i>PCSK2</i>	rs111925767	20	17,331,621	T/G	0.23	0.05	0.01	1.6×10^{-9}
<i>RNF6</i>	rs10507349	13	26,781,528	G/A	0.78	0.05	0.01	1.9×10^{-9}
<i>PAM</i>	rs75457267	5	102,658,770	C/T	0.96	0.10	0.02	2.2×10^{-9}
<i>SLC7A14</i>	rs56252324	3	170,334,547	A/C	0.87	0.06	0.01	5.4×10^{-9}
<i>WIPI1</i>	rs2302783	17	66,447,073	C/T	0.72	0.04	0.01	1.1×10^{-8}
<i>NKX6-3/ANK1</i>	rs13266210	8	41,533,514	G/A	0.21	0.05	0.01	2.1×10^{-8}
<i>FAM185A</i>	rs10228495	7	102,440,184	C/T	0.45	0.04	0.01	2.9×10^{-8}
<i>JARID2</i>	rs16876519	6	15,496,122	A/G	0.85	0.05	0.01	3.5×10^{-8}
Previously reported loci								
<i>STARD10</i>	rs77464186	11	72,460,398	C/A	0.19	0.26	0.01	3.7×10^{-202}
<i>MADD</i>	rs10501320	11	47,293,799	G/C	0.76	0.21	0.01	1.3×10^{-165}
<i>PCSK1</i>	rs13169290	5	95,729,406	A/G	0.28	0.12	0.01	3.3×10^{-59}
<i>CDC4A/B</i>	rs11856307	15	62,399,093	A/C	0.54	0.09	0.01	6.4×10^{-40}
<i>TCF7L2</i>	rs7903146	10	114,758,349	T/C	0.26	0.10	0.01	1.9×10^{-39}
<i>SLC30A8</i>	rs4300038	8	118,217,915	G/A	0.66	0.09	0.01	4.1×10^{-39}
<i>LARP6</i>	rs113350503	15	71,111,437	G/A	0.57	0.06	0.01	6.5×10^{-18}
<i>DDX31</i>	rs368476	9	135,456,552	A/G	0.65	0.07	0.01	7.6×10^{-21}
<i>SNX7</i>	rs6702126	1	99,199,954	G/A	0.65	0.04	0.01	8.7×10^{-10}
<i>SGSM2</i>	rs61741902	17	2,282,779	A/G	0.01	0.47	0.03	5.8×10^{-49}
<i>TBC1D30</i>	rs150781447	12	65,224,220	T/C	0.02	0.30	0.04	9.1×10^{-17}
<i>KANK1</i>	rs146375546	9	727,176	G/A	0.03	0.26	0.04	4.3×10^{-11}
Loci in model without BMI adjustment								
<i>SLC2A10</i>	rs3091537	20	45,332,200	A/C	0.64	0.04	0.01	3.9×10^{-8}
<i>BCL11A</i>	rs243018	2	60,586,707	G/C	0.45	0.04	0.01	2.4×10^{-8}

Chr, chromosome; EA, effect allele; NEA, non-effect allele; EAF, effect allele frequency; Std Err, SE of beta. Loci are labeled by one or more nearby candidate genes.

WIPI1, *JARID2*, *SLC2A10*, *ELAPOR1*, and *PCSK2*) were not colocalized with T2D or any glycemic trait.

We obtained the direction of allelic effect of the 30 lead proinsulin leads on fasting glucose²⁷ and more than 30 other related glycemic traits including proinsulin levels after an oral glucose challenge¹⁰ (Figure 1, Tables S2 and S10). The allele associated with higher glucose was associated with higher proinsulin for half the lead variants (15 of 30) and associated with lower proinsulin for the other half.

Putative candidate genes

To identify potential candidate genes for each signal, we identified nearby genes, obtained their iESI deciles, and performed colocalization and SMR analyses with eQTL data (Tables S11–S14).^{38,39} Genes with high expression levels in islets, particularly those that are not highly expressed in other tissues, represent strong candidate genes for influencing the proinsulin to insulin processing pathway. These genes that are highly and specifically expressed in islets will have high iESI values (defined as iESI

Table 2. Six conditionally distinct proinsulin signals

Locus	Marginal associations						Conditional associations			LD with primary (r^2)
	rs ID	EA/NEA	EAF	Beta	Std Err	p value	bC	bC_se	pC	
<i>STARD10</i>	rs481206	C/T	0.69	0.12	0.01	3.8×10^{-62}	0.06	0.01	1.0×10^{-16}	0.068
<i>MADD</i>	rs35233100	C/T	0.94	0.35	0.02	1.9×10^{-104}	0.23	0.02	3.0×10^{-46}	0.154
<i>MADD</i>	rs1449626	A/C	0.78	0.01	0.01	4.8×10^{-1}	0.06	0.01	7.0×10^{-15}	0.068
<i>PCSK1</i>	rs2117141	C/T	0.41	0.06	0.01	4.0×10^{-16}	0.07	0.01	1.9×10^{-24}	0.008
<i>SGSM2</i>	rs2447103	C/A	0.51	0.07	0.01	3.5×10^{-26}	0.07	0.01	5.3×10^{-22}	0.004
<i>DDX31</i>	rs7864386	G/A	0.56	0.03	0.01	1.6×10^{-6}	0.04	0.01	1.8×10^{-10}	0.027

Conditionally distinct signals identified with GCTA-COJO and the eMERGE reference panel. EA, effect allele; NEA, non-effect allele; EAF, effect allele frequency; Std Err, SE of beta; bC, conditional beta; bC_se, conditional SE of beta; pC, conditional p value. Results for both *MADD* signals are from the analyses conditioning on the other two *MADD* signals.

decile > 7).³⁷ Most (29/36) proinsulin signals fell within 100 kb of at least one gene with a high iESI (Table S11). Top iESI genes included well-documented beta-cell genes, such as *MADD*, *PCSK1*, and *PCSK2*,^{56–58} as well as genes at loci not previously described in glycemic trait studies: *ELAPOR1* and *SLC7A14*.

To identify additional candidate genes underlying the proinsulin association signals, we colocalized them with eQTL signals^{38,39} (Tables S12 and S13). Through colocalization with eQTLs in pancreatic islets from the Inspire consortium,³⁸ we identified 11 proinsulin signals that colocalized with eQTL signals for 17 genes (Table S12); six proinsulin signals colocalized with eQTLs for more than one gene. The alleles associated with higher proinsulin were associated with higher expression of eight genes (*MADD*, *RNF6*, *CDK8*, *SLC2A10*, *SNX7*, *ARAP1*, *STARD10*, and *TCF7L2*) and lower expression of nine protein-coding genes or noncoding transcripts (*SIX3*, *SIX2*, *RP11-89K21.1*, *AC012354.6*, *ARSG*, *WIP1I*, *SLC7A14*, *FAM46C*, and *LA RP6*). All 17 colocalizations also passed the experiment-wide significance threshold for SMR (p value < 0.0029). Using HEIDI, we detected heterogeneity for just one gene at p value < 0.0029: *STARD10*. While this may indicate the correlation is due to linkage rather than pleiotropy, the result may also be due to the complicated structure of this locus, which may violate the assumption of only one causal variant in the eQTL region.

Signal colocalization at the *NKX6-3/ANK1* locus provided additional data with which to interpret this complex locus. The locus includes two T2D signals^{24,26}: one colocalized with the *NKX6-3* eQTL in islets²⁴ and the other colocalized with an *ANK1* eQTL in adipose and muscle.^{26,59} *NKX6-3* is highly and specifically expressed in islets (iESI decile = 10), while *ANK1* is not (iESI decile = 2). The T2D risk alleles for the two signals were associated with lower islet *NKX6-3* expression and higher *ANK1* expression in adipose and muscle, suggesting that the signals affect T2D risk in different tissues. We observed only one proinsulin association signal at this locus. While we might have expected it to align with the proposed islet *NKX6-3* eQTL signal, it instead colocalized with the adipose *ANK1*

eQTL signal (Figures 2 and S8, Table S13). The proinsulin lead variant rs13266210 is in strong LD with the *ANK1* eQTL (rs3802315, $r^2 = 0.84$) and the East Asian AGEN T2D lead variant (rs62508166, $r^2 = 0.92$), and HyPrColoc shows strong evidence of colocalization across all three studies (PPFC = 0.92). The A allele of rs13266210 is associated with increased T2D risk, higher *ANK1* expression in adipose, and lower proinsulin. At this proinsulin signal, proxy variant rs6989203 (LD $r^2 = 0.84$ with rs13266210) overlaps with an islet beta-cell single nucleus ATAC peak⁵² and is in high LD with the *ANK1* eQTL site ($r^2 = 0.93$). Of the two T2D signals at the *ANK1/NKX6-3* locus previously proposed to act in different tissues on different genes, the proinsulin signal colocalizes with the adipose *ANK1* signal rather than the expected colocalization with islet *NKX6-3*.

Credible sets and variant annotation and function

We built a credible set of putative causal variants for each of the 36 signals. These 36 sets together contained 814 variants (Table S15). We extended the credible sets to include 276 additional variants exhibiting LD $r^2 \geq 0.8$ (1000 Genome European-ancestry reference) with the lead variants, including 142 variants that were unavailable in the meta-analysis and therefore could not have been included in the Bayesian credible set. Three signals had one variant in the extended credible set (*SGSM2*, *ELAPOR1*, and the second signal in *DDX31*) and 14 signals (39%) had ten variants or fewer.

The extended credible sets for 17 proinsulin signals contained coding variants (Table S16). Across all credible sets, we observed one nonsense, 18 missense, and 31 synonymous variants. The credible sets for 13 proinsulin signals contained at least one missense variant: seven signals in previously identified proinsulin loci (*TBC1D30*, *PCSK1*, *KANK1*, *FAM185A*, the first and second signals at *SGSM2*, and the third signal in *MADD*), four in loci known in other glycemic trait GWASs (*SLC30A8*, *GIPR*, *FAM46C*, and *PAM*), and two that are not known proinsulin or glycemic trait genes (*ELAPOR1* and *WIP1I*). The lead variant rs74920406 at the *ELAPOR1* locus, a missense variant of low




Figure 1. Direction of allelic effect of fasting glucose versus fasting proinsulin
Standardized effect sizes for lead variants are shown from this study compared to fasting glucose from Chen et al. (2021).²⁷ Left of the vertical line, alleles associated with higher fasting glucose and lower proinsulin; right of the vertical line, alleles associated with higher fasting glucose and higher proinsulin.

frequency (p.His55Tyr, MAF = 0.04), was not previously associated with proinsulin or other glycemic traits but was associated with low-density lipoprotein (LDL) (Table S17).⁶⁰ This variant is conserved across species^{48,61,62} and has a probably damaging effect on the protein.⁴⁶ *ELAPOR1* encodes endosome-lysosome associated apoptosis and autophagy regulator 1 and inhibits beta-cell insulin signaling by accelerating endocytosis of the insulin receptor and insulin-like growth factor receptors.⁶³ The credible set for *WIPI1* contained a coding missense variant (p.Thr31Ile; rs883541). *WIPI1* is a phosphatidylinositol-2-phosphate effector gene, which encodes a component of the autophagy machinery; skeletal muscle from severely insulin-resistant individuals with T2D displayed decreased expression of autophagy-related genes, including *WIPI1*.⁶⁴

Among the 1,090 variants in the extended credible sets for all signals, 62 overlapped with an active enhancer in islets and 76 overlapped with an islet cell type single-nucleus ATAC-seq peak (Table S18). We thus examined regulatory annotations of proinsulin-associated credible sets. The variants were enriched in islet active enhancers (Figure 3, fold enrichment = 8.8, p value = 4.6×10^{-12}). Among islet single-nucleus ATAC-seq peaks, beta-cell peaks were most enriched (fold enrichment = 2.9, p value = 5.1×10^{-10}).

To further investigate plausible allelic effects of one variant located in an annotated ATAC-seq peak, we examined the regulatory function of lead variant rs10501320, at *MADD*, in transcriptional reporter assays. *MADD* is a well-documented proinsulin locus associated with proinsulin-to-insulin conversion.⁶⁵ Compared to a negative control, a genomic fragment spanning rs10501320 and the surrounding ATAC-seq peak showed ~3-fold increased transcriptional activity in rat insulinoma 832/13 cells and a

~4-fold increase in transcriptional activity in mouse insulinoma MIN6 cells, consistent with a role as an enhancer (Figures 3 and S9). The rs10501320-G allele showed 1.3- to 1.6-fold greater transcriptional activity than the C allele (p value < 0.0001); the G allele was associated with higher proinsulin in this GWAS meta-analysis and higher fasting glucose previously.²⁷ The direction of effect was consistent with the *MADD* nonsense mutation rs35

233100, which has been predicted to cause a loss of function and was associated with decreased proinsulin (Figure S9). These data suggest that rs10501320 may contribute to allele-specific differences in *MADD* transcriptional activity in islets. The direction of effect was consistent with the *MADD* nonsense mutation rs35233100, which has been predicted to cause a loss of function and was associated with decreased proinsulin (Figure S9).¹⁰ These data suggest that rs10501320 may contribute to allele-specific differences in *MADD* transcriptional activity in islets and further suggest that *MADD* is a causal transcript at this multi-gene locus.^{10,66}

Discussion

These genetic analyses of circulating proinsulin levels, based on large GWAS meta-analyses, identified 36 signals at 30 loci. We identified 12 previously reported proinsulin loci and 18 additional proinsulin loci. We replicate associations with low-frequency variants at *TBC1D30*, *SGSM2*, and *MADD*, loci that had previously been reported in an exome array analysis in a single cohort.¹⁰ The only previously described proinsulin locus that our study did not replicate was one reported as a cohort-specific signal near *SV2B* (p value = 0.17).¹¹ Characterization of these loci through eQTL colocalization, coding and regulatory annotation, and nearby gene function (Tables S11–S14) provided candidate genes that may influence insulin processing and secretion.

Understanding how glycemic trait signals influence proinsulin can help elucidate potential pathways by which the variants may ultimately influence T2D. We identified five plausible broad groups of encoded proteins: prohormone




Figure 2. The ANK1/NKX6-3 locus associations with proinsulin, T2D, and adipose ANK1 expression

The proinsulin signal at this locus colocalizes with the second AGEN T2D signal and the METSIM adipose ANK1 eQTL signal (HyPrColoc PPFC = 0.92). We used approximate conditional analysis results for the AGEN second signal in HyPrColoc as well as for the plot shown above. AGEN results colored by ASN 1000 Genomes LD reference.

convertases, beta-cell transcription, G-protein modulators, regulation of cytoskeleton dynamics, and lysosomal maturation/endosome recycling (Tables S11 and S14). In the first

group, we include genes *PCSK1* and *PCSK2*, encoding the prohormone convertases PCSK1/3 and PCK2 that are respectively responsible for cleaving the B-chain and




Figure 3. Candidate variants may influence regulatory activity

(A) Regulatory element enrichment analyses using enhancers, accessible chromatin, and other data from islets, skeletal muscle, and adipose. Proinsulin variants are enriched in islet active enhancers and accessible chromatin, especially in beta cells.

(B) The *MADD* locus in proinsulin, lead variant rs10501320. The *MADD* region is an area of extensive LD—the full locus is shown in Figure S4.

(C) The lead variant of the primary *MADD* signal is located in an intron of *MADD* and is in accessible chromatin in islets and an enhancer state and a region conserved across species.

(D) A 411-bp genomic element spanning the lead variant rs10501320 showed strong enhancer activity in a transcriptional reporter assay in two beta cell lines: MIN6 and 832/13. EV, empty vector; G/C, alleles at the lead variant rs10501320. In the eQTL and GWAS data, the G allele at rs10501320 that showed higher transcriptional activity showed higher *MADD* expression levels in islets and is associated with higher proinsulin. Bars show standard errors; p values correspond to two-sided t tests.

A-chain from the C-peptide during proinsulin processing to insulin. While targeted studies have implicated an association between genetic variants in *PCSK2* and glucose homeostasis and T2D,⁶⁷ the association had not yet reached significance in a GWAS with T2D or other glycemic traits, and one study had suggested that *PCSK2* did not significantly impact the beta cells' ability to produce mature insulin.⁶⁸ We now demonstrate that the association reaches genome-wide significance in proinsulin, supporting a significant role for *PCSK2* in beta cells during the processing of proinsulin to insulin. The second group includes candidate genes implicated in beta-cell differentiation (*BARHL1* at the *DDX31* locus, *JARID2*, *NKX6-3*, *SIX2*, and *SIX3*) or the activation and maintenance of beta-cell transcription (*BCL11A*, *C2CD4B*, *TCF7L2*, and *TLE1*). For example,

JARID2 has been shown to play a role in pancreatic and endocrine cell differentiation and beta-cell mass in mouse embryos.^{69–71} The third group consists of genes mediating vesicle translocation and membrane fusion events by affecting the activity of small G proteins, such as Rab and Rho GTPases. *DLC1*, at the *DLC1* locus, encodes a GTPase-activating protein that promotes actin polymerization through regulating the Rho/Rock1 and is modulated by insulin-responsive pathways.^{72,73} The three remaining loci in this group are established proinsulin loci whose nearby genes have been described previously (*MADD*, *SGSM2*, and *TBC1D30*).¹⁰ The fourth group is comprised of genes affecting the cytoskeleton, which undergoes dynamic changes during the processing and secretion of proinsulin at basal and stimulated states: *ANK1*, *KANK1*,

LRR49, and *RNF6*. *KANK1* promotes exocytotic events by mediating actin polymerization;⁷⁴ *LRR49* at the *LARP6* locus is a member of the tubulin polyglutamylase complex;⁷⁵ and *RNF6* is an E3 ubiquitin-protein ligase that regulates actin remodeling.^{76,77} Finally, the fifth group includes genes (*ELAPOR1*, *SNX7*, *STX16*, *TPD52*, *WIPI1*, and *ARSG*) implicated in endosome recycling and lysosomal maturation. In the beta cells, proinsulin is degraded in autophagosome-derived lysosomes via an endocytotic pathway that contributes to the tight regulation of insulin secretion and glucose homeostasis.^{78,79} Both *SNX7* (encoding a sorting nexin⁸⁰) and *WIPI1* (encoding a WD40 repeat protein) play a role in forming autophagosome and transiting autophagosome to early endosome.^{81,82} *STX16* encodes a t-SNARE involved in secretory vesicle membrane fusion and endosome recycling in the Golgi.^{83,84} These genes might help further elucidate the mechanisms for insulin synthesis, processing, and secretion.

Previously proposed clusters of T2D loci included two related to insulin deficiency that differed on the basis of the direction of effect of the T2D risk allele on circulating proinsulin levels.^{6–9} The allele associated with higher glucose was associated with higher proinsulin for half the lead variants, including all variants located near genes involved in beta-cell dysfunction and transcriptional regulation (Tables S10, S11, and S14). For the remaining proinsulin loci, the alleles associated with higher glucose were associated with lower proinsulin; many of these variants are located near genes involved in cytoskeleton dynamics, lysosomal maturation, or endosome recycling (e.g., *WIPI1*, *ELAPOR1*, and *RNF6*). Thus, the directions of allelic effect on proinsulin relative to glucose can help distinguish between clusters of T2D loci.^{6–9}

As another approach to identify potential causal genes, we integrated GWAS signals with islet eQTLs through colocalization and SMR analyses. This approach identified four potential candidate genes at three loci that have not previously been reported in proinsulin or any of the T2D and glycemic studies: *SLC2A10*, *SLC7A14*, *WIPI1*, and *ARSG*. Loci that colocalized with eQTL signals of more than one gene, such as *SIX3* and *WIPI1*, could correspond to allelic effects on more than one gene, sequential effects, or effects on both genes for which only one gene is physiologically relevant to the trait. Our eQTL colocalization analyses also showed that the proinsulin signal at the *NKX6-3/ANK1* locus does not colocalize with the primary AGEN T2D signal and *NKX6-3* in islets but rather with the secondary AGEN T2D signal and the *ANK1* eQTL in adipose.^{26,38,39} Larger eQTL datasets and further characterization of their conditionally distinct signals may be valuable to better interpret colocalization with GWAS signals. Together, the several GWAS traits and eQTL colocalizations at this locus suggest that the underlying mechanisms are not yet fully understood. While we attempt to offer plausible candidate genes for all our proinsulin signals, the genes identified through physical proximity to the lead variant, coding variants in the credible

set, islet expression, and literature searches (Tables S11–S14) are predictions; functional work is invaluable to elucidate genes' roles in the proinsulin.

The *SIX3* proinsulin locus was described previously as a T2D and glucose signal in East Asians.^{26,27,85} Both *SIX3* and *SIX2* are highly and specifically expressed in islets, with an iESI score of 10 for both genes. *SIX3* regulates beta-cell development coordinately with *SIX2*, and knock-down of either gene impairs insulin secretion.^{86,87} Despite a common allele frequency (MAF > 0.13 for all 1000 Genomes ancestries) across ancestries and evidence that the lead variant affects transcriptional factor binding and transcriptional activity,⁸⁵ GWAS meta-analyses of T2D and fasting glucose have failed to date to identify an association at p value < 5×10^{-8} in European-ancestry individuals.^{24,27} Our proinsulin results demonstrate that the glycemic associations at this *SIX3* signal are not specific to East Asians (Figure S10).

The primary *STARD10* signal, which colocalized with a T2D^{24–26} signal, also colocalized with both the *STARD10* and *ARAP1* lead islet eQTL signals (Figure S11). The proinsulin-decreasing allele at the *STARD10* lead variant (rs77464186) was associated with decreased expression of both *STARD10* and *ARAP1*. Although the strength of association was stronger with *STARD10* expression (eQTL p value with rs77464186 for *STARD10* expression = 5×10^{-34} versus *ARAP1* expression = 6×10^{-7}), the evidence for colocalization was stronger with *ARAP1* (*ARAP1* r^2 = 0.99, PPFC = 0.9) versus *STARD10* (r^2 = 0.93, PPFC = 0.60). Both *STARD10* and *ARAP1* are highly expressed in islets, with iESI scores of 9 and 7, respectively. The strength and direction of association between proinsulin and *STARD10* were consistent with the evidence that *STARD10* influences insulin granule biosynthesis and insulin processing by binding to phosphatidylinositides; beta-cell deletion of *Stard10* in mice led to impaired insulin secretion while overexpression of *Stard10* improved glucose tolerance in high-fat-fed animals.^{88,89}

Approximate conditional analysis software such as GCTA requires use of a large LD reference panel representative of the study participants. Even among single-ancestry analyses such as this European-only proinsulin meta-analysis, use of different LD reference panels of the same broad European ancestry can result in strikingly different signals. This issue is particularly noticeable in regions with at least one strongly significant signal. For example, at the *MADD* locus (p = 1.4×10^{-165}), GCTA analyses identified nine, 12, or 22 conditionally distinct signals, depending on which reference panel we employed (Table S6). The discrepancy in results led us to report a signal only when we observed it in at least two of three reference panels, reducing the total number of signals in the *MADD* locus to three—all of which had been previously reported to be associated with proinsulin, adding further confidence to the validity of these signals. While identifying conditionally distinct signals with meta-analysis summary results is invaluable, caution in interpretation of signals is warranted.

To identify potential causal variants driving our observed signals that would have been missed in the regular credible sets built by the Bayesian fine-mapping approach from the association results alone, we defined an extended credible set as the union of variants in the Bayesian credible set and variants in high LD with the lead variant ($r^2 > 0.8$ in 1000 Genomes European). This approach recognizes that standard fine-mapping approaches may be mis-calibrated when applied to meta-analyses,⁹⁰ that variants may have been excluded from the meta-analysis because of analytic or technical factors (e.g., indels are not imputed by the Haplotype Reference Consortium or variants with MAF < 0.5%), and that there were variants that were poorly represented in our meta-analysis as a result of factors such as low sample size. The extended credible set approach added 276 variants, including 142 variants that were not included in the meta-analysis and therefore could not have been included in the Bayesian credible set. The extended credible set identified an additional missense variant in *PCSK1* (rs6234), 15 variants that overlap active enhancers in islets, and 24 variants that overlap islet single-nucleotide ATAC-seq cluster peaks. The extended credible sets provide a more comprehensive pool of candidate variants for mechanistic studies.

Integration of proinsulin loci with complementary glyce-mic traits, expression data in trait-relevant tissues, and functional follow-up provide candidate genes for T2D and hypotheses on potential avenues of mechanism for known T2D loci. While these proinsulin meta-analyses include a large sample size, the difficulty and cost in obtaining proin-sulin measurements limits the sample size compared to studies of many other glyce-mic traits. Future research into genetic contributors to proinsulin will benefit from more and more diverse cohorts. Nonetheless, these findings may help accelerate our understanding of T2D disease pa-thology and promote translation into new therapeutics.

Data and code availability

Upon publication, GWAS summary statistics will be available on the MAGIC Investigators website, <https://magicinvestigators.org/downloads/>, and through the Common Metabolic Diseases knowledge portal, <https://hugeamp.org/>.

Supplemental information

Supplemental information can be found online at <https://doi.org/10.1016/j.ajhg.2023.01.002>.

Acknowledgments

The authors thank the studies' investigators, staff members, and participants for their contributions. T2D data were accessed from <https://diagram-consortium.org>, <https://blog.nus.edu.sg/agen/>, and through dbGaP under accession number dbGaP: phs001672.v3.pl. Data on glyce-mic traits have been contributed by MAGIC investigators and have been downloaded from www.magicinvestigators.org.

www.magicinvestigators.org. We are grateful for the eMERGE network for use of their data as an LD reference panel. Assistance with phenotype harmonization and genotype data cleaning was provided by the eMERGE Administrative Coordinating Center (U01HG004603) and the National Center for Biotechnology Information (NCBI). The eMERGE datasets used for the analyses described in this manuscript were obtained from dbGaP at <http://www.ncbi.nlm.nih.gov/gap> through dbGaP accession number dbGaP: phs000888.v1.p1. Full acknowledgments, including participating cohort and personal acknowledgments, can be found in the [supplemental information](#).

Declaration of interests

J.B.M. is an academic associate for Quest Diagnostics Endocrine R&D. M.E.K. is employed by SYNLAB Holding Deutschland GmbH. C.M.L. receives grants from Bayer Ag and Novo Nordisk and her husband works for Vertex. B.Z. is employed at the Swedish Medical Products Agency, SE-751 03 Uppsala, Sweden; the views expressed in this paper are the personal views of the authors and not necessarily the views of the Swedish government agency. B.Z. has not received any funding or benefits from any sponsor for the present work. J.C.F. receives consulting honoraria from Goldfinch Bio and AstraZeneca and speaker honoraria from Novo Nordisk, AstraZeneca, and Merck for research lectures over which he had full control on content. D.A.L. has received support from Medtronic Ltd and Roche Diagnostics for research unrelated to this paper. W.M. reports grants and personal fees from Siemens Diagnostics, grants and personal fees from Aegerion Pharmaceuticals, grants and personal fees from AMGEN, grants and personal fees from AstraZeneca, grants and personal fees from Danone Research, grants and personal fees from Sanofi, personal fees from Hoffmann LaRoche, personal fees from MSD, grants and personal fees from Pfizer, personal fees from Synageva, grants and personal fees from BASF, grants from Abbott Diagnostics, and grants and personal fees from Numares, outside the submitted work. W.M. is employed by Synlab Holding Deutschland GmbH. R.W. reports lecture fees from Novo Nordisk and Sanofi and served on an advisory board for Akcea Therapeutics, Daiichi Sankyo, Sanofi, and Novo Nordisk. E.W. is now an employee of AstraZeneca.

Received: August 3, 2022

Accepted: January 3, 2023

Published: January 23, 2023

References

1. Porte, D. (1991). Banting lecture 1990. Beta-cells in type II diabetes mellitus. *Diabetes* *40*, 166–180.
2. Ward, W.K., Bolgiano, D.C., McKnight, B., Halter, J.B., and Porte, D. (1984). Diminished B cell secretory capacity in patients with noninsulin-dependent diabetes mellitus. *J. Clin. Invest.* *74*, 1318–1328.
3. Mezza, T., Ferraro, P.M., Sun, V.A., Moffa, S., Cefalo, C.M.A., Quero, G., Cinti, F., Sorice, G.P., Pontecorvi, A., Folli, F., et al. (2018). Increased β -cell workload modulates proinsulin-to-insulin ratio in humans. *Diabetes* *67*, 2389–2396.
4. Liu, M., Weiss, M.A., Arunagiri, A., Yong, J., Rege, N., Sun, J., Haataja, L., Kaufman, R.J., and Arvan, P. (2018). Biosynthesis, structure, and folding of the insulin precursor protein. *Diabetes, Obes. Metab.* *20*, 28–50.

5. Strawbridge, R.J., Dupuis, J., Prokopenko, I., Barker, A., Ahlqvist, E., Rybin, D., Petrie, J.R., Travers, M.E., Bouatia-Naji, N., Dimas, A.S., et al. (2011). Genome-wide association identifies nine common variants associated with fasting proinsulin levels and provides new insights into the pathophysiology of type 2 diabetes. *Diabetes* 60, 2624–2634.
6. Udler, M.S., Kim, J., von Grotthuss, M., Bonàs-Guarch, S., Cole, J.B., Chiou, J., Christopher D Anderson on behalf of METASTROKE and the ISGC, Boehnke, M., Laakso, M., Atzmon, G., et al. (2018). Type 2 diabetes genetic loci informed by multi-trait associations point to disease mechanisms and subtypes: A soft clustering analysis. *PLoS Med.* 15, e1002654.
7. Mansour Aly, D., Dwivedi, O.P., Prasad, R.B., Käräjämäki, A., Hjort, R., Thangam, M., Åkerlund, M., Mahajan, A., Udler, M.S., Florez, J.C., et al. (2021). Genome-wide association analyses highlight etiological differences underlying newly defined subtypes of diabetes. *Nat. Genet.* 53, 1534–1542.
8. DiCorpo, D., LeClair, J., Cole, J.B., Sarnowski, C., Ahmadizar, F., Bielak, L.F., Blokstra, A., Bottinger, E.P., Chaker, L., Chen, Y.-D.I., et al. (2022). Type 2 diabetes partitioned polygenic scores associate with disease outcomes in 454, 193 individuals across 13 cohorts. *Diabetes Care* 45, 674–683.
9. Wesolowska-Andersen, A., Brorsson, C.A., Bizzotto, R., Mari, A., Tura, A., Koivula, R., Mahajan, A., Vinuela, A., Tajes, J.F., Sharma, S., et al. (2022). Four groups of type 2 diabetes contribute to the etiological and clinical heterogeneity in newly diagnosed individuals: An IMI DIRECT study. *Cell Rep. Med.* 3, 100477.
10. Huyghe, J.R., Jackson, A.U., Fogarty, M.P., Buchkovich, M.L., Stančáková, A., Stringham, H.M., Sim, X., Yang, L., Fuchsberger, C., Cederberg, H., et al. (2013). Exome array analysis identifies new loci and low-frequency variants influencing insulin processing and secretion. *Nat. Genet.* 45, 197–201.
11. Strawbridge, R.J., Silveira, A., Hoed, M.d., Gustafsson, S., Luan, J., Rybin, D., Dupuis, J., Li-Gao, R., Kavousi, M., Dehghan, A., et al. (2017). Identification of a novel proinsulin-associated SNP and demonstration that proinsulin is unlikely to be a causal factor in subclinical vascular remodelling using Mendelian randomisation. *Atherosclerosis* 266, 196–204.
12. 1000 Genomes Project Consortium, Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P., Kang, H.M., Korbel, J.O., Marchini, J.L., McCarthy, S., McVean, G.A., and Abecasis, G.R. (2015). A global reference for human genetic variation. *Nature* 526, 68–74.
13. Chien, L.-C. (2020). A rank-based normalization method with the fully adjusted full-stage procedure in genetic association studies. *PLoS One* 15, e0233847.
14. Kang, H.M., Sul, J.H., Service, S.K., Zaitlen, N.A., Kong, S.Y., Freimer, N.B., Sabatti, C., and Eskin, E. (2010). Variance component model to account for sample structure in genome-wide association studies. *Nat. Genet.* 42, 348–354.
15. Zhan, X., Hu, Y., Li, B., Abecasis, G.R., and Liu, D.J. (2016). RVTESTS: An efficient and comprehensive tool for rare variant association analysis using sequence data. *Bioinformatics* 32, 1423–1426.
16. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J., Sklar, P., de Bakker, P.I.W., Daly, M.J., and Sham, P.C. (2007). PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* 81, 559–575.
17. Winkler, T.W., Day, F.R., Croteau-Chonka, D.C., Wood, A.R., Locke, A.E., Mägi, R., Ferreira, T., Fall, T., Graff, M., Justice, A.E., et al. (2014). Quality control and conduct of genome-wide association meta-analyses. *Nat. Protoc.* 9, 1192–1212.
18. Willer, C.J., Li, Y., and Abecasis, G.R. (2010). METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 26, 2190–2191.
19. Yang, J., Lee, S.H., Goddard, M.E., and Visscher, P.M. (2011). GCTA: a tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* 88, 76–82.
20. Yang, J., Genetic Investigation of ANthropometric Traits GIANT Consortium, Ferreira, T., Morris, A.P., Montgomery, G.W., Weedon, M.N., DIAbetes Genetics Replication And Meta-analysis (DIAGRAM) Consortium, Medland, S.E., Madden, P.A.F., Heath, A.C., and Martin, N.G. (2012). Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nat. Genet.* 44, 369–375.
21. Stančáková, A., Paananen, J., Soininen, P., Kangas, A.J., Bonnycastle, L.L., Morken, M.A., Collins, F.S., Jackson, A.U., Boehnke, M.L., Kuusisto, J., et al. (2011). Effects of 34 risk loci for type 2 diabetes or hyperglycemia on lipoprotein subclasses and their composition in 6, 580 nondiabetic Finnish men. *Diabetes* 60, 1608–1616.
22. Rolfe, E.D.L., Loos, R.J.F., Druet, C., Stolk, R.P., Ekelund, U., Griffin, S.J., Forouhi, N.G., Wareham, N.J., and Ong, K.K. (2010). Association between birth weight and visceral fat in adults. *Am. J. Clin. Nutr.* 92, 347–352.
23. McCarty, C.A., Chisholm, R.L., Chute, C.G., Kullo, I.J., Jarvik, G.P., Larson, E.B., Li, R., Masys, D.R., Ritchie, M.D., Roden, D.M., et al. (2011). The eMERGE Network: A consortium of biorepositories linked to electronic medical records data for conducting genomic studies. *BMC Med. Genomics* 4, 13.
24. Mahajan, A., Taliun, D., Thurner, M., Robertson, N.R., Torres, J.M., Rayner, N.W., Payne, A.J., Steinthorsdottir, V., Scott, R.A., Grarup, N., et al. (2018). Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. *Nat. Genet.* 50, 1505–1513.
25. Mahajan, A., Spracklen, C.N., Zhang, W., Ng, M.C.Y., Petty, L.E., Kitajima, H., Yu, G.Z., Rieger, S., Speidel, L., Kim, Y.J., et al. (2022). Multi-ancestry genetic study of type 2 diabetes highlights the power of diverse populations for discovery and translation. *Nat. Genet.* 54, 560–572.
26. Spracklen, C.N., Horikoshi, M., Kim, Y.J., Lin, K., Bragg, F., Moon, S., Suzuki, K., Tam, C.H.T., Tabara, Y., Kwak, S.-H., et al. (2020). Identification of type 2 diabetes loci in 433, 540 East Asian individuals. *Nature* 582, 240–245.
27. Chen, J., Spracklen, C.N., Marenne, G., Varshney, A., Corbin, L.J., Luan, J., Willems, S.M., Wu, Y., Zhang, X., Horikoshi, M., et al. (2021). The trans-ancestral genomic architecture of glycemic traits. *Nat. Genet.* 53, 840–860.
28. Foley, C.N., Staley, J.R., Breen, P.G., Sun, B.B., Kirk, P.D.W., Burgess, S., and Howson, J.M.M. (2021). A fast and efficient colocalization algorithm for identifying shared genetic risk factors across multiple traits. *Nat. Commun.* 12, 764.
29. Giambartolomei, C., Vukcevic, D., Schadt, E.E., Franke, L., Hingorani, A.D., Wallace, C., and Plagnol, V. (2014). Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* 10, e1004383.
30. Carey, V. (2021). *ldblock: data structures for linkage disequilibrium measures in populations.* R Packag. Version 1.24.0.
31. Yin, X., Chan, L.S., Bose, D., Jackson, A.U., VandeHaar, P., Locke, A.E., Fuchsberger, C., Stringham, H.M., Welch, R., Yu, K., et al. (2022). Genome-wide association studies of

- metabolites in Finnish men identify disease-relevant loci. *Nat. Commun.* *13*, 1644.
32. Ghossaini, M., Mountjoy, E., Carmona, M., Peat, G., Schmidt, E.M., Hercules, A., Fumis, L., Miranda, A., Carvalho-Silva, D., Buniello, A., et al. (2021). Open Targets Genetics: systematic identification of trait-associated genes using large-scale genetics and functional genomics. *Nucleic Acids Res.* *49*, D1311–D1320.
 33. Mountjoy, E., Schmidt, E.M., Carmona, M., Schwartzentruber, J., Peat, G., Miranda, A., Fumis, L., Hayhurst, J., Buniello, A., Karim, M.A., et al. (2021). An open approach to systematically prioritize causal variants and genes at all published human GWAS trait-associated loci. *Nat. Genet.* *53*, 1527–1533.
 34. Buniello, A., MacArthur, J.A.L., Cerezo, M., Harris, L.W., Hayhurst, J., Malangone, C., McMahon, A., Morales, J., Mountjoy, E., Sollis, E., et al. (2019). The NHGRI-EBI GWAS Catalog of published genome-wide association studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res.* *47*, D1005–D1012.
 35. Zhou, W., Nielsen, J.B., Fritsche, L.G., Dey, R., Gabrielsen, M.E., Wolford, B.N., LeFaive, J., VandeHaar, P., Gagliano, S.A., Gifford, A., et al. (2018). Efficiently controlling for case-control imbalance and sample relatedness in large-scale genetic association studies. *Nat. Genet.* *50*, 1335–1341.
 36. Kurki, M.I., Karjalainen, J., Palta, P., Sipilä, T.P., Kristiansson, K., Donner, K., Reeve, M.P., Laivuori, H., Aavikko, M., Kainisto, M.A., et al. (2022). FinnGen: Unique genetic insights from combining isolated population and national health register data. Preprint at medRxiv.
 37. Varshney, A., Scott, L.J., Welch, R.P., Erdos, M.R., Chines, P.S., Narisu, N., Albanus, R.D., Orchard, P., Wolford, B.N., Kursawe, R., et al. (2017). Genetic regulatory signatures underlying islet gene expression and type 2 diabetes. *Proc. Natl. Acad. Sci. USA.* *114*, 2301–2306.
 38. Viñuela, A., Varshney, A., van de Bunt, M., Prasad, R.B., Asplund, O., Bennett, A., Boehnke, M., Brown, A.A., Erdos, M.R., Fadista, J., et al. (2020). Genetic variant effects on gene expression in human pancreatic islets and their implications for T2D. *Nat. Commun.* *11*, 4912.
 39. Raulerson, C.K., Ko, A., Kidd, J.C., Currin, K.W., Brotman, S.M., Cannon, M.E., Wu, Y., Spracklen, C.N., Jackson, A.U., Stringham, H.M., et al. (2019). Adipose tissue gene expression associations reveal hundreds of candidate genes for cardiometabolic traits. *Am. J. Hum. Genet.* *105*, 773–787.
 40. Pruim, R.J., Welch, R.P., Sanna, S., Teslovich, T.M., Chines, P.S., Gliedt, T.P., Boehnke, M., Abecasis, G.R., Willer, C.J., and Frisman, D. (2010). LocusZoom: Regional visualization of genome-wide association scan results. *Bioinformatics* *26*, 2336–2337.
 41. Zhu, Z., Zhang, F., Hu, H., Bakshi, A., Robinson, M.R., Powell, J.E., Montgomery, G.W., Goddard, M.E., Wray, N.R., Visscher, P.M., and Yang, J. (2016). Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* *48*, 481–487.
 42. Wellcome Trust Case Control Consortium, Maller, J.B., McVean, G., Byrnes, J., Vukcevic, D., Palin, K., Su, Z., Howson, J.M.M., Auton, A., Myers, S., et al. (2012). Bayesian refinement of association signals for 14 loci in 3 common diseases. *Nat. Genet.* *44*, 1294–1301.
 43. Neph, S., Kuehn, M.S., Reynolds, A.P., Haugen, E., Thurman, R.E., Johnson, A.K., Rynes, E., Maurano, M.T., Vierstra, J., Thomas, S., et al. (2012). BEDOPS: high-performance genomic feature operations. *Bioinformatics* *28*, 1919–1920.
 44. McLaren, W., Pritchard, B., Rios, D., Chen, Y., Flicek, P., and Cunningham, F. (2010). Deriving the consequences of genomic variants with the Ensembl API and SNP effect predictor. *Bioinformatics* *26*, 2069–2070.
 45. Ng, P.C., and Henikoff, S. (2003). SIFT: Predicting amino acid changes that affect protein function. *Nucleic Acids Res.* *31*, 3812–3814.
 46. Adzhubei, I.A., Schmidt, S., Peshkin, L., Ramensky, V.E., Gerasimova, A., Bork, P., Kondrashov, A.S., and Sunyaev, S.R. (2010). A method and server for predicting damaging missense mutations. *Nat. Methods* *7*, 248–249.
 47. Rentzsch, P., Schubach, M., Shendure, J., and Kircher, M. (2021). CADD-Splice-improving genome-wide variant effect prediction using deep learning-derived splice scores. *Genome Med.* *13*, 31.
 48. Kircher, M., Witten, D.M., Jain, P., O’Roak, B.J., Cooper, G.M., and Shendure, J. (2014). A general framework for estimating the relative pathogenicity of human genetic variants. *Nat. Genet.* *46*, 310–315.
 49. Reva, B., Antipin, Y., and Sander, C. (2011). Predicting the functional impact of protein mutations: application to cancer genomics. *Nucleic Acids Res.* *39*, e118.
 50. Varshney, A., Kyono, Y., Elangovan, V.R., Wang, C., Erdos, M.R., Narisu, N., Albanus, R.D., Orchard, P., Stitzel, M.L., Collins, F.S., et al. (2021). A transcription start site map in human pancreatic islets reveals functional regulatory signatures. *Diabetes* *70*, 1581–1591.
 51. Cannon, M.E., Currin, K.W., Young, K.L., Perrin, H.J., Vadlamudi, S., Safi, A., Song, L., Wu, Y., Wabitsch, M., Laakso, M., et al. (2019). Open chromatin profiling in adipose tissue marks genomic regions with functional roles in cardiometabolic traits. *G3 (Bethesda)* *9*, 2521–2533.
 52. Rai, V., Quang, D.X., Erdos, M.R., Cusanovich, D.A., Daza, R.M., Narisu, N., Zou, L.S., Didion, J.P., Guan, Y., Shendure, J., et al. (2020). Single-cell ATAC-Seq in human pancreatic islets and deep learning upscaling of rare cells reveals cell-specific type 2 diabetes regulatory signatures. *Mol. Metab.* *32*, 109–121.
 53. Miguel-Escalada, I., Bonàs-Guarch, S., Cebola, I., Ponsa-Cobas, J., Mendieta-Esteban, J., Atla, G., Javierre, B.M., Rolando, D.M.Y., Farabella, I., Morgan, C.C., et al. (2019). Human pancreatic islet three-dimensional chromatin architecture provides insights into the genetics of type 2 diabetes. *Nat. Genet.* *51*, 1137–1148.
 54. Schmidt, E.M., Zhang, J., Zhou, W., Chen, J., Mohlke, K.L., Chen, Y.E., and Willer, C.J. (2015). GREGOR: evaluating global enrichment of trait-associated variants in epigenomic features using a systematic, data-driven approach. *Bioinformatics* *31*, 2601–2606.
 55. Fogarty, M.P., Cannon, M.E., Vadlamudi, S., Gaulton, K.J., and Mohlke, K.L. (2014). Identification of a regulatory variant that binds FOXA1 and FOXA2 at the CDC123/CAMK1D type 2 diabetes GWAS locus. *PLoS Genet.* *10*, e1004633.
 56. Imai, A., Ishida, M., Fukuda, M., Nashida, T., and Shimomura, H. (2013). MADD/DENN/Rab3GEP functions as a guanine nucleotide exchange factor for Rab27 during granule exocytosis of rat parotid acinar cells. *Arch. Biochem. Biophys.* *536*, 31–37.
 57. Bailyes, E.M., Shennan, K.I., Seal, A.J., Smeekens, S.P., Steiner, D.F., Hutton, J.C., and Docherty, K. (1992). A member of the eukaryotic subtilisin family (PC3) has the enzymic properties

- of the type 1 proinsulin-converting endopeptidase. *Biochem. J.* 285, 391–394.
58. Davidson, H.W., Rhodes, C.J., and Hutton, J.C. (1988). Intraorganellar calcium and pH control proinsulin cleavage in the pancreatic beta cell via two distinct site-specific endopeptidases. *Nature* 333, 93–96.
 59. Scott, L.J., Erdos, M.R., Huyghe, J.R., Welch, R.P., Beck, A.T., Wolford, B.N., Chines, P.S., Didion, J.P., Narisu, N., Stringham, H.M., et al. (2016). The genetic regulatory signature of type 2 diabetes in human skeletal muscle. *Nat. Commun.* 7, 11764.
 60. Klimentidis, Y.C., Arora, A., Newell, M., Zhou, J., Ordovas, J.M., Renquist, B.J., and Wood, A.C. (2020). Phenotypic and genetic characterization of lower LDL cholesterol and increased type 2 diabetes risk in the UK biobank. *Diabetes* 69, 2194–2205.
 61. Garber, M., Guttman, M., Clamp, M., Zody, M.C., Friedman, N., and Xie, X. (2009). Identifying novel constrained elements by exploiting biased substitution patterns. *Bioinformatics* 25, i54–i62.
 62. Huber, C.D., Kim, B.Y., and Lohmueller, K.E. (2020). Population genetic models of GERP scores suggest pervasive turnover of constrained sites across mammalian evolution. *PLoS Genet.* 16, e1008827.
 63. Ansarullah, Jain, C., Far, F.F., Homberg, S., Wißmiller, K., von Hahn, F.G., Raducanu, A., Schirge, S., Sterr, M., Bilekova, S., et al. (2021). Inceptor counteracts insulin signalling in β -cells to control glycaemia. *Nature* 590, 326–331.
 64. Møller, A.B., Kampmann, U., Hedegaard, J., Thorsen, K., Nordentoft, I., Vendelbo, M.H., Møller, N., and Jessen, N. (2017). Altered gene expression and repressed markers of autophagy in skeletal muscle of insulin resistant patients with type 2 diabetes. *Sci. Rep.* 7, 43775.
 65. Wagner, R., Dudziak, K., Herzberg-Schäfer, S.A., Machicao, F., Stefan, N., Staiger, H., Häring, H.U., and Fritsche, A. (2011). Glucose-raising genetic variants in MADD and ADCY5 impair conversion of proinsulin to insulin. *PLoS One* 6, e23639.
 66. Cornes, B.K., Brody, J.A., Nikpoor, N., Morrison, A.C., Chu, H., Ahn, B.S., Wang, S., Dauriz, M., Barzilay, J.I., Dupuis, J., et al. (2014). Association of levels of fasting glucose and insulin with rare variants at the chromosome 11p11.2-MADD locus: cohorts for heart and aging research in genomic epidemiology (CHARGE) consortium targeted sequencing study. *Circ. Cardiovasc. Genet.* 7, 374–382.
 67. Chang, T.-J., Chiu, Y.-F., Sheu, W.H.-H., Shih, K.-C., Hwu, C.-M., Quertermous, T., Jou, Y.-S., Kuo, S.-S., Chang, Y.-C., and Chuang, L.-M. (2015). Genetic polymorphisms of PCSK2 are associated with glucose homeostasis and progression to type 2 diabetes in a Chinese population. *Sci. Rep.* 5, 14380.
 68. Ramzy, A., Asadi, A., and Kieffer, T.J. (2020). Revisiting proinsulin processing: evidence that human β -cells process proinsulin with prohormone convertase (PC) 1/3 but Not PC2. *Diabetes* 69, 1451–1462.
 69. Cervantes, S., Fontcuberta-PiSunyer, M., Servitja, J.-M., Fernandez-Ruiz, R., García, A., Sanchez, L., Lee, Y.-S., Gomis, R., and Gasa, R. (2017). Late-stage differentiation of embryonic pancreatic β -cells requires Jarid2. *Sci. Rep.* 7, 11643.
 70. Soyer, J., Flasse, L., Raffelsberger, W., Beucher, A., Orvain, C., Peers, B., Ravassard, P., Vermot, J., Voz, M.L., Mellitzer, G., and Gradwohl, G. (2010). Rfx6 is an Ngn3-dependent winged helix transcription factor required for pancreatic islet cell development. *Development* 137, 203–212.
 71. White, P., May, C.L., Lamounier, R.N., Brestelli, J.E., and Kaestner, K.H. (2008). Defining pancreatic endocrine precursors and their descendants. *Diabetes* 57, 654–668.
 72. Liao, Y.-C., and Lo, S.H. (2008). Deleted in liver cancer-1 (DLC-1): a tumor suppressor not just for liver. *Int. J. Biochem. Cell Biol.* 40, 843–847.
 73. Hers, I., Wherlock, M., Homma, Y., Yagisawa, H., and Tavaré, J.M. (2006). Identification of p122RhoGAP (deleted in liver cancer-1) Serine 322 as a substrate for protein kinase B and ribosomal S6 kinase in insulin-stimulated cells. *J. Biol. Chem.* 281, 4762–4770.
 74. Rafiq, N.B.M., Nishimura, Y., Plotnikov, S.V., Thiagarajan, V., Zhang, Z., Shi, S., Natarajan, M., Viasnoff, V., Kanchanawong, P., Jones, G.E., and Bershadsky, A.D. (2019). A mechano-signalling network linking microtubules, myosin IIA filaments and integrin-based adhesions. *Nat. Mater.* 18, 638–649.
 75. Wang, L., Paudyal, S.C., Kang, Y., Owa, M., Liang, F.-X., Spector, A., Knaut, H., Sánchez, I., and Dynlacht, B.D. (2022). Regulators of tubulin polyglutamylation control nuclear shape and cilium disassembly by balancing microtubule and actin assembly. *Cell Res.* 32, 190–209.
 76. Liu, L., Zhang, Y., Wong, C.C., Zhang, J., Dong, Y., Li, X., Kang, W., Chan, F.K.L., Sung, J.J.Y., and Yu, J. (2018). RNF6 promotes colorectal cancer by activating the Wnt/ β -catenin pathway via ubiquitination of TLE3. *Cancer Res.* 78, 1958–1971.
 77. Tursun, B., Schlüter, A., Peters, M.A., Viehweger, B., Ostendorff, H.P., Soosairajah, J., Drung, A., Bossenz, M., Johnsen, S.A., Schweizer, M., et al. (2005). The ubiquitin ligase Rnf6 regulates local LIM kinase 1 levels in axonal growth cones. *Genes Dev.* 19, 2307–2319.
 78. Riahi, Y., Wikstrom, J.D., Bachar-Wikstrom, E., Polin, N., Zucker, H., Lee, M.-S., Quan, W., Haataja, L., Liu, M., Arvan, P., et al. (2016). Autophagy is a major regulator of beta cell insulin homeostasis. *Diabetologia* 59, 1480–1491.
 79. Zhou, Y., Liu, Z., Zhang, S., Zhuang, R., Liu, H., Liu, X., Qiu, X., Zhang, M., Zheng, Y., Li, L., et al. (2020). RILP restricts insulin secretion through mediating lysosomal degradation of proinsulin. *Diabetes* 69, 67–82.
 80. Antón, Z., Betin, V.M.S., Simonetti, B., Traer, C.J., Attar, N., Cullen, P.J., and Lane, J.D. (2020). A heterodimeric SNX4–SNX7 SNX-BAR autophagy complex coordinates ATG9A trafficking for efficient autophagosome assembly. *J. Cell Sci.* 133, jcs246306.
 81. Velikkakath, A.K.G., Nishimura, T., Oita, E., Ishihara, N., and Mizushima, N. (2012). Mammalian Atg2 proteins are essential for autophagosome formation and important for regulation of size and distribution of lipid droplets. *Mol. Biol. Cell* 23, 896–909.
 82. Tsuyuki, S., Takabayashi, M., Kawazu, M., Kudo, K., Watanabe, A., Nagata, Y., Kusama, Y., and Yoshida, K. (2014). Detection of WIPI1 mRNA as an indicator of autophagosome formation. *Autophagy* 10, 497–513.
 83. Hastoy, B., Clark, A., Rorsman, P., and Lang, J. (2017). Fusion pore in exocytosis: More than an exit gate? A β -cell perspective. *Cell Calcium* 68, 45–61.
 84. Mallard, F., Tang, B.L., Galli, T., Tenza, D., Saint-Pol, A., Yue, X., Antony, C., Hong, W., Goud, B., and Johannes, L. (2002). Early/recycling endosomes-to-TGN transport involves two SNARE complexes and a Rab6 isoform. *J. Cell Biol.* 156, 653–664.

85. Spracklen, C.N., Shi, J., Vadlamudi, S., Wu, Y., Zou, M., Raulerson, C.K., Davis, J.P., Zeynalzadeh, M., Jackson, K., Yuan, W., et al. (2018). Identification and functional analysis of glycaemic trait loci in the China Health and Nutrition Survey. *PLoS Genet.* *14*, e1007275.
86. Velazco-Cruz, L., Goedegebuure, M.M., Maxwell, K.G., Augsornworawat, P., Hoglebe, N.J., and Millman, J.R. (2020). SIX2 regulates human β cell differentiation from stem cells and functional maturation in vitro. *Cell Rep.* *31*, 107687.
87. Bevacqua, R.J., Lam, J.Y., Peiris, H., Whitener, R.L., Kim, S., Gu, X., Friedlander, M.S.H., and Kim, S.K. (2021). SIX2 and SIX3 coordinately regulate functional maturity and fate of human pancreatic β cells. *Genes Dev.* *35*, 234–249.
88. Carrat, G.R., Haythorne, E., Tomas, A., Haataja, L., Müller, A., Arvan, P., Piunti, A., Cheng, K., Huang, M., Pullen, T.J., et al. (2020). The type 2 diabetes gene product STARD10 is a phosphoinositide-binding protein that controls insulin secretory granule biogenesis. *Mol. Metab.* *40*, 101015.
89. Carrat, G.R., Hu, M., Nguyen-Tu, M.-S., Chabosseau, P., Gaulton, K.J., van de Bunt, M., Siddiq, A., Falchi, M., Thurner, M., Canouil, M., et al. (2017). Decreased STARD10 expression is associated with defective insulin secretion in humans and mice. *Am. J. Hum. Genet.* *100*, 238–256.
90. Tsuo, K., Zhou, W., Wang, Y., Kanai, M., Namba, S., Gupta, R., Majara, L., Nkambule, L.L., Morisaki, T., Okada, Y., et al. (2022). Meta-analysis fine-mapping is often miscalibrated at single-variant resolution. *Cell Genomics* *2*, 100212.