

Efficient Segmental Isotope Labeling of Integral Membrane Proteins for High-Resolution NMR Studies

Melina Daniilidis, Laura E. Sperl, Benedikt S. Müller, Antonia Babl, and Franz Hagn*

Cite This: *J. Am. Chem. Soc.* 2024, 146, 15403–15410


Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information



ABSTRACT: High-resolution structural NMR analyses of membrane proteins are challenging due to their large size, resulting in broad resonances and strong signal overlap. Among the isotope labeling methods that can remedy this situation, segmental isotope labeling is a suitable strategy to simplify NMR spectra and retain high-resolution structural information. However, protein ligation within integral membrane proteins is complicated since the hydrophobic protein fragments are insoluble, and the removal of ligation side-products is elaborate. Here, we show that a stabilized split-intein system can be used for rapid and high-yield protein trans-splicing of integral membrane proteins under denaturing conditions. This setup enables segmental isotope labeling experiments within folded protein domains for NMR studies. We show that high-quality NMR spectra of markedly reduced complexity can be obtained in detergent micelles and lipid nanodiscs. Of note, the nanodisc insertion step specifically selects for the ligated and correctly folded membrane protein and simultaneously removes ligation byproducts. Using this tailored workflow, we show that high-resolution NMR structure determination is strongly facilitated with just two segmentally isotope-labeled membrane protein samples. The presented method will be broadly applicable to structural and dynamical investigations of (membrane-) proteins and their complexes by solution and solid-state NMR but also other structural methods where segmental labeling is beneficial.

INTRODUCTION

Membrane proteins are challenging systems for high-resolution structural analyses. These proteins are difficult to crystallize, and cryo-EM is particularly suited for large membrane protein systems.¹ Small- to medium-sized membrane proteins can be investigated by solution- or solid-state NMR.² However, adverse relaxation properties and limited long-term stability render NMR studies of larger membrane proteins very challenging. The use of high-level isotope-labeled protein and a suitable native-like membrane mimetic, such as lipid nanodiscs, can in part remedy this situation.^{3,4} Despite these important advancements, severe NMR signal overlap with larger proteins still markedly reduces spectral resolution and impedes the data analysis for resonance assignment and structure determination.⁵ This issue is particularly limiting for solid-state NMR experiments, where ¹H line widths are markedly larger.⁶ Amino acid-selective isotope labeling⁷ has been used to simplify the NMR resonance assignment procedure. However, this approach requires the production of multiple samples which can be expensive regarding time and reagent costs. Segmental isotope labeling has turned out to be

effective in simplifying NMR spectra.⁸ For segmental isotope labeling, various protein ligation methods have been reported, such as enzymatic approaches using trans peptidases,⁹ expressed protein ligation,^{10,11} and split-intein mediated trans-splicing.^{12–14} Typically, expressed protein ligation, or native chemical ligation,¹⁵ requires a C-terminal thioester and an N-terminal cysteine. The ligation yield can be further enhanced by using selenium cysteine at the ligation site.¹⁶ However, for expressed protein ligation, at least one fragment is typically of synthetic origin. A major advantage of intein-mediated trans-splicing is the direct use of recombinant protein fragments without the need for specific chemical modifications or synthetic educts for the ligation reaction.¹⁷ While trans-splicing of soluble (multi-) protein domains can be done in

Received: March 6, 2024

Revised: May 10, 2024

Accepted: May 10, 2024

Published: May 24, 2024






Figure 1. Intein-mediated protein trans-splicing for segmental isotope labeling of membrane proteins for NMR studies. (a) Strategy to obtain properly folded membrane proteins in detergent micelles and nanodiscs by protein trans-splicing. (b) Prototype membrane proteins OmpX (2m06.pdb⁴) and MPV17 (Alphafold²⁰ model) with a β -barrel and α -helical topology, respectively. The N- and C-exteins and the splicing sites are indicated. MSP, membrane scaffold protein; MP, membrane protein.




Figure 2. High-yield production of membrane proteins by intein-mediated protein trans-splicing. (a) SDS-PAGE of the splicing reaction of OmpX. (b) ESI-MS data of the splicing reaction indicating the generation of full-length OmpX. (c) Thermal stability of spliced OmpX refolded into DPC micelles. (d) same as in (a) but with the α -helical membrane protein MPV17. (e) ESI-MS data of the intein fragments and the spliced MPV17 product. (f) Same as in (c) but with MPV17 in DPC micelles. (a,d) 0* indicates the ~ 10 s time point right after mixing the two OmpX or MPV17 fragments.

very good yields and thus turned out to be very useful for structural studies by NMR,⁸ the ligation within a compactly folded integral membrane protein remains difficult for various reasons, such as solubility, sample purity as well as ligation efficacy and product yield.¹⁸ Due to these obstacles, it was so far very difficult to produce sufficient amounts of segmentally isotope labeled membrane proteins by intein trans-splicing for high-resolution NMR studies.

Here, we used a highly stable DnaE split-intein system¹⁹ for high-yield protein trans splicing of integral membrane proteins. We showcase this method by performing segmental isotope labeling of an α -helical and a β -barrel integral membrane

protein for high-resolution NMR studies. The obtained NMR spectra are markedly simplified, facilitating NMR spectral analysis and resonance assignment work. In addition, the insertion into lipid nanodiscs can serve as a quality control filter, rendering the time-consuming removal of ligation side products unnecessary. With this optimized workflow, NMR structure determination of integral membrane proteins is markedly facilitated, contributing to expanding the size limit of membrane proteins that are accessible to solution- and solid-state NMR. In addition, this workflow will be beneficial for other methods that rely on the selective integration of labels or chemical probes.

RESULTS

First, we designed suitable protein constructs for intein trans-splicing, where the N-terminal membrane protein fragment is fused to the N-intein and the C-terminal part to the C-intein (Figures 1a, S1, Supporting Information Materials and Methods section). Since the membrane protein fragments are predominantly insoluble in aqueous solution, the ligation reaction needs to be conducted under denaturing conditions, leading to a reduction in the activity of the intein. The Cfa intein used in this study¹⁹ is still very active at high urea or guanidine hydrochloride concentrations and is thus compatible with our ligation strategy. This setup allows for intein-mediated trans-splicing within a compactly folded membrane protein. The ligated full-length membrane protein can then be refolded into detergent micelles and, if desired, subsequently inserted into lipid bilayer nanodiscs of a suitable size.^{4,21} To demonstrate that this approach is feasible for membrane proteins of different secondary structures and topologies, we used the bacterial β -barrel membrane protein OmpX and the human α -helical inner mitochondrial membrane protein MPV17 as model systems (Figure 1b). The trans-splicing activity was reported to be dependent on the C-extein residues directly flanking the intein.¹⁷ To address this requirement, we here placed the splicing sites within loops to be able to incorporate additional residues that are beneficial for high splicing activity without perturbing the protein fold. If the structure of the membrane protein is not known, loop regions can be identified by secondary structure or AlphaFold²⁰ predictions or using secondary chemical shift information from experimental NMR backbone resonance assignment data.²² Here, we incorporated the motif CFN at the C-extein positions 1–3, which is found in the native extein sequence of the naturally occurring split intein DnaE from *Nostoc punctiforme*.²³ Furthermore, we added short linkers next to the exteins that have previously been shown to favor efficient splicing (Figure S1).^{19,23} If the length of a loop cannot be expanded, it is also possible to replace existing loop residues with this optimized amino acid stretch. In addition, it is recommended to evaluate the biophysical and functional properties of the desired protein construct as well as its refolding properties in advance using a full-length protein modified accordingly. For designing suitable OmpX constructs, we used the previously obtained NMR structure in lipid nanodiscs (2m06.pdb⁴). For MPV17, the AlphaFold²⁰ model together with NMR backbone chemical shift information²⁴ was used (Figures 1b, and S1). Each split-intein fusion protein was produced in *E. coli* and purified from inclusion bodies under denaturing conditions. The protein ligation reaction was performed in 6 M urea at 30 °C (Figure 2). With OmpX, the trans-splicing reaction was already close to completion after 15 to 30 min, with a maximum conversion of ~75% determined after 120 min by SDS-PAGE (Figures 2a, and S2). The use of the Cfa intein led to a strongly increased yield and markedly reduced reaction times compared to a native split intein that has been previously used for the ligation of the bacterial outer membrane protein OmpF.¹⁸ Since the bands in the SDS-PAGE of the C-terminal Intein-OmpX construct (Int_C-OmpX_C) and the spliced full-length OmpX product almost completely overlapped, we used mass spectrometry to probe the mass (17 kDa) of the desired full-length OmpX product (Figures 2b, and S3).

Full-length OmpX obtained by intein trans-splicing (Figure S4a) shows identical thermal stability as wild-type OmpX, indicating that the slight sequence modifications required for efficient ligation did not perturb the protein structure (Figures 2c, and S4b). For the inner mitochondrial membrane protein MPV17, the splicing reaction was completed after 60 min, as probed by SDS-PAGE, resulting in a conversion of ~50% after 120 min reaction time (Figures 2d, and S5), which was still enough to produce milligram quantities of the spliced protein. The spliced full-length MPV17 as well as the educts and the cleaved intein fragments were identified by mass spectrometry (Figure 2e). During the splicing reaction, full-length MPV17 precipitated in 6 M urea, whereas the educts remained soluble, which contributed to the high product yield and resulted in a high product purity. Since the soluble fraction was used for SDS-PAGE, the intensity of the band for full-length MPV17 consequently decreased with increasing reaction time (Figures 2d, and S6). As described above for OmpX, the introduction of a short linker for optimal ligation efficiency did not perturb the protein structure, as evident from a similar thermal melting point of intein-ligated MPV17 in DPC micelles compared to wild-type MPV17²⁴ (Figure 2f).

Since some protocols utilize SDS for the solubilization of membrane proteins prior to refolding²⁵ we wondered whether the Cfa-intein trans-splicing reaction can also be performed in 0.5% SDS, a typical concentration that is used for solubilization. As shown in Figure S7, product formation could not be observed, which can be rationalized by unfolding of the intein by this harsh detergent. However, this limitation does not prevent the solubilization of ligated membrane proteins in SDS for subsequent refolding experiments. To achieve this, the urea in the ligation reaction just needs to be removed by dialysis, followed by solubilization of the resulting protein precipitate in SDS. In contrast to intein trans-splicing, expressed protein ligation, not relying on the folding state of an intein, has been shown to be compatible with a harsh detergent environment.¹⁰

A common side product of intein trans-splicing reactions is the C-terminal extein without the C-intein fragment,²⁶ together with unreacted educts. In our setup, we also observed the presence of the C-terminal OmpX fragment without the intein (OmpX_C). His-tagged educts and inteins could be successfully removed by IMAC (Figure S8a). The OmpX_C fragment lacks a His-tag and was consequently copurified with the spliced OmpX product (Figure S8b). Finally, the removal of this unwanted fragment was achieved by elaborate size exclusion chromatography under denaturing conditions (Figure S9).

Next, we applied this tailored and optimized protocol to perform segmental isotope labeling of both membrane proteins. For this, we produced OmpX and MPV17 samples where each extein was either unlabeled (¹H,¹⁴N) or isotope labeled (²H,¹⁵N) and visible in NMR (see Figure 1b). OmpX samples were inserted into size-optimized lipid nanodiscs^{4,21,27} containing a DMPC/DMPG (3:1) lipid blend (Figure S10). MPV17 forms large oligomers in lipids²⁴ which prevented its insertion into lipid nanodiscs. Thus, the purification (Figure S11) and the NMR experiments were instead conducted in DPC micelles as previously described.²⁴

To our positive surprise, the nanodisc assembly step with ligated OmpX served as a protein quality control filter since only compactly folded full-length protein inserted into the lipid environment of a nanodisc (Figures 3a, and S12). This




Figure 3. Removal of ligation side products by insertion into lipid nanodiscs. (a) After intein splicing and refolding, properly folded full-length and a misfolded OmpX_C fragment are copurifying in DPC detergent micelles, giving rise to a 2D-^[15N, 1H]-TROSY spectrum containing signals of both species (blue spectrum in panel b). The black spectrum in (b) is a reference with folded U-^[2H, 15N]-labeled OmpX in DPC micelles. Insertion into lipid nanodiscs selects for the properly folded species and efficiently removes the OmpX_C fragment, indicated by SDS-PAGE and a 2D-^[15N, 1H]-TROSY spectrum lacking signals in the unfolded region (c). NMR spectra were recorded at 318 K and at 950 MHz ¹H frequency with a sample where the C-extein of OmpX (Figure 1b) is labeled with ²H and ¹⁵N (blue spectra). The N-extein is unlabeled and not visible in NMR. Trp sc: tryptophane NHε side chain signals.




Figure 4. 2D-NMR analysis of the segmentally isotope-labeled integral membrane proteins OmpX and MPV17. (a) 2D-^[15N, 1H]-TROSY spectra of uniformly labeled OmpX (gray) and segmentally labeled OmpX (red and blue). (b) Chemical shift perturbations (CSPs) within OmpX calculated from the spectra shown in (a). (c) CSPs mapped onto the structure of OmpX (2m06.pdb⁴). The splicing site is indicated by the two green spheres. (d–f) Same as in (a–c) but with segmentally isotope labeled MPV17 in DPC micelles. The structural model of MPV17 was obtained with AlphaFold.²⁰

approach allowed for skipping the elaborate size exclusion chromatography step under denaturing conditions (Figure S9)

to remove OmpX_C. NMR spectra of spliced OmpX containing ²H,¹⁵N-labeled OmpX_C in detergent micelles (Figure 3b) and




Figure 5. Segmental isotope labeling of membrane proteins facilitates NMR structure determination. (a) Structural interface between $^2\text{H},^{15}\text{N}$ -labeled OmpX-N-extein and unlabeled OmpX-C-extein with observed NOE distance restraints (shown in (b)) within the labeled or to the unlabeled part. (c) and (d), same as in (a) and (b) but with an inverse isotope labeling pattern, i.e., $^2\text{H},^{15}\text{N}$ -labeled OmpX-C-extein and unlabeled OmpX-N-extein. (e) 20 lowest energy structures obtained with the two segmentally labeled OmpX samples in lipid nanodiscs, showing a root mean square deviation (rmsd) of backbone atoms in ordered secondary structure elements of 0.29 Å. The assignment of NOE contacts was facilitated by the lower complexity and signal overlap in the individual spectra.

lipid nanodiscs (Figure 3c) show that the undesired set of signals corresponding to the unfolded OmpX_C fragment in DPC micelles are completely absent in lipid nanodiscs. Thus, the nanodisc insertion procedure can be considered a general and convenient way for efficient product purification, rendering more elaborate purification strategies obsolete.

With this optimized ligation and purification workflow, we next recorded 2D- $^{15}\text{N},^1\text{H}$ -TROSY spectra of segmentally isotope-labeled OmpX (*vide supra*) in lipid nanodiscs (Figure 4a) and compared them with spectra of uniformly $^2\text{H},^{15}\text{N}$ -labeled OmpX. In these spectra, the signals of segmentally labeled OmpX overlaid almost perfectly with the corresponding signals of uniformly labeled OmpX. Slight chemical shift perturbations were only observed in the region around the splicing site and the neighboring loop region (Figure 4b,c).

Overall, the complexity of the spectra is markedly reduced due to a lower number of NMR signals, which is most pronounced in the central region of the spectrum where unfolded protein stretches usually lead to heavy signal overlap. NMR spectra of similar quality were obtained with segmentally labeled MPV17 in detergent micelles (Figure 4d). For this α -helical membrane protein, where signal dispersion is less pronounced leading to strong overlap, the benefit of segmental labeling is even more apparent. In line with the very similar thermal stability data (Figure 2f), the NMR spectra are only slightly perturbed around the splicing site (Figure 4e,f), indicating that the structure of MPV17 is not altered.

Due to the simplification of the NMR spectra of segmentally isotope-labeled proteins, NMR structure determination is considered to be markedly facilitated.^{13,14} Typically, this strategy requires the production of two orthogonally labeled

samples, e.g., only one segment is labeled with NMR-active nuclei with the other one only ^1H labeled and *vice versa*. Here, we used a $^2\text{H},^{15}\text{N}$ versus $^1\text{H},^{14}\text{N}$ labeling pattern (see Figure 4a), which is sufficient for the structure determination of smaller β -barrel membrane proteins⁴ using backbone NOE contacts. However, any isotope labeling pattern on each side can be used if necessary, including selective ^{13}C methyl labeling²⁸ in both fragments to enable the detection of specific contacts within larger integral membrane proteins. For extracting distance restraints, we recorded 3D- ^{15}N -edited-TROSY- $^{15}\text{N},^1\text{H}$ -NOESY NMR experiments with the two segmentally labeled OmpX samples in lipid nanodiscs. As expected, we could observe NOE-connections within the isotope-labeled segment in each sample (Figure 5a–d), e.g., G8-N25 or A67-V83 amide proton NOEs.

In addition, at the edges of each isotope-labeled segment, NOE contacts are visible between an ^{15}N -isotope-labeled amide and any proton in the adjacent unlabeled (^1H) β -stand, e.g., G7-Y146, G7-V144 or G68-I40, G68-V39. This pattern further facilitated the assignment of NOE contacts since not only the amide of a cross- β -stand residue is detected but also its side chain protons. Moreover, additional NOE restraints can be used for structure determination, providing additional structural information in these regions.

With this strategy, we could collect 180 NOE distance restraints to determine a high-resolution structure of OmpX in lipid nanodiscs (Figure 5e) with a backbone atom root mean squared deviation (rmsd) of 0.29 Å (Table 1), which is slightly better than our previous structure based on backbone amide NOEs only.⁴ It can be further envisioned that this approach will be beneficial to unambiguously probe distances across β -

Table 1. Structural Statistics of Segmentally Labeled OmpX in Phospholipid Nanodiscs^a

structural information	
NOEs (amide and side-chain contacts)	180
Hydrogen bond restraints	77
Dihedral angle restraints (TALOS ³⁰)	291
Backbone rmsd in β -strands (Å) ^b	0.29 ± 0.05
Backbone rmsd for all residues (Å)	1.12 ± 0.29
Ramachandran map analysis ^c	
Most favored regions	93.4%
Additionally allowed regions	4.1%
Generously allowed regions	2.5%
Disallowed regions	0.0%
Deviations from restraints and idealized geometry	
Distance restraints (Å)	0.11 ± 0.03
Dihedral angle restraints (°)	0.19 ± 0.02
Bonds (Å)	0.0024 ± 0.00008
Angles (°)	0.66 ± 0.01
Impropers (°)	1.91 ± 0.13

^aAnalysis of the 20 lowest-energy structures. ^bOnly ordered secondary structure elements were used for structural superimposition: 3–14, 20–30, 38–48, 60–71, 78–90, 104–115, 122–132, 135–147; rmsd values are calculated relative to a nonminimized average structure of each ensemble. ^cRamachandran analysis with PROCHECK-NMR³¹ was performed on the lowest-energy structure.

barrels or α -helical bundles between selectively labeled protein fragments using orthogonal isotope labeling patterns in larger membrane proteins. Side-chain methyl group spectra typically show heavy overlap but are essential to determine accurate tertiary structures, even for β -barrel membrane proteins.²⁹

DISCUSSION

This study provides the first example where high-resolution NMR structure determination was conducted with a segmental isotope labeled integral membrane protein. Segmental isotope labeling for NMR has been used for decades but is limited to soluble proteins or, e.g., attaching unfolded tails to membrane-associated domains.¹⁴ Here, we use the exceptional properties of the Cfa intein¹⁹ to perform protein ligation by intein-splicing within an integral membrane protein. In comparison to a previous study,¹⁸ the exceptionally fast reaction rate and high yield of the protein trans-splicing with the Cfa intein under denaturing conditions enables the production of milligram quantities of segmental isotope-labeled membrane proteins and renders this approach feasible for a wide range of membrane protein systems where refolding is possible. To obtain optimal protein ligation yields we here used naturally occurring sequence elements between the inteins and the exteins. This strategy requires the positioning of the splicing sites in loop regions of the membrane protein. However, recent improvements of the Cfa intein³² render traceless splicing more efficient with markedly reduced sequence requirements at the splicing site. If a particular membrane protein system requires segmental isotope labeling of an internal fragment, orthogonal split inteins can in principle be used to ligate three segments.³³ However, since every additional ligation step reduces the overall protein yield, the design of fast and efficient orthogonal split inteins that are active under denaturing conditions will be necessary to make this approach usable for NMR sample preparation.

NMR investigations of membrane proteins require not only high-field instrumentation and optimized experimental setup but also cutting-edge biochemical sample production methods. The use of any isotope-labeling strategy that simplifies the NMR spectral signature and renders the NMR resonance assignment procedure less elaborate is essential for enabling investigations of the structure, interactions, and dynamics of membrane proteins of increasing complexity. An important class of integral membrane proteins are G-protein coupled receptors (GPCRs), where in some cases the production in *E. coli* and refolding was shown to be possible³⁴ although high-level isotope labeling and NMR resonance assignments are still challenging.³⁵ Thus, our approach might be used to specifically visualize functional elements in such larger membrane proteins to probe their conformational states and interactions with small molecules and partner proteins.

While we here used solution-state NMR, we anticipate that our method will also be highly beneficial for solid-state NMR studies of membrane proteins. At ultra-fast magic angle spinning conditions and at ultra-high magnetic field, solid-state NMR is in principle size-independent,^{6,36} enabling investigations of very large systems. However, the increasing number of resonances in large proteins impedes the spectral analysis due to heavy signal overlap. The ability to segmentally label selected regions of interest will thus also be essential for structural studies of larger membrane proteins by solid-state NMR. In addition to NMR studies, the presented approach can be further used to selectively attach spin labels to desired protein segments within a folded (membrane) protein domain for electron paramagnetic resonance (EPR) spectroscopy. This procedure renders mutagenesis of surface-exposed reactive amino acid residues in parts of the protein where no labeling is desired obsolete.³⁷ Furthermore, segmental isotope labeling of membrane proteins might be useful for small-angle neutron scattering (SANS) experiments,³⁸ where contrast matching can be employed to selectively observe conformational changes of a particular structural element within a folded membrane protein. Hence, we believe that the presented method will be broadly applicable to the investigation of the structure, dynamics, and function of integral membrane proteins and other large protein systems.

CONCLUSIONS

We showed that intein-based trans-splicing can be conducted at high yields with integral membrane proteins. This setup opens the possibility for segmental isotope labeling for high-resolution NMR studies, where signal overlap is a main issue that prevents a detailed analysis of larger and more challenging systems. We show that lipid nanodisc insertion provides an efficient platform for selecting the correctly ligated and properly folded membrane protein species. This workflow facilitates the high-resolution structure determination of membrane proteins in a native lipid environment and will be a versatile tool to selectively study the structure and dynamics of functionally important parts of membrane proteins by NMR and other structural methods.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.4c03294>.

Additional experimental details, materials, and methods, including figures on split-intein fusion protein construct design, reaction yields, mass spectrometry, purification of the protein constructs and ligation products, and Supporting Information references (PDF)

AUTHOR INFORMATION

Corresponding Author

Franz Hagn – Bavarian NMR Center, Department of Bioscience, School of Natural Sciences, Technical University of Munich, 85748 Garching, Germany; Institute of Structural Biology, Helmholtz Munich, 85764 Neuherberg, Germany; orcid.org/0000-0002-1315-459X; Email: franz.hagn@tum.de

Authors

Melina Daniilidis – Bavarian NMR Center, Department of Bioscience, School of Natural Sciences, Technical University of Munich, 85748 Garching, Germany; orcid.org/0000-0001-7662-3325

Laura E. Sperl – Bavarian NMR Center, Department of Bioscience, School of Natural Sciences, Technical University of Munich, 85748 Garching, Germany

Benedikt S. Müller – Bavarian NMR Center, Department of Bioscience, School of Natural Sciences, Technical University of Munich, 85748 Garching, Germany

Antonia Babl – Bavarian NMR Center, Department of Bioscience, School of Natural Sciences, Technical University of Munich, 85748 Garching, Germany

Complete contact information is available at:

<https://pubs.acs.org/10.1021/jacs.4c03294>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by funds of the Deutsche Forschungsgemeinschaft (CRC1035, project number 201302640, project B13) and the Technical University of Munich and Helmholtz Munich. We want to thank Drs Gerd Gemmecker and Sam Asami for general NMR support and the Bavarian NMR Center (www.bnmrz.org) for access to high-field NMR instruments. Juliane Scheer (TUM) is gratefully acknowledged for initial experiments.

ABBREVIATIONS

DPC, dodecylphosphocholine; MSP, membrane scaffold protein; MS, mass spectrometry; NMR, nuclear magnetic resonance; SEC, size exclusion chromatography; IMAC, immobilized metal ion affinity chromatography; SDS, sodium dodecyl sulfate; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis

REFERENCES

- (1) Cheng, Y. Membrane protein structural biology in the era of single particle cryo-EM. *Curr. Opin. Struct. Biol.* **2018**, *52*, 58–63.
- (2) Liang, B.; Tamm, L. K. NMR as a tool to investigate the structure, dynamics and function of membrane proteins. *Nat. Struct. Mol. Biol.* **2016**, *23* (6), 468–474.
- (3) Markus, M. A.; Dayie, K. T.; Matsudaira, P.; Wagner, G. Effect of deuteration on the amide proton relaxation rates in proteins. Heteronuclear NMR experiments on villin 14T. *J. Magn. Reson. B* **1994**, *105* (2), 192–195. Daniilidis, M.; Brandl, M. J.; Hagn, F. The Advanced Properties of Circularized MSP Nanodiscs Facilitate High-resolution NMR Studies of Membrane Proteins. *J. Mol. Biol.* **2022**, *434* (24), No. 167861.
- (4) Hagn, F.; Eitzkorn, M.; Raschle, T.; Wagner, G. Optimized phospholipid bilayer nanodiscs facilitate high-resolution structure determination of membrane proteins. *J. Am. Chem. Soc.* **2013**, *135* (5), 1919–1925.
- (5) Wider, G.; Wuthrich, K. NMR spectroscopy of large molecules and multimolecular assemblies in solution. *Curr. Opin. Struct. Biol.* **1999**, *9* (5), 594–601.
- (6) Reif, B.; Ashbrook, S. E.; Emsley, L.; Hong, M. Solid-state NMR spectroscopy. *Nat. Rev. Methods Primers* **2021**, *1*, 2 DOI: [10.1038/s43586-020-00002-1](https://doi.org/10.1038/s43586-020-00002-1).
- (7) Fiaux, J.; Bertelsen, E. B.; Horwich, A. L.; Wuthrich, K. Uniform and residue-specific ¹⁵N-labeling of proteins on a highly deuterated background. *J. Biomol. NMR* **2004**, *29* (3), 289–297. Trbovic, N.; Klammt, C.; Koglin, A.; Lohr, F.; Bernhard, F.; Dotsch, V. Efficient strategy for the rapid backbone assignment of membrane proteins. *J. Am. Chem. Soc.* **2005**, *127* (39), 13504–13505. Tugarinov, V.; Kanelis, V.; Kay, L. E. Isotope labeling strategies for the study of high-molecular-weight proteins by solution NMR spectroscopy. *Nat. Protoc.* **2006**, *1* (2), 749–754. Michel, E.; Skrisovska, L.; Wuthrich, K.; Allain, F. H. Amino acid-selective segmental isotope labeling of multidomain proteins for structural biology. *Chembiochem* **2013**, *14* (4), 457–466.
- (8) Liu, D.; Xu, R.; Cowburn, D. Segmental isotopic labeling of proteins for nuclear magnetic resonance. *Methods Enzymol* **2009**, *462*, 151–175.
- (9) Refaei, M. A.; Combs, A.; Kojetin, D. J.; Cavanagh, J.; Caperelli, C.; Rance, M.; Sapiro, J.; Tsang, P. Observing selected domains in multi-domain proteins via sortase-mediated ligation and NMR spectroscopy. *J. Biomol. NMR* **2011**, *49* (1), 3–7. Freiburger, L.; Sonntag, M.; Hennig, J.; Li, J.; Zou, P.; Sattler, M. Efficient segmental labeling of multi-domain proteins using Sortase A. *J. Biomol. NMR* **2015**, *63* (1), 1–8. Raltchev, K.; Pipercevic, J.; Hagn, F. Production and Structural Analysis of Membrane-Anchored Proteins in Phospholipid Nanodiscs. *Chemistry* **2018**, *24* (21), 5493–5499. Mikula, K. M.; Krumwiede, L.; Pluckthun, A.; Iwai, H. Segmental isotopic labeling by asparaginyl endopeptidase-mediated protein ligation. *J. Biomol. NMR* **2018**, *71* (4), 225–235. Williams, F. P.; Milbradt, A. G.; Embrey, K. J.; Bobby, R. Segmental Isotope Labelling of an Individual Bromodomain of a Tandem Domain BRD4 Using Sortase A. *PLoS One* **2016**, *11* (4), No. e0154607. Boyko, K. V.; Rosenkranz, E. A.; Smith, D. M.; Miers, H. L.; Oued Es Cheikh, M.; Lund, M. Z.; Young, J. C.; Reardon, P. N.; Okon, M.; Smirnov, S. L.; et al. Sortase-mediated segmental labeling: A method for segmental assignment of intrinsically disordered regions in proteins. *PLoS One* **2021**, *16* (10), No. e0258531.
- (10) Valiyaveetil, F. I.; MacKinnon, R.; Muir, T. W. Semisynthesis and folding of the potassium channel KcsA. *J. Am. Chem. Soc.* **2002**, *124* (31), 9113–9120.
- (11) Vitali, F.; Henning, A.; Oberstrass, F. C.; Hargous, Y.; Auweter, S. D.; Erat, M.; Allain, F. H. Structure of the two most C-terminal RNA recognition motifs of PTB using segmental isotope labeling. *EMBO J.* **2006**, *25* (1), 150–162. Schubeis, T.; Yuan, P.; Ahmed, M.; Nagaraj, M.; van Rossum, B. J.; Ritter, C. Untangling a Repetitive Amyloid Sequence: Correlating Biofilm-Derived and Segmentally Labeled Curli Fimbriae by Solid-State NMR Spectroscopy. *Angew. Chem., Int. Ed. Engl.* **2015**, *54* (49), 14669–14672.
- (12) Otomo, T.; Ito, N.; Kyogoku, Y.; Yamazaki, T. NMR observation of selected segments in a larger protein: central-segment isotope labeling through intein-mediated ligation. *Biochemistry* **1999**, *38* (49), 16040–16044. Muona, M.; Aranko, A. S.; Raulinaitis, V.; Iwai, H. Segmental isotopic labeling of multi-domain and fusion proteins by protein trans-splicing in vivo and in vitro. *Nat. Protoc.* **2010**, *5* (3), 574–587.
- (13) Yagi, H.; Tsujimoto, T.; Yamazaki, T.; Yoshida, M.; Akutsu, H. Conformational change of H⁺-ATPase beta monomer revealed on segmental isotope labeling NMR spectroscopy. *J. Am. Chem. Soc.* **2004**, *126* (50), 16632–16638. Skrisovska, L.; Allain, F. H. Improved

segmental isotope labeling methods for the NMR study of multidomain or large proteins: application to the RRM of Npl3p and hnRNP L. *J. Mol. Biol.* **2008**, *375* (1), 151–164.

(14) Gupta, S.; Tycko, R. Segmental isotopic labeling of HIV-1 capsid protein assemblies for solid state NMR. *J. Biomol NMR* **2018**, *70* (2), 103–114. Wiegand, T.; Cadalbert, R.; von Schroetter, C.; Allain, F. H.; Meier, B. H. Segmental isotope labelling and solid-state NMR of a 12 × 59 kDa motor protein: identification of structural variability. *J. Biomol NMR* **2018**, *71* (4), 237–245. Ciragan, A.; Backlund, S. M.; Mikula, K. M.; Beyer, H. M.; Samuli Ollila, O. H.; Iwai, H. NMR Structure and Dynamics of TonB Investigated by Scar-Less Segmental Isotopic Labeling Using a Salt-Inducible Split Intein. *Front Chem.* **2020**, *8*, 136.

(15) Agouridas, V.; El Mahdi, O.; Diemer, V.; Cargoet, M.; Monbaliu, J. M.; Melnyk, O. Native Chemical Ligation and Extended Methods: Mechanisms, Catalysis, Scope, and Limitations. *Chem. Rev.* **2019**, *119* (12), 7328–7443.

(16) Liu, J.; Chen, Q.; Rozovsky, S. Utilizing Selenocysteine for Expressed Protein Ligation and Bioconjugations. *J. Am. Chem. Soc.* **2017**, *139* (9), 3430–3437.

(17) Muralidharan, V.; Muir, T. W. Protein ligation: an enabling technology for the biophysical analysis of proteins. *Nat. Methods* **2006**, *3* (6), 429–438.

(18) Brenzel, S.; Cebi, M.; Reiss, P.; Koert, U.; Mootz, H. D. Expanding the scope of protein trans-splicing to fragment ligation of an integral membrane protein: towards modulation of porin-based ion channels by chemical modification. *Chembiochem* **2009**, *10* (6), 983–986.

(19) Stevens, A. J.; Brown, Z. Z.; Shah, N. H.; Sekar, G.; Cowburn, D.; Muir, T. W. Design of a Split Intein with Exceptional Protein Splicing Activity. *J. Am. Chem. Soc.* **2016**, *138* (7), 2162–2165.

(20) Jumper, J.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Zidek, A.; Potapenko, A.; et al. Highly accurate protein structure prediction with AlphaFold. *Nature* **2021**, *596* (7873), 583–589.

(21) Gunsel, U.; Hagn, F. Lipid Nanodiscs for High-Resolution NMR Studies of Membrane Proteins. *Chem. Rev.* **2022**, *122* (10), 9395–9421. Hagn, F.; Nasr, M. L.; Wagner, G. Assembly of phospholipid nanodiscs of controlled size for structural studies of membrane proteins by NMR. *Nat. Protoc* **2018**, *13* (1), 79–98.

(22) Wishart, D. S.; Sykes, B. D.; Richards, F. M. The chemical shift index: a fast and simple method for the assignment of protein secondary structure through NMR spectroscopy. *Biochemistry* **1992**, *31* (6), 1647–1651.

(23) Iwai, H.; Zuger, S.; Jin, J.; Tam, P. H. Highly efficient protein trans-splicing by a naturally split DnaE intein from *Nostoc punctiforme*. *FEBS Lett.* **2006**, *580* (7), 1853–1858.

(24) Sperl, L. E.; Hagn, F. NMR Structural and Biophysical Analysis of the Disease-Linked Inner Mitochondrial Membrane Protein MPV17. *J. Mol. Biol.* **2021**, *433* (15), No. 167098.

(25) Schmidt, P.; Bender, B. J.; Kaiser, A.; Gulati, K.; Scheidt, H. A.; Hamm, H. E.; Meiler, J.; Beck-Sickingler, A. G.; Huster, D. Improved in Vitro Folding of the Y(2) G Protein-Coupled Receptor into Bicelles. *Front Mol. Biosci* **2018**, *4*, 100.

(26) Aranko, A. S.; Zuger, S.; Buchinger, E.; Iwai, H. In vivo and in vitro protein ligation by naturally occurring and engineered split DnaE inteins. *PLoS One* **2009**, *4* (4), No. e5185. Chong, S.; Shao, Y.; Paulus, H.; Benner, J.; Perler, F. B.; Xu, M. Q. Protein splicing involving the *Saccharomyces cerevisiae* VMA intein. The steps in the splicing pathway, side reactions leading to protein cleavage, and establishment of an in vitro splicing system. *J. Biol. Chem.* **1996**, *271* (36), 22159–22168.

(27) Klopfer, K.; Hagn, F. Beyond detergent micelles: The advantages and applications of non-micellar and lipid-based membrane mimetics for solution-state NMR. *Prog. Nucl. Magn. Reson. Spectrosc.* **2019**, *114–115*, 271–283.

(28) Kurauskas, V.; Schanda, P.; Sounier, R. Methyl-Specific Isotope Labeling Strategies for NMR Studies of Membrane Proteins. In *Membrane Protein Structure and Function Characterization Methods in*

Molecular Biology. Lacapere, J. J., Ed.; Humana Press, New York, NY, vol 1635, 2017; https://doi.org/10.1007/978-1-4939-7151-0_6.

(29) Hagn, F.; Wagner, G. Structure refinement and membrane positioning of selectively labeled OmpX in phospholipid nanodiscs. *J. Biomol NMR* **2015**, *61* (3–4), 249–260.

(30) Shen, Y.; Delaglio, F.; Cornilescu, G.; Bax, A. TALOS+: a hybrid method for predicting protein backbone torsion angles from NMR chemical shifts. *Journal of biomolecular NMR* **2009**, *44* (4), 213–223.

(31) Laskowski, R. A.; Rullmann, J. A.; MacArthur, M. W.; Kaptein, R.; Thornton, J. M. AQUA and PROCHECK-NMR: programs for checking the quality of protein structures solved by NMR. *J. Biomol. NMR* **1996**, *8* (4), 477–486.

(32) Stevens, A. J.; Sekar, G.; Shah, N. H.; Mostafavi, A. Z.; Cowburn, D.; Muir, T. W. A promiscuous split intein with expanded protein engineering applications. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114* (32), 8538–8543.

(33) Pinto, F.; Thornton, E. L.; Wang, B. An expanded library of orthogonal split inteins enables modular multi-peptide assemblies. *Nat. Commun.* **2020**, *11* (1), 1529.

(34) Casiraghi, M.; Damian, M.; Lescop, E.; Point, E.; Moncoq, K.; Morellet, N.; Levy, D.; Marie, J.; Guittet, E.; Baneres, J. L.; et al. Functional Modulation of a G Protein-Coupled Receptor Conformational Landscape in a Lipid Bilayer. *J. Am. Chem. Soc.* **2016**, *138* (35), 11170–11175. Schrottke, S.; Kaiser, A.; Vortmeier, G.; Els-Heindl, S.; Worm, D.; Bosse, M.; Schmidt, P.; Scheidt, H. A.; Beck-Sickingler, A. G.; Huster, D. Expression, Functional Characterization, and Solid-State NMR Investigation of the G Protein-Coupled GHS Receptor in Bilayer Membranes. *Sci. Rep* **2017**, *7*, 46128.

(35) Mohamadi, M.; Goricanec, D.; Wagner, G.; Hagn, F. NMR sample optimization and backbone assignment of a stabilized neurotensin receptor. *J. Struct. Biol.* **2023**, *215* (2), No. 107970.

(36) Bonaccorsi, M.; Le Marchand, T.; Pintacuda, G. Protein structural dynamics by Magic-Angle Spinning NMR. *Curr. Opin Struct. Biol.* **2021**, *70*, 34–43. Le Marchand, T.; Schubeis, T.; Bonaccorsi, M.; Paluch, P.; Lalli, D.; Pell, A. J.; Andreas, L. B.; Jaudzems, K.; Stanek, J.; Pintacuda, G. (1)H-Detected Biomolecular NMR under Fast Magic-Angle Spinning. *Chem. Rev.* **2022**, *122* (10), 9943–10018.

(37) Becker, C. F.; Lauecker, K.; Balog, M.; Kalai, T.; Hideg, K.; Steinhoff, H. J.; Engelhard, M. Incorporation of spin-labelled amino acids into proteins. *Magn. Reson. Chem.* **2005**, *43* (Spec no.), S34–39.

(38) Sonntag, M.; Jagtap, P. K. A.; Simon, B.; Appavou, M. S.; Geerlof, A.; Stehle, R.; Hennig, J.; Sattler, M. Segmental, Domain-Selective Perdeuteration and Small-Angle Neutron Scattering for Structural Analysis of Multi-Domain Proteins. *Angew. Chem., Int. Ed. Engl.* **2017**, *56* (32), 9322–9325.