



Kaempferol 3-O-rhamnoside-7-O-rhamnoside is an endogenous flavonol inhibitor of polar auxin transport in Arabidopsis shoots

Ruohe Yin^{1,4}, Kerstin Han¹, Werner Heller¹, Andreas Albert², Petre I. Dobrev³, Eva Zažímalová³ and Anton R. Schäffner¹

¹Institute of Biochemical Plant Pathology, Helmholtz Zentrum München, 85764 Neuherberg, Germany; ²Research Unit Environmental Simulation, Helmholtz Zentrum München, 85764 Neuherberg, Germany; ³Institute of Experimental Botany, Academy of Sciences of the Czech Republic, Prague 6, Czech Republic; ⁴Present address: Department of Botany and Plant Biology, University of Geneva, Sciences III, 1211 Geneva 4, Switzerland

Authors for correspondence: Ruohe Yin Tel: +41 22 379 3653 Email: ruohe.yin@gmail.com

Anton R. Schäffner Tel: +49 89 31872930 Email: schaeffner@helmholtz-muenchen.de

Received: 14 August 2013 Accepted: 15 September 2013

New Phytologist (2013) doi: 10.1111/nph.12558

Key words: Arabidopsis thaliana, flavonol biosynthesis, flavonol glycoside, flavonol glycosyltransferases, plant growth, polar auxin transport.

Introduction

The phytohormone auxin, represented predominantly by indole-3-acetic acid (IAA), plays a crucial role in plant growth and development. Auxin needs to be transported from the sites of synthesis, mainly in the apices and young leaves, to the distal part of the plant to exert its function (Berleth et al., 2007). Auxin transporters, including ABCB proteins, AUX1/LAX family members and PIN proteins (Noh et al., 2001; Friml, 2003; Petrasek et al., 2006; Yang et al., 2006; Zažímalová et al., 2010; Peer et al., 2011), are responsible for the auxin fluxes and patterning in plants (Friml et al., 2003). Flavonoids, phenylpropanoic secondary metabolites, have been implicated in the blocking of auxin transport (Peer & Murphy, 2007). The supply of flavonols to detached zucchini hypocotyls resulted in decreased polar auxin transport (PAT) (Jacobs & Rubery, 1988). In Arabidopsis, PAT

was increased in transparent testa4 (tt4), a flavonoid-deficient mutant defective in the first step of flavonoid production (Shirley et al., 1995; Buer & Muday, 2004; Peer et al., 2004) (Fig. 1). Accordingly, *tt4* roots exhibited delayed gravitropism, which was reversed by chemical complementation by naringenin, an intermediate of flavonoid biosynthesis (Buer & Muday, 2004). By

Summary

• Polar auxin transport (PAT) plays key roles in the regulation of plant growth and development. Flavonoids have been implicated in the inhibition of PAT. However, the active flavonoid derivative(s) involved in this process in vivo has not yet been identified. Here, we provide evidence that a specific flavonol bis-glycoside is correlated with shorter plant stature and reduced PAT.

• Specific flavonoid-biosynthetic or flavonoid-glycosylating steps were genetically blocked in Arabidopsis thaliana. The differential flavonol patterns established were analyzed by highperformance liquid chromatography (HPLC) and related to altered plant stature. PAT was monitored in stem segments using a radioactive $[^{3}H]$ -indole-3-acetic acid tracer.

• The flavonoid 3-O-glucosyltransferase mutant ugt78d2 exhibited a dwarf stature in addition to its altered flavonol glycoside pattern. This was accompanied by reduced PAT in ugt78d2 shoots. The ugt78d2-dependent growth defects were flavonoid dependent, as they were rescued by genetic blocking of flavonoid biosynthesis. Phenotypic and metabolic analyses of a series of mutants defective at various steps of flavonoid formation narrowed down the potentially active moiety to kaempferol 3-O-rhamnoside-7-O-rhamnoside. Moreover, the level of this compound was negatively correlated with basipetal auxin transport.

• These results indicate that kaempferol 3-O-rhamnoside-7-O-rhamnoside acts as an endogenous PAT inhibitor in Arabidopsis shoots.

> contrast, PAT was reduced in the flavonol over-production mutant #3 defective in dihydroflavonol reductase (Fig. 1), consistent with an inhibitory role of flavonols in PAT (Peer et al., 2004). Despite these and other substantial pieces of evidence supporting a role of flavonols in the modulation of auxin transport (Kuhn et al., 2011; Lewis et al., 2011; Grunewald et al., 2012), neither specific flavonol aglycones nor their conjugates active in this process in vivo have been identified so far.

> The difficulty in relating specific flavonols to auxin transport modulation is, in part, a result of the complex flavonol modification in planta. Flavonol aglycones are intensively modified by UDP-dependent glycosyltransferases (UGTs), which include UGT78D2, UGT78D3, UGT73C6 UGT78D1, and UGT89C1 in the model plant Arabidopsis thaliana (Fig. 1) (Jones et al., 2003; Tohge et al., 2005; Yonekura-Sakakibara et al., 2007, 2008). The glycosides are distributed in an organspecific manner. In contrast with the complex flavonol profile in flowers, it is rather simple in inflorescence stems (Yonekura-Sakakibara et al., 2008; Stracke et al., 2010b). Thus, the inflorescence stem, which is implicated in basipetal auxin movement, is the optimal organ for searching for the flavonol derivative(s) active in auxin transport modulation.



Here, we show that the loss of the flavonoid 3-O-glucosyltransferase UGT78D2 resulted in an altered flavonol glycoside pattern and reduced PAT in shoots, which was accompanied by a reduced plant height and increased branching. Blocking of flavonoid biosynthesis and/or glycosylation at specific steps clearly related the enhanced accumulation of kaempferol 3-O-rhamnoside-7-O-rhamnoside (k1) to the growth defects of *ugt78d2*. Through analyses of auxin transport in several genotypes, which contained different levels of k1, an inverse correlation between basipetal auxin transport and k1 level was identified. Therefore, we propose that k1 acts as an endogenous auxin transport inhibitor in *Arabidopsis* shoots.

Materials and Methods

Plant material and growth conditions

The ugt78d1 (SAIL_568_F08), ugt78d2 (SALK_049338), ugt 89c1 (SALK_071113), tt4 (SALK_020583), tt7 (GK_349F05) and tt18 (SALK_028793) mutants are based on Arabidopsis thaliana (L.) Heynh., accession Columbia (Col-0). These mutant lines have been described in previous studies (Abrahams et al.,

Fig. 1 Flavonoid biosynthesis pathway in Arabidopsis thaliana. (a) Scheme of flavonoid biosynthesis. CHS (TT4), chalcone synthase; F3'H (TT7), flavonoid 3'hydroxylase; DFR (TT3), dihydroflavonol 4-reductase; FLS, flavonol synthase; ANS (TT18), anthocyanidin synthase; UGT78D1, flavonol 3-O-rhamnosyltransferase; UGT78D2, flavonoid 3-Oglucosyltransferase; UGT89C1, flavonol 7-O-rhamnosyltransferase. As shown by Yin et al. (2012), the combined loss of UGT78D1 and UGT78D2 does not imply an accumulation of flavonol aglycones because of a feedback inhibition of flavonol biosynthesis. (b) Glycosylation reactions catalyzed by UGT78D1, UGT78D2 and UGT89C1 (Jones et al., 2003; Tohge et al., 2005; Yonekura-Sakakibara et al., 2007). Abbreviations: Kaempferol (k); rhamnoside (rha); glucoside (glu). k1, k-3-O-rha-7-Orha; k2, k-3-O-glu-7-O-rha; k3, k-3-O-[rha (1->2 glu)]-7-O-rha; q1, q2 and q3 are guercetins structurally equivalent to k1, k2 and k3, respectively.

2003; Alonso *et al.*, 2003; Jones *et al.*, 2003; Rosso *et al.*, 2003; Tohge *et al.*, 2005; Yonekura-Sakakibara *et al.*, 2007; Stracke *et al.*, 2010b). The *tt3-1* (NASC_N84; L*er* background) and *fls1* (Riken_PST16145; Nö background) mutants have different genetic backgrounds (Shirley *et al.*, 1995; Ito *et al.*, 2005).

tt4 ugt78d2, tt7 ugt78d2, ugt78d1 ugt78d2, ugt89c1 ugt78d2, tt18 ugt78d2 and fls1 ugt78d2 double mutants were generated by genetic crossing and confirmed by PCR genotyping. Plants were grown in a sun simulator chamber, which provides a photobiological environment very close to natural global solar radiation (Seckmeyer & Payer, 1993; Thiel *et al.*, 1996). Plants were exposed to photosynthetically active radiation (PAR, 400– 700 nm) of 200–230 μ mol m⁻² s⁻¹ with a 16 h : 8 h, light : dark regime at a temperature of 22°C and a relative humidity of 70%. Similar phenotypes were observed when grown in a glasshouse with higher natural light intensities at temperatures of 20–25°C (European springtime).

A DR5:: β -glucuronidase (DR5::GUS) line (Col background; Sabatini *et al.*, 1999) was introgressed into *ugt78d2*; plants were grown on half-strength Murashige and Skoog (MS) medium agar plates supplemented with 1.5% sucrose for 9 d and stained for β -glucuronidase expression for 1 h at 37°C (Deruère *et al.*, 1999).

Genetic complementation of *ugt78d2*

A genomic fragment containing the *UGT78D2* gene was amplified from Col-0 genomic DNA using the primers 5'-GGGGAC AAGTTTGTACAAAAAAGCAGGCTTTCGGTCCAAAGGA TTTCAG-3' and 5'-GGACCACTTTGTACAAGAAAGCTG GGTAGATTTTCTGAGCCGTGCAT-3'. The PCR product was cloned into vector pDONR221 using GATEWAYTM (Invitrogen, Karlsruhe, Germany) recombination, confirmed by DNA sequencing and further recombined into pBGW (Karimi *et al.*, 2002). The resulting binary vector was used to transform *ugt78d2* knockout mutant plants by the floral dip method (Clough & Bent, 1998). Transgenic plants with a single insertion of the transgene were selected by segregation analysis for experiments.

Analyses of flavonols

For flavonol extraction, the lower 2.5-cm stem segment of 4-wk-old plants was harvested and immediately frozen in liquid nitrogen. The plant material was thoroughly ground in liquid nitrogen using a mortar and pestle, and further homogenized after the addition of 1 ml of methanol per 100 mg of fresh material in a Douncer on ice. The suspension was incubated for 1 h with moderate rotation at 4°C. The extracts were then clarified by centrifugation at $14\,000\,g$ for 10 min with a table-top centrifuge at 4°C. One-third volume of distilled water was added to the supernatant, vortexed and centrifuged as above. The cleared extract was analyzed using a reversephase high-performance liquid chromatography (HPLC) system (Gemini C18 Phenyl, 5 μ m, 150 \times 4.6 mm; Phenomenex, Aschaffenburg, Germany). Solvent A was 1% acetic acid and solvent B consisted of 89% methanol with 1% acetic acid. The elution gradient program followed a linear gradient from 20% solvent B to 100% solvent B (75 min) at room temperature. The quantification of k1 was referred to a kaempferol aglycone authentic standard (Carl Roth, Karlsruhe, Germany).

Inflorescence stem gravitropic response assay

Arabidopsis plants were grown on commercial soil substrate (Einheitserde; Einheitserde- und Humuswerke, Sinntal-Altengronau, Germany) for 4 wk in a phytochamber (light intensity, 120 μ mol m⁻²s⁻¹; 16 h : 8 h, light : dark cycle; relative humidity, 60%; 23°C). Plants were adapted in darkness for 2 h before gravitropic stimulation. Gravitropic stimulation was begun by turning plants by 90° at 23°C in darkness. Except for the 5 min needed for the measurement of the bending angles at each time point, the whole gravitropic response assay was conducted in darkness.

Examination of parenchyma cell size of inflorescence stems

Seven-week-old inflorescence stems were fixed in ethanol: glycerol: water (36:1:10, v/v/v), hand sectioned along the longitudinal axis, stained with astral blue and safranin, and examined under a microscope in bright field mode. Cell length was measured using Cell^P Professional Imaging Software (Olympus Europe, Hamburg, Germany).

PAT assay

The auxin transport assays were performed as described previously with slight modifications (Okada *et al.*, 1991). Segments (length, 2.5 cm) from the basal part of the primary inflorescence stem were inserted into 30 µl of $1 \times MS$ medium (pH 5.5) with $10.5 \,\mu$ M [³H]-IAA (460 kBq ml⁻¹). Both basipetal and acropetal auxin transport were assayed. One end of the stem fragment (apical end for basipetal auxin transport and basal end for acropetal auxin transport) was dipped into the assay mixture. After 6 h of incubation at room temperature in the dark in a closed vial, a 5-mm segment was cut off from the non-submerged end and measured by scintillation counting. For the inhibition assay, $10 \,\mu$ M of *N*-1-naphthylphthalamic acid (NPA) was added. The measurements without NPA treatment were compared between each genotype by a paired *t*-test.

IAA quantification

Col-0 and *ugt78d2* plants were grown for 28 d before the inflorescence stems were harvested. The whole inflorescence stem was cut into three segments of equal length, namely the apex, middle and basal segment. The extraction and quantification of free IAA were carried out as described previously, with the endogenous IAA determined by isotope dilution followed by liquid chromatography-mass spectrometry (Dobrev *et al.*, 2005; Dobrev & Vankova, 2012).

Results

Shoot architecture of *ugt78d2* plants is altered

A loss-of-function mutant of the flavonoid 3-O-glucosyltransferase UGT78D2 showed a pronounced shoot phenotype (Fig. 2). ugt78d2 mutant plants exhibited a significantly shorter stature soon after bolting (Fig. 2). At a later growth stage, the mutants developed more branches with an obvious loss of apical dominance (Supporting Information Table S1). To elucidate the mechanism underlying this dwarfism, the cellular architectures of inflorescence stems were examined. In ugt78d2, the parenchymal cell length was reduced significantly when compared with the wild-type. In contrast with stems, other organs were not obviously affected. The lengths of hypocotyls and siliques of ugt78d2 were similar to those of the wild-type (Table S1). Young seedlings and root systems developed in the same way in both genotypes (Fig. S1) Transformation of ugt78d2 with a genomic fragment covering the complete UGT78D2 gene rescued plants from growth disturbances and restored the wild-type phenotype (Fig. S2). Thus, the lesion in UGT78D2 was responsible for the observed shoot growth defects.

4 Research



Fig. 2 Phenotypes of flavonoid biosynthesisrelated mutants. Shoots of *Arabidopsis thaliana* wild-type and the indicated mutants were imaged after 28 d of growth. Bar, 5 cm.

The ugt78d2 growth phenotype is flavonol dependent

Flavonoids are probable candidates responsible for the ugt78d2 phenotype, as UGT78D2 has been characterized as a flavonoid 3-O-glucosyltransferase and the ugt78d2 mutant exhibits an altered flavonoid pattern (Lee et al., 2005; Tohge et al., 2005). Moreover, transformation of ugt78d2 with the UGT78D2 gene did not only re-establish the wild-type growth phenotype, but also restored its flavonol pattern (Fig. 3a). A flavonoid-deficient tt4 ugt78d2 double mutant was generated to further check this effect. The growth defects of ugt78d2 were indeed eliminated in tt4 ugt78d2, indicating that they were flavonoid dependent. Moreover, this observation in the flavonoid-deficient background indicated that the growth defects were not related to the repression of specific flavonol-3-O-glucoside derivatives (k2, k3, q2 and q3) in ugt78d2 (Figs 2, 3a). In addition to flavonols, UGT78D2 also recognizes anthocyanidins as substrates (Lee et al., 2005; Tohge et al., 2005). To test the possible involvement of anthocyanins in the development of the ugt78d2 growth phenotype, an anthocyanin-deficient *tt18 ugt78d2* double mutant was generated. However, loss of the anthocyanidin synthase in the tt18 ugt78d2 line did not rescue the growth defects of ugt78d2 (Figs 1, 2). The flavonol glycoside pattern of tt18ugt78d2 was similar to that of ugt78d2 (Fig. 3a). By contrast, the introgression of fls1 (accession Nö) into ugt78d2, which specifically blocks the biosynthesis of the flavonol moieties, restored wild-type-like growth (Fig. S3). The impact of flavonols was further supported by the slightly repressed growth of the tt3 mutant in comparison with its genetic background Ler; the loss of dihydroflavonol reductase in the tt3 mutant leads to enhanced k1 levels (Fig. S4; Peer *et al.*, 2001). Collectively, these observations indicated that the ugt78d2 growth phenotype was flavonol dependent and that anthocyanins were not responsible for the ugt78d2 growth defects.

The *ugt78d2* phenotype is correlated with kaempferol 3-O-rhamnoside-7-O-rhamnoside

To identify the flavonols responsible for the *ugt78d2* phenotype, a genetic approach using mutants defective in flavonol biosynthetic and/or conjugating steps was employed. In concert with published data (Yonekura-Sakakibara *et al.*, 2008; Stracke *et al.*, 2010b), six flavonol glycosides were detected in wild-type stems. Three kaempferol glycosides (k1–k3) were the most abundant

Fig. 3 Flavonol glycoside pattern in Arabidopsis thaliana mutants affecting flavonoid biosynthesis and conjugation. (a) Flavonols extracted from stems (lower region of 2.5 cm) of the wild-type and flavonoid metabolism-related mutants. The scale of all representative high-performance liquid chromatograms is identical. Peaks u1-u8 represent unknown flavonoids. SM, sinapoyl malate. (b) Quantification of k1 (Fig. 1b) in the wild-type and flavonoid metabolism-related mutants. The means of $k1 \pm SD$ from four biological samples (a pool of three to five individual stem segments for one sample) are shown. k1 was quantified using kaempferol aglycone as a standard. Letters above the error bars represent a significant difference between genotypes by a paired *t*-test (P < 0.01).



flavonols, whereas three quercetin glycosides (q1-q3) were present at much lower levels (Fig. 3a). No flavonol aglycones were detected in the wild-type and *ugt78d2*, which is in accordance with previous analyses of the single 3-O-glycosylation mutants *ugt78d1* and *ugt78d2* (Tohge *et al.*, 2007; Yin *et al.*, 2012) (Fig. 1). Furthermore, even the double mutant *ugt78d1 ugt78d2*, which blocks the major enzymes performing the initial 3-O-glycosylation of the flavonol moiety, did not accumulate aglycones (Yin *et al.*, 2012). In accordance with the loss of 3-O-glucosyl-transferase activity in *ugt78d2*, the 3-O-glucoside derivatives (k2, k3, q2 and q3) were strongly reduced, whereas the 3-O-rhamnoside derivatives (k1 and q1) were elevated, when compared with the wild-type counterpart (Fig. 3a). Therefore, both k1 and q1 were positively correlated with the *ugt78d2* growth defects.

A *tt7 ugt78d2* double mutant was generated in order to further distinguish whether the increases in k1 and/or q1 levels were

related to the *ugt78d2* growth phenotype. The conversion of kaempferol to quercetin is blocked in *tt7* (Fig. 1). As expected, *tt7 ugt78d2* was devoid of q1 and accompanied by an even larger amount of k1 than in *ugt78d2* (Fig. 3b). *tt7 ugt78d2* was more severely dwarfed than *ugt78d2*, indicating that a further enhanced k1 level led to a more pronounced phenotype (Fig. 2). These observations suggested that the development of the *ugt78d2* growth defects was independent of quercetin derivatives, and that k1 was sufficient to induce the *ugt78d2* phenotype.

k1 contains 3-O-rhamnosyl and 7-O-rhamnosyl residues (Fig. 1b). To further substantiate that k1 was responsible for the observed *ugt78d2* phenotype, additional mutants were employed to selectively inhibit the attachment of 3-O- or 7-O-rhamnosyl residues to the k1 backbone. The attachment of these residues to the k1 backbone is catalyzed by UGT78D1 and UGT89C1, respectively (Jones *et al.*, 2003; Yonekura-Sakakibara *et al.*,

2007). As expected, k1 could not be detected in *ugt78d2 ugt89c1*, but unknown flavonol derivatives accumulated (Fig. 3a). Importantly, the *ugt78d2* phenotype reverted to a wild-type stature in this double mutant (Fig. 2). Likewise, no *ugt78d2*-like phenotype was observed in *ugt78d1 ugt78d2*, which was also devoid of k1 (Figs 2, 3a). All of these observations clearly indicated that k1 was indeed the flavonol metabolite inducing the *ugt78d2* phenotype.

Basipetal auxin transport is reduced in ugt78d2 plants

The dwarf phenotype of ugt78d2 resembled that of the abcb1 abcb19 double mutant, which was greatly impaired in auxin transport (Noh et al., 2001). As flavonols have been postulated to be endogenous auxin transport inhibitors, we reasoned that the increase in k1 suppressed auxin transport in inflorescence stems, resulting in the growth defects. Therefore, polar transport of [³H]-IAA was measured in inflorescence stem segments. NPAsensitive, basipetal auxin transport in ugt78d2 was reduced significantly to c. 45% of the wild-type level, whereas acropetal auxin transport remained unchanged (Figs 4a, S5). Importantly, auxin transport activity further decreased to c. 25% of the wild-type level in the tt7 ugt78d2 double mutant, which accumulated even greater amounts of k1 (Figs 3b, 4a). There was a dose-dependent, linear relationship between the k1 level and PAT, when comparing the wild-type, ugt78d2, tt7 and tt7 ugt78d2 (Fig. 4b). The regulation did not affect the transcript level of genes encoding



Fig. 4 Basipetal auxin transport in inflorescence stems. (a) Basipetal auxin transport in *Arabidopsis thaliana* wild-type, *ugt78d2*, *tt7* and *tt7 ugt78d2* basal stem segments. Data represent radioactivity accumulated in basal segments (means \pm SE obtained from 10 individual plants without *N*-1-naphthylphthalamic acid (NPA) treatment and from two individual plants for NPA treatment). Closed bars, +NPA; open bars, –NPA. Letters above the error bars represent a significant difference between genotypes by a paired *t*-test (*P* < 0.05). (b) Basipetal auxin transport in relation to k1 levels. Mean value and standard error are plotted. A linear relationship was observed (*R*² = 0.9911).

major, putative auxin export proteins, as mRNA levels of *PIN1*, *PIN3*, *PIN7*, *ABCB1* and *ABCB19* were not significantly different in wild-type and *ugt78d2* stems (Fig. S6).

The altered PAT also raised the question of whether there was a general difference in auxin responsiveness. However, several auxin-inducible marker genes were up-regulated by exogenous IAA application in the same manner in *ugt78d2* and the wild-type (Fig. S7).

Free IAA level is reduced in ugt78d2 inflorescence stems

To examine whether the reduced auxin transport influenced the concentration of free IAA in ugt78d2 stems, the free IAA contents of three consecutive stem segments of ugt78d2 and wild-type plants were compared. The endogenous levels of free IAA in the apical inflorescence segments were not different from those of the wild-type (P > 0.05), whereas the middle and basal stem segments of ugt78d2 contained lower steady-state IAA levels than the wild-type counterparts (P < 0.05; Table 1). This suggested that the decreased IAA content in the lower parts of the inflorescence stem of ugt78d2 was caused by the reduced basipetal transport of auxin from sites of its biosynthesis in the apex.

Gravitropism of the inflorescence stem is delayed in *ugt78d2* plants

The reduction of PAT in the inflorescence stems of *ugt78d2* could also affect tropic responses. Therefore, gravitropism was analyzed in the mutant. Four-week-old plants were laid down horizontally in the dark to exert a gravistimulus, which would lead to an upward bending of the inflorescence stems. The bending angle of the inflorescence stem was monitored at different time points after the gravistimulus. *ugt78d2* exhibited a strongly delayed gravitropic response, indicating that the repressed auxin transport was correlated with a reduced gravitropism in *ugt78d2* inflorescence stems (Fig. 5).

Discussion

Over the past decade, strong evidence, both *in vitro* and *in vivo*, has accumulated implicating flavonols in plant growth regulation and auxin transport modulation (Jacobs & Rubery, 1988; Buer & Muday, 2004; Peer *et al.*, 2004; Ringli *et al.*, 2008; Santelia

 Table 1
 Free indole-3-acetic acid (IAA) quantification in Arabidopsis

 thaliana inflorescence stem segments
 \$\$\$

	Apical segment	Middle segment	Basal segment
Wild-type	$\begin{array}{c} 207\pm50\\ 194\pm17 \end{array}$	253 ± 23	247 ± 27
ugt78d2		118 \pm 14*	$142 \pm 18*$

Free IAA was measured in three consecutive stem segments (pmol g⁻¹ FW). The mean values \pm SE of three independent biological samples (one sample is a pool of four to six individual stem segments) are displayed. Statistical analyses were performed by a paired *t*-test for uneven variance for each part of the stem between the two genotypes (*, *P* < 0.05).



Fig. 5 Gravitropic response of wild-type and *ugt78d2* inflorescence stems. Four-week-old *Arabidopsis thaliana* plants were laid down from an upright to a horizontal position in darkness. The curvature of the primary inflorescence stems was measured at different time points to assess the gravitropic response. The angles refer to a horizontal positioning of the tip of the inflorescence (0°) and a fully upright position of the tangential approximation of the bent tip (90°). Wild-type, blue line; *ugt78d2*, red line. Means and SE are shown (n = 15).

et al., 2008; Kuhn et al., 2011; Lewis et al., 2011; Grunewald et al., 2012; Buer et al., 2013). However, the physiologically active flavonol derivative(s) could not be pinpointed unequivocally. In vivo feeding studies were hampered by the possible modification and metabolism of the exogenously added compounds. Similarly, genetic studies affecting either flavonol biosynthesis or flavonol conjugation did not generate effects that could be traced back to a single flavonol moiety (Buer & Muday, 2004; Peer et al., 2004; Ringli et al., 2008; Buer et al., 2013). Nevertheless, the genetic identification of the kaempferol derivative k1 as an endogenous inhibitor of PAT in this work is in agreement with previous studies. These have indicated activity associated with exogenously applied flavonols that could be metabolized to k1 (Jacobs & Rubery, 1988; Mathesius et al., 1998). Furthermore, a release from PAT suppression in flavonol-free tt4 was observed, whereas PAT is further repressed in tt3, which exhibits an enhanced flavonol content at the expense of anthocyanins, and, importantly, in tt7, which shows specifically increased kaempferol glycoside levels at the expense of quercetins (Buer & Muday, 2004; Peer et al., 2004). Furthermore, the formation of k1 is dependent on the 3-O-rhamnosyltransferase UGT78D1, which is relatively abundant at the shoot apex in comparison with other tissues and almost absent from roots in agreement with a low k1 level in roots (Jones et al., 2003) (Figs S1, S8). Thus, the expression pattern of UGT78D1 correlates with an in situ activity of k1 in modulating auxin transport and affecting the shoot phenotype.

Nevertheless, the identification of k1 as an active compound does not exclude an impact of other flavonol derivatives on auxin fluxes. Flavonol biosynthesis is under developmental and organ-specific regulation by MYB and WRKY transcription factors (Stracke *et al.*, 2007, 2010b; Grunewald *et al.*, 2012). Accordingly, the active compounds could also be dependent on the developmental stage and/or on the organ. WRKY23 has been revealed to be part of a feedback loop of auxin to repress its own transport in roots, as it was induced by auxin and enhanced the biosynthesis of flavonols, in particular of quercetins. Thus, quercetins have been suggested to be active agents in roots (Grunewald et al., 2012; Buer et al., 2013). This finding was supported by the analysis of the quercetin-less tt7, which indicated that no kaempferol derivatives, but rather quercetins, were involved in the suppression of basipetal auxin flux in roots (Lewis et al., 2011; Grunewald et al., 2012). However, Buer et al. (2013) also found a repressed root PAT in both quercetin-accumulating tt3 and quercetin-deficient tt7. Nevertheless, these data and the very low k1 level in roots, as well as the lack of obvious ugt78d2related phenotypes in hypocotyls and roots, including a similar auxin accumulation in mutant and wild-type root tips monitored by a DR5::β-glucuronidase reporter line, strongly support the notion that different flavonol compounds may affect PAT in root and shoot (Figs S1, S9; Table S1).

Suppression of rhamnose biosynthesis in *rol1-2* strongly altered the flavonol profile and induced, for example, hyponastic growth of cotyledons and aberrant leaf cell development. This was related to enhanced auxin accumulation and a repressed auxin (non-IAA) efflux from mesophyll protoplasts (Ringli *et al.*, 2008; Kuhn *et al.*, 2011). As *rol1-2* growth defects were rescued in *tt4 rol1*, but retained in *tt7 rol1*, they were attributed to kaempferols. However, k1 was not the causal metabolite for this particular phenotype, as *rol1 ugt78d1* lacking k1 retained the *rol1* cotyledon phenotype (Ringli *et al.*, 2008).

Several possible mechanisms have been reported on how flavonols might affect auxin transport. Flavonol biosynthetic mutants led to an altered expression, subcellular localization and perhaps modified dynamics in the plasma membrane of some PIN proteins, which could have been caused by a direct or indirect impact of flavonols on these proteins (Peer et al., 2004; Santelia et al., 2008). At the transcriptional level, there was no difference in mRNA abundance of several PIN as well as ABCB1 and ABCB19 genes in ugt78d2 and wild-type plants (Fig. S6). ABCB1 and ABCB19 are ABC transporters which primarily function in the long-distance auxin transport streams and the movement of auxin out of apical tissues (Bandyopadhyay et al., 2007). Therefore, these ABCB transporters are potential targets of k1 in planta. Indeed, ABCB1/19 proteins are able to bind the quercetin aglycone (Murphy et al., 2002), and inhibition of ABCB1 auxin transport activity by the quercetin aglycone has been demonstrated in Arabidopsis protoplasts (Geisler et al., 2005). These observations are in agreement with previous in vitro experiments showing that flavonol aglycones can compete with the synthetic auxin transport inhibitor NPA for binding to isolated microsomal vesicles, although these results may not necessarily reflect an in vivo relevance (Jacobs & Rubery, 1988; Murphy et al., 2002). Nevertheless, these studies indicate that flavonol derivatives, including k1, could interact directly with and/or inhibit auxin transport proteins.

Despite these *in vitro* studies, a possible involvement of flavonol aglycones in auxin transport inhibition *in vivo* and, in

particular, in the ugt78d2 growth phenotype appears to be unlikely. This notion is based on two lines of evidence: first, to the best of our knowledge, so far only two of numerous studies have detected flavonol aglycones in Arabidopsis by either HPLC (Peer et al., 2001) or mass spectrometry (Buer et al., 2013); however, the latter authors also discussed inconsistencies with respect to the occurrence and absence of aglycones in different tt mutants. Although flavonol aglycones exist at least as biosynthetic intermediates, only minute amounts of the hydrophobic molecules might be tolerated in living cells, and therefore free flavonols are not consistently detected; however, this would not preclude a regulatory role (Yin et al., 2012). The important second line of evidence stems from loss-of-function mutants affecting the two major UGTs in Arabidopsis, UGT78D1 and UGT78D2, which perform the initial 3-O-flavonol glycosylation (Fig. 1). The lack of either UGT is compensated by the remaining one, and results in a shift of the flavonol glycoside pattern with no aglycones being detected (this work; Jones et al., 2003; Yonekura-Sakakibara et al., 2008; Yin et al., 2012). The ugt78d1 ugt78d2 double mutant strongly enhances the chance of flavonol aglycone accumulation, yet no free flavonols were detected (Yin et al., 2012) and, more importantly, the ugt78d2-dependent growth retardation was reversed in the double mutant (Fig. 2).

Flavonols have also been shown to exert their function by modulating the interaction between TWD1, a regulatory protein, and ABCB1 transporters, thus suggesting another possible scenario (Bailly *et al.*, 2008). Furthermore, it has been reported that PINOID modulates ABCB1-mediated auxin transport through its kinase activity, and that this effect is reversed by direct quercetin binding, that is, the flavonol acts as an endogenous kinase inhibitor to modulate PAT (Henrichs *et al.*, 2012). In summary, these mechanistic studies on auxin transport inhibition allow for both different active flavonol moieties and multiple target proteins. The discovery of k1 as an active molecule *in planta* may serve as a valuable tool to obtain further insight into the detailed mechanism of flavonol-dependent auxin transport inhibition.

Apart from the genetic designation of k1, the dose-dependent PAT repression by k1 further supports its identification as an endogenous PAT inhibitor in *Arabidopsis* shoots (Fig. 4). Interestingly, there is also almost the same quantitative relationship in *ugt78d2* relative to the wild-type between the concentrations of k1 (*c.* two-fold higher in the mutant), endogenous auxin (*c.* two-fold lower in the lower inflorescence stem segments of mutant plants) and basipetal auxin transport (*c.* two-fold lower in the basal part of the inflorescence stem of the mutant) (Figs 3b, 4a; Table 1). A further increase in k1 in *tt7 ugt78d2* resulted in more severe growth defects. However, the complete lack of k1 (in *tt4, ugt78d1, ugt89c1*) did not result in a higher stature of the *A. thaliana* plants, suggesting a threshold for k1-dependent growth repression or an independent limitation of growth of these mutants.

In plants, a minor increase in k1 levels would only lead to moderate PAT inhibition, which might not lead to obvious phenotypic changes (as seen in the shoots of *abcb1* and *abcb*19 single mutants; Noh *et al.*, 2001). However, under harsh natural growth conditions, including high light, UV-B irradiation, extreme temperatures and drought, k1 as well as other flavonol glycosides could be strongly increased (Hannah *et al.*, 2006; Rausher, 2006; Caldwell *et al.*, 2007; Korn *et al.*, 2008; Götz *et al.*, 2010; Stracke *et al.*, 2010a). Thus, regulation of flavonol glycoside levels may constitute a means to adjust plant growth and stature by environmental factors and in ecological contexts.

Acknowledgements

We thank R. Stracke for providing *tt4* and *ugt89c1* seeds, J. Durner for helpful suggestions and R. Ulm, G. Bahnweg and L. Lin for critical reading of the manuscript. We are also grateful to two anonymous reviewers whose remarks helped to improve this study. B. Geist and S. Stich provided excellent technical assistance; S. Dräxl assisted with the auxin transport experiments. The work of P.I.D. and E.Z. was supported by the Grant Agency of the Czech Republic, project P305/11/0797.

References

- Abrahams S, Lee E, Walker AR, Tanner GJ, Larkin PJ, Ashton AR. 2003. The *Arabidopsis TDS4* gene encodes leucoanthocyanidin dioxygenase (LDOX) and is essential for proanthocyanidin synthesis and vacuole development. *Plant Journal* 35: 624–636.
- Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R et al. 2003. Genome-wide insertional mutagenesis of Arabidopsis thaliana. Science 301: 653–657.
- Bailly A, Sovero V, Vincenzetti V, Santelia D, Bartnik D, Koenig BW, Mancuso S, Martinoia E, Geisler M. 2008. Modulation of P-glycoproteins by auxin transport inhibitors is mediated by interaction with immunophilins. *Journal of Biological Chemistry* 283: 21817–21826.
- Bandyopadhyay A, Blakeslee JJ, Lee OR, Mravec J, Sauer M, Titapiwatanakun B, Makam SN, Bouchard R, Geisler M, Martinoia E *et al.* 2007. Interactions of PIN and PGP auxin transport mechanisms. *Biochemical Society Transactions* 35: 137–141.
- Berleth T, Scarpella E, Prusinkiewicz P. 2007. Towards the systems biology of auxin-transport-mediated patterning. *Trends in Plant Science* 12: 151–159.
- Buer CS, Kordbacheh F, Truong TT, Hocart CH, Djordjevic MA. 2013. Alteration of flavonoid accumulation patterns in *transparent testa* mutants disturbs auxin transport, gravity responses, and imparts long-term effects on root and shoot architecture. *Planta* 238: 171–189.
- Buer CS, Muday GK. 2004. The transparent testa4 mutation prevents flavonoid synthesis and alters auxin transport and the response of Arabidopsis roots to gravity and light. *Plant Cell* 16: 1191–1205.
- Caldwell MM, Bornman JF, Ballare CL, Flint SD, Kulandaivelu G. 2007. Terrestrial ecosystems, increased solar ultraviolet radiation, and interactions with other climate change factors. *Photochemical & Photobiological Sciences* 6: 252–266.
- Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant Journal 16: 735–743.
- Deruère J, Jackson K, Garbers C, Soll D, Delong A. 1999. The RCN1-encoded A subunit of protein phosphatase 2A increases phosphatase activity *in vivo*. *Plant Journal* 20: 389–399.
- Dobrev PI, Havlicek L, Vagner M, Malbeck J, Kaminek M. 2005. Purification and determination of plant hormones auxin and abscisic acid using solid phase extraction and two-dimensional high performance liquid chromatography. *Journal of Chromatography A* 1075: 159–166.
- Dobrev PI, Vankova R. 2012. Quantification of abscisic acid, cytokinin, and auxin content in salt-stressed plant tissues. *Methods in Molecular Biology* 913: 251–261.

Friml J. 2003. Auxin transport – shaping the plant. *Current Opinion in Plant Biology* 6: 7–12.

Friml J, Vieten A, Sauer M, Weijers D, Schwarz H, Hamann T, Offringa R, Jurgens G. 2003. Efflux-dependent auxin gradients establish the apical–basal axis of *Arabidopsis. Nature* 426: 147–153.

Geisler M, Blakeslee JJ, Bouchard R, Lee OR, Vincenzetti V, Bandyopadhyay A, Titapiwatanakun B, Peer WA, Bailly A, Richards EL *et al.* 2005. Cellular efflux of auxin catalyzed by the *Arabidopsis* MDR/PGP transporter AtPGP1. *Plant Journal* 44: 179–194.

Götz M, Albert A, Stich S, Heller W, Scherb H, Krins A, Langebartels C, Seidlitz HK, Ernst D. 2010. PAR modulation of the UV-dependent levels of flavonoid metabolites in *Arabidopsis thaliana* (L.) Heynh. leaf rosettes: cumulative effects after a whole vegetative growth period. *Protoplasma* 243: 95–103.

Grunewald W, De Smet I, Lewis DR, Lofke C, Jansen L, Goeminne G, Vanden Bossche R, Karimi M, De Rybel B, Vanholme B *et al.* 2012. Transcription factor WRKY23 assists auxin distribution patterns during *Arabidopsis* root development through local control on flavonol biosynthesis. *Proceedings of the National Academy of Sciences, USA* 109: 1554–1559.

Hannah MA, Wiese D, Freund S, Fiehn O, Heyer AG, Hincha DK. 2006. Natural genetic variation of freezing tolerance in *Arabidopsis. Plant Physiology* 142: 98–112.

Henrichs S, Wang B, Fukao Y, Zhu J, Charrier L, Bailly A, Oehring SC, Linnert M, Weiwad M, Endler A *et al.* 2012. Regulation of ABCB1/ PGP1-catalysed auxin transport by linker phosphorylation. *EMBO Journal* 31: 2965–2980.

Ito T, Motohashi R, Kuromori T, Noutoshi Y, Seki M, Kamiya A, Mizukado S, Sakurai T, Shinozaki K. 2005. A resource of 5,814 dissociation transposon-tagged and sequence-indexed lines of *Arabidopsis* transposed from start loci on chromosome 5. *Plant and Cell Physiology* 46: 1149–1153.

Jacobs M, Rubery PH. 1988. Naturally occurring auxin transport regulators. *Science* 241: 346–349.

Jones P, Messner B, Nakajima J, Schäffner AR, Saito K. 2003. UGT73C6 and UGT78D1, glycosyltransferases involved in flavonol glycoside biosynthesis in *Arabidopsis thaliana*. *Journal of Biological Chemistry* 278: 43910–43918.

Karimi M, Inze D, Depicker A. 2002. GATEWAY vectors for Agrobacterium-mediated plant transformation. Trends in Plant Science 7: 193–195.

Korn M, Peterek S, Mock HP, Heyer AG, Hincha DK. 2008. Heterosis in the freezing tolerance, and sugar and flavonoid contents of crosses between *Arabidopsis thaliana* accessions of widely varying freezing tolerance. *Plant, Cell* & *Environment* 31: 813–827.

Kuhn BM, Geisler M, Bigler L, Ringli C. 2011. Flavonols accumulate asymmetrically and affect auxin transport in *Arabidopsis. Plant Physiology* 156: 585–595.

Lee Y, Yoon HR, Paik YS, Liu JR, Chung WI, Choi G. 2005. Reciprocal regulation of *Arabidopsis* UGT78D2 and BANYULS is critical for regulation of the metabolic flux of anthocyanidins to condensed tannins in developing seed coats. *Journal of Plant Biology* 48: 356–370.

Lewis DR, Ramirez MV, Miller ND, Vallabhaneni P, Ray WK, Helm RF, Winkel BS, Muday GK. 2011. Auxin and ethylene induce flavonol accumulation through distinct transcriptional networks. *Plant Physiology* 156: 144–164.

Mathesius U, Schlaman HR, Spaink HP, Of Sautter C, Rolfe BG, Djordjevic MA. 1998. Auxin transport inhibition precedes root nodule formation in white clover roots and is regulated by flavonoids and derivatives of chitin oligosaccharides. *Plant Journal* 14: 23–34.

Murphy AS, Hoogner KR, Peer WA, Taiz L. 2002. Identification, purification, and molecular cloning of N-1-naphthylphthalamic acid-binding plasma membrane-associated aminopeptidases from *Arabidopsis. Plant Physiology* 128: 935–950.

Noh B, Murphy AS, Spalding EP. 2001. *Multidrug resistance*-like genes of *Arabidopsis* required for auxin transport and auxin-mediated development. *Plant Cell* **13**: 2441–2454.

Okada K, Ueda J, Komaki MK, Bell CJ, Shimura Y. 1991. Requirement of the auxin polar transport system in early stages of *Arabidopsis* floral bud formation. *Plant Cell* **3**: 677–684.

Peer WA, Brown DE, Tague BW, Muday GK, Taiz L, Murphy AS. 2001. Flavonoid accumulation patterns of transparent testa mutants of Arabidopsis. *Plant Physiology* 126: 536–548.

Peer WA, Bandyopadhyay A, Blakeslee JJ, Makam SN, Chen RJ, Masson PH, Murphy AS. 2004. Variation in expression and protein localization of the PIN family of auxin efflux facilitator proteins in flavonoid mutants with altered auxin transport in *Arabidopsis thaliana*. *Plant Cell* **16**: 1898–1911.

Peer WA, Blakeslee JJ, Yang H, Murphy AS. 2011. Seven things we think we know about auxin transport. *Molecular Plant* 4: 487–504.

Peer WA, Murphy AS. 2007. Flavonoids and auxin transport: modulators or regulators? *Trends in Plant Science* 12: 556–563.

Petrasek J, Mravec J, Bouchard R, Blakeslee JJ, Abas M, Seifertova D, Wisniewska J, Tadele Z, Kubes M, Covanova M *et al.* 2006. PIN proteins perform a rate-limiting function in cellular auxin efflux. *Science* **312**: 914–918.

Rausher MD. 2006. The evolution of flavonoids and their genes. In: Grotewold E, ed. *The science of flavonoids*. New York, NY, USA: Springer, 175–211.

Ringli C, Bigler L, Kuhn BM, Leiber RM, Diet A, Santelia D, Frey B, Pollmann S, Klein M. 2008. The modified flavonol glycosylation profile in the *Arabidopsis rol1* mutants results in alterations in plant growth and cell shape formation. *Plant Cell* 20: 1470–1481.

Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B. 2003. An Arabidopsis thaliana T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics. *Plant Molecular Biology* 53: 247–259.

Sabatini S, Beis D, Wolkenfelt H, Murfett J, Guilfoyle T, Malamy J, Benfey P, Leyser O, Bechtold N, Weisbeek P *et al.* 1999. An auxin-dependent distal organizer of pattern and polarity in the *Arabidopsis* root. *Cell* 99: 463–472.

Santelia D, Henrichs S, Vincenzetti V, Sauer M, Bigler L, Klein M, Bailly A, Lee Y, Friml J, Geisler M et al. 2008. Flavonoids redirect PIN-mediated polar auxin fluxes during root gravitropic responses. *Journal of Biological Chemistry* 283: 31218–31226.

Seckmeyer G, Payer HD. 1993. A new sunlight simulator for ecological research on plants. *Journal of Photochemistry and Photobiology B: Biology* 21: 175–181.

Shirley BW, Kubasek WL, Storz G, Bruggemann E, Koornneef M, Ausubel FM, Goodman HM. 1995. Analysis of *Arabidopsis* mutants deficient in flavonoid biosynthesis. *Plant Journal* 8: 659–671.

Stracke R, Favory JJ, Gruber H, Bartelniewoehner L, Bartels S, Binkert M, Funk M, Weisshaar B, Ulm R. 2010a. The *Arabidopsis* bZIP transcription factor HY5 regulates expression of the *PFG1/MYB12* gene in response to light and ultraviolet-B radiation. *Plant, Cell & Environment* 33: 88–103.

Stracke R, Ishihara H, Huep G, Barsch A, Mehrtens F, Niehaus K, Weisshaar B. 2007. Differential regulation of closely related R2R3-MYB transcription factors controls flavonol accumulation in different parts of the *Arabidopsis thaliana* seedling. *Plant Journal* 50: 660–677.

Stracke R, Jahns O, Keck M, Tohge T, Niehaus K, Fernie AR, Weisshaar B. 2010b. Analysis of PRODUCTION OF FLAVONOL GLYCOSIDES-dependent flavonol glycoside accumulation in *Arabidopsis thaliana* plants reveals MYB11-, MYB12- and MYB111-independent flavonol glycoside accumulation. *New Phytologist* 188: 985–1000.

Thiel S, Dohring T, Kofferlein M, Kosak A, Martin P, Seidlitz HK. 1996. A phytotron for plant stress research: how far can artificial lighting compare to natural sunlight? *Journal of Plant Physiology* 148: 456–463.

Tohge T, Nishiyama Y, Hirai MY, Yano M, Nakajima J, Awazuhara M, Inoue E, Takahashi H, Goodenowe DB, Kitayama M *et al.* 2005. Functional genomics by integrated analysis of metabolome and transcriptome of *Arabidopsis* plants over-expressing an MYB transcription factor. *Plant Journal* 42: 218–235.

Tohge T, Yonekura-Sakakibara K, Niida R, Watanabe-Takahashi A, Saito K. 2007. Phytochemical genomics in *Arabidopsis thaliana*: a case study for functional identification of flavonoid biosynthesis genes. *Pure Applied Chemistry* 79: 811–823.

Yang Y, Hammes UZ, Taylor CG, Schachtman DP, Nielsen E. 2006. High-affinity auxin transport by the AUX1 influx carrier protein. *Current Biology* 16: 1123–1127.

Yin R, Messner B, Faus-Kessler T, Hoffmann T, Schwab W, Hajirezaei MR, von Saint Paul V, Heller W, Schäffner AR. 2012. Feedback inhibition of the general phenylpropanoid and flavonol biosynthetic pathways upon a compromised flavonol-3-*O*-glycosylation. *Journal of Experimental Botany* **63**: 2465–2478.

- Yonekura-Sakakibara K, Tohge T, Matsuda F, Nakabayashi R, Takayama H, Niida R, Watanabe-Takahashi A, Inoue E, Saito K. 2008. Comprehensive flavonol profiling and transcriptome coexpression analysis leading to decoding gene–metabolite correlations in *Arabidopsis*. *Plant Cell* 20: 2160–2176.
- Yonekura-Sakakibara K, Tohge T, Niida R, Saito K. 2007. Identification of a flavonol 7-O-rhamnosyltransferase gene determining flavonoid pattern in *Arabidopsis* by transcriptome coexpression analysis and reverse genetics. *Journal of Biological Chemistry* 282: 14932–14941.
- Zažímalová E, Murphy AS, Yang HB, Hoyerová K, Hošek P. 2010. Auxin transporters – why so many? *Cold Spring Harbor Perspectives in Biology* 2: a001552.

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 Morphology and flavonol content of young seedlings.

Fig. S2 Genetic complementation of the *ugt78d2*-dependent shoot growth phenotype.

Fig. S3 Growth phenotype and k1 levels of *fls1 ugt78d2*.

Fig. S4 Growth phenotype and k1 levels of tt3-1 and Ler.

Fig. S5 Acropetal auxin transport activity in inflorescence stems of wild-type and *ugt78d2*.

Fig. S6 Expression of auxin transporter genes in *ugt78d2* stem segments relative to wild-type.

Fig. S7 Auxin-responsive gene expression in *ugt78d2* and wild-type.

Fig. S8 Analysis of UGT78D1 gene expression in plant organs.

Fig. S9 Expression of auxin reporter *DR5::GUS* in wild-type and *ugt78d2* roots.

Table S1 Phenotypic analysis of Col-0 and ugt78d2 plants

Please note: Wiley Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.

About New Phytologist

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged.
 We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* our average time to decision is <25 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@ornl.gov)
- For submission instructions, subscription and all the latest information visit www.newphytologist.com