

Molecular architecture of thylakoid membranes within intact spinach chloroplasts

Reviewed Preprint

v1 • March 25, 2025

Not revised

Wojciech Wietrzynski , Lorenz Lamm, William HJ Wood, Matina-Jasemi Loukeri, Lorna Malone, Tingying Peng, Matthew P Johnson , Benjamin D Engel 

Biozentrum, University of Basel, Basel, Switzerland • Helmholtz AI, Helmholtz Zentrum München, Neuherberg, Germany • Plants, Photosynthesis and Soil, School of Biosciences, University of Sheffield, Sheffield, United Kingdom • Department of Structural Cell Biology, Max Planck Institute of Biochemistry, Martinsried, Germany

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

The macromolecular organization of photosynthetic complexes within the thylakoids of higher plant chloroplasts has been a topic of significant debate. Using in situ cryo-electron tomography, this study reveals the native thylakoid architecture of spinach thylakoid membranes with single-molecule precision. The experimental methods are unique and **compelling**, providing **important** information for understanding the structural features that impact photosynthetic regulation in vascular plants and addressing several long-standing questions about the organization and regulation of photosynthesis.

<https://doi.org/10.7554/eLife.105496.1.sa2>

Abstract

Thylakoid membranes coordinate the light reactions of photosynthesis across multiple scales, coupling the architecture of an elaborate membrane network to the spatial organization of individual protein complexes embedded within this network. Previously, we used *in situ* cryo-electron tomography (cryo-ET) to reveal the native thylakoid architecture of the green alga *Chlamydomonas reinhardtii* [1] and then map the molecular organization of these thylakoids with single-molecule precision [2]. However, it remains to be shown how generalizable this green algal blueprint is to the thylakoids of vascular plants, which possess distinct membrane architecture subdivided into grana stacks interconnected by non-stacked stromal lamellae. Here, we continue our cryo-ET investigation to reveal the molecular architecture of thylakoids within intact chloroplasts isolated from spinach (*Spinacia oleracea*). We visualize the fine ultrastructural details of grana membranes, as well as interactions between thylakoids and plastoglobules. We apply and further develop AI-based computational approaches for automated membrane segmentation and membrane protein picking [3], enabling us to quantify the organization of photosynthetic complexes within the plane of the thylakoid membrane and across adjacent stacked membranes. Our analysis reveals that, despite different 3D architecture, the molecular organization of thylakoid membranes in vascular plants and green algae is strikingly similar. In contrast to isolated plant thylakoids,

where semi-crystalline arrays of photosystem II (PSII) appear to hold some membranes together, we find in intact chloroplasts that PSII is non-crystalline and has uniform concentration both within the membrane plane and across stacked grana membranes. Similar to *C. reinhardtii*, we observe strict lateral heterogeneity of PSII and PSI at the boundary between appressed and non-appressed thylakoid domains, with no evidence for a distinct grana margin region where these complexes have been proposed to intermix. Based on these measurements, we support a simple two-domain model for the molecular organization of thylakoid membranes in both green algae and plants.

Introduction

Biological membranes and the proteins embedded within them are interwoven components of a dynamic multiscale system. Thylakoid membranes present one of the best examples of such a system, with interconnected relationships between membrane architecture, molecular organization, and physiological function [4, 5]. Therefore, to fully understand the light-dependent reactions of photosynthesis, it is crucial to describe thylakoid architecture with high precision in its most native state.

Thylakoids are the basic unit of oxygenic photosynthesis (with the exception of *Gloeobacteria*, which completely lack them [6, 7]). They form a variety of architectures in cyanobacteria and within the chloroplasts of plants and algae [8]. In vascular plants, thylakoids assemble into one of the most intricate membrane networks in living organisms: one continuous cisternae-like compartment that is folded and flattened into an interconnected network [9–11]. Thylakoid shape and packing within plant chloroplasts are optimized to maximize the surface area for light absorption and the coupled electron and proton transfer reactions conducted by photosynthetic electron transport chain complexes. The thylakoid network is divided into domains of appressed, stacked membranes (called grana) and non-appressed, free thylakoids (called stromal lamellae) that interconnect the grana. This architecture is crucial for the spatial separation of photosystem I (PSI) and photosystem II (PSII) and their associated light harvesting antenna complexes, LHCI and LHCII, thereby preventing the ‘spillover’ (and thus loss) of excitation energy from the short wavelength trap of PSII (680 nm) to the longer wavelength trap of PSI (700 nm) [12, 13]. The existence of two thylakoid domains and their dynamics has been shown to underpin a variety of regulatory mechanisms that control linear electron transfer (LET), cyclic electron transfer (CET) [14, 15], the PSII repair cycle [16], non-photochemical quenching (NPQ) [17], and the balance of excitation energy distributed to PSI and PSII [18, 19]. Despite over half a century of efforts using a multitude of techniques, including freeze-fracture electron microscopy (EM), negative-stain EM, atomic force microscopy, and fluorescence microscopy, our understanding of the organizational principles of thylakoid networks remains limited by resolution and the requirement for intactness of the system [19–25]. Indeed, numerous controversies remain, including the nature and organization of the grana margins and grana end membranes [26], the architecture of the connections between grana and stromal lamellae [9, 22, 27, 28], and the exact distribution of protein complexes such as ATP synthase (ATPsyn) and cytochrome *b₆f* (cyt_{b₆f}) [29, 30].

A decade ago, we combined cryo-focused ion beam (FIB) milling [31, 32] with cryo-electron tomography (cryo-ET) [33, 34] to provide a detailed view of thylakoid architecture *in situ*, inside the chloroplasts of *Chlamydomonas reinhardtii* cells [1]. We then built upon this by leveraging advances in direct electron detector cameras and the contrast-enhancing Volta phase plate to understand the distribution of photosynthetic complexes within the native thylakoid network of *C. reinhardtii* cells [2]. Now, we apply *in situ* cryo-ET to the more challenging system of plant chloroplasts, utilizing an artificial intelligence (AI)-assisted approach [3, 35] to identify individual photosynthetic complexes and quantify their organization within networks of grana and stromal lamellae membranes.

Results and Discussion

We isolated chloroplasts from six-week-old spinach plants then plunged them onto EM grids, thinned them by cryo-FIB milling, and finally imaged them by cryo-ET with defocus contrast (no Volta phase plate). To enhance contrast of the defocus imaging, we used AI-based denoising algorithms [36, 37]. The resulting tomograms provide a glimpse into the native organization of the spinach plastid (**Fig. 1A**, Sup. Fig. 1). The intact double-membrane envelope encloses a protein-dense stroma (note the high concentration of Rubisco throughout the chloroplast, e.g., **Fig. 2A**, **Fig. 3A**). Suspended within the stroma, the thylakoid network is the main architectural feature of the chloroplast (**Fig. 1A-C**). The thylakoids form a discrete network that is discontinuous from the surrounding chloroplast envelope. However, we did observe rare contact sites between the thylakoid and envelope membranes (Sup. Fig. 4), consistent with previous cryo-ET observations in *C. reinhardtii* [1, 38]. Spherical plastoglobules [39] decorated in putative protein densities were often seen interacting with thylakoid surfaces (**Fig. 1B-C**, **Fig. 2C**). We noticed holes through the thylakoid sheets precisely at the plastoglobule contact sites (**Fig. 2D-E**). At those sites, plastoglobules attach to thylakoids along an extended stretch of high curvature thylakoid membrane, as opposed to making a single point contact to a flat thylakoid sheet (**Fig. 2F**). This may help facilitate exchange between the two compartments.

Segmentation of the membranes highlights the nature of the folded thylakoid network (**Fig. 1C-D**, Sup. Fig. 2). The grana form well defined stacks of appressed membranes, which when viewed from the top, were revealed to be irregular cylinders with wavy edges (**Fig. 3A**). Sometimes, thylakoid layers change length and extend laterally beyond the stack, breaking the grana's vertical order (e.g., Sup. Fig. 2A). Distances between grana differ, and in some instances they almost merge with one another (**Fig. 5A-B**; Sup. Fig. 2A). In our dataset, the number of thylakoids per grana stack varied from 4 to 22, with a mean of 10 (**Fig. 1E** left). Every granum connects with the stromal lamellae membranes that spiral around each stack, in accordance with the helical staircase model (**Fig. 1C-D**, **Fig. 3A**) [9, 24]. Stromal lamellae extend from the stacks at variable angles, and after some distance, join other grana or rarely dead-end in the stroma (**Fig. 1C-D**, Sup. Fig. 2A). Our cryo-ET volumes reveal the fine structural details of native membrane bilayers at high-resolution (e.g., **Fig. 4** left), allowing us to precisely quantify the morphometric parameters of thylakoids with sub-nanometer precision. We measured a membrane thickness of 5.1 ± 0.3 nm, a stromal gap of 3.2 ± 0.3 nm, a luminal thickness of 10.8 ± 2.0 nm, and a total thylakoid thickness (including two membranes plus the enclosed lumen) of 21.1 ± 1.8 nm (**Fig. 4**) (for comparison see [1, 2, 30, 40]).

It is important to note that individual tomograms represent only a small slice of the chloroplast volume (approximately $1200 \times 1200 \times 120$ nm). This makes sampling of the larger micrometer-scale features less statistically relevant (see [41] and references within for alternative approaches). For example, grana diameter measured in our cryo-ET volumes is inherently underestimated due to FIB milling yielding only a narrow slice through the grana stack (**Fig. 1E** middle). We therefore complemented these measurements with TEM overviews of FIB-milled lamellae, where many top views of grana were visible (Sup. Fig. 1B), facilitating more accurate diameter measurements (**Fig. 1E** middle). Similarly, we are able to report the exact ratio between appressed and non-appressed membrane surface areas in our dataset (**Fig. 1E** right, F), but we are aware that this ratio may not be representative of the entire chloroplast volume (see Limitations and Future Perspectives).

Thylakoid membranes are populated primarily by the photosynthetic complexes. Their distribution follows the principle of lateral heterogeneity [42]: PSII complexes are localized to appressed membranes, whereas PSI and ATPsyn complexes reside in the non-appressed regions [43–46]. The size of the stroma-exposed domains of PSI and ATPsyn (~ 3.5 nm and ~ 12 nm,




Figure 1.

Overview of intact spinach chloroplasts visualized by cryo-ET.

A) Tomographic slice through an intact spinach chloroplast. **B)** Corresponding 3D segmentation of the volume. Characteristic elements are labeled. Thylakoid membranes (green) are organized into stacks of appressed thylakoids (grana) and individual thylakoids connecting the grana (stromal lamellae). Note the intact chloroplast envelope (blue) and single thylakoid surrounded by two plastoglobules (orange). Stroma particles (grey) are a mix of different complexes but predominantly Rubisco. **C)** The same thylakoid network as in B, here shown from the opposite side. Note the inclined stromal lamellae spiraling around the granum on the left and a smaller plastoglobule sandwiched between the membranes. **D)** Close-up on the margin of a granum (from a different tomogram) showing the individual thylakoids transitioning into inclined stromal lamellae. In the first transition at the top of the image, note the hole in the thylakoid where the stromal lamella meets the granum. **E)** Box plots showing the number of thylakoids per granum (left), grana diameter measured in TEM overviews and in tomograms (middle), and the ratio of appressed to non-appressed membrane surfaces measured in selected tomograms. Box: 75 percentile, yellow lines: mean, whiskers: 95 percentile. **F)** Representative segmentation and classification of appressed (teal) and non-appressed (yellow) thylakoid membranes (quantification shown in E).




Figure 2.

Fine details of plastoglobules interacting with thylakoids.

A) Tomographic slice showing two plastoglobules: dense, spherical droplets that have surfaces covered with protein densities. **B)** 3D render of the surfaces of three plastoglobules. Note the ordered, small densities. Purple arrows in A and B indicate the protein coat. **C)** Gallery of tomographic slices showing plastoglobules in close proximity of thylakoids and the chloroplast envelope (second panel in the gallery of plastoglobules). **D-E)** Zoom-in on a segmentation of thylakoids with visible fenestrations in stromal lamellae. These holes may be caused by interactions with adjacent plastoglobules. **F)** Two examples of plastoglobules potentially interacting with thylakoids. Each example shows three consecutive tomographic z-slices. Note the protein-membrane interactions in example 2, with an array of small protein densities (potentially coming from the plastoglobule coat) mediating the contact between compartments.




Figure 3.

Top view of the interconnections between two grana.

A) Averaged image of 40 tomographic slices (56 nm total averaged volume) from a tomogram denoised with the DeepDeWedge neural network [37] (high contrast, missing wedge inpainting), showing top views of two neighboring grana. Note the irregular shape of the grana, with wavy edges (traced with white line). Three stromal lamellae connect the two grana stacks. PSII particles are visible in the stacks. **B)** Close up of a region from A, showing stromal lamellae bridging two grana. Blue arrowheads: PSII particles in grana membranes.




Figure 4.

Thylakoid spacing and thickness measurements.

Left: a zoom into thylakoids at the granum edge, showing the level of details observed in defocus cryo- tomograms denoised with cryo-CARE. Notice how the thylakoid tips stick to the appressed membrane, making them non-symmetrical. Right: quantification of thylakoid morphometric parameters. Membrane (1, blue) is the mean thickness of the lipid bilayer; stromal gap (2, green) is the mean distance between stacked thylakoids; lumen (3, yellow) is the mean thylakoid lumen width; thylakoid (4, red) is mean thickness of the thylakoid including two membrane bilayers plus the enclosed lumen (measured separately from the individual measurements of those features). Errors: SEM. Note the larger SEM values for lumen and thylakoid. This reflects the lumen variability between rigid appressed thylakoids of the granum and more labile stromal lamellae.




Figure 5.

Lateral heterogeneity of PSII and PSI between appressed and non-appressed membranes.

A) Segmentation of the thylakoid network showing two close grana. Three membrane pieces are highlighted: M1, an appressed membrane of the granum (teal) transitions into a stromal lamellae membrane (yellow) and then into an end membrane of the second granum (yellow); M12, an appressed membrane in the middle of the granum; M23, a membrane that spans two grana with a small unstacked stromal lamella in the middle (zoomed-in and rotated in green inset panel A') and at the right end. The eye with the arrow indicates the viewing direction in membranograms. **B)** Slice through the corresponding tomographic volume. **C-E)** Membranograms showing both the luminal and stromal sides of the three membranes highlighted in A. Color strips indicate the membrane domain (blue: appressed, yellow: non-appressed). Stromal lamellae and end membrane are annotated above membranogram. All membranograms show densities 2 nm above the membrane surface (see F for reference). There are sharp boundaries between densities in the appressed and non-appressed regions. PSII densities are seen only on the luminal side of the appressed membranes, whereas PSI densities are seen only on the stromal side of non-appressed membranes. **F)** Top: Diagram showing side views of structures (blue: PSII, orange: *cytb₆f*, green: PSI, pink: ATPsyn) in a thylakoid membrane (grey). Bottom: membrane surface views showing densities that protrude from each structure 2 nm into the thylakoid lumen or stroma. **G)** Cartoon representation of the stromal and luminal surfaces of the membrane in E, with particle footprints pasted in (colored as in F).

respectively) prevents them from entering the narrow stromal gap between appressed thylakoids (~3.2 nm, **Fig. 4**). We were able to visualize particle distributions in appressed and non-appressed regions by projecting the tomogram volumes onto both the stromal and luminal surfaces of segmented membranes (**Fig. 5A**; Sup. Fig. 3) to generate “membranograms” (**Fig. 5C-E**, **Fig. 6A**). Similar to *C. reinhardtii* [2] and as observed in other systems [28, 47, 48], large densities corresponding to the protruding oxygen evolving complexes (OEC) of PSII are visible on the luminal side of the appressed membranes (e.g., **Fig. 3**, **Fig. 5C-D**; Sup. Fig. 5D). These OEC densities mark the center positions of large membrane-embedded supercomplexes of PSII bound to trimers of LHCII [8, 49, 50], which do not extend from the membrane and thus are not visible in the membranograms. Interspersed between the PSII densities, we observe smaller dimeric particles that are consistent with the size of the lumen-exposed domains of *cytb₆f*, confirming the long-debated presence of this complex in appressed membranes of vascular plant thylakoids (e.g., [51]). In contrast, the stromal surface of the appressed membranes is generally smooth and featureless, consistent with the exclusion of PSI and ATPsyn, as well as the lack of large protrusions on the stromal faces of PSII, LHCII or *cytb₆f*. The PSII complexes abruptly disappear from the luminal face at the junction between appressed and non-appressed thylakoids (teal-to-yellow transitions in **Fig. 5C, E**; Sup. Fig. 3A). Instead, both the luminal and stromal surfaces of the non-appressed thylakoids are covered by high number of smaller densities (**Fig. 5C**; Sup. Fig. 3). The identity of these densities is difficult to assign with confidence, but a majority of them likely correspond to PSI (on the stromal side) and *cytb₆f* (on the luminal side) (see diagram in **Fig. 5F**).

Biochemical isolation experiments often sub-divide the non-appressed membranes into stromal lamellae, grana margins, and grana end membranes on the basis of detergent solubilization and mechanical fragmentation, and have reported different protein compositions for the regions (for references see e.g., [26, 52]). In contrast, we observe little difference in the appearance and distribution of the densities on non-appressed membranes irrespective of whether they are immediately adjacent to grana (margins), extend away from them (stromal lamellae), or cap the grana stacks (grana end membranes) (**Fig. 5C, E**; Sup. Fig. 3). Rather, we observe a simpler two-domain organization of photosynthetic complexes segregated into appressed and non-appressed regions, which corresponds to the original lateral heterogeneity model put forward by Andersson and Anderson [42, 53, 54]. It should be noted that the edges (tips) of the appressed membranes are difficult to visualize with membranograms, although their highly curved nature makes it unlikely that they could accommodate large photosynthetic complexes (see zoomed-in views of thylakoid tips in **Fig. 1D**, **Fig. 4** and Sup. Fig. 2B-D).

Using AI-assisted approaches [3, 35] confirmed by manual inspection, we detected essentially all particles populating appressed spinach membranes ($[All_{Spin}] = 2160 \text{ particles}/\mu\text{m}^2$). Dimeric PSII complexes were clearly identified in the tomograms ($[PSII_{Spin}] = 1415 \text{ particles}/\mu\text{m}^2$), enabling us to assign their center positions and orientations (**Fig. 6A, C**) [55]. In the best quality tomograms, we were also able to manually assign dimeric *cytb₆f* particles in the appressed regions ($[cytb_{6f}_{Spin}] = 446 \text{ particles}/\mu\text{m}^2$) (**Fig. 6A**). Note that the total particle concentration, but not the PSII and *cytb₆f* subclasses, includes unknown densities as well as clipped particles at the edges of membrane patches selected for analysis. The PSII concentration and mean center-to-center distance between nearest neighbor complexes ($PSII-NN_{Spin} = 21.2 \pm 3.1 \text{ nm}$) (**Fig. 6B**) is similar to what was reported in *Arabidopsis thaliana* by freeze fracture EM [56]. In contrast, our previous cryo-ET measurements of *C. reinhardtii* showed a slightly lower PSII concentration and correspondingly longer nearest neighbor distances within appressed thylakoid regions ($[PSII_{Chlamy}] = 1122 \text{ PSII}/\mu\text{m}^2$, $PSII-NN_{Chlamy} = 24.4 \pm 5.2 \text{ nm}$) [2]. This discrepancy is in agreement with the relative sizes of the PSII-light harvesting complex II (PSII-LHCII) supercomplexes isolated from the respective species, with $C_2S_2M_2L_2$ supercomplexes from *C. reinhardtii* having a larger footprint in the membrane than $C_2S_2M_2$ supercomplexes from vascular plants [57–59]. The concentration of PSII complexes is consistent between membranes (**Fig.**




Figure 6.

Distribution and concentration of PSII in grana membranes.

A) Top: membranogram view of an appressed membrane piece with characteristic densities of dimers of PSII and *cytb₆f*. Bottom: same membrane with models of PSII and *cytb₆f* complexes fitted into the EM densities. **B)** Nearest neighbor distance plot of all the particles from the appressed membranes. Blue: PSII, orange: *cytb₆f*, grey: all particles. Note that "all" also includes unassigned particles of unknown identity. The dotted line labeled PSII-NN_{spin} indicates the mean ($\pm$ SEM) center-to-center distance between neighboring PSII cores. Nearest neighbor distances are much more variable between two *cytb₆f* than between two PSII complexes. Identification of *cytb₆f* particles was possible only in the best resolved membranes; therefore, particle number and distribution estimation are not as reliable. **C)** Segmentation of a granum with all detected PSII particles pasted in. Note that we did not analyze PSII distribution at the very tips of thylakoids; this is why the tips look unoccupied. **D)** Plot showing PSII nearest neighbor center-to-center distances (light blue) and concentration (dark blue) as a function of the particle position from the edge of the granum, in 10 nm bins (diagrammed above the plot). Dot size represents number of counts in each bin. **E)** Plot of the lengths of grana thylakoids selected for the analysis in D. Thylakoids of significantly different lengths were selected to minimize any potential effect of the membrane size on the PSII distribution.

6B) and is independent of distance to the edge of the granum and thylakoid position in the stack (**Fig. 6E** shows lengths of thylakoids analyzed in **Fig. 6D**; we sampled membranes with variable lengths to depict the entire range of thylakoid sizes).

Using the observable orientations of OEC densities within the membrane plane, we placed footprints into the segmented membranes corresponding to structures of the C_2 dimeric PSII core complex, the C_2S_2 PSII-LHCII supercomplex (isolated from spinach [58], PDB: 3JCU), and the $C_2S_2M_2$ PSII-LHCII supercomplex (isolated from *A. thaliana* [59], PDB: 7OUI). When considering the PSII cores alone (C_2), only rare events of complexes touching were observed (**Fig. 7A** top row, D). Placing in the C_2S_2 supercomplexes resulted in a small in-plane clash between some particles (~2.1% overlap of total footprint area, **Fig. 7A** second row, D). The steric hindrance increased to ~8.2% overlap when larger $C_2S_2M_2$ supercomplexes were used (**Fig. 7A** third row, D). Depending on which footprint was mapped in, the total membrane surface coverage by the PSII cores and the PSII-LHCII supercomplexes ranged from ~20% to ~55% (**Fig. 7B**), leaving ample space in the membrane for additional free LHCII trimers and *cytb₆f*. Our observation of some clash between $C_2S_2M_2$ supercomplexes suggests that different supercomplex variants probably coexist within the same membrane. Cumulative coverage of C_2 , C_2S_2 , and $C_2S_2M_2$ footprints when overlaying sequential membranes across the granum (known as the “granum cross-section”) resulted in complete cross-section coverage after minimum of 6 membranes (3 consecutive thylakoids) in the case of the biggest supercomplex (**Fig. 7C**).

We do not observe instances where PSII complexes directly face each other across the luminal gap (**Fig. 8A-B**). Rather, PSII complexes interdigitate despite the thylakoid lumen being wide enough to accommodate two directly apposed complexes (**Fig. 4**). Indeed, the luminal overlap is negligible between all visible densities (including *cytb₆f* and unknown particles) for the two membranes of an appressed grana thylakoid (e.g., M19-M20 and M21-M22 membrane pairs in **Fig. 8B**). Similarly, across the stromal gap, the overlap of all densities is only slightly higher (~4%) (e.g., M20-M21 membrane pair in **Fig. 8B**). Performing this stromal overlap quantification with supercomplex footprints (**Fig. 7E**) showed that PSII cores have 9% of their surface area overlapping, whereas C_2S_2 and $C_2S_2M_2$ supercomplexes show 22% and 38% overlap of their area, respectively (**Fig. 7F**). Within intact chloroplasts, we did not observe patterns of PSII supercomplexes aligning across appressed membranes (**Fig. 8A-B**), contrary to the ordered lattices of PSII previously observed in thylakoids isolated from pea [28]. However, in tomograms of chloroplasts ruptured during blotting (e.g., Sup. Fig. 1F), we did observe that PSII densities sometimes align into semi-crystalline lattices with strong overlap between the PSII complexes from adjacent appressed membranes (**Fig. 8C-D**). It is therefore plausible that extraction of thylakoids away from the chloroplast environment can induce the formation of PSII lattices between appressed membranes, some of which may have undergone restacking.

Conclusions and Speculation

Our cryo-ET study of intact spinach chloroplasts provides insights that span scales from the level of local membrane architecture (fine 3D ultrastructure of individual grana and connecting stromal lamellae) to the level of how individual protein complexes are organized within these membranes.

The architecture of spinach thylakoids visualized here is consistent with the previously proposed helical staircase models [24, 60, 61], although the structural details we observe are perhaps less stringent. The grana stacks of appressed membranes are not completely cylindrical: they have a wavy circumference (when viewed from the top, e.g., **Fig. 3**), vary in diameter along the stack, and sometimes nearly merge with one another. We observe a variety of membrane architectures at the interconnections between appressed and non-appressed regions. Sometimes, the appressed domains shift smoothly into the non-appressed domains without architectural contortions (**Fig. 5**; Sup. Fig. 3C), similar to what was observed in *C. reinhardtii*. However, much




Figure 7.

Occupancy of PSII and PSII-LHCII supercomplexes in grana membranes.

A) Top panel: cartoon representation of an analyzed membrane patch (M4, fourth membrane in a granum) with footprints of all PSII core complexes (C_2) placed into their respective positions. Middle panels: the same membrane patch with footprints of C_2S_2 and $C_2S_2M_2$ PSII-LHCII supercomplexes placed in. Bottom panel: red outlines highlight possible in-plane clashes between different types of supercomplexes within the same membrane (outlines carried down from panels above). **B)** Quantification of the membrane coverage, depending on the type of supercomplexes placed in (the area of S_2 LHCII trimers in supercomplexes is also shown). Reproduced results from the *C. reinhardtii* study (Chlamy) shown for comparison [24]. Error bars: SD. **C)** Cumulative membrane coverage of PSII supercomplexes throughout the stack. Each line in the plot represents the percentage of area occupied by PSII or different PSII-LHCII supercomplexes when adding membranes across the stack (from the first stacked membrane until the end of the granum). Line: mean, shading: SEM. Membranes from 15 grana were used for the analysis. **D)** Quantification of the in-plane clashes between PSII cores or different supercomplexes. PSII-PSII clash: 0.0002% (within error range); these might be complexes in direct contact like the example in A. **E)** Top: cartoon representation of the fifth membrane (M5) from the same granum with all C_2S_2 supercomplexes in place. Bottom, overlay of M4 (blue) and M5 (magenta) showing the overlap (white) between PSII-LHCII in appressed membranes separated by a stromal gap. **F)** Quantification of the stromal overlap between PSII cores, LHCII S_2 trimers, and different supercomplex variants. For all plots, data was generated from 3 chloroplast preparations, using the best resolved membranes ($n=173$).




Figure 8.

Overlap of densities across stacked membranes: intact vs. broken chloroplasts.

A) Tomographic slice showing a typical granum inside an intact chloroplast. White box marks the stack of membranes visualized in B. **B)** Above: segmentation of the piece of a granum from A with four membranes highlighted (M19-M22). Odd numbered membranes in magenta, even numbered in green. Below: corresponding membranogram overlays of the adjacent membrane pairs. Coloring represents all EM densities protruding ~2 nm from the luminal side of each membrane. The three examples show two luminal overlaps (top and bottom) and an overlap across a stromal gap (middle). White: overlap between EM densities. **C)** Tomographic slice showing a part of a broken chloroplast, containing individual membranes and potentially re-stacked thylakoids. White box marks the pieces of two membranes from adjacent thylakoids. **D)** Top: membranogram from a piece of one appressed membrane in C, showing densities protruding ~2 nm from the membrane's luminal surface. Note the arrays of PSII complexes and absence of any other type of particles. Bottom: membranograms from the pair of appressed membranes (green and magenta) overlayed with each other. White: overlap between EM densities.

more complex transitions were also observed, with narrow membrane bridges or sharp forks (Sup. Fig. 2). At this level of membrane architecture, we also observed thylakoid contacts with plastoglobules (Fig. 2) and the chloroplast envelope (Sup. Fig. 4). The pristine sample preservation inside vitrified chloroplasts, combined with the high resolution of our cryo-ET imaging, allowed us to measure thylakoid morphometrics (bilayer thickness, stromal and luminal gaps) with sub-nanometer precision (Fig. 4).

At the level of molecular organization, the appressed and non-appressed membranes have markedly different composition, clearly visualized by the protruding protein densities in membranograms (Fig. 5; Sup. Fig. 3). Similar to *C. reinhardtii* [2], there is a precise segregation between the two domains, with PSII restricted to the appressed membranes and PSI restricted to the non-appressed membranes. *Cytb_f* is present in both domains. In contrast to previous biochemical studies, no obvious difference was seen between sub-regions of the non-appressed membranes (stromal lamellae, grana margins, grana end membranes) (Fig. 5C, E; Sup. Fig. 3). Similarly, our analysis of PSII distribution in the appressed grana membranes showed that these complexes are uniformly distributed through the stack (Fig. 6, Fig. 7), with no changes in concentration observed across the grana layers nor close to the granum edge, where it meets the non-appressed membrane region. This confirms that, at least on the level of PSII distribution, there are no organizational differences across the granum.

Together, these observations support a simple two-domain model for thylakoid lateral heterogeneity *in vivo*, which states that 1) the appressed and non-appressed domains each have a homogeneous mix of their respective proteins throughout, and 2) there is an abrupt and discrete transition between the domains, with no visible mixing between the PSI and PSII complexes. Nonetheless, the domain boundary (grana margin) must be sufficiently fluid to facilitate the exchange of protein complexes during dynamic processes such as PSII repair and state transitions.

In the intact chloroplasts, we did not observe ordered lattices of PSII that were previously reported in less intact samples from various species [28, 47, 62, 63]. However, we did see semi-crystalline arrangements of PSII complexes in chloroplasts that had ruptured on the EM grid (Fig. 8). Rearrangement of PSII into ordered arrays may therefore be triggered by exposure of the thylakoid network to a non-native environment with different buffering properties [64–66]. It remains to be determined whether PSII lattices could also have a physiological function *in vivo* [67] (to the best of our knowledge, the only *in situ* report of PSII arrays is in the dehydrated state of the resurrection plant *Craterostigma pumilum* [68]).

In situ cryo-ET only provides snapshots of living systems, but perhaps we can infer hints about the dynamic nature of thylakoids from these frozen moments in time. Our results show that PSII-LHCII supercomplexes do not align with each other under standard growth conditions. This distribution is consistent with supercomplexes mixing freely within the appressed domains without persistent static interactions with other supercomplexes. Based on evidence that LHCII plays an important role in thylakoid stacking [28, 44, 69–72], we hypothesize that while LHCII trimers can make relatively stable in-plane interactions to form PSII-LHCII supercomplexes, the other interactions within an appressed membrane and across the stromal gap to an adjacent stacked membrane are more transient. LHCII trimers are intrinsically multivalent and make electrostatic interactions with each other across the stromal gap using their disordered, positively-charged N-terminus [72]. Such a network of weak and multivalent interactions is reminiscent of the organizing principles of liquid-liquid phase separation [73, 74]. We speculate that stacked thylakoids in both green algae and plants are a condensation of membrane-embedded PSII-LHCII supercomplexes and free LHCII proteins. This condensation is driven by an ensemble of interactions both within the membrane plane and between stacked membranes (the latter via the N-terminus of LHCII), which are modulated by macromolecular crowding effects and membrane surface charge, respectively [66, 75, 76]. In contrast to canonical globular condensates, we propose that thylakoid stacks are multiple interacting layers of planar condensates. In addition to

being a main driving force for the condensation of LHCII and PSII, these interactions between membrane layers also likely provide the mechanism for sterically excluding ATPsyn and PSI from the condensate, while permitting entry of *cytb₆f* because it lacks a large stromal density. An important prediction of this stacking condensation model is that PSII, LHCII, and *cytb₆f* are mobile within the appressed regions; however, we lack the methodology to directly track movements of single protein complexes within a native grana membrane.

Limitations and Future Perspectives

Plant cells are generally difficult to vitrify and are too large for conventional FIB milling workflows. We opted for chloroplast extraction and plunge-freezing (Sup. Fig. 1), which facilitated the preparation, however not without disadvantages. The minutes-long isolation protocol can lead to relaxation of the physiological state of the chloroplasts, especially if it is temperature- or light-dependent. Moreover, plunge-freezing on EM grids can rupture the chloroplast envelope (Sup. Fig. 1B, F) and promote preferential orientation of the plastids — most often the entire thylakoid network lies flat on the surface of the grid support. Flat membranes, orthogonal to the electron beam during cryo-ET acquisition, provide top views (**Fig. 3**, Sup. Fig. 5) that suffer from missing-wedge effects and are difficult to segment and analyze. We collected most of the tomograms towards the edges of the chloroplasts where this effect is less pronounced (more frequent side views), but this approach could induce region-specific bias.

Although more technically demanding, recent advances in cryo-FIB lift-out of high-pressure-frozen multicellular samples [77, 78] open future possibilities to perform cryo-ET on chloroplasts within native leaf tissue. Still, the relatively low throughput of cryo-ET and intrinsically small volume imaged by this technique present challenges for answering micron-scale questions about the overall membrane ultrastructure of the chloroplast; those questions are being successfully tackled by other microscopy techniques that image larger sample volumes at the trade-off of resolution (e.g., [22–24, 27, 60, 79, 80]). A more complete picture of thylakoid networks across length and time scales will require integrating the native molecular views provided by cryo-ET with complementary structural and cellular techniques [81]. Thylakoids are dynamic systems that adapt their molecular architecture to modulate their function in response to environmental changes. A future challenge is to accurately replicate this complexity with mesoscale modeling that incorporates parameters from diverse experiments to bridge thylakoid structure and function.

Materials and Methods

Plant growth

Spinacia oleracea WT plants were germinated and grown in a growth chamber for 6 weeks with a 12 h day/night cycle at 21–23°C and white-light illumination (150 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$). One day before the experiment, the night cycle was extended to 20 h to minimize starch content in the chloroplasts. Prior to chloroplast isolation, plants were transferred from darkness to illumination for 1 h to acclimate them to light. This experiment was performed in three independent batches.

Chloroplast isolation

Approximately 80 grams of fresh leaf mass was collected and immediately blended with 160 mL of isolation buffer (0.45 M Sorbitol, 20 mM Tricine-KOH, 10 mM EDTA, 10 mM NaHCO_3 , 5 mM MgCl_2 , 0.1% BSA, 0.2% D-Ascorbate) on ice. The slurry was filtered twice through cotton and a double layer of Miracloth to remove debris. The filtrate was then deposited on a 50% percoll cushion in isolation buffer and centrifuged for 8–10 mins at 7500 rpm. The green band in between percoll

layers was collected and diluted in the isolation buffer at least 3 times (dilution depended on the efficiency of purification and type of the EM grids to be used). Chloroplast were plunge-frozen immediately afterwards.

Plunge-freezing

Plunging was performed using a Vitrobot Mark 4 (Thermo Fisher Scientific). 4 μL of chloroplasts solution was placed onto R2/1 or R1.2/1.3 carbon-coated 200-mesh copper EM grids (Quantifoil Micro Tools), blotted for 7-8 seconds from the back side (front side pad covered with a Teflon sheet) with blot-force 10, and plunge-frozen in a liquid ethane/propane mixture cooled with liquid nitrogen. Grids were clipped into “autogrid” support rings modified with a cut-out on one side (FEI, Thermo Fisher Scientific) and stored in plastic boxes in liquid nitrogen until used for cryo-FIB milling.

Focused Ion Beam milling

Cryo-FIB milling was performed as described previously [31] using an Aquilos dual-beam FIB/SEM instrument (Thermo Fisher Scientific). Grids were screened for good chloroplast coverage and coated with a layer of organometallic platinum using a gas injection system to protect the sample surface. The sample was milled with a gallium ion beam to produce $\sim 100\text{--}150\text{ nm}$ -thick lamellae. Some batches of lamellae were additionally sputter-coated with platinum at the end of the milling process. Grids were then transferred in liquid nitrogen to the TEM microscope for tomogram acquisition.

Cryo-ET data acquisition

Tomographic data was collected on a 300 kV Titan Krios TEM instrument (Thermo Fisher Scientific), equipped with a post-column energy filter (Quantum, Gatan), and a direct detector camera (K2 Summit, Gatan). Tilt series were acquired using SerialEM software [82] with a dose symmetric scheme [83] starting at the pretilt of $10\text{--}10^\circ$ (depending on the loading direction of the grids), with 2° increment between tilts and spanning a total of 120° . Individual tilts were recorded in counting mode with an imaging rate of 8-12 frames per second, at a pixel size of $3.52\text{ \AA}$ and target defocus in the range of $2.5\text{--}5\text{ }\mu\text{m}$. The total accumulated dose for the tilt-series was kept at approximately $120\text{ e}/\text{\AA}^2$.

Tomogram reconstruction and processing

All tilt-series were processed using the TOMOMAN pipeline [84]. Raw frames were aligned using MotionCor2 (v.1.4.0) [85]. Bad tilts were manually removed, and cleaned tilt-series were dose-weighted. The assembled tilt-series were aligned with patch tracking using the Etomo program from the IMOD (v.4.11.1) package [86, 87]. Final tomograms were reconstructed by weighted back projection in IMOD. To enhance contrast, the cryo-CARE denoising was applied [36]. For the views in Fig. 3, we instead performed denoising with DeepDeWedge, which additionally attempts to fill in missing wedge information [37]. All images of tomograms (tomographic slices) shown were acquired using the IMOD 3dmod viewer.

Membrane segmentation

Membrane segmentation was performed using AI-assisted approaches as described in [3]. Briefly, in order to generate membrane segmentations, we initially segmented the tomograms using the TomoSegMemTV segmentation protocol [88], and extracted patches of 160^3 voxels of tomogram-segmentation pairs. After manually cleaning these segmented patches, we used these pairs to train a neural network (nnU-Net) to segment the entire thylakoid lattice. We predicted segmentations using the trained network, extracted and corrected new membrane patches, merged these into our training set, and started a new round of training. We followed this iterative approach until we were satisfied with the performance of our segmentation model. This model is

now available in the MemBrain-seg package [3]. All segmentations shown in the figures were generated using this model, then cleaned in Amira (Thermo Fisher Scientific) and rendered in UCSF ChimeraX.

Preparation of single-membrane segmentations

For all analyses requiring single membrane instances or stacks of individual membranes (e.g., particle organization in the membrane and overlays between membranes), the separate segmentations were made by cutting the full-tomogram segmentations from MemBrain-seg in Amira and exporting them as separate volume files (.mrc) and triangulated surface meshes.

Measurement of appressed vs. non-appressed areas

For the quantification of appressed vs. non-appressed regions (shown in Fig. 1E-F), we first segmented all membrane networks in all our tomograms using our trained MemBrain-seg model and then manually cleaned the output segmentations in Amira to remove tomogram edge artifacts. Afterwards, we converted the segmentations into triangular meshes using PyVista's marching cubes algorithm [89], followed by Laplacian smoothing, giving normal vectors for each triangle on the mesh. For each triangle, we traced the distance to the next triangle along the normal vector. This next triangle along the normal belongs to potentially neighboring membranes, and by thresholding the distance, we divided the meshes into “appressed” and “non-appressed” triangles. To account for noisy distance predictions, we applied a majority vote among geodesically neighboring triangles to receive a smooth representation of appressed and non-appressed regions. Furthermore, by computing the principal curvature at each triangle, we excluded the membrane segmentation edges (e.g., clipped by the tomogram volume or ending due to mis-segmentation), as these have normal vectors pointing away from the stack and therefore would distort the metrics towards more non-appressed regions. By summing up the triangle areas of all remaining mesh triangles for both classes, we received the total areas of appressed and non-appressed membranes.

Thylakoid spacing and thickness calculations

In order to measure distances between different membrane layers, stromal and luminal gaps, as well as entire thylakoid thickness (shown in Fig. 4), we utilized membrane flattening functionalities from blik [90] to process our membrane dataset. The goal of this step was to receive a “stack” of 2D images for each membrane, where each 2D image in the z-stack corresponds to the flattened front view of the membrane at a different distance along the membrane normal vector. To perform the flattening, we first sampled a grid of points on each 3D membrane segmentation. This grid allowed blik to generate the 2D image stacks along the normal vectors. By averaging the intensities of each 2D image in these stacks, we received a single value for each z-coordinate, leading to a 1D plot along the z-axis, i.e., the membrane normal vector. From this 1D plot, we extracted local intensity minima and maxima corresponding to center points of lipid bilayers. After manual cleaning of outlier plots (i.e., plots where the automated assignment of keypoints failed), this representation allowed us to estimate the distances between different membranes and gaps.

Membrane protein ground truth positions

For training MemBrain, our membrane protein localization network [3], we required ground truth positions for the PSII proteins. To generate them, we imported the single membrane .obj files into Membranorama (<https://github.com/dtegunov/membranorama>) to visualize the membranes and their embedded proteins by projecting the tomogram densities onto the surface mesh. In this program, we were able to manually annotate PSII positions and orientations in 45 membrane patches from seven different tomograms, which we then used as training data for the MemBrain model.

MemBrain training

We utilized our membrane protein ground truth data to train a MemBrain model. To do this, we split our full dataset consisting of 45 membranes into a training set (28 membranes), a validation set (7 membranes), and a test set (10 membranes). As a required first step in the MemBrain workflow, we manually defined the side of each membrane facing toward the thylakoid lumen as the side of interest. Then, we trained MemBrain using its default parameters to predict “heatmaps” depicting each membrane point’s distance to the closest PSII. Using our ground truth annotations (validation set), we further tuned the best bandwidth parameter for the subsequent Mean Shift clustering, which we determined to be 21 bin4 voxels, corresponding to 29.6 nm

Application of trained MemBrain model

After training and tuning MemBrain, we applied it to the full dataset, consisting of 455 discrete membrane segmentations from the nine best quality tomograms. This gave us good initial predictions for positions and orientations of PSII complexes in all membranes. In order to confirm their validity, we inspected all 455 membranes again, discarded 152 membranes due to bad quality, and manually corrected false MemBrain predictions. The remaining 303 membranes were used for the analysis. Wherever image quality allowed, we also annotated *cytb₆f* positions. This gave us an accurate membrane protein dataset, consisting of 14433 PSII, 937 *cytb₆f*, and 5934 positions of unknown particles (non-identifiable densities).

Center-to-center distances

In order to analyze the nearest neighbor distances between our particle positions, we first converted each voxel-wise membrane segmentation into a surface mesh representation using PyVista’s marching cubes algorithm, followed by Laplacian surface smoothing (applied to individual single membrane instances). This allowed us to compute exact geodesic distances between two positions on the mesh using The Virtual Brain’s (<https://github.com/the-virtual-brain>) implementation of the discrete geodesic distance algorithm [91]. For each particle location, we computed the geodesic distance to the center position of the nearest neighbor from the respective target particle class (e.g., PSII-NN_{Spin}, *cytb₆f*-NN_{Spin}, all-NN_{Spin}; shown in Fig. 6B).

Distance to appressed region edge

To analyze the change of protein distribution with respect to distance from the edge of appressed membrane regions, we first annotated these membrane edges manually using Membranorama. We then divided the Euclidean distance to the membrane edge into 10nm- sized bins and computed the total area of mesh triangles within each bin. By dividing the number of particle positions by the triangle area within each bin, we obtain average protein complex concentrations per bin. Analogously, for each bin, we computed the average nearest neighbor distances. Plotting these concentrations against the respective bin distances provided the graphs displayed in Fig. 6D.

Overlap analysis

To analyze the overlap between PSII-LHCII supercomplexes in neighboring membranes (Fig. 7E and F) or cumulative occupancy in membranes across the stack (Fig. 7C), we projected protein positions and orientations onto a common membrane surface. We calculated their overlaps by comparing the overlaying triangles on the segmented membrane. To process a stack of membranes, we calculated a connection vector between the first and last membrane of the stack by computing the average membrane normal vector for the starting membrane. Then, for each membrane, we projected the corresponding membrane protein positions along the normal vector onto the starting membrane mesh. Mapping the protein complex footprint (based on its 3D structure filtered to 14 Å resolution) into the correct orientation in this new position allowed us to


calculate the area on the membrane surface that was occupied by the respective complexes. Doing this for all membranes gave us a set of occupied areas per membrane, which we then analyzed for overlapping areas. To avoid bias due to membrane misalignment (e.g., stack being inclined in the tomogram slab resulting in incomplete overlap between membranes), we also projected the segmentations of all membranes onto the starting mesh and only analyzed the areas that were shared by all projected membrane segmentations.

Data availability

The raw cryo-ET data will be available at the Electron Microscopy Public Image Archive (EMPIAR), accession code: EMPIAR-XXXXX. Denoised tomograms and segmentations shown in the figures will be deposited in the Electron Microscopy Data Bank (EMDB): EMD-XXXXX.

Acknowledgements


We thank the Plant and Greenhouse facility of LMU Munich and especially Anja Schneider for help with the cultivation of plants. We thank Jürgen Plitzko and Wolfgang Baumeister at the Max Planck Institute of Biochemistry for access to electron microscopes. We thank Sebastian Ziegler and Fabian Isensee from the Applied Computer Vision Lab (ACVL) at the German Cancer Research Center (DKFZ) for their support in designing the membrane segmentation framework. L.L. also thanks Julia Schnabel from the Institute of Machine Learning in Biomedical Imaging (IML) at Helmholtz Munich for advice and supervision. Cryo-ET analysis was performed at the sciCORE (<http://scicore.unibas.ch/>) scientific computing center at the University of Basel. B.D.E. and W.W. acknowledge funding from a Human Frontier Science Program (HFSP) research grant (award number RGP0005/2021) and European Research Council (ERC) consolidator grant “cryOcean” (fulfilled by the Swiss State Secretariat for Education, Research and Innovation, award number M822.00045). M.P.J. was supported the grants from Biotechnology and Biological Sciences Research Council (BBSRC) (award number BB/V006630/1) and the Leverhulme Trust (award numbers RPG-2019-045 and RPG-2021-345). L.M. was supported by a BBSRC White Rose DTP studentship in Mechanistic Biology. L.L. acknowledges support from the Munich School for Data Science (MUDS) and a fellowship from the Boehringer Ingelheim Fonds.



Supplemental Figure 1.

Sample preparation, tomogram selection on the FIB-milled lamellae, and types of tomograms collected.


A) Simplified schematic of the chloroplast preparation protocol, which aimed to minimize the time between chloroplast isolation and freezing. **B)** An overview of a typical FIB-milled lamella containing chloroplasts as seen in the TEM microscope. Chloroplasts can break and sometimes re-close during the isolation and plunging procedures. We tried to identify intact, spheroidal chloroplast sections in the lamellae and focused tomogram collection in those regions. White arrows point the grana stacks, visible in the overviews. **C-F)** Examples of tomograms collected in different regions of an intact chloroplast (C-E) as well as in a region containing a broken chloroplast (F).



Supplemental Figure 2.

Ultrastructure of grana sides and transition to stromal lamellae.


A) Segmentation of thylakoid membranes from two grana in close proximity. Note the irregular stacks and shifted thylakoids. **B)** View of the margin region on one side. Grana-forming thylakoids are flat and straight, whereas stromal lamellae are more undulating. **C)** View of the other side. Note the convoluted shape of stromal lamellae connecting with two other lamellae spanning from the same granum. The top part of the membrane segmentation was set to high transparency to improve visibility. **D)** Serial perpendicular slices through the granum, showing appressed membranes transitioning into stromal lamellae. Membranes merge and form bridges between each other.



Supplemental Figure 3.

Protein densities on two domains of thylakoids.


A) A 3D rendering of a thylakoid granum and stromal lamellae with tomographic densities projected onto the membrane surfaces (membranograms). The first membrane of the granum (M1) is separated from the stack to show the PSII and *cyt.*b₆f** densities in the appressed region of M2 below. Note that all non-appressed membranes (including grana margins, end membranes, and stromal lamellae) are populated with similar densities. **B)** and **C)** Zoom-in views of stromal lamellae extending from the side of the granum. The eye icon shows the viewing direction in C. All panels show membranes at the same threshold value and the same color scale.



Supplemental Figure 4.

Thylakoid-chloroplast envelope contacts.

Rare instances of thylakoids and thylakoid-like membranes approaching and contacting the inner envelope membrane inside intact chloroplasts. **A-C)** Tomographic slices showing overviews of the contact sites (white arrowheads). **D-F)** Zoom-ins of the contacts from respective images. Dashed box in B corresponds to the enlarged view in E.



Supplemental Figure 5.

Top view of a granum.

Tomographic slices from a tomogram denoised with DeepDeWedge program [37] (high contrast, missing wedge inpainting) showing the chloroplast stroma 15 nm above the granum (A), 7 nm above the granum (B), and 3 nm above the top membrane (C), as well as the luminal space between the top and second membranes of the granum (D). Note the high number of globular particles (mostly Rubisco) in A. Arrowheads indicate complexes appearing in the slices (Yellow: ribosomes, purple: ATPsyn, Green: putative PSI, Blue: PSII). Note the similar size of Rubisco and the ATPsyn F_1 subunit, which complicates assignment. The tomographic volume was rotated so the top membrane was approximately parallel to the xy-plane.

References

1. Engel B.D., et al. (2015) **Native architecture of the Chlamydomonas chloroplast revealed by in situ cryo-electron tomography** *eLife* 4
2. Wietrzynski W., et al. (2020) **Charting the native architecture of Chlamydomonas thylakoid membranes with single-molecule precision** *eLife* 9
3. Lamm L., et al. (2024) **MemBrain v2: an end-to-end tool for the analysis of membranes in cryo-electron tomography** *bioRxiv*
4. Wietrzynski W., Engel B.D. (2023) **Chapter 23 - Supramolecular organization of chloroplast membranes** In: Grossman A.R.Wollman F.-A., editors. *The Chlamydomonas Sourcebook* London: Academic Press pp. 763–785
5. Ostermeier M., et al. (2024) **Structure, biogenesis, and evolution of thylakoid membranes** *The Plant Cell* 36:4014–4035
6. Rippka R., Waterbury J., Cohen-Bazire G. (1974) **A cyanobacterium which lacks thylakoids** *Archives of microbiology* 100:419–436
7. Rahmatpour N., et al. (2021) **A novel thylakoid-less isolate fills a billion-year gap in the evolution of Cyanobacteria** *Current Biology* 31:2857–2867
8. Perez-Boerema A., Engel B.D., Wietrzynski W. (2024) **Evolution of Thylakoid Structural Diversity** *Annual Review of Cell and Developmental Biology* 40:169–193
9. Austin J.R., Staehelin L.A. (2011) **Three-Dimensional Architecture of Grana and Stroma Thylakoids of Higher Plants as Determined by Electron Tomography** *Plant Physiology* 155:1601–1611
10. Nevo R., et al. (2012) **Composition, architecture and dynamics of the photosynthetic apparatus in higher plants** *The Plant Journal* 70:157–176
11. Gounaris K., Barber J., Harwood J.L. (1986) **The thylakoid membranes of higher plant chloroplasts** *Biochemical Journal* 237:313–326
12. Kitajima M., Butler W.L. (1975) **Quenching of chlorophyll fluorescence and primary photochemistry in chloroplasts by dibromothymoquinone** *Biochim Biophys Acta* 376:105–15
13. Murata N (1969) **Control of excitation transfer in photosynthesis. II. Magnesium ion-dependent distribution of excitation energy between two pigment systems in spinach chloroplasts** *Biochimica et biophysica acta* 189:171–181
14. Garty Y., et al. (2024) **Thylakoid membrane stacking controls electron transport mode during the dark-to-light transition by adjusting the distances between PSI and PSII** *Nat Plants* 10:512–524
15. Wood W.H.J., et al. (2018) **Dynamic thylakoid stacking regulates the balance between linear and cyclic photosynthetic electron transfer** *Nat Plants* 4:116–127

16. Puthiyaveetil S., et al. (2014) **Compartmentalization of the protein repair machinery in photosynthetic membranes** *Proceedings of the National Academy of Sciences* **111**:15839–15844
17. Johnson M.P., et al. (2011) **Photoprotective Energy Dissipation Involves the Reorganization of Photosystem II Light-Harvesting Complexes in the Grana Membranes of Spinach Chloroplasts** *The Plant Cell* **23**:1468–1479
18. Kyle D.J., Staehelin L.A., Arntzen C.J. (1983) **Lateral mobility of the light-harvesting complex in chloroplast membranes controls excitation energy distribution in higher plants** *Archives of Biochemistry and Biophysics* **222**:527–541
19. Wood W.H.J., et al. (2018) **Dynamic thylakoid stacking regulates the balance between linear and cyclic photosynthetic electron transfer** *Nature Plants* **4**:116–127
20. Staehelin L.A. (2003) **Chloroplast structure: from chlorophyll granules to supra-molecular architecture of thylakoid membranes** *Photosynth Res* **76**:185–96
21. Staehelin L.A., Paolillo D.J. (2020) **A brief history of how microscopic studies led to the elucidation of the 3D architecture and macromolecular organization of higher plant thylakoids** *Photosynthesis Research* **145**:237–258
22. Shimoni E., et al. (2005) **Three-Dimensional Organization of Higher-Plant Chloroplast Thylakoid Membranes Revealed by Electron Tomography** *The Plant Cell* **17**:2580
23. Bos P.R., Berentsen J., Wientjes E. (2023) **Expansion microscopy resolves the thylakoid structure of spinach** *Plant Physiol*
24. Mustardy L., et al. (2008) **The three-dimensional network of the thylakoid membranes in plants: quasihelical model of the granum-stroma assembly** *Plant Cell* **20**:2552–7
25. Ruban A.V., Johnson M.P. (2015) **Visualizing the dynamic structure of the plant photosynthetic membrane** *Nature Plants* **1**:15161
26. Albertsson P.-Å (2001) **A quantitative model of the domain structure of the photosynthetic membrane** *Trends in Plant Science* **6**:349–354
27. Bussi Y., et al. (2019) **Fundamental helical geometry consolidates the plant photosynthetic membrane** *Proc Natl Acad Sci U S A* **116**:22366–22375
28. Daum B., et al. (2010) **Arrangement of Photosystem II and ATP Synthase in Chloroplast Membranes of Spinach and Pea** *The Plant Cell* **22**:1299–1312
29. Johnson M.P., et al. (2014) **Nanodomains of Cytochrome b6f and Photosystem II Complexes in Spinach Grana Thylakoid Membranes** *The Plant Cell* **26**:3051
30. Kirchhff H., Li M., Puthiyaveetil S. (2017) **Sublocalization of Cytochrome b(6)f Complexes in Photosynthetic Membranes** *Trends Plant Sci* **22**:574–582
31. Schaffer M., et al. (2017) **Optimized cryo-focused ion beam sample preparation aimed at in situ structural studies of membrane proteins** *Journal of Structural Biology* **197**:73–82
32. Lam V., Villa E. (2021) **Practical Approaches for Cryo-FIB Milling and Applications for Cellular Cryo-Electron Tomography** *Methods Mol Biol* **2215**:49–82

33. Nogales E., Mahamid J. (2024) **Bridging structural and cell biology with cryo-electron microscopy** *Nature* **628**:47–56
34. Young L.N., Villa E. (2023) **Bringing Structure to Cell Biology with Cryo-Electron Tomography** *Annu Rev Biophys* **52**:573–595
35. Lamm L., et al. (2022) **MemBrain: A deep learning-aided pipeline for detection of membrane proteins in Cryo-electron tomograms** *Comput Methods Programs Biomed* **224**:106990
36. Buchholz T.O., et al. (2019) **Cryo-CARE: Content-Aware Image Restoration for Cryo-Transmission Electron Microscopy Data** In: *2019 IEEE 16th International Symposium on Biomedical Imaging (ISBI 2019)*
37. Wiedemann S., Heckel R. (2024) **A deep learning method for simultaneous denoising and missing wedge reconstruction in cryogenic electron tomography** *Nature Communications* **15**:8255
38. Gupta T.K., et al. (2021) **Structural basis for VIPP1 oligomerization and maintenance of thylakoid membrane integrity** *Cell* **184**:3643–3659
39. Shanmugabalaji V., et al. (2022) **Chapter Three - Plastoglobules: A hub of lipid metabolism in the chloroplast** In: Rébeillé F., Maréchal E., editors. *Advances in Botanical Research* Academic Press pp. 91–119
40. Li M., et al. (2020) **Measuring the dynamic response of the thylakoid architecture in plant leaves by electron microscopy** *Plant Direct* **4**:e00280
41. Mazur R., Mostowska A., Kowalewska Ł. (2021) **How to Measure Grana – Ultrastructural Features of Thylakoid Membranes of Plant Chloroplasts** *Frontiers in Plant Science* **12**
42. Andersson B., Anderson J.M. (1980) **Lateral heterogeneity in the distribution of chlorophyll-protein complexes of the thylakoid membranes of spinach chloroplasts** *Biochim Biophys Acta* **593**:427–40
43. Miller K.R., Staehelin L.A. (1976) **Analysis of the thylakoid outer surface. Coupling factor is limited to unstacked membrane regions** *Journal of Cell Biology* **68**:30–47
44. Armond P.A., Staehelin L.A., Arntzen C.J. (1977) **Spatial relationship of photosystem I, photosystem II, and the light-harvesting complex in chloroplast membranes** *The Journal of cell biology* **73**:400–418
45. Vallon O., Wollman F.A., Olive J. (1986) **Lateral distribution of the main protein complexes of the photosynthetic apparatus in Chlamydomonas reinhardtii and in spinach: an immunocytochemical study using intact thylakoid membranes and a PS II enriched membrane preparation** *Photobiochemistry and Photobiophysics* **12**:203–220
46. Armond P.A., Arntzen C.J. (1977) **Localization and Characterization of Photosystem II in Grana and Stroma Lamellae** *Plant Physiol* **59**:398–404
47. Levitan O., et al. (2019) **Structural and functional analyses of photosystem II in the marine diatom Phaeodactylum tricornutum** *Proc Natl Acad Sci U S A* **116**:17316–17322

48. Li M., et al. (2021) **In situ cryo-ET structure of phycobilisome-photosystem II supercomplex from red alga** *eLife* **10**:e69635
49. Croce R., van Amerongen H. (2020) **Light harvesting in oxygenic photosynthesis: Structural biology meets spectroscopy** *Science* **369**
50. Grieco M., et al. (2015) **Light-harvesting II antenna trimers connect energetically the entire photosynthetic machinery — including both photosystems II and I** *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1847**:607–619
51. Pribil M., Labs M., Leister D. (2014) **Structure and dynamics of thylakoids in land plants** *Journal of Experimental Botany* **65**:1955–1972
52. Koochak H., et al. (2019) **The structural and functional domains of plant thylakoid membranes** *The Plant Journal* **97**:412–429
53. Anderson J.M., Andersson B. (1982) **The architecture of photosynthetic membranes: lateral and transverse organization** *Trends in Biochemical Sciences* **7**:288–292
54. Anderson J.M (1981) **Consequences of spatial separation of photosystem 1 and 2 in thylakoid membranes of higher plant chloroplasts** *FEBS Letters* **124**:1–10
55. Lamm L., et al. (2022) **MemBrain: A deep learning-aided pipeline for detection of membrane proteins in Cryo-electron tomograms** *Computer Methods and Programs in Biomedicine* **224**:106990
56. Goral T.K., et al. (2012) **Light-harvesting antenna composition controls the macrostructure and dynamics of thylakoid membranes in Arabidopsis** *The Plant Journal* **69**:289–301
57. Sheng X., et al. (2019) **Structural insight into light harvesting for photosystem II in green algae** *Nature Plants* **5**:1320–1330
58. Wei X., et al. (2016) **Structure of spinach photosystem II-LHCII supercomplex at 3.2 Å resolution** *Nature* **534**:69–74
59. Graça A.T., et al. (2021) **High-resolution model of Arabidopsis Photosystem II reveals the structural consequences of digitonin-extraction** *Scientific Reports* **11**:15534
60. Paolillo D.J. (1970) **The three-dimensional arrangement of intergranal lamellae in chloroplasts** *Journal of Cell Science* **6**:243–253
61. Bussi Y., et al. (2019) **Fundamental helical geometry consolidates the plant photosynthetic membrane** *Proceedings of the National Academy of Sciences* **116**:22366–22375
62. Boekema E.J., et al. (2000) **Arrangement of photosystem II supercomplexes in crystalline macrodomains within the thylakoid membrane of green plant chloroplasts** Edited by M F. Moody. *Journal of Molecular Biology* **301**:1123–1133
63. Simpson D.J (1983) **Freeze-fracture studies on barley plastid membranes. VI. Location of the P700-chlorophyll a-protein 1** *Eur J Cell Biol* **31**:305–14
64. Staehelin L.A (1976) **Reversible particle movements associated with unstacking and restacking of chloroplast membranes in vitro** *The Journal of cell biology* **71**:136–158

65. Tsvetkova N.M., et al. (1995) **Factors influencing PS II particle array formation in *Arabidopsis thaliana* chloroplasts and the relationship of such arrays to the thermostability of PS II** *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1228**:201–210
66. Ojakian G.K., Satir P. (1974) **Particle movements in chloroplast membranes: quantitative measurements of membrane fluidity by the freeze-fracture technique** *Proc Natl Acad Sci U S A* **71**:2052–6
67. Eckardt N.A., et al. (2024) **Lighting the way: Compelling open questions in photosynthesis research** *The Plant Cell* **36**:3914–3943
68. Charuvi D., et al. (2016) **Studying the Supramolecular Organization of Photosynthetic Membranes within Freeze-fractured Leaf Tissues by Cryo-scanning Electron Microscopy** *J Vis Exp* **112**
69. McDonnell A., Staehelin L.A. (1980) **Adhesion between liposomes mediated by the chlorophyll a/b light-harvesting complex isolated from chloroplast membranes** *J Cell Biol* **84**:40–56
70. Albanese P., et al. (2020) **How paired PSII-LHCII supercomplexes mediate the stacking of plant thylakoid membranes unveiled by structural mass-spectrometry** *Nat Commun* **11**:1361
71. Guardini Z., et al. (2022) **Thylakoid grana stacking revealed by multiplex genome editing of LHCII encoding genes** *bioRxiv*
72. Standfuss J., et al. (2005) **Mechanisms of photoprotection and nonphotochemical quenching in pea light-harvesting complex at 2.5 Å resolution** *The EMBO Journal* **24**:919–928
73. Mittag T., Pappu R.V. (2022) **A conceptual framework for understanding phase separation and addressing open questions and challenges** *Mol Cell* **82**:2201–2214
74. Snead W.T., Gladfelter A.S. (2019) **The Control Centers of Biomolecular Phase Separation: How Membrane Surfaces, PTMs, and Active Processes Regulate Condensation** *Mol Cell* **76**:295–305
75. Izawa S., Good N.E. (1966) **Effect of salts and electron transport on the conformation of isolated chloroplasts. I. Light-scattering and volume changes** *Plant Physiol* **41**:533–43
76. Kim E.H., et al. (2005) **Entropy-assisted stacking of thylakoid membranes** *Biochim Biophys Acta* **1708**:187–95
77. Schiøtz O.H., et al. (2024) **Serial Lift-Out: sampling the molecular anatomy of whole organisms** *Nat Methods* **21**:1684–1692
78. Nguyen H.T.D., et al. (2024) **Serialized on-grid lift-in sectioning for tomography (SOLIST) enables a biopsy at the nanoscale** *Nat Methods* **21**:1693–1701
79. Iwai M., Roth M.S., Niyogi K.K. (2018) **Subdiffraction-resolution live-cell imaging for visualizing thylakoid membranes** *Plant J* **96**:233–243
80. Pipitone R., et al. (2021) **A multifaceted analysis reveals two distinct phases of chloroplast biogenesis during de-etiolation in *Arabidopsis*** *eLife* **10**

81. McCafferty C.L., et al. (2024) **Integrating cellular electron microscopy with multimodal data to explore biology across space and time** *Cell* **187**:563–584
82. Mastronarde D.N. (2005) **Automated electron microscope tomography using robust prediction of specimen movements** *J Struct Biol* **152**:36–51
83. Hagen W.J.H., Wan W., Briggs J.A.G. (2017) **Implementation of a cryo-electron tomography tilt-scheme optimized for high resolution subtomogram averaging** *J Struct Biol* **197**:191–198
84. Khavnekar S., Erdmann P.S., Wan W. (2024) **TOMOMAN: a software package for large scale cryo-electron tomography data preprocessing, community data sharing, and collaborative computing** *bioRxiv*
85. Zheng S.Q., et al. (2017) **MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy** *Nat Methods* **14**:331–332
86. Kremer J.R., Mastronarde D.N., McIntosh J.R. (1996) **Computer visualization of three-dimensional image data using IMOD** *J Struct Biol* **116**:71–6
87. Mastronarde D.N., Held S.R. (2017) **Automated tilt series alignment and tomographic reconstruction in IMOD** *J Struct Biol* **197**:102–113
88. Martinez-Sanchez A., et al. (2014) **Robust membrane detection based on tensor voting for electron tomography** *Journal of Structural Biology* **186**:49–61
89. Sullivan C.B., Kaszynski A. (2019) **PyVista: 3D plotting and mesh analysis through a streamlined interface for the Visualization Toolkit (VTK)** *Journal of Open Source Software* **4**:1450
90. Gaifas L., et al. (2024) **Blik is an extensible 3D visualisation tool for the annotation and analysis of cryo-electron tomography data** *PLoS Biol* **22**:e3002447
91. Mitchell J.S.B., Mount D.M., Papadimitriou C.H. (1987) **The Discrete Geodesic Problem** *SIAM Journal on Computing* **16**:647–668

Author information

Wojciech Wietrzynski*

Biozentrum, University of Basel, Basel, Switzerland
ORCID iD: [0000-0001-8898-2392](https://orcid.org/0000-0001-8898-2392)

For correspondence: wojciech.wietrzynski@unibas.ch

*These authors contributed equally.

Lorenz Lamm*

Biozentrum, University of Basel, Basel, Switzerland, Helmholtz AI, Helmholtz Zentrum München, Neuherberg, Germany
ORCID iD: [0000-0003-0698-7769](https://orcid.org/0000-0003-0698-7769)

*These authors contributed equally.

William HJ Wood

Plants, Photosynthesis and Soil, School of Biosciences, University of Sheffield, Sheffield, United Kingdom

ORCID iD: [0000-0003-0683-8085](https://orcid.org/0000-0003-0683-8085)

Matina-Jasemi Loukeri

Department of Structural Cell Biology, Max Planck Institute of Biochemistry, Martinsried, Germany

Lorna Malone[†]

Plants, Photosynthesis and Soil, School of Biosciences, University of Sheffield, Sheffield, United Kingdom

ORCID iD: [0000-0002-7625-7545](https://orcid.org/0000-0002-7625-7545)

[†]Current address: Electron Bio-imaging Centre, Diamond Light Source, Didcot, United Kingdom

Tingying Peng

Helmholtz AI, Helmholtz Zentrum München, Neuherberg, Germany

ORCID iD: [0000-0002-7881-1749](https://orcid.org/0000-0002-7881-1749)

Matthew P Johnson

Plants, Photosynthesis and Soil, School of Biosciences, University of Sheffield, Sheffield, United Kingdom

ORCID iD: [0000-0002-1663-0205](https://orcid.org/0000-0002-1663-0205)

For correspondence: matt.johnson@sheffield.ac.uk

Benjamin D Engel

Biozentrum, University of Basel, Basel, Switzerland

ORCID iD: [0000-0002-0941-4387](https://orcid.org/0000-0002-0941-4387)

For correspondence: ben.engel@unibas.ch

Editors

Reviewing Editor

David Kramer

Michigan State University, East Lansing, United States of America

Senior Editor

Felix Campelo

Institute of Photonic Sciences, Barcelona, Spain

Reviewer #1 (Public review):

Summary:

In this study, authors utilized in situ cryo-electron tomography (cryo-ET) to uncover the native thylakoid architecture of spinach chloroplasts and mapped the molecular organization of these thylakoids with single-molecule resolution. The obtained images show the detailed ultrastructural features of grana membranes and highlight interactions between thylakoids and plastoglobules. Interestingly, despite the distinct three-dimensional architecture of vascular plant thylakoids, their molecular organization closely resembles that of green algae. The pronounced lateral segregation of PSII and PSI was observed at the interface between appressed and non-appressed thylakoid regions, without evidence of a specialized grana margin zone where these complexes might intermix. Furthermore, unlike isolated thylakoid membranes, photosystem II (PSII) did not form a semi-crystalline array and distributed uniformly within the membrane plane and across stacked grana membranes in intact chloroplasts. Based on the above observations, the authors propose a simplified two-domain model for the molecular organization of thylakoid membranes that can apply to both green algae and vascular plants. This study suggests that the general understanding of the functional separation of thylakoid membranes in vascular plants should be reconsidered.

Strengths:

By employing and refining AI-driven computational tools for the automated segmentation of membranes and identification of membrane proteins, this study successfully quantifies the spatial organization of photosynthetic complexes both within individual thylakoid membranes and across neighboring stacked membranes.

Weaknesses:

This study's weakness is that it requires the use of chloroplasts isolated from leaves and the need to freeze them on a grid for observation, so it is unclear to what extent the observations reflect physiological conditions. In particular, the mode of existence of the thylakoid membrane complexes seems to be strongly influenced by the physicochemical environment surrounding the membranes, as indicated by the different distribution of PSII between intact chloroplasts and those with ruptured envelope membranes.

<https://doi.org/10.7554/eLife.105496.1.sa1>

Reviewer #2 (Public review):

Summary:

For decades, the macromolecular organization of photosynthetic complexes within the thylakoids of higher plant chloroplasts has been a topic of significant debate. Using focused ion beam milling, cryo-electron tomography, and advanced AI-based image analysis, the authors compellingly demonstrate that the macromolecular organization in spinach thylakoids closely mirrors the patterns observed in their earlier research on *Chlamydomonas reinhardtii*. Their findings provide strong evidence challenging long-standing assumptions about the existence of a 'grana margin'-a region at the interface between grana and stroma lamellae domains that was thought to contain intermixed particles from both areas. Instead, the study establishes that this mixed zone is absent and reveals a distinct, well-defined boundary between the grana and stroma lamellae.

Strengths:

By situating high-resolution structural data within the broader cellular context, this work contributes valuable insights into the molecular mechanisms governing the spatial organization of photosynthetic complexes within thylakoid membranes.

