



Nonredundant Function of Soluble LT α_3 Produced by Innate Lymphoid Cells in Intestinal Homeostasis

Andrey A. Kruglov *et al.*

Science **342**, 1243 (2013);

DOI: 10.1126/science.1243364

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of December 9, 2013):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/342/6163/1243.full.html>

Supporting Online Material can be found at:

<http://www.sciencemag.org/content/suppl/2013/12/05/342.6163.1243.DC1.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/342/6163/1243.full.html#related>

This article **cites 37 articles**, 12 of which can be accessed free:

<http://www.sciencemag.org/content/342/6163/1243.full.html#ref-list-1>

Nonredundant Function of Soluble LT α_3 Produced by Innate Lymphoid Cells in Intestinal Homeostasis

Andrey A. Kruglov,^{1,2*} Sergei I. Grivennikov,³ Dmitry V. Kuprash,^{2,4} Caroline Winsauer,¹ Sandra Prepens,¹ Gitta Maria Selezniek,⁵ Gerard Eberl,⁶ Dan R. Littman,⁷ Mathias Heikenwalder,^{5,8} Alexei V. Tumanov,⁹ Sergei A. Nedospasov^{1,2,4*}

Immunoglobulin A (IgA) production at mucosal surfaces contributes to protection against pathogens and controls intestinal microbiota composition. However, mechanisms regulating IgA induction are not completely defined. We show that soluble lymphotoxin α (sLT α_3) produced by ROR γ ⁺ innate lymphoid cells (ILCs) controls T cell–dependent IgA induction in the lamina propria via regulation of T cell homing to the gut. By contrast, membrane-bound lymphotoxin β (LT $\alpha_1\beta_2$) produced by ROR γ ⁺ ILCs is critical for T cell–independent IgA induction in the lamina propria via control of dendritic cell functions. Ablation of LT α in ROR γ ⁺ cells abrogated IgA production in the gut and altered microbiota composition. Thus, soluble and membrane-bound lymphotoxins produced by ILCs distinctly organize adaptive immune responses in the gut and control commensal microbiota composition.

Production of immunoglobulin A (IgA) at mucosal surfaces contributes to host defense against intestinal pathogens and governs quantitative and qualitative control of commensal microbiota composition by the host (1, 2). IgA can be induced by distinct T cell–dependent or T cell–independent pathways. T cell–dependent regulation of IgA production takes place mainly in Peyer’s patches and requires the formation of germinal centers and the interaction of B cells with follicular helper T cells (3). T cell–independent mucosal IgA is produced both in isolated lymphoid follicles (ILFs) and in the lamina propria aided by exposure of B cells to various cytokines and growth factors, without the formation of germinal centers (4, 5).

Lymphotoxin α (LT α) and lymphotoxin β (LT β) are trimeric cytokines of the tumor necrosis factor (TNF) superfamily that are expressed in either soluble (sLT α_3) or membrane-bound (LT $\alpha_1\beta_2$) forms by T and B cells, as well as by retinoic acid–related orphan receptor positive (ROR γ ⁺) innate lymphoid cells (ILCs) (6). Soluble lymphotoxin is a TNF-like cytokine and its signaling is me-

diated via both TNFR1 and TNFR2, whereas membrane-bound lymphotoxin signals via LT β R (6). Ablation of the surface lymphotoxin–driven pathway via inactivation of the genes encoding LT α , LT β , or LT β R results in block of lymphoid organ development and in diminished IgA plasma cell numbers in mucosal tissues (7, 8), indicating that membrane-bound lymphotoxin is critical for intestinal IgA production. However, the possible contribution of soluble lymphotoxin to this process is not known.

ILCs were recently described as an important subset of innate immune cells that lack specific antigen receptors but are able to produce a range of effector cytokines (9). They are predominantly located in mucosal tissues and provide the first line of defense against various mucosal pathogens (9–14). In particular, ROR γ ⁺ ILCs, via LT production, induce the development of gut-associated lymphoid tissues such as lymph nodes, Peyer’s patches, and isolated lymphoid follicles (8, 15, 16) and are critical for protection against intestinal pathogens (10, 11, 17), for maintenance of the epithelial barrier, and for the prevention of systemic dissemination of commensal microbiota (16, 18). However, the molecular mechanisms that mediate host control of commensals by ROR γ ⁺ ILCs remain largely unknown.

We used mice lacking LT α or LT β production by ROR γ ⁺-expressing cells (LT α ^{ΔILC,T} and LT β ^{ΔILC,T} mice, respectively) (19, 20) (fig. S1). Because the transcription factor ROR γ is expressed in double positive (CD4⁺CD8⁺) thymocytes, these mice also exhibit LT gene deletion in all $\alpha\beta$ T cells (21) in addition to ROR γ ⁺ ILCs. Therefore, to define the role of LT expressed specifically by ROR γ ⁺ ILCs, we included mice with T cell ablation of LT α and LT β , using CD4-Cre transgenic mice (LT α ^{ΔT} and LT β ^{ΔT}, respectively) as controls in all our analyses (fig. S1). Mice lacking expression of either the LT α or LT β gene in T cells showed no

developmental defects in their secondary lymphoid tissues (20), whereas LT β ^{ΔILC,T} mice lacked Peyer’s patches, isolated lymphoid follicles, and all peripheral lymph nodes except mesenteric (fig. S1), recapitulating the anatomical phenotype of complete LT β ablation (22). LT α ^{ΔILC,T} mice lacked Peyer’s patches, isolated lymphoid follicles, and all lymph nodes (fig. S1). Together, these data demonstrated the role of LT produced by ROR γ ⁺ ILCs during embryogenesis for secondary lymphoid organ development (15).

Membrane-bound lymphotoxin produced by ROR γ ⁺ ILCs is implicated as one of the critical cytokines required for generation of mucosal IgA through the formation of ILFs (8). However, LT β ^{ΔILC,T} animals that lacked ILFs exhibited normal fecal IgA levels and only slightly diminished blood IgA levels relative to wild-type controls (Fig. 1, A and B). By contrast, concomitant inactivation of surface and soluble lymphotoxins via deletion of the LT α gene, in both ROR γ ⁺ ILCs and $\alpha\beta$ T cells (but not in $\alpha\beta$ T cells alone), led to a striking decrease in both blood and fecal IgA levels and was essential for the presence of IgA⁺ cells in the lamina propria (Fig. 1, A to C). By contrast, LT α expression by T cells was not required either for IgA production or for the recruitment of major immune cell subsets in the small intestine (Fig. 1, A to C, and fig. S2). To rule out a possible contribution from mesenteric lymph nodes that are present in LT β ^{ΔILC,T} animals (but not in LT α ^{ΔILC,T} mice), we established bone marrow transfers into lethally irradiated LT α -deficient recipients that lacked gut-associated lymphoid tissue. In contrast to wild-type, LT α ^{ΔT}, and LT β ^{ΔILC,T} bone marrow cells, transfer of LT α ^{ΔILC,T} bone marrow cells failed to induce IgA in recipient mice (Fig. 1D); this result implies a direct role of LT α produced by ROR γ ⁺ ILCs in this process, irrespective of the presence of mesenteric lymph nodes and consistent with previous findings in LT α -deficient mice (7, 23).

One of the known functions of IgA in mucosal tissues is to contain and control the composition of commensal microbiota (24). Deep sequencing analysis of ileal commensal microflora in wild-type and LT α ^{ΔILC,T} animals, and further real-time polymerase chain reaction (PCR) of selected intestinal commensals, revealed a marked expansion of segmented filamentous bacteria and a reduction in Bacteroidetes in mice lacking LT α expression by ROR γ ⁺ cells (Fig. 1, E and F), implicating LT expression by ROR γ ⁺ ILCs in the control of gut microbiota.

Although wild-type, LT α ^{ΔT}, LT β ^{ΔILC,T}, LT α ^{ΔILC,T}, and LT α ^{−/−} animals showed similar ILC numbers in the lamina propria (figs. S3 and S4), c-Kit and CCR6 expression by the small intestinal lymphoid tissue inducer-like (LTi) (CD45⁺Thy1.2⁺c-Kit^{high}IL-7R α ⁺CCR6⁺) cells was controlled by membrane-bound lymphotoxin produced by ROR γ ⁺ ILCs (fig. S3). However, comparable phenotypic changes were found in adult LTi cells from LT β ^{ΔILC,T}, LT α ^{ΔILC,T}, and LT α ^{−/−} mice (fig. S4); therefore, this cannot explain the lack

¹German Rheumatism Research Center (DRFZ), a Leibniz Institute, Berlin 10117, Germany. ²Belozersky Institute of Physico-Chemical Biology and Biological Faculty, Lomonosov Moscow State University, Moscow 119991, Russia. ³Cancer Prevention and Control Program, Fox Chase Cancer Center, Philadelphia, PA 19111, USA. ⁴Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow 119991, Russia. ⁵Department of Pathology, Institute of Clinical Pathology, University Hospital Zürich, CH 8091 Zürich, Switzerland. ⁶Institut Pasteur and Centre National de la Recherche Scientifique, Unité de Recherche Associée 1961, 75724 Paris, France. ⁷Howard Hughes Medical Institute and Molecular Pathogenesis Program, Kimmel Center for Biology and Medicine, Skirball Institute, New York University School of Medicine, New York, NY 10016, USA. ⁸Institute of Virology, Helmholtz Zentrum München, Technische Universität München, D-81675 München, Germany. ⁹Trudeau Institute, Saranac Lake, NY 12983, USA.

*Corresponding author. E-mail: andrey_krugloff@mail.ru (A.A.K.); sergei@nedos.net (S.A.N.)

of intestinal IgA observed in the two latter gene-deficient animals.

Signaling via both TNFR1 and TNFR2 contributed to intestinal IgA production, whereas complete genetic ablation of TNF did not affect fecal IgA levels (fig. S5), consistent with the role for soluble LT produced by ROR γ ⁺ ILCs. Furthermore, both TNFR1 and TNFR2 expressed by non-hematopoietic cells contributed to IgA induction, as revealed by reciprocal bone marrow transfer experiments (fig. S5). Therefore, soluble lymphotoxin acts via TNFR1 and TNFR2 expressed by lamina propria stromal cells to promote IgA production.

In the context of intestinal immunity, LT β R signaling is important for B cell homing to the gut (7). We found significantly reduced expression of chemokine and adhesion molecules such as CXCL13, VCAM-1, and CCL20 in the small intestine upon ablation of LT α , but not of LT β , from ROR γ ⁺ cells, whereas MAdCAM-1 and CCL21 expression remained unaffected (fig. S5).

Fig. 1. Soluble LT α_3 produced by ROR γ ⁺ ILCs regulates IgA production and microbiota composition in the gut. (A) Fecal IgA levels in naïve wild-type (WT), LT α ^{ΔT}, LT α ^{ΔILC,T}, LT β ^{ΔILC,T}, and LT α ^{-/-} animals. **(B)** Serum IgA levels in naïve WT, LT α ^{ΔT}, LT α ^{ΔILC,T}, LT β ^{ΔILC,T}, and LT α ^{-/-} animals. **(C)** Immunofluorescence analysis of IgA expression in the small intestine in naïve mice lacking LT α and LT β expression by ROR γ ⁺ cells. Scale bars, 80 μ m. **(D)** Fecal IgA levels in LT α ^{-/-} recipients, reconstituted with WT, LT α ^{ΔT}, LT α ^{ΔILC,T}, and LT β ^{ΔILC,T} bone marrow. Feces were collected 2 months after bone marrow transfer; IgA fecal levels were measured as described (30). **(E)** Deep sequencing analysis of microbiota composition from terminal ileum of LT α ^{ΔILC,T} mice and littermate WT controls. Representative microbiota composition in WT and LT α ^{ΔILC,T} ileum is presented ($n = 2$ mice per group). **(F to I)** Real-time PCR analysis of microbiota composition in terminal ileum of naïve WT, LT α ^{ΔT}, LT α ^{ΔILC,T}, and LT β ^{ΔILC,T} animals. Error bars, SEM; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ (Student's t test). All data, except deep sequencing analysis, are representative of two or more independent experiments with $n \geq 3$ mice.


Moreover, both soluble and membrane-bound lymphotoxins expressed by ROR γ ⁺ ILCs facilitated the homing of lamina propria IgM⁺ B cells (Fig. 2, A and B).

Because intestinal IgA plasma cells can develop from peritoneal IgM⁺ B cells recruited to the gut (25), we assessed the influence of LT α and LT β expression by ROR γ ⁺ ILCs in the peritoneal B cell compartment. Indeed, peritoneal cavity exudate cells from both LT α ^{ΔILC,T} and LT β ^{ΔILC,T} mice contained increased numbers of B1 and B2 B cells (fig. S6), whereas numbers of B cells in the spleen and bone marrow were normal (fig. S6). Additionally, B cells from LT α ^{ΔILC,T} mice were able to undergo class switching toward IgA in vitro, ruling out a B cell-intrinsic defect in class switch recombination (fig. S6).

Unexpectedly, when LT β ^{ΔILC,T} mice were crossed onto a T cell receptor (TCR) $\alpha\beta$ -deficient background, we found reduced IgA levels both in the feces and blood (Fig. 2, C and D), which

correlated with the absence of IgA⁺ plasma cells in the lamina propria (Fig. 2, E to G). T cell deficiency in LT β ^{ΔILC,T} mice did not further affect homing of B cells to the lamina propria (Fig. 2F and fig. S7). B cell proliferation, activation, and expression of activation-induced cytidine deaminase (AID) are all required for class switch recombination in B cells (26), and, indeed, no AID mRNA was detected in freshly isolated lamina propria lymphocytes from LT β ^{ΔILC,T} TCR $\alpha\beta$ ^{-/-} mice, whereas lamina propria lymphocytes from LT β ^{ΔILC,T} mice showed AID expression (fig. S7). Moreover, lamina propria B cells in LT β ^{ΔILC,T} mice were actively proliferating as revealed by KI67 staining (fig. S7). Taken together, these findings indicate that in the absence of surface lymphotoxin expression by ROR γ ⁺ ILCs, IgA class switching can occur in the lamina propria, and that $\alpha\beta$ T cells are crucial for this process.

LT expression by ROR γ ⁺ ILCs did not affect the development of various dendritic cell (DC) subsets



in mesenteric lymph nodes and in the small intestine (fig. S8), but $LT\alpha_1\beta_2$ expression by $ROR\gamma^+$ cells did control inducible nitric oxide synthase (iNOS) expression by mesenteric lymph node $CD11c^+$ DCs (Fig. 2H), which is known to be critical for IgA induction (27, 28). Moreover, $CD11c^+$ DCs isolated from mesenteric lymph nodes of $LT\beta^{\Delta ILCT}$ mice were less potent in inducing IgA in vitro (Fig. 2I). Together, these data indicated that $LT\alpha_1\beta_2$ may control T cell-independent IgA production via regulation of iNOS expression by DCs. Furthermore, our analysis revealed a reduction of CD40L mRNA levels in the small intestine of $LT\beta^{\Delta ILCT}$ mice after T cell ablation (Fig. 3A). When the T cell compartment in $LT\beta^{\Delta ILCT}$ $TCR\beta^{-/-}$ animals was reconstituted with wild-type or $CD40L^{-/-}$ $\alpha\beta$ T cells, we found that wild-type, but not $CD40L^{-/-}$, T cells could induce IgA production (Fig. 3, B and C). Interestingly, $LT\alpha$, but


not $LT\beta$, expressed by $ROR\gamma^+$ cells controlled T cell numbers in the lamina propria (Fig. 3, D and E). However, $LT\alpha$ produced by $ROR\gamma^+$ cells did not affect the numbers of gut-homing T cells on the periphery (fig. S8).

Consistently, we found significant reduction of CD40L mRNA expression in $LT\alpha^{\Delta ILCT}$ animals relative to littermate controls (Fig. 3F), which led us to hypothesize that soluble lymphotoxin may regulate IgA induction via control of T cell homing to the lamina propria. Indeed, forced activation of CD40 signaling by agonistic antibody in $LT\alpha^{\Delta ILCT}$ mice resulted in IgA induction (Fig. 3, G and H) without affecting T and B cell homing to the small intestine (fig. S9), further implying that T cells may contribute to the regulation of IgA switching in the lamina propria via the CD40-CD40L pathway. Notably, the induction of unspecific intestinal inflammation, such as dex-

tran sodium sulfate-induced colitis, failed to induce generation of IgA plasma cells (fig. S9). Collectively, these data indicated that s $LT\alpha_3$ derived from ILCs may control IgA induction via regulation of T cell homing to the lamina propria.

Induction of fully competent adaptive immune responses in the intestinal tract is a host defense mechanism directed against potential pathogens, which also allows control of commensal microbiota by the host. Here, we delineated distinct functions for membrane-bound and soluble lymphotoxins expressed by $ROR\gamma^+$ ILCs in the induction of IgA in the lamina propria (fig. S10). We found that production of membrane-bound lymphotoxin by ILCs regulates T cell-independent IgA induction via iNOS production by $CD11c^+$ DCs. We further demonstrated that ILC-derived soluble LT regulates the

Fig. 2. Membrane-bound $LT\alpha_1\beta_2$ produced by $ROR\gamma^+$ cells controls T cell-independent IgA induction in the lamina propria in the absence of organized gut-associated lymphoid tissue. (A) B cell frequencies in the lamina propria of WT, $LT\beta^{\Delta ILCT}$, and $LT\alpha^{\Delta ILCT}$ mice. (B) Numbers of $CD45^+CD19^+IgM^+$ cells in the lamina propria of WT, $LT\beta^{\Delta ILCT}$, and $LT\alpha^{\Delta ILCT}$ mice. (C) Fecal IgA levels in WT, $LT\beta^{\Delta ILCT}$, $TCR\alpha\beta^{-/-}$, and $LT\beta^{\Delta ILCT}$ $TCR\alpha\beta^{-/-}$ mice. (D) Serum IgA levels in $TCR\alpha\beta^{-/-}$ and $LT\beta^{\Delta ILCT}$ $TCR\alpha\beta^{-/-}$ mice. (E) Immunofluorescence analysis of IgA expression in the small intestine in WT, $LT\beta^{\Delta ILCT}$, $TCR\alpha\beta^{-/-}$, and $LT\beta^{\Delta ILCT}$ $TCR\alpha\beta^{-/-}$ mice. Scale bar, 80 μ m. (F and G) Representative fluorescence-activated cell sorter (FACS) dot plots (F) and frequencies (G) of $CD45^+IgA^+$ cells in lamina propria of $LT\beta^{\Delta ILCT}$ and $LT\beta^{\Delta ILCT}$ $TCR\alpha\beta^{-/-}$ mice. (H) iNOS mRNA expression levels in $CD11c^+$ cells sorted from mesenteric lymph nodes. (I) IgA levels in 5-day cultures of WT splenic IgM^+ B cells together with $CD11c^+$ DCs isolated from mesenteric lymph nodes of WT or $LT\beta^{\Delta ILCT}$ mice. All data are representative of two or more independent experiments with $n \geq 3$ mice. Error bars, SEM; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ (Student's t test).






Fig. 3. Regulation of T cell-dependent IgA production by soluble LT α_3 produced by ROR γ^t ILCs. (A) CD40L mRNA levels in jejunum of naive WT, LT $\beta^{\Delta ILCT}$, and LT $\beta^{\Delta ILCT}$ TCR $\beta^{-/-}$ mice. (B) Immunofluorescence analysis of IgA expression in the small intestine of naive TCR $\beta^{-/-}$, LT $\beta^{\Delta ILCT}$ TCR $\beta^{-/-}$ mice, as well as in LT $\beta^{\Delta ILCT}$ TCR $\beta^{-/-}$ mice, reconstituted with WT or CD40L $^{-/-}$ $\alpha\beta$ T cells. Scale bar, 100 μ m. (C) Frequency of CD45 $^+$ IgA $^+$ cells in the lamina propria of LT $\beta^{\Delta ILCT}$ TCR $\beta^{-/-}$ mice 2 weeks after adoptive transfer of WT or CD40L $^{-/-}$ $\alpha\beta$ T cells. (D and E) Numbers of CD45 $^+$ CD3 $^+$ cells (D) and CD45 $^+$ CD3 $^+$ CD4 $^+$ cells (E) in the lamina propria of WT mice and mice with LT α or LT β ablation in ROR γ^t cells. (F) CD40L mRNA levels in jejunum of WT and LT $\alpha^{\Delta ILCT}$ animals. (G) In vivo induction of IgA in LT $\alpha^{\Delta ILCT}$ mice by agonistic antibody to CD40. (H) CD45 $^+$ IgA $^+$ cells in the lamina propria of LT $\alpha^{\Delta ILCT}$ mice after anti-CD40 treatment. All data are representative of two or more independent experiments with $n \geq 3$ mice. Error bars, SEM; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ (Student's t test).

T cell-dependent pathway of IgA production via control of T cell homing to the gut. Through these processes, lymphotoxins produced by ROR γ^t ILCs control microbiota composition in the host and may influence various pathophysiological processes. Our findings highlight a rare nonredundant

function of soluble lymphotoxin and may be relevant for anti-TNF therapy using etanercept, as this drug can block not only TNF but also soluble lymphotoxin (29), and thus the effects of such treatment may affect IgA levels and gut microbiota in patients.

References and Notes

1. K. Honda, D. R. Littman, *Annu. Rev. Immunol.* **30**, 759–795 (2012).
2. A. Cerutti, K. Chen, A. Chorny, *Annu. Rev. Immunol.* **29**, 273–293 (2011).
3. S. Fagarasan, S. Kawamoto, O. Kanagawa, K. Suzuki, *Annu. Rev. Immunol.* **28**, 243–273 (2010).
4. A. J. Macpherson *et al.*, *Science* **288**, 2222–2226 (2000).
5. J. R. Mora *et al.*, *Science* **314**, 1157–1160 (2006).
6. C. F. Ware, *Annu. Rev. Immunol.* **23**, 787–819 (2005).
7. H. S. Kang *et al.*, *Nat. Immunol.* **3**, 576–582 (2002).
8. M. Tsuji *et al.*, *Immunity* **29**, 261–271 (2008).
9. H. Spits, T. Cupedo, *Annu. Rev. Immunol.* **30**, 647–675 (2012).
10. G. F. Sonnenberg, L. A. Monticelli, M. M. Elloso, L. A. Fouser, D. Artis, *Immunity* **34**, 122–134 (2011).
11. Y. Wang *et al.*, *Immunity* **32**, 403–413 (2010).
12. L. A. Zenewicz *et al.*, *Immunity* **29**, 947–957 (2008).
13. N. Satoh-Takayama *et al.*, *Immunity* **29**, 958–970 (2008).
14. M. R. Hepworth *et al.*, *Nature* **498**, 113–117 (2013).
15. G. Eberl *et al.*, *Nat. Immunol.* **5**, 64–73 (2004).
16. D. Bouskra *et al.*, *Nature* **456**, 507–510 (2008).
17. A. V. Tumanov *et al.*, *Cell Host Microbe* **10**, 44–53 (2011).
18. G. F. Sonnenberg *et al.*, *Science* **336**, 1321–1325 (2012).
19. D. J. Liepinsh *et al.*, *Mol. Cell. Biol.* **26**, 4214–4225 (2006).
20. A. Tumanov *et al.*, *Immunity* **17**, 239–250 (2002).
21. G. Eberl, D. R. Littman, *Science* **305**, 248–251 (2004).
22. M. B. Alimzhanov *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **94**, 9302–9307 (1997).
23. R. D. Newberry, J. S. McDonough, K. G. McDonald, R. G. Lorenz, *J. Immunol.* **168**, 4988–4997 (2002).
24. K. Suzuki *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **101**, 1981–1986 (2004).
25. F. G. Kroese *et al.*, *Int. Immunol.* **1**, 75–84 (1989).
26. M. Muramatsu *et al.*, *Cell* **102**, 553–563 (2000).
27. H. Tezuka *et al.*, *Nature* **448**, 929–933 (2007).
28. J. H. Fritz *et al.*, *Nature* **481**, 199–203 (2012).
29. D. Tracey, L. Klareskog, E. H. Sasso, J. G. Salfeld, P. P. Tak, *Pharmacol. Ther.* **117**, 244–279 (2008).
30. See supplementary materials on Science Online.

Acknowledgments: We thank S. Fillatreau, J. Gommerman, A. Rudensky, and F. Melchers for useful discussions and critiques; K. Holden-Dye for editing of the manuscript; L. Drutskaya, K. Horn, M. Drutskaya, and D. Schlieznz for expert technical support throughout the project; H. Schäfer, S. Gruczek, and M. Ohde for excellent animal husbandry; the members of the German Rheumatism Research Center Flow Cytometry Core Facility (T. Kaiser, J. Kirsch) for help with FACS analysis and cell sorting; H. Hecker-Kia, H. Schliemann, and T. Geske for antibody preparation; and J. Hirscher for help with figure preparation. Supported by Deutsche Forschungsgemeinschaft grants (SFB633 A13 and NE1466/2-1) and the MCB Program of the Russian Academy of Sciences (S.A.N. and D.V.K.), and the Crohn's and Colitis Foundation (A.V.T.). Bacterial 16S rRNA pyrosequencing reads have been deposited in the NCBI Short Read Archive database with accession numbers SRX362155, SRX362156, SRX362157, and SRX362158. Author contributions: A.A.K. and S.A.N. designed research; A.A.K., D.V.K., S.I.G., S.P., C.W., G.M.S., M.H., and A.V.T. performed experiments; D.R.L. and G.E. provided ROR γ^t -Cre animals and discussed the data; and A.A.K. and S.A.N. wrote the manuscript.

Supplementary Materials

www.sciencemag.org/content/342/6163/1243/suppl/DC1
Materials and Methods
Figs. S1 to S10
Table S1
References (31–38)

17 July 2013; accepted 24 October 2013
10.1126/science.1243364