



OPEN ACCESS

APPROVED BY
Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*CORRESPONDENCE
Martin Hrabe de Angelis
✉ martin.hrabedeangelis@helmholtz-
munich.de

[†]These authors have contributed equally
to this work

RECEIVED 21 January 2026
REVISED 26 January 2026
ACCEPTED 04 February 2026
PUBLISHED 23 February 2026

CITATION

Oestereicher MA, Ward CS, Schneltzer E,
Marschall S, Fuchs H, Gailus-Durner V,
About GB, Selloum M, Meziane H,
Stewart M, Teboul L, Norris C, Pimm D,
Kan M, Gómez FL, Wilson R, Monroy M,
Pasha S, Zabrodzka E, Prochazka J,
Reguera DP, Nichtova Z, Herault Y,
Wells S, Parkinson H, Heaney JD,
Sedlacek R, Gao X, de Angelis MH and
Spielmann N (2026) Correction:
Establishing standardized transthoracic
echocardiography reference ranges for
mouse models: insights into the impact
of anesthesia, sex, and age.
Front. Cardiovasc. Med. 13:1792916.
doi: 10.3389/fcvm.2026.1792916

COPYRIGHT

© 2026 Oestereicher, Ward, Schneltzer,
Marschall, Fuchs, Gailus-Durner, About,
Selloum, Meziane, Stewart, Teboul,
Norris, Pimm, Kan, Gómez, Wilson,
Monroy, Pasha, Zabrodzka, Prochazka,
Reguera, Nichtova, Herault, Wells,
Parkinson, Heaney, Sedlacek, Gao, de
Angelis and Spielmann. This is an open-
access article distributed under the
terms of the [Creative Commons
Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other
forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which does
not comply with these terms.

Correction: Establishing standardized transthoracic echocardiography reference ranges for mouse models: insights into the impact of anesthesia, sex, and age

Manuela A. Oestereicher^{1†}, Christopher S. Ward^{2†},
Elida Schneltzer¹, Susan Marschall¹, Helmut Fuchs¹,
Valerie Gailus-Durner¹, Ghina Bou About³, Mohammed Selloum³,
Hamid Meziane³, Michelle Stewart⁴, Lydia Teboul⁴, Clare Norris⁴,
Dale Pimm⁴, Marina Kan⁵, Federico López Gómez⁵,
Robert Wilson⁵, Mayra Monroy⁶, Sheraz Pasha⁶, Eva Zabrodzka⁷,
Jan Prochazka⁷, David Pajuelo Reguera⁷, Zuzana Nichtova⁷,
Yann Herault³, Sara Wells⁴, Helen Parkinson⁵, Jason D. Heaney⁸,
Radislav Sedlacek⁷, Xiang Gao⁹, Martin Hrabe de Angelis^{1,10,11*} and
Nadine Spielmann¹

¹Institute of Experimental Genetics and German Mouse Clinic, Helmholtz Center Munich, German
Research Center for Environmental Health, Neuherberg, Germany, ²Department of Integrative
Physiology, Baylor College of Medicine, Houston, TX, United States, ³Université de Strasbourg, CNRS,
INSERM, Institute de la Clinique de la Souris, CELPHEDIA, PHENOMIN, Illkirch, France, ⁴The Mary Lyon
Centre and MRC Harwell, Harwell Campus, Oxfordshire, United Kingdom, ⁵European Molecular Biology
Laboratory-European Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton,
Cambridgeshire, United Kingdom, ⁶Advanced Technology Cores, Baylor College of Medicine, Houston,
TX, United States, ⁷Czech Centre for Phenogenomics, Institute of Molecular Genetics of the Czech
Academy of Sciences, Prague, Czechia, ⁸Department of Molecular and Human Genetics, Baylor
College of Medicine, Houston, TX, United States, ⁹SKL of Pharmaceutical Biotechnology and Model
Animal Research Center, Collaborative Innovation Center for Genetics and Development, Nanjing
Biomedical Research Institute, Nanjing University, Nanjing, China, ¹⁰Chair of Experimental Genetics,
School of Life Science Weihenstephan, Technische Universität München, Freising, Germany, ¹¹German
Center for Diabetes Research (DZD), Neuherberg, Germany

KEYWORDS

anesthesia effects on cardiac function, c57BL/6N mouse model, cardiovascular
phenotyping, mouse cardiac reference ranges, transthoracic echocardiography

A Correction on

[Establishing standardized transthoracic echocardiography reference ranges
for mouse models: insights into the impact of anesthesia, sex, and age](#)

By Oestereicher MA, Ward CS, Schneltzer E, Marschall S, Fuchs H, Gailus-Durner V, About GB,
Selloum M, Meziane H, Stewart M, Teboul L, Norris C, Pimm D, Kan M, Gómez FL, Wilson R,
Monroy M, Pasha S, Zabrodzka E, Prochazka J, Reguera DP, Nichtova Z, Herault Y, Wells S,
Parkinson H, Heaney JD, Sedlacek R, Gao X, de Angelis MH and Spielmann N (2025). *Front.
Cardiovasc. Med.* 12:1695034. doi: 10.3389/fcvm.2025.1695034.

Affiliations 2–8 were in the wrong order. Affiliations 2–8 have been reordered.
The original version of this article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.