

Table S11: Rate of new atopic dermatitis cases with number of children with completed assessment and incidence rate (IR) in the first year of life for each outcome by intervention group in the subgroup of infants with parental AD

AD outcome and assessment time points (in months)	Subgroup of infants with parental AD	
	WGF % (n/N) / IR (n/person-years)	CF % (n/N) / IR (n/person-years)
AD_{Primary}		
4	2.6% (2/77)	5.3% (5/94)
6	4.1% (3/74)	7.1% (6/84)
12	5.8% (4/69)	14.3% (11/77)
Incidence rate	13.2 (9/68.0)	27.8 (22/79.2)
AD_{UKWP}		
4	2.6% (2/78)	5.2% (5/97)
6	3.9% (3/77)	6.7% (6/89)
12	5.6% (4/72)	15% (12/80)
Incidence rate	13.1 (9/69.0)	28.3 (23/81.4)
AD_{Doctor}		
4	3.8% (3/78)	8.2% (8/97)
6	2.6% (2/76)	7% (6/86)
12	5.6% (4/72)	18.2% (14/77)
Incidence rate	13.2 (9/68.2)	36.9 (28/76.0)
AD_{Any}		
4	3.8% (3/78)	10.3% (10/97)
6	5.3% (4/76)	10.7% (9/84)
12	7.1% (5/70)	20.8% (15/72)
Incidence rate	17.9 (12/67.1)	46.0 (34/73.9)

AD: Atopic dermatitis; WGF: Whole goat milk formula; CF: Cow milk formula; UKWP: United Kingdom Working Party; (n): new AD cases per time point; (N): observed number of infants per time point; IR: Incidence rate per 100 person-years