

CORRECTION

Open Access



Correction: Circulating nitric oxide pathway metabolites in heart failure with preserved ejection fraction: a sex-stratified cross-sectional analysis

Sophia Marie-Theres Dinges^{1,2,3*}, Edzard Schwedhelm^{4,5}, Flavia Baldassarri^{1,2}, Bernhard Haller⁶, Andreas B. Gevaert^{7,8}, Rainer Böger^{4,5}, Ephraim B. Winzer⁹, Frank Edelmann^{10,11,12}, Ulrik Wisløff¹³, Volker Adams⁹, Burkert Pieske¹⁴, Emeline M. Van Craenenbroeck^{7,8}, Martin Halle^{1,2}, Susanna M. Hofmann^{3,15,16†} and Stephan Mueller-Stegner^{1,2†}

Correction: *Biology of Sex Differences* (2026) 17:120
<https://doi.org/10.1186/s13293-026-00940-7>

Following publication of the original article [1], the authors reported a typesetting error, whereby the footnote for Table 3 was erroneously omitted.

The footnote for Table 3 is given below and the original article [1] has been corrected. The publishers apologise for this error.

Results show the regression coefficient β for sex and the corresponding 95% confidence interval.

- β coefficient: lower values in women.

+ β coefficient: higher values in women.

Model 1: unadjusted.

Model 2: Model 1 + age.

Model 3: Model 2 + %-predicted VO₂peak.

Model 4: Model 3 + BMI, HDL cholesterol.

[†]Susanna M. Hofmann and Stephan Mueller-Stegner contributed equally to this work.

The original article can be found online at <https://doi.org/10.1186/s13293-026-00940-7>.

*Correspondence:

Sophia Marie-Theres Dinges
Sophia.Dinges@mri.tum.de

¹TUM School of Medicine and Health, Department for Preventive Sports Medicine and Sports Cardiology, Technical University of Munich, TUM University Hospital, Munich, Germany

²DZHK (German Centre for Cardiovascular Research), Partner Site Munich Heart Alliance, Munich, Germany

³Research Group "Women and Diabetes", Institute for Diabetes and Regeneration, Helmholtz Center Munich - German Research Center for Environmental Health, Neuherberg, Germany

⁴Institute of Clinical Pharmacology and Toxicology, University Medical Center Hamburg- Eppendorf, Hamburg, Germany

⁵DZHK (German Centre for Cardiovascular Research), Partner Site North, Hamburg, Germany

⁶TUM School of Medicine and Health, Institute of AI and Informatics in Medicine, Technical University of Munich, Munich, Germany

⁷Department of Cardiology, Antwerp University Hospital, Edegem, Belgium

⁸Research Group Cardiovascular Diseases, GENCOR Department, University of Antwerp, Antwerp, Belgium

⁹Department of Internal Medicine/Cardiology, Heart Center, University Clinic, Technische Universität Dresden, Dresden, Germany

¹⁰Department of Cardiology, Angiology and Intensive Care Medicine, Deutsches Herzzentrum der Charité (DHZC) - Medical Heart Centre of Charité and German Heart Institute Berlin, Berlin, Germany

¹¹Charité - Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt Universität zu Berlin, Berlin, Germany

¹²DZHK (German Centre for Cardiovascular Research), Partner Site Berlin, Berlin, Germany

¹³Department of Circulation and Medical Imaging, Faculty of Medicine and Health Sciences, Division of Cardiology, Department of Internal Medicine, Norwegian University of Science and Technology, University Medicine Rostock, Trondheim, Norway

¹⁴Division of Cardiology, Department of Internal Medicine, University Medicine Rostock, Rostock, Germany

¹⁵Department of Medicine IV, University Hospital, LMU Munich, Munich, Germany

¹⁶German Center for Diabetes Research (DZD), Neuherberg, Germany



Model 5: Model 4 + comorbidities (diabetes, hypertension, coronary artery disease, atrial fibrillation, eGFR).

Model 6: Model 5 + ACE inhibitors, angiotensin receptor blocker, diuretics, and statins.

Published online: 06 July 2026

sex-stratified cross-sectional analysis. *Biol Sex Differ.* 2026;17:120. <https://doi.org/10.1186/s13293-026-00940-7>.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Reference

1. Dinges SMT, Schwedhelm E, Baldassarri F, et al. Circulating nitric oxide pathway metabolites in heart failure with preserved ejection fraction: a