

SOLUTION STRUCTURE AND MOLECULAR INTERACTIONS OF THE LAMIN B RECEPTOR TUDOR DOMAIN

Stamatis Liokatis¹, Christian Edlich², Katerina Soupsana^{3,4}, Ioannis Giannios^{3,4}, Parthena Panagiotidou¹, Konstantinos Tripsianes^{5,6}, Michael Sattler^{2,5,6}, Spyros D. Georgatos^{3,4} and Anastasia S. Politou^{1,4,*}

From the Laboratories of Biological Chemistry¹ and Biology³, School of Medicine, University of Ioannina, GR-45110 Ioannina, Greece,
Structural and Computational Biology Unit², European Molecular Biology Laboratory, D-69012 Heidelberg, Germany,
Biomedical Research Institute⁴, Foundation for Research and Technology (BRI-FORTH), GR-45110 Ioannina, Greece,
Institute of Structural Biology⁵, Helmholtz Zentrum München, 85764 Neuherberg, Germany, and
Munich Center for Integrated Protein Science (CiPSM)⁶, Department Chemie, Technische Universität München, 85747 Garching, Germany
Running head: Structure/interactions of the LBR Tudor domain

Address correspondence to: Anastasia S. Politou, Laboratory of Biological Chemistry, University of Ioannina, School of Medicine, GR-45110 Ioannina, Greece. Fax: 30 26510 07868; E-mail: apolitou@cc.uoi.gr

Capsule

Background: Lamin B Receptor (LBR) is an integral nuclear envelope protein and contains a Tudor domain.

Results: The NMR structure of LBR-Tudor was determined and its interactions with nuclear proteins, histones and nucleosomes were explored.

Conclusion: LBR-Tudor is not involved in recognition of methylated histones and binds free H3.

Significance: Tudor domains may act as histone chaperone-like platforms.

LBR is a polytopic protein of the nuclear envelope thought to connect the inner nuclear membrane with the underlying nuclear lamina and peripheral heterochromatin. To better understand the function of this protein, we have examined in detail its nucleoplasmic region, which is predicted to harbor a Tudor domain (LBR-TD). Structural analysis by multi-dimensional NMR spectroscopy establishes that LBR-TD indeed adopts a classical β -barrel Tudor fold in solution, which however features an incomplete aromatic cage. Removal of LBR-TD renders LBR more mobile

at the plane of the nuclear envelope, but the isolated module does not bind to nuclear lamins, heterochromatin proteins (MeCP2) and nucleosomes, nor does it associate with methylated Arg/Lys residues through its aromatic cage. Instead, LBR-TD exhibits tight and stoichiometric binding to the “histone fold” region of unassembled, free histone H3, suggesting an interesting role in histone assembly. Consistent with such a role, robust binding to native nucleosomes is observed when LBR-TD is extended towards its carboxy-terminus, to include an area rich in

Ser-Arg residues (RS). The RS region, alone or in combination with LBR-TD, binds both unassembled and assembled H3/H4 histones, suggesting that the TD/RS interface may operate as a “histone chaperone-like platform”.

Tudor domains are 50-70 amino acid modules, named after the synonymous *Drosophila* protein, which harbors 11 such copies in its molecule (1). Along with the chromodomain, PWWP, MBT and *Agenet*, these modules comprise a structural superfamily, the so-called ‘Royal family’. The members of the Royal family occur in a variety of chromatin-associated proteins and are thought to originate from a common ancestor (2). Originally, Tudor domains were thought to be RNA-binding motifs, because they were first identified in RNA-binding proteins or ribonucleoprotein particles (3). However, subsequent structural and biochemical studies involving the survival motor neuron (SMN) protein suggested that Tudor domains may associate with symmetrically dimethylated Arg residues in spliceosomal Sm proteins (4-7). This hypothesis has received further support from more recent studies with a variety of proteins (8-13) and it is now clear that Tudor domains can bind either methylated Lys residues in H3 and H4 histone tails (14-18), or methylated arginines usually flanked by glycine residues (11-13, 19). These interactions involve the methylated side chains and a cluster of aromatic residues that constitute the so-called aromatic cage, present in many Tudor and chromodomains (20-22).

A putative Tudor domain has been recently identified by inspection of the amino acid sequence of the Lamin B Receptor (LBR). LBR is a ubiquitous integral protein of the nuclear envelope (NE) and was initially characterized by virtue of its ability to associate with nuclear lamin B (23). It is now thought to participate in a variety of nuclear functions, including tethering of the nuclear lamina to the inner nuclear membrane and “transient trapping” of nuclear components that are involved in chromatin remodeling and transcriptional inactivation (24-30).

The putative Tudor domain of LBR (hereon referred to as LBR-TD) is accommodated within the amino-terminal part of the protein (LBR-NT), which faces the nucleoplasm and has been shown to mediate binding to the nuclear lamina and

peripheral heterochromatin. Sequence analyses suggest that LBR-NT contains three distinct regions (see scheme in Fig. 1A): (a) the LBR-TD, which spans the first 60 residues; (b) a highly charged 40-residue hinge region that is rich in Arg-Ser (RS) dipeptide motifs; and (c) a 110-amino acid segment (SGD) with no apparent sequence kinship to other proteins. The middle segment has features of a “natively disordered” protein and exhibits multiple SRPK1 phosphorylation sites (29,31,32).

Biochemical studies have implicated LBR-NT in lamin B interactions (33). It is also likely that this part of the protein participates in other interactions, such as binding to histone H3/H4 oligomers (30), methyl-CpG binding protein MeCP2 (34) and Heterochromatin Protein 1 (HP1) (32). LBR-NT has also been implicated in binding linker DNA (35) and in LBR-LBR interactions (28), but the significance of these findings and the involvement of LBR-TD have not been precisely determined.

In an attempt to elucidate the interactions of LBR at the molecular level and to further understand its *in situ* organization at the nuclear envelope, we have begun to dissect the LBR-NT into structurally/functionally relevant domains, starting from LBR-TD. Here, we report the solution structure of chicken LBR-TD and its interactions with other proteins and cellular components. Our biochemical data suggest that instead of binding to methylated arginine or lysine residues the LBR Tudor fold may have a histone chaperone-like activity, thus extending the range of functional roles of the Tudor family.

EXPERIMENTAL PROCEDURES

Expression and purification of recombinant proteins- Chicken LBR cDNAs encoding the LBR sub-domains Tudor (residues 1-62), RS (residues 63-100), SGD (residues 101-208), Tudor-RS (residues 1-100) and RSSGD (residues 63-208) were inserted into the NcoI/NotI sites of a modified pET24d (Novagen) expression vector encoding an amino terminal His₆-GST tag, followed by a TEV protease cleavage site, a generous gift from G. Stier (EMBL). The correct cDNA sequences of the expression clones were confirmed by DNA sequencing. Fusion proteins were expressed in BL21 (DE3) cells according to

standard procedures (36) and purified from bacterial lysates using Ni-NTA affinity chromatography. A uniformly ^{15}N , ^{13}C -labeled polypeptide corresponding to LBR-TD was also prepared by growing the *E.coli* strain BL21(DE3)plysS overexpressing LBR-TD in a minimal medium containing $^{15}\text{NH}_4\text{Cl}$ and ^{13}C -glucose. The N-terminal tag was removed after TEV digestion and the pure Tudor domain was collected after passing through a second Ni-NTA column. NMR samples were dialyzed into 20 mM sodium phosphate buffer (pH 6.9), 100 mM NaCl and concentrated using an Amicon ultrafiltration device to a final concentration of 0.8 mM. The recombinant Tudor protein used for the NMR experiments in addition to residues 1-62 of chicken LBR carries four N-terminal residues (GAMG) from the TEV cleavage site.

Native-calf thymus histones were purchased from Roche. A pET3a expression vector carrying the cDNAs encoding H3 and histone H3 core region (aa 27-135) from *Xenopus laevis* was a kind gift from K. Luger, University of Colorado. Recombinant H3 and tail-less H3 were expressed in *E.coli* BL21 (DE3) cells and purified under denaturing conditions using SP-sepharose chromatography, as previously described (37). After removal of urea with extensive dialysis, the samples were lyophilized, dissolved in water and their concentration was adjusted to 1mg/ml.

The yeast H3 tail region (1-46) was expressed in *E.coli* BL21 (DE3) cells as a fusion protein with a N-terminal GST tag from a pGEX2T expression vector kindly provided by M. Grunstein, University of California, and was purified as previously described (38).

The recombinant human N-terminal region of LBR (1-201) was expressed from a pET15b expression vector encoding an amino terminal His₁₀ tag. The fusion protein was expressed in BL21 (DE3) cells according to standard procedures (36) and purified from bacterial lysates under denaturing conditions using Ni-NTA affinity chromatography followed by protein refolding on the column (39).

Human MeCP2 (1-486) was expressed as a fusion protein with an amino terminal 6xHis tag from a pET30a expression vector, which was kindly provided by G. Badaracco, University of Insubria (34). The fusion protein was expressed in BL21 (DE3) cells according to standard

procedures (36) and was purified under native conditions by using Ni-NTA agarose beads.

Preparation of nuclear envelope extracts- Turkey erythrocytes were obtained from whole blood and their nuclei were isolated with standard methods (28,30,40). Isolated nuclei were digested with MNase (100units/ml digestion buffer) for 10min at 37°C (in 20mM HEPES-KOH buffer, pH 7.4, 5mM MgCl₂, 0.025% Triton, 1mM CaCl₂, 1mM DTT, 1mM PMSF, protease inhibitors – leupeptin, pepstatin, antipain, aprotinin at 2µg/ml). The reaction was stopped with 2mM EDTA and, after centrifugation at 10,000g, the resulting nuclear envelopes were washed with digestion buffer containing 2mM EDTA. Nuclear extract was prepared with resuspension of the nuclear envelope pellet in 300mM NaCl, 20mM Tris-HCl pH 7.5, 250mM sucrose, 2mM MgCl₂, 1mM EGTA, 1mM DTT, 1mM PMSF and protease inhibitors, followed by sonication. After ultracentrifugation for 30min at 4°C and 300,000g, the soluble extract was collected and used in pull-down assays.

Isolation of native lamin B- Nuclei were isolated from rat liver according to standard methods (41). Isolated nuclei (5ml) were resuspended in 0.1mM MgCl₂, 1mM DTT, 0.5mM PMSF (10ml). 15ml of 10% sucrose, 20mM Tris-HCl pH 8.5, 0.1mM MgCl₂, 1mM DTT, 0.5mM PMSF were added to the suspension. The preparation was digested with 500µl DNaseI (2mg/ml) for 15min at ambient temperature under rotation and the suspension centrifuged at 10,000g and 4°C for 15min. The pellet was resuspended in 10% sucrose, 20mM Tris-HCl pH 7.5, 1mM MgCl₂, 0.5mM PMSF (10ml) and 15ml of 30% sucrose, 20mM Tris-HCl pH 7.5, 0.1mM MgCl₂, 1mM DTT 0.5mM PMSF were added. Another round of DNaseI digestion was performed as above. The pellet was resuspended in 1M KCl, 50mM Tris-HCl pH 7.5, 1mM EDTA, 1mM DTT, 0.5mM PMSF (10ml) and was centrifuged as above. The final pellet was washed with ice cold ddH₂O and was centrifuged at 10,000g and 4°C for 45min. The nuclear envelope pellet was extracted with 8M urea, 10mM Tris-HCl pH 7.6, 4mM EDTA, 1mM DTT, 0.5mM PMSF followed by sonication. The soluble material was collected after 35min of ultracentrifugation at 300,000g and 18°C. Soluble lamin B was isolated from the extract by using DE53 ion exchange

chromatography. All buffers contained protease inhibitors – leupeptin, pepstatin, antipain, aprotinin – to a final concentration of 2µg/ml. Urea was removed from the desired elutions with extensive dialysis against 25mM Tris-HCl pH 8.5, 150mM NaCl, 1mM EGTA, 0.1mM DTT, 1mM PMSF. Antibodies against lamin B were prepared as previously described (42).

Isolation of native H3-H4 tetramers and interaction assays using sucrose gradients- Native H3-H4 tetramers were isolated using hydroxylapatite chromatography (43). Briefly, nuclear extract from MNase digested turkey erythrocyte nuclei was dialyzed against 10mM phosphate buffer pH6.8, 1mM DTT and subsequently passed through an equilibrated hydroxylapatite column (Biorad). Different histone pairs were eluted from the column using increasing ionic strength buffers (the H3-H4 tetramer is eluted with 2M NaCl).

For the sucrose gradient runs, GST-TD or Tudor protein samples were mixed with equimolar amounts of H3-H4 tetramer. The mixtures were dialyzed against 20mM Tris pH7.5, 300mM NaCl, 5% sucrose, 2mM MgCl₂, 0.1mM EDTA, 1mM DTT, 1mM PMSF and subsequently concentrated to a total protein concentration of 1mg/ml. 300µl samples were coated on the top of a 5-20% sucrose gradient made in the same buffer in a total volume of 11ml. Samples were spun at 40,000rpm in a SW40 Beckman rotor for 20h at 4°C. Fractions of 500µl were collected and run in a SDS-PAGE gel. Protein bands were visualized with Coomassie Brilliant Blue G-250.

Pull-down assays- All reactions were carried out in Eppendorf tubes coated with 1% boiled/filtered fish skin gelatin. GST-fusion proteins and GST alone as control (10-20µg) were attached to glutathione-agarose beads. The beads were combined with nuclear extract or 15-30µg native/recombinant proteins in 300mM NaCl buffer and incubated for 1h at room temperature. The low-speed sediments were subsequently washed five times with the 300mM NaCl buffer and once with isotonic buffer. Bound proteins were eluted with hot SDS sample buffer and run in a SDS-PAGE gel. Protein bands were visualized with Coomassie Brilliant Blue G-250 and by western blot in the case of lamin-B. Pull-down assays were repeated at high ionic strengths (up to 1M).

The semi-quantitative experiments were performed in 400, 750, 1000, 2000, 5000, 10000 µl buffer with a constant amount of GST-TD and H3. Bands were quantified using the software Quantity-1 (Biorad). All experiments were repeated at least three times.

NMR spectroscopy- NMR spectra of LBR-TD were recorded at 25°C on Bruker DRX 500 and DRX 600 NMR spectrometers equipped with triple resonance cryoprobes and pulsed field gradients. Multi-dimensional NMR spectra were processed with NMRPipe (44) and analyzed with NMRView (45). Backbone and side chain ¹H, ¹⁵N and ¹³C resonances were assigned using a set of triple resonance experiments (46). The ¹H, ¹⁵N and ¹³C chemical shifts of the backbone resonances were obtained from sensitivity-enhanced three-dimensional HNCA, CBCA(CO)NH and CBCANH experiments. The side chain signals were assigned from three-dimensional HCCH-TOCSY and aromatic Cβ-Hδ/ε correlation experiments (47).

For NMR titrations, ¹H, ¹⁵N heteronuclear single quantum correlation (HSQC) experiments were recorded at 600 MHz proton frequency using a 50 µM ¹⁵N-labeled Tudor domain sample and an excess of the ligand in 20 mM sodium phosphate buffer (pH 6.9), 100 mM NaCl. Mono-, di- and trimethylated Lys, as well as symmetrically and asymmetrically dimethylated Arg amino acids were added up to a molar ratio of 1:25, while native and recombinant histone H3 were added stepwise up to a molar ratio of 1:3 (LBR-TD:H3).

Structure calculation- Distance restraints were derived from ¹⁵N- and ¹³C-edited 3D NOESY experiments. Hydrogen bond restraints were also derived from identification of slow-exchanging amide protons. Combined NOE cross-peak assignment and 3D protein structure calculation were performed using the program CYANA (48). Torsion angle restraints were derived from TALOS (49). The final ensemble of structures was refined in a box of solvent molecules as described (50). The quality of the structure ensemble was evaluated using WHAT IF (51) and PROCHECK-NMR (52). Molecular images were generated with PyMol (53). Structure similarity searches were performed using the DALI (54) server (www.ebi.ac.uk/dali). Coordinates have been deposited in the Protein Data Bank (accession code 2L8D). Chemical shift assignments and NOE

peak lists have been deposited in the BioMagResBank (accession code 17402).

FRAP experiments- Fluorescence recovery after photobleaching (FRAP) assays were performed on a Leica laser scanning confocal microscope (SP5) using suitable software and the 488-nm line of an argon laser. GFP-transfected cells were grown on special Petri dishes with coverslips attached and visualized in phenol-free culture medium buffered with Hepes-KOH. FRAP was performed with a bleach pulse of 6.5 seconds and initiated after 5 pre-bleach images. Post-bleach images (512 times 512 pixels) were collected for 307s at low laser power (24%). Data were corrected for fluorescence quench and recovery observed in the entire cell and in the background.

RESULTS

Overview of the structure and similarity to the Royal family domains. The protein analyzed, LBR-TD, comprised residues 1–62 of chicken LBR (see schematic diagram in Fig. 1A) and was monomeric in solution, as assessed by analytical ultracentrifugation (data not shown). The solution structure of LBR-TD was determined by heteronuclear multidimensional NMR spectroscopy with distance restraints derived from three-dimensional ¹⁵N- and ¹³C-edited NOE spectra. Experimental restraints and structural statistics over the 10 lowest energy structures are summarized in Table S1 (Supplementary data). An ensemble of the 10 lowest-energy NMR structures and a ribbon representation of the average structure are presented in Fig. 1B, C.

The NMR data show that residues 4-58 of LBR-TD form a well-defined tertiary structure, with an RMSD of 0.53Å for backbone atoms. All residues fall in the allowed regions of the Ramachandran plot. The amino-terminal residues 1-3 and the carboxy-terminal residues 59-62 are apparently disordered, as indicated by the paucity of inter-residue nuclear Overhauser effects (NOEs). The structured part adopts a β -barrel-like fold, consisting of five antiparallel β -strands. Strands β 1-4 are connected by short turns, while strands β 4 and β 5 are linked by a short 3_{10} -helix. A β -bulge (residues 28-29) allows one strand (β 2) to span both sides of the domain. The structure is stabilized by a well-defined hydrophobic core that

consists of residues Tyr6, Val12, Val25, Val27, Tyr37, Val39, Leu49 and Ile54 (Fig. 1D). These residues are conserved in most of the TD sequences known (2).

The side chains of the highly conserved amino acids Trp16 (loop 1), Tyr23 (strand β 2), Tyr41 (strand β 3) and Asp43 form a cluster on the surface of the domain (Fig. 1E). The location of these residues is highly reminiscent of the aromatic cages found in chromo and Tudor domains, which are thought to mediate recognition of methylated Lys or Arg side chains (4, 10, 12-15, 17-22) (Figure 2A). The conformation of the side chains in the aromatic residues is well-defined by a large number of NOE restraints in this area (Supplementary data, Fig. S1).

Structural homology searches using DALI (54) showed that the structure of the chicken LBR-TD is very similar to the TD of the survival motor neuron (SMN) protein (PDB 1g5v, Z score=7.9, rmsd=1.8Å) and the first of the two hybrid Tudor domains of the JMJD2A demethylase (PDB 2gfa, Z score=7.6, rmsd=1.7Å), although the sequence identity with these two homologs did not exceed 19% (Fig. 2B). Extensive similarity (Z score=8.0, rmsd=1.3) was also observed with a structure that corresponds to the Tudor domain of human LBR. This structure has been recently determined by NMR in the context of a Structural Genomics project by the RIKEN consortium and deposited in the database (PDB 2dig), but has not been further analyzed.

The amino acid composition of the LBR aromatic cage is very similar to that of the second hybrid Tudor domain of JMJD2A (HTD2), which is to date the only Tudor domain known to interact with trimethyllysine residues through a binding pocket that consists of only three aromatic residues and an aspartate (Fig. 2A). However, a more detailed comparison reveals that the aromatic cage of LBR-TD more strongly resembles the aromatic cage of the first hybrid Tudor domain of JMJD2A (HTD1) (Fig. 2B,C), which is not involved in binding trimethylated lysine residues (15, 17). Given that methyllysine or methylarginine recognition by Tudor domains does not involve appreciable structural rearrangements of the aromatic cage upon binding (Supplementary data, Fig. S2), it is therefore likely that LBR-TD is similar to HTD1 and does not recognize methylation marks.

Molecular interactions and functional role of LBR-TD. To explore the role of LBR-TD under *in vivo* conditions, we compared the properties of full-length LBR (FL-GFP) and a truncated LBR form lacking the entire LBR-TD module (Δ TD-GFP) in transiently transfected HeLa cells. As documented in Fig. 3A, FL-GFP and Δ TD-GFP exhibited a similar distribution, partitioning with the nuclear envelope (NE) and the endoplasmic reticulum (ER). Furthermore, when a segment corresponding to half the NE rim was photobleached (Fig. 3B, gallery), the two proteins recovered to a similar extent, yielding mobile fractions in the order of 0.6 and indicating that at steady-state a large fraction of the corresponding subunits does not exchange. However, assessing the relative recovery rates, we noticed that the median $t_{1/2}$ calculated from plateau fluorescence was greater for FL-LBR (14.0s) than for the mutant missing the LBR-TD (8.7s). Moreover, when the rates of fluorescence recovery in the NE and the ER were compared, the ratio of $NE_{t_{1/2}}/ER_{t_{1/2}}$ was 1.15 for FL-LBR and 0.94 for the mutant (Fig. 3B, Table). From these data it can be inferred that the absence of the Tudor domain renders NE-associated LBR relatively more mobile, presumably because the truncated protein fails to interact with some of the LBR partners.

Interactions of LBR-TD in vitro. As has been observed with other Tudor domains, no binding of LBR-TD to DNA and/or RNA could be detected by electrophoretic mobility-shift assays (data not shown). Furthermore, when increasing amounts of mono-, di- and tri- methylated Lys, or symmetrically and asymmetrically dimethylated Arg, were added to ^{15}N -labeled LBR-TD (up to a molar ratio of 1:25), there was no appreciable chemical shift or intensity changes in the ^1H - ^{15}N HSQC spectra (data not shown), suggesting that LBR-TD does not recognize free and methylated amino acids.

To better understand the role of LBR-TD at the molecular level, we studied its interactions with other nuclear proteins under *in vitro* conditions employing pull-down assays. For these purposes, we designed and expressed five recombinant proteins in bacteria, i.e. GST-TD, GST-RS, GST-SGD, GST-TDRS and GST-RSSGD, covering the entire amino-terminal part of LBR (for a schematic diagram see Fig. 1A). When the various LBR derivatives were

co-incubated with a NE-peripheral heterochromatin extract (for details and characterization, see 28), LBR-TD did not appear to bind any of the components present in the extract. However, GST-RS and all proteins containing the RS motif co-precipitated core histones and a 68kDa polypeptide corresponding to nuclear lamin B (Fig. 4A). Lamin binding could be directly confirmed by co-incubating each recombinant protein with purified, rat liver lamin B and probing the corresponding precipitate with specific anti-lamin B antibodies (Fig. 4B). Although a band with appropriate Mr was not immediately obvious in the initial precipitates, specific binding was also detected when GST-RS and GST-TDRS were co-incubated with purified MeCP2, a chromatin protein that has been shown to interact with LBR (Fig. 4C). Under the same conditions GST-TD failed to associate with this polypeptide.

LBR has also been shown to bind the isolated histones H3 and H4 (30). Based on this and taking into consideration the suggested role of the Tudor domains as potential “histone code readers”, we examined the *in vitro* interactions of LBR-TD, GST-RS and GST-TDRS with native core histones isolated from calf thymus. As shown in Fig. 4D, LBR-TD bound efficiently to histone H3, but did not associate with histones H4, H2A, or H2B. On the other hand, both GST-RS and GST-TDRS bound specifically to histones H3 and H4.

To ascertain that GST-TD binding to isolated H3 was specific, we repeated the assay in a quantitative fashion and under stringent ionic conditions, using both recombinant and native histones. As shown in Fig. 5A, binding was nearly stoichiometric, saturable, and could still occur in 1M salt. Furthermore, these observations could be confirmed by NMR-based titration experiments, in which increasing amounts of calf thymus H3 were added to ^{15}N -labeled LBR-TD and spectral changes monitored by recording ^1H - ^{15}N HSQC spectra. Extensive broadening and disappearance of several amide resonances upon H3 addition indicated that a LBR-TD-H3 complex was formed (Fig. 6A), presumably with an off-rate corresponding to an “intermediate exchange” binding regime (55). Histone interactions with their partners, such as chaperones or other chromatin-related proteins, are known to be

transient and often accompanied by conformational changes upon binding (56). Therefore, their interface is highly dynamic and this is reflected in the NMR spectra of their complexes, although with the data at hand we cannot differentiate line-broadening due to on/off binding exchange and conformational dynamics of the bound state. Residues perturbed by the interaction were identified based on the percentage of reduced signal intensity of amide resonances in LBR-TD upon addition of H3 at 1:3 molar ratio. These residues seemed to cover an extended area on one face of the LBR-TD surface, were not localized in the aromatic cluster or its vicinity (Fig. 6B) and are highly conserved in LBR sequences from different species (Fig. 6C). Notably, the same binding behavior was observed when LBR-TD was titrated with recombinant H3, reinforcing the idea that post-translational modifications of H3 were not required for this interaction (data not shown).

To get some insight on the type of contacts that could stabilize the LBR-TD/H3 interaction, we also examined the charge and the hydrophobic character of the binding surface (Fig. 6D). Based on this analysis, we can safely conclude that the binding is not solely dependent on electrostatic interactions, since there is a distinctly charged patch only in one part of the interaction surface, while the central part of the interface seems to be highly hydrophobic (as a result of the solvent exposed side-chains of Val20, Leu21 and Tyr23). These observations imply that the binding between LBR-TD and H3 is mediated by both electrostatic and hydrophobic interactions and is probably conserved in different organisms.

To identify the region of the histone H3 that was responsible for LBR-TD binding, we utilized different H3 fragments and assessed binding of LBR-TD to either H3 tails or tail-less H3. As shown in Fig. 5B, LBR-TD did not bind to the former peptide, but exhibited robust binding to the latter. Therefore, the association between the two proteins seems to be mediated by a region of the H3 molecule located in the so-called “histone fold” domain.

Puzzled by the fact that LBR-TD is able to associate with unassembled histone H3 but fails to bind core histone octamers present in NE-peripheral heterochromatin extracts, we repeated the experiments using an assortment of

chromatin particles and sub-particles. The results of these experiments were consistently negative (data not shown), except for H3-H4 tetramers isolated from salt-dissociated chromatin and hydroxylapatite column chromatography. As shown in Fig. 7A, both GST-TD and unfused TD obtained after cleavage with TEV protease exhibited mobility shifts when combined with H3-H4 tetramers and analyzed by rate zonal centrifugation in sucrose gradients. However, when an analogous experiment with H3-H4 tetramers was done in a column chromatography format (passing the histone oligomers through a glutathione-GST-TD column), we discovered that LBR-TD, instead of binding stably to H3-H4 tetramers, was apparently “stealing” H3 from the sub-particles (Fig. 7B). From these results it would appear that binding of histone H3 to LBR-TD is antagonistic to binding to histone H4. This interpretation is graphically presented in the model shown in Fig. 8 and discussed in detail below.

DISCUSSION

In this study we have determined the tertiary structure of chicken LBR-TD in solution and examined its interactions with a variety of substrates and potential binding partners. As expected, LBR-TD was found to adopt the characteristic β -barrel fold of other Tudor domains and was apparently required for normal LBR dynamics in living cells. However, in contrast to a previously published report (33), our results showed that LBR-TD was not sufficient for lamin B binding *in vitro*, and that this interaction also required the adjacent Arg/Ser (RS) region.

Unlike other Tudor domains, LBR-TD did not associate with unmodified or modified Arg/Lys residues. Thus, no binding to free or methylated aminoacids, to the LBR-RS, to Lys and Arg-rich histone tails or to intact nucleosomal core particles was observed *in vitro*. This rules out the possibility that LBR-TD might operate as an “RS-trap” or as a “K/R modification reader”, in the fashion that other Tudor domains bind to Arg- and Lys-methylated proteins and peptides, Arg-Gly rich sequences or free amino acids (4-19,57). At a first glance this may seem paradoxical, because LBR-TD *does* contain an aromatic cage, similar to the second hybrid Tudor domain of JMJD2A, the chromodomain of HP1 and other Royal family

domains that can accommodate methylated Lys residues (2,20-22). However, recognition of symmetrically dimethylated arginine by Tudor domains has been shown to require four aromatic residues in the binding pocket (12-13 and Figs. 2A, S3), while the aromatic cage of LBR-TD comprises only three aromatic residues (Figs. 1E, 2B) and may therefore be incapable of accommodating a methylated guanidium group (Supplementary data, Fig. S3). Furthermore, based on the structural data that are available, the same argument holds for dimethyl-lysine recognition, since 53BP1 utilizes four aromatic residues to line the binding pocket (14, 59). In a recent, comprehensive review article on LBR (26), the authors compared the structure and the aromatic cage of human LBR-TD with the ones from other Tudor domains and suggested that LBR-TD and 53BP1-TD, being structurally similar, might recognize the same ligand, i.e. H4K20me2. However, based on the structure of the 53BP1/H4-K20me2 complex, the binding pocket of 53BP1 has been described before as a distinctly compact aromatic cage with four aromatic residues coming in direct contact with the methylated side chains and a fifth aromatic residue packing tightly against the ligand peptide backbone (14). The same mode of dimethyllysine recognition by 53BP1-TD has also been identified in the complex it forms with a p53-K382me2 peptide (58). Therefore, the aromatic cage of LBR-TD differs significantly from that of 53BP1 (see also Supplementary data, Fig. S3). The only exception in the Tudor tendency to construct pockets with four aromatic residues for caging the methylated marks are the two hybrid Tudor domains of JMJD2A that form an interdigitated structure (15,17). Although in both hybrid domains of JMJD2A the aromatic cages consists of three aromatic residues and an aspartate residue like the one found in LBR-TD, only the second of the two hybrid lobes (HTD2) is able to bind peptides containing a trimethyllysine residue (because the aromatic side chains are positioned orthogonally with respect to each other, which in turn generates sufficient space for accommodating the large trimethylammonium group). In the first hybrid lobe of JMJD2A (HTD1) the histidine sidechain is oriented towards the inside of the cage-like enclosure and occupies a position that occludes caging of any methylated Lys. In

LBR-TD, Trp16 has the same orientation as the histidine residue of HTD1-JMJD2A (Supplementary data, Fig. S3) and (most likely) precludes LBR-TD from binding methylation marks. It should be noted that the relative disposition of the aromatic sidechains of LBR-TD is well defined due to the large number of NOEs detected (Supplementary data, Fig. S1). In addition, the arrangement of these residues in the chicken LBR-TD structure presented here and in the human LBR-TD structure previously determined is very similar (Supplementary data, Fig. S4). In short, the aromatic pocket of LBR is too “open” to allow efficient caging of Arg or Lys dimethylated side chains and too restrictive for the trimethylated Lys side chain to fit in.

Invariably, recognition of methylated ligands by the Royal family of domains depends largely on electrostatic stabilization mediated by cation- π interactions. Therefore, the surface of the binding pocket has a negatively charged electrostatic potential to attract the cationic moiety of the Lys or Arg residue. In LBR-TD the surface surrounding the cluster of aromatic residues is not as negatively charged as that of Tudor domains known to bind methylated marks (Supplementary data, Fig. S5). Taken together, despite the overall fold similarity with other Tudor domains, these differences in the spatial arrangement of the aromatic rings that build the binding pocket in LBR-TD prohibit stable binding of the so-far characterized Tudor ligands.

Interestingly, a protein-array approach for identifying novel methyl-lysine-dependent interactions of several Royal family domains, in agreement with our experimental observations and detailed structural analysis, also failed to detect any interaction between LBR-TD and variably methylated H3 or H4 tails (16,59). This work was based on a protein-domain microarray screening and tested, among others, human LBR-TD as one of several chromatin-associated domains that might specifically recognize histone H3 and H4 tail peptides methylated to varying degrees on specific lysine or arginine residues. The results of this investigation revealed that LBR-TD failed to bind to any of the peptides used as baits, among them the H4-K20me2, but also H3-K79me2, H3-K4me1,2,3, H3K9-me1,2,3 and H4K20-me1,2,3. On the contrary, the Tudor-domain of 53BP1 was found to recognize

H3 and H4 peptides bearing dimethyl-groups on H4-K20, H3-K4 and H3-K9. Nevertheless, we cannot exclude the possibility that LBR-TD could recognize a larger peptide ligand in the context of the full-length LBR protein as has been observed for methyl-lysine binding PHD-containing proteins (60).

Instead of mediating binding to methylated amino acids, we found that LBR-TD binds tightly and selectively to isolated histone H3 through a patch on its surface distinct from the aromatic cage and comprised of residues highly conserved in LBR sequences from different species. The binding is mediated by both electrostatic and hydrophobic interactions and is probably conserved in different organisms. In addition, our data indicate that H3-H4 oligomers and fully assembled nucleosomes interact with the RS region of LBR, which is physically contiguous with LBR-TD. Based on these findings, we propose a model (Fig. 8), whereby LBR-TD may in fact operate as a transient docking site or as a storage chaperone for newly imported histone H3 molecules, as, for example, CIA/ASF1 (56,60-61). In this scenario, H3-H4 binding may occur at a later phase, following chaperoning by LBR-TD. The assembly of H3-H4 tetramers might trigger their release from the Tudor domain and allow the deposition of nucleosome assembly intermediates (or complete histone octamers) to the adjacent RS region. Interestingly, an antiparallel β -sheet motif similar to that of LBR-TD has been recognized before as a recurring theme in histone chaperones and as a scaffold for histone binding elements (61-62). Certainly, at this point and in the lack of sound experimental evidence, this is only a hypothesis and needs to be validated in a concrete

biological context. However, this hypothesis highlights an alternative scenario for the functional role of LBR in the nuclear envelope, which remains elusive, despite its experimentally documented association with chromatin and nuclear lamina (26). This scenario is compatible with the presumed role of nuclear envelope as a platform for peripheral heterochromatin assembly and chromatin remodeling, based on a large body of experimental evidence (24-27).

Tudor domains are found in many proteins and serve distinct functions. To some extent, these diverse binding properties match the various different arrangements of the Tudor domains, which occur as single (SMN and LBR), or as tandem units (53BP1 and UHRF1). An interdigitated arrangement of tandem Tudor domains (JMJD2A), or an insertion of the Tudor domain into another fold (SND1), have also been observed. In addition, there are proteins that contain multiple repeats of such domains. For example, the *Drosophila* Tudor protein has eleven copies of the Tudor domain, whereas in the TDRD family the numbers range from one to eight. Based on their primary sequences and the general mechanisms used to recognise methylated ligands, it is not expected that all Tudor domains represent genuine methyl-binders. Instead, it is likely that the stable Tudor fold might serve architectural purposes and can be used as a platform for orienting other functional domains in the full-length protein. Such a role may apply to LBR-TD, because this domain adopts a stable globular structure and may thus assist in organizing the rest of the molecule and coordinating its interactions with chromatin particles.

REFERENCES

1. Talbot, K., Miguel-Aliaga, I., Mohaghegh, P., Ponting, C. P., and Davies, K. E. (1998) *Hum. Mol. Genet.*, **7**, 2149-2156
2. Maurer-Stroh, S., Dickens, N. J., Hughes-Davies, L., Kouzarides, T., Eisenhaber, F., and Ponting, C. P. (2003) *Trends Biochem. Sci.*, **28**, 69-74
3. Ponting, C. P. (1997) *Trends Biochem. Sci.*, **22**, 51-52
4. Selenko, P., Sprangers, R., Stier, G., Buhler, D., Fischer, U., and Sattler, M. (2001) *Nat. Struct. Biol.*, **8**, 27-31
5. Sprangers, R., Groves, M. R., Sinning, I., and Sattler, M. (2003) *J. Mol. Biol.*, **327**, 507-520
6. Bühler, D., Raker, V., Lührmann, R., and Fischer, U. (1999). *Hum. Mol. Genet.*, **8**, 2351-2357
7. Brahm, H., Meheus, L., Brabandere, V., Fischer, U., and Lührmann, R. (2001). *RNA*, **7**, 1531-1542
8. Cote, J., Richard, S. (2005) *J. Biol. Chem.*, **280**, 28476-28483

9. Shaw, N., Zhao, M., Cheng, C., Xu, H., Saarikettu, J., Li, Y., Da, Y., Yao, Z., Silvennoinen, O., Yang, J. et al. (2007) *Nat. Struct. Mol. Biol.*, **14**, 779-784
10. Friberg, A., Corsini, L., Mourao, A., and Sattler, M. (2009) *J. Mol. Biol.*, **387**, 921-934
11. Kirino, Y., Vourekas, A., Sayed, N., de Lima Alves, F., Thomson, T., Lasko, P., Rappsilber, J., Jongens, T.A., and Mourelatos, Z. (2010) *RNA*, **16**, 70-78
12. Liu, K., Chen, C., Guo, Y., Lam, R., Bian, C., Xu, C., Zhao, D.Y., Jin, J., MacKenzie, F., Pawson, T., and Min, J. (2010) *Proc. Natl. Acad. Sci. USA*, **107**, 18398-18403
13. Liu, H., Wang, J. H. S., Huang, Y., Li, Z., Gong, W., Lehmann, R., and Xu, R.M. (2010) *Genes Dev.*, **24**, 1876-1881
14. Botuyan, M. V., Lee, J., Ward, I. M., Kim, J. E., Thompson, J. R., Chen, J. and Mer, G. (2006) *Cell*, **127**, 1361-1373
15. Huang, Y., Fang, J., Bedford, M. T., Zhang, Y., and Xu, R. M. (2006) *Science*, **312**, 748-751
16. Kim, J., Daniel, J., Espejo, A., Lake, A., Krishna, M., Xia, L., Zhang, Y., and Bedford, M. T. (2006) *EMBO Rep.*, **7**, 397-403
17. Lee, J., Thompson, J. R., Botuyan, M. V., and Mer, G. (2008) *Nat. Struct. Mol. Biol.*, **15**, 109-111
18. Ramos, A., Hollingworth, D., Adinolfi, S., Castets, M., Kelly, G., Frenkiel, T. A., Bardoni, B., and Pastore, A. (2006) *Structure*, **14**, 21-31
19. Charier, G., Couprie, J., Alpha-Bazin, B., Meyer, V., Quemeneur, E., Guerois, R., Callebaut, I., Gilquin, B., and Zinn-Justin, S., (2004) *Structure*, **12**, 1551-1562
20. Corsini, L., and Sattler, M. (2007) *Nat. Struct. Mol. Biol.*, **14**, 98-99
21. Jacobs, S. A., and Khorasanizadeh, S. (2002) *Science*, **295**, 2080-2083
22. Nielsen, P. R., Nietlispach, D., Mott, H. R., Callaghan, J., Bannister, A., Kouzarides, T., Murzin, A. G., Murzina, N. V., and Laue, E. D. (2002) *Nature*, **416**, 103-107
23. Worman, H. J., Yuan, J., Blobel, G., and Georgatos, S. D. (1988) *Proc. Natl. Acad. Sci. USA*, **85**, 8531-8534
24. Wilson, K. L., and Foisner, R. (2010) *Cold Spring Harb. Perspect. Biol.*, **2**, a000554
25. Georgatos, S. D. (2001) *EMBO J.*, **20**, 2989-2994
26. Olins, A. L., Rhodes, G., Welch, D. B., Zwerger, M., and Olins, D. E. (2010) *Nucleus*, **1**, 53-70
27. Schirmer, E. C., and Foisner, R. (2007) *Exp. Cell Res.*, **313**, 2167-79
28. Makatsori, D., Kourmouli, N., Polioudaki, H., Shultz, L. D., McLean, K., Theodoropoulos, P. A., Singh, P. B., and Georgatos, S. D. (2004) *J. Biol. Chem.*, **279**, 25567-25573
29. Nikolakaki, E., Simos, G., Georgatos, S. D., and Giannakouros, T. (1996) *J. Biol. Chem.*, **271**, 8365-8372
30. Polioudaki, H., Kourmouli, N., Drosou, V., Bakou, A., Theodoropoulos, P. A., Singh, P. B., Giannakouros, T., and Georgatos, S. D. (2001) *EMBO Rep.*, **2**, 920-925
31. Nikolakaki, E., Drosou, V., Sanidas, I., Peidis, P., Papamarcaki, T., Iakoucheva, L. M., and Giannakouros, T. (2008) *Biochim. Biophys. Acta*, **1780**, 214-225
32. Ye, Q., Callebaut, I., Pezhman, A., Courvalin, J. C., and Worman, H. J. (1997) *J. Biol. Chem.*, **272**, 14983-14989
33. Lin, F., Noyer, C. M., Ye, Q., Courvalin, J. C., and Worman, H. J. (1996) *Hepatology*, **23**, 57-61
34. Guarda, A., Bolognese, F., Bonapace, I. M., and Badaracco, G. (2009) *Exp. Cell. Res.*, **315**, 1895-903
35. Ye, Q., and Worman, H. J. (1994) *J. Biol. Chem.*, **269**, 11306-11311
36. Sambrook, J., Fritsch, E. F., and Maniatis, T. M. (1989) *In Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
37. Luger, K., Rechsteiner, T. J., Flaus, A. J., Waye, M. M., and Richmond, T. J. (1997) *J. Mol. Biol.*, **272**, 301-311
38. Ling, X., Harkness, T. A., Schultz, M. C., Fisher-Adams, G., and Grunstein, M. (1996) *Genes Dev.*, **10**, 686-699
39. Oganessian, N., Kim, S. H., and Kim, R. (2005) *J. Struct. Funct. Genomics*, **6**, 177-182
40. Georgatos, S. D., and Blobel, G. (1987) *J. Cell Biol.*, **105**, 105-115

41. Blobel, G., and Potter, V. R. (1966) *Science*, **154**, 1662-1665
42. Maison, C., Horstmann, H., and Georgatos, S. D. (1993) *J. Cell Biol.*, **123**, 1491-1505
43. Simon, R. H., and Felsenfeld, G. (1979) *Nucleic Acids Res.*, **6**, 689-696
44. Delaglio, F., Grzesiek, S., Vuister, G. W., Zhu, G., Pfeifer, J., and Bax, A. (1995) *J. Biomol. NMR*, **6**, 277-293
45. Johnson, B. A. (2004) *Methods Mol. Biol.*, **278**, 313-352
46. Sattler, M., Schleucher, J., and Griesinger, C. (1999) *Prog. NMR Spectrosc.*, **34**, 93-158
47. Yamazaki, T., Forman-Kay, J. D., and Kay, L. E. (1993) *J. Am. Chem. Soc.*, **115**, 11054-11055
48. Herrmann, T., Guntert, P., and Wüthrich, K. (2002) *J. Biomol. NMR*, **24**, 171-189
49. Cornilescu, G., Delaglio, F., and Bax, A. (1999) *J. Biomol. NMR*, **13**, 289-302
50. Brunger, A. T., Adams, P. D., Clore, G. M., DeLano, W. L., Gros, P., Grosse-Kunstleve, R. W., Jiang, J. S., Kuszewski, J., Nilges, N., Pannu, N. S., Read, R. J., Rice, L. M., Simonson, T., and Warren, G. L. (1998) *Acta Cryst. D*, **54**, 905-921
51. Vriend, G. (1990) *J. Mol. Graph.*, **8**, 52-56.
52. Laskowski, R. A., Rullmann, J. A., MacArthur, M. W., Kaptein, R., and Thornton, J. M. (1996) *J. Biomol. NMR*, **8**, 477-486
53. Delano, W. L. (2002) *The Pymol Molecular Graphics System*, Delano Scientific, San Carlos, CA.
54. Holm, L., and Sander, C. (1995) *Trends Biochem. Sci.*, **20**, 478-480
55. Matsuo, H., Walters, K. J., Teruya, K., Tanaka, T., Gasser, G. T., Lippard, S. J., Kyogoku, Y., and Wagner, G. (1999) *J. Am. Chem. Soc.*, **121**, 9903-9904
56. Mousson, F., Lautrette, A., Thuret, J. Y., Agez, M., Courbeyrette, R., Amigues, B., Becker, E., Neumann, J. M., Guerois, R., Mann, C., and Ochsenbein, F. (2005) *Proc. Natl. Acad. Sci. USA*, **102**, 5975-80
57. Grenon, M., Costelloe, T., Jimeno, S., O'Shaughnessy, A., Fitzgerald, J., Zgheib, O., Degerth, L., and Lowndes, N. F. (2007) *Yeast*, **24**, 105-119
58. Ro, S., Musselman, C. A., Kachirskaia, I., Hayashi, R., Glass, K.C., Nix, J. C., Gozani, O., Appella, E., and Kutateladze, T. G. (2010) *J. Mol. Biol.*, **398**, 489-496
59. Yang, Y., Lu, Y., Espejo, A., Wu, J., Xu, W., Liang, S., and Bedford, M. T. (2010) *Mol. Cell*, **40**, 1016-1023
60. Horton, J. R., Upadhyay, A. K., Qi, H. H., Zhang, X., Shi, Y., and Cheng, X. (2010) *Nat. Struct. Mol. Biol.*, **17**, 38-43
61. Natsume, R., Eitoku, M., Akai, Y., Sano, N., Horikoshi, M., and Senda, T. (2007) *Nature*, **446**, 338-41
62. Park, Y. J., and Luger, K. (2006) *Proc. Natl. Acad. Sci. USA*, **103**, 1248-1253

FOOTNOTES

We would like to thank G. Stier (EMBL) for the generous gift of the pETM-30 plasmid, K. Luger (University of Colorado) for H3 plasmids, M. Grunstein (University of California) for H3-tail plasmids, G. Badaracco (University of Insubria) for MeCP2 plasmids and antibodies, E. Nikolakaki (Aristotle University of Thessaloniki) for help with cloning experiments at the early stages of this work, Y. Markaki and D. Makatsori (University of Ioannina) for providing advice with the nuclear envelope extract isolation and GST pull-down experiments, B. Simon (EMBL) for help with NMR titration experiments, A. Lustig (Biozentrum, University of Basel) for Analytical Ultracentrifugation experiments and T. Papamarcaki (University of Ioannina) for materials. We thank the confocal laser microscope facility of the University of Ioannina for use of the Leica TCS-SP scanning confocal microscope. This work was supported by the Greek Ministry of Education and the European Commission under the program "Pythagoras II", by the Greek General Secretariat for Research and Development under the program "PENED2003" and by the "IKYDAD 2004" Programme of scientific exchange and cooperation between Greece and Germany.

The abbreviations used are: LBR, Lamin B Receptor; LBR-TD, LBR Tudor domain; FRAP, Fluorescence recovery after photobleaching; RS, Arg/Ser motifs; SMN, survival motor neuron protein; NE, nuclear envelope; LBR-NT, amino-terminal segment of LBR; TD, Tudor domain; HP1, Heterochromatin Protein 1; NOE, nuclear Overhauser effect; FL-GFP, full-length LBR-GFP; Δ TD, truncated LBR form lacking the entire LBR-TD; ER, endoplasmic reticulum; SGD, second globular domain

FIGURE LEGENDS

Fig. 1. Structure of the LBR Tudor domain (LBR-TD). (A) Layout of LBR molecule. (B) Stereo view of the backbone atoms (N, C α , C') for residues of a NMR ensemble of 10 (out of 100 computed) lowest energy structures of LBR-TD. Secondary structure elements are colored cyan for β -strands and red for the 3_{10} -helix connecting β 4 and β 5. (C) Ribbon diagram of the closest to the mean LBR-TD structure. Secondary structure elements colored as above. (D) Rotated view of the Tudor domain structure showing the side chains of hydrophobic core residues in green. (E) Close view of the aromatic cage discussed in the text. The side chains of the three residues (W16, Y23, Y41) that form the aromatic cluster are shown in green.

Fig. 2. Comparison of the structures of the closest structural homologues of LBR-TD. (A) Structure-based sequence alignment of the LBR-TD, the interdigitated Tudor domains of JMJD2A, and the Tudor domains of 53BP1, *Drosophila* TUDOR, SMN and SND1 proteins. Residues that form the aromatic cage in each domain are boxed and in colored background. Other invariant residues among the LBR and JMJD2A domains are bold. Notice that only the second domain of JMJD2A binds trimethyl-lysine (Kme3), 53BP1 binds dimethyl-lysine (Kme2), whereas the SMN, *Drosophila* TUDOR, and SND1 Tudor domains specifically recognise symmetrically dimethylated arginine (sDMA). Due to the interdigitated structure of JMJD2A the sequences for the superimposed structures are: HTD1 T900..E925 - I983..V1003, HTD2 A958..P982 - T926..I946. (B) Ribbon diagrams of LBR-TD structural homologues. The structural homologues were identified by DALI and are displayed in green for LBR, magenta for SMN (PDB code: 1g5v), red and pink for JMJD2A Tudor domain 1 (HTD1) and 2 (HTD2), respectively (PDB code: 2gfa). (C) Close-up view of the superimposed aromatic cages of the three domains. Coloring scheme same as in B.

Fig. 3. (A) Steady-state localization of full length LBR (FL-GFP) and Tudor truncated LBR (Δ TD-GFP) in transiently transfected HeLa cells. (B) (upper panel) FRAP data from transfected HeLa cells expressing FL-GFP and Δ TD-GFP LBR. (bottom panel) The table shows the median $t_{1/2}$ when half of the nuclear envelope was bleached ($t_{1/2}$ NE), the ratio of median $t_{1/2}$ between NE and ER and the number of independent experiments (n). Data shown correspond to 41 independent FRAP assays for the full length LBR and 17 independent experiments for the Δ TD mutant.

Fig. 4. GST pull-down experiments with several regions of N-terminal LBR. (A) GST-fused LBR-TD, LBR-RS, LBR-SGD, LBR-TDRS, LBR-RSSGD were incubated with nuclear envelope extracts, produced after MNase digestion of turkey erythrocyte nuclei. Bound proteins were eluted from GST-beads with SDS sample buffer and visualized with Coomassie staining. (B) GST-fused LBR-TD, LBR-RS, LBR-TDRS were also incubated with: (B) Lamin B, (C) recombinant MeCP2 and (D) purified native core histones.

Fig. 5. Interaction of LBR-TD with histone H3. (A) Semiquantitative pull-down experiment using constant amounts of GST-Tudor and native H3 in serial dilutions with buffer, as detailed in Materials and Methods. Experiments were repeated in the same manner with recombinant H3. Inset gels shows 20% of input native H3/recombinant H3. (B) Interaction of LBR-TD with core and tail parts of H3. GST pull-down assays with GST-fused LBR-TD and recombinant tail-less H3 (left panel) and with LBR-TD and GST-fused H3 N-terminal tail (right panel). Panels show Coomassie Blue-stained SDS-PAGE gels and 50% of input tailless H3 (left) or LBR-TD (right).

Fig. 6. Features of the cLBR-TD-H3 interface. (A) NMR-based titration experiment using ^{15}N -labeled LBR Tudor and increasing amounts of native or recombinant histone H3. A representative overlay of HSQC spectra for LBR-TD (blue) and for equimolar amounts of H3 and LBR-TD (red) is shown. (B) Mapping the interaction surface of histone H3 onto LBR's LBR-TD structure. Surface representations of the LBR-TD structure that differ by a 180° rotation around the y axis. The side chains of residues with perturbed amide resonances upon titration with H3 are shown in magenta and the LBR-TD surface in light blue. (C) Sequence alignment of the LBR-TD from several species (from top to bottom: chicken, human, Pongo abelii, bovine, rat, mouse, zebrafis, xenopus laevis). The conservation of residues across species is shown under the alignment. Symbols denote identical residues (*), highly similar residues (:) and similar residues (.). Arrows point to the aminoacids comprising the interaction surface between cLBR-TD and H3 (identified by the NMR-chemical shift perturbation experiments). (D) Sequence conservation, hydrophobicity and electrostatic potential of the H3 binding surface of cLBR-TD. Mapping on the surface of (from left to right): the residues of cLBR-TD comprising the interaction surface with H3, the sequence conservation, the hydrophobicity and the charge separation.

Fig. 7. Interaction of LBR-TD with H3-H4 tetramers. LBR-TD either fused with GST (left panel) or unfused (right panel) was mixed with H3-H4 tetramers and run in a 5-20% sucrose gradients. Fractions were collected and analyzed in SDS-PAGE gels stained with Coomassie Brilliant Blue. (B) GST-fused LBR was incubated with isolated H3-H4 tetramer. Bound proteins were eluted from GST-beads with SDS sample buffer and analyzed in a SDS-PAGE gel, stained with Coomassie Brilliant Blue.

Fig. 8. A hypothetical model for the “chaperone-like” function of LBR-TD. According to this model, LBR-TD operates as a transient docking site for “free” H3 (I). Subsequent H3-H4 binding (II) triggers release of H3-H4 tetramers and deposition of nucleosome assembly intermediates to the adjacent RS region (III).

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6





Figure 7

A.



B.




Figure 8

