

Article type: Other

**Genome-wide exploration identifies sex-specific genetic effects of alleles upstream *NPY*
to increase the risk of severe periodontitis in men**

Sandra Freitag-Wolf¹, Henrik Dommisch², Christian Graetz³, Yvonne Jockel-Schneider⁴,
Inga Harks⁵, Ingmar Staufienbiel⁶, Joerg Meyle⁷, Peter Eickholz⁸, Barbara Noack⁹, Corinna
Bruckmann¹⁰, Christian Gieger¹¹, Søren Jepsen², Wolfgang Lieb¹², Stefan Schreiber¹³, Inke
R. König^{14*}, Arne S. Schaefer^{13*}

¹Institute of Medical Informatics and Statistics, Christian-Albrechts-University Kiel,
Germany

²Department of Periodontology, Operative and Preventive Dentistry, Rheinische-Friedrichs-
Wilhelm-University, Bonn, Germany

³University Medical Center Schleswig-Holstein, Campus Kiel, Department of Operative
Dentistry and Periodontology, Kiel, Germany

⁴Department of Periodontology, Clinic of Preventive Dentistry and Periodontology,
University Medical Center of the Julius-Maximilians-University, Würzburg, Germany

⁵Center of Periodontology, Operative and Preventive Dentistry, University Medical Center
Münster, Germany

⁶Department of Conservative Dentistry, Periodontology and Preventive Dentistry, Hannover
Medical School, Germany

⁷Department of Periodontology, University Medical Center Giessen and Marburg, Germany

This article has been accepted for publication and undergone full peer review but has not
been through the copyediting, typesetting, pagination and proofreading process, which may
lead to differences between this version and the Version of Record. Please cite this article as
doi: 10.1111/jcpe.12317

This article is protected by copyright. All rights reserved.

⁸Department of Periodontology, Centre for Dental, Oral, and Maxillofacial Medicine (Carolinum), Johann Wolfgang Goethe-University, Frankfurt am Main, Germany

⁹University Medical Center Carl Gustav Carus der Technischen Universität Dresden, Center of Periodontology, Operative and Preventive Dentistry, Clinic of Preventive Dentistry, Dresden, Germany

¹⁰Department of Conservative Dentistry and Periodontology, Bernhard Gottlieb University Clinic of Dentistry, Vienna, Austria

¹¹Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

¹²Institute of Epidemiology and Biobank popgen, University Medical Center Schleswig-Holstein, Germany

¹³Institute for Clinical Molecular Biology, Christian-Albrechts-University Kiel, Germany

¹⁴Institute for Medical Biometry and Statistics, University of Lübeck, University Hospital Schleswig-Holstein, Lübeck, Germany

*both authors contributed equally

Running title: Sex-specific effects of *NPY* alleles

Keywords: periodontitis; *NPY*, sex, interaction, genetic association

Correspondence address: Arne S. Schaefer

Christian-Albrechts-University Kiel, Institute of Clinical Molecular Biology,
Arnold-Heller-Str. 3, 24105 Kiel, Germany

Tel.: +49-(0)431-597-3882, Fax: +49-(0)431-597-1842

This article is protected by copyright. All rights reserved.

Email: a.schaefer@ikmb.uni-kiel.de

Inke R. König, Prof., Ph.D.

Institute for Medical Biometry and Statistics,

University Hospital Schleswig-Holstein, Campus Lübeck,

Maria-Goeppert-Str. 1, 23562 Lübeck, Germany

Tel.: +49-(0)451-500 5580, Fax: +49-(0)451-500 2999. E-Mail: inke.koenig@imbs.uni-luebeck.de

Conflict of interest and source of funding statement

Author AS and SJ are funded by grants of the Deutsche Forschungsgemeinschaft (SCHA 1582/2-1, KFO208); The popgen 2.0 network is supported by a grant from the German Ministry for Education and Research (01EY1103). The popgen 2.0 network is supported by a grant from the German Ministry for Education and Research (01EY1103) and by grants from the Deutsche Forschungsgemeinschaft Excellence Cluster “Inflammation at Interfaces” (EXC306, EXC306/2). The KORA research platform (KORA, Cooperative Research in the Region of Augsburg) was initiated and is financed by the Helmholtz Zentrum München - German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research and by the State of Bavaria.

Abstract

Aim: Periodontitis (PD) is influenced by genetic as well as life-style and socio-economic factors. Epidemiological studies show that men are at greater risk of severe forms of PD, suggesting interplay between sex and genetic factors. We aimed to systematically analyze patients with aggressive periodontitis (AgP) for gene-sex interactions.

Materials and Methods: 329 German AgP cases and 983 controls were genotyped with Affymetrix 500K Arrays and were analyzed by logistic regression analysis. The most significant gene-sex interaction was replicated in an independent sample of 382 German/Austrian AgP cases and 489 controls.

Results: Ten single nucleotide polymorphisms (SNPs) in strong linkage disequilibrium ($r^2 > 0.85$) upstream the gene neuropeptide Y (*NPY*) suggested gene-sex interaction ($p < 5 \times 10^{-5}$). SNP rs198712 showed the strongest association in interaction with sex ($p = 5.4 \times 10^{-6}$) with odds ratios in males and females of 1.63 and 0.69, respectively. In the replication, interaction of sex with rs198712 was verified with $p = 0.022$ (pooled $p = 4.03 \times 10^{-6}$) and similar genetic effects. Analysis of chromatin elements from ENCODE data revealed tissue specific transcription at the associated non-coding region.

Conclusion: The current study is the first to observe a sexually dimorphic role of alleles at *NPY* in humans and support previous genome-wide findings of a role of *NPY* in severe PD.

Clinical Relevance

Scientific rationale for the study: Epidemiological studies showed that men are at greater risk of severe forms of periodontitis suggesting interplay between sex and genetic factors.

Principal findings: Ten alleles upstream the gene neuropeptide Y (*NPY*) suggested gene-sex interaction ($p < 5 \times 10^{-5}$). rs198712 showed the strongest association ($p = 5.4 \times 10^{-6}$) with odds ratios in males and females of 1.63 and 0.69, respectively. In the replication, interaction of sex with rs198712 was verified with $p = 0.022$.

Practical implications: The data provide evidence of a sexually dimorphic role of alleles at *NPY* in humans and support previous genome-wide findings of a role of *NPY* in PD.

Introduction

Periodontitis (PD) is a chronic inflammatory disease of the oral cavity. The inflammation is elicited by the oral microbial biofilm that leads to gingival bleeding, pocket formation, attachment and bone loss and eventually to tooth loss as final outcome (Buchwald et al., 2013). PD affects human populations worldwide at prevalence rates of 11 % for the severe forms (Marcenes et al., 2013). It is largely classified into the sub-forms chronic periodontitis (CP) and aggressive periodontitis (AgP). Whereas CP is mostly observed in adults and is characterized by slow progression of the disease, AgP is found in young individuals and is diagnosed based on rapid attachment loss and destruction of the alveolar bone. A pathogenic microflora is the causing agent of PD, however, factors such as smoking, diabetes and obesity can contribute to the disease risk (Buchwald et al., 2013, Mealey & Oates, 2006, Sangwan et al., 2013, Schenkein & Loos, 2013). Related to these risks, socio-economic factors such as poor educational attainment and low income are also associated with the progression of PD and tooth loss. (Buchwald et al., 2013) For CP, epidemiologic studies provide broad-based evidence that men are at greater risk of developing severe periodontal disease than women (McGrath & Bedi, 2003, Shiau & Reynolds, 2010). A recent study estimated the prevalence, severity, and extent of periodontitis in the adult U.S. population with data from the 2009 and 2010 National Health and Nutrition Examination Survey (NHANES) cycle and reported that after adjustment for the effect of age, total periodontitis was significantly higher in men than in women aged 30 years and older (males= 57%; females= 39%), with males showing a 3%, 7%, and 8% higher prevalence of mild, moderate, and severe PD, respectively, compared with women. (Eke et al., 2012) Findings like these suggest interplay between genetics, sex

and possibly other environmental factors in PD. For AgP, an increased prevalence in males or females is currently not supported in the literature (Hart et al., 1991). However, the few available studies are small, which decreased their ability for generalization.

A genetic basis of PD has been known by formal genetic studies for many years (Corey et al., 1993, Marazita et al., 1994, Michalowicz et al., 1991, Michalowicz et al., 2000) but compared to other complex diseases a relatively low number of risk alleles have been identified that provide evidence of association by repeated replication in independent, large case-control populations (Schaefer et al., 2013).

In common with other complex diseases, there is a gap between statistical modeling and biological phenomena, which is notoriously understudied. In general, genetic susceptibility regions were analyzed for the average proband, but further subject characteristics such as sex were not taken into account. A limitation of this approach is that relevant influential factors can be overlooked simply because they exert their effect only against specific patterns of expression of other factors.

We hypothesized interactions between specific SNPs and sex that influence the disease risk of PD. To test this hypothesis, the genetic constitution and the interaction term of sex was analyzed on a genome-wide level in a case-control sample of the early onset phenotype AgP. A focus on this phenotype allowed a better control of confounding effects such as age and accumulating life-style factors and due to the strong severity at a young age, it is assumed that genetic factors play a prominent role in disease susceptibility. The main result was verified in an independent AgP case-control replication sample.

Material and Methods

Study population

Cases and controls that were genotyped with Affymetrix 500K Arrays were described before (Schaefer et al., 2010). In brief, inclusion criteria for the 329 German AgP cases (131 males, 189 females) of the GWAS panel were ≥ 7 teeth with 50% alveolar bone loss from the cemento-enamel junction to the tooth apex in subjects who were under the age of 35 years and whose parents and grandparents were born in Germany. To limit a potential bias in the diagnosis of the disease phenotype by different examiners, a set of full-mouth dental radiographs was used. Copies were mailed for independent confirmative periodontal bone scoring to the study center and the initial diagnosis was confirmed by a second specialist in periodontology who has, without exception, been located and educated at the study center. The use of radiographs allowed a reproducible and quantifiable diagnosis of the disease phenotype. The controls were population representative individuals from the region of Kiel, Germany (N = 500) and blood donors from the University-Clinic Schleswig-Holstein, Kiel, Germany (N = 500) collected by the biobank popgen (Krawczak et al., 2006). The inclusion criteria for the 382 German/Austrian cases of the replication panel (174 males, 208 females) were ≥ 7 teeth with 30% bone loss under the age of 35. The parents and grandparents of the German cases were born in Germany. The parents and grandparents of the Austrian cases (N=69) were born in Austria with German family names. These cases were described in (Schaefer et al., 2013). The controls of the German replication sample were 489 population representative controls from South-Germany (246 males, 243 females), provided by the Cooperative Health Research in the Region of Augsburg Study (KORA), Bavaria, Germany (Wichmann et al., 2005). Population representative controls and blood donors were generally regarded to be free of AgP, as the prevalence of AgP is very low with an estimated occurrence of $<0.1\%$.

The study was approved by each institute's own ethical review board and all participants provided written informed consent.

DNA Isolation and Genotyping

Genomic DNA was extracted from frozen blood samples. All DNA samples were quality controlled on agarose gels. Genotyping of the GWAS was performed with the Affymetrix Gene Chip Human Mapping 500K Array Set for patients and controls (Schaefer et al., 2010). Genotypes were assigned using the BRLMM-p algorithm. In the replication, SNP rs198712 was genotyped with the TaqMan Assay hCV9946741 (Applied Biosystems, Foster City, USA) on an automated platform, employing TECAN Freedom EVO and 96-well and 384-well TEMO liquid handling robots (TECAN, Männedorf, Switzerland).

Statistical Analysis

SNPs with a genotype call rate <90% and a MAF >5% were excluded from the study. Markers were tested for deviations from Hardy-Weinberg equilibrium in controls before inclusion into the analyses ($\alpha = 0.05$). From a total of 500,568 SNPs, 65% passed the quality criteria (287,224 SNPs). Possible statistical interactions between remaining SNPs and sex on AgP were assessed by logistic regression analysis at a significance threshold of $p < 0.05$. The number of alleles, specified as a continuous influence factor (additive model) and sex were included in the model equation as main effects. From these models, we report p values and sex specific odds ratios (ORs) that were calculated for the explorative GWAS data, for the replication data and for both data pooled. Model-based multifactor dimensionality reduction (Calle et al., 2010) was performed using MB-MDR (version 4.1) for a binary phenotype, fixing sex as interaction partner for all individual SNPs. Statistical analyses were conducted

using PLINK v2.049 (Purcell et al. 2007) and the R software package, version 2.12.2 (<http://www.r-project.org>).

***In-silico* Analysis of Chromatin State Segmentation**

Chromatin state segmentation of human cell lines were produced by ChIP-seq data (Chromatin In-situ Precipitation) generated by Broad Institute, (Bernstein laboratory, Massachusetts General Hospital/Harvard Medical School) and produced in Manolis Kellis's Computational Biology group at the Massachusetts Institute of Technology. Chromatin states were learned from this binarized data using a multivariate Hidden Markov Model (HMM) as previously described (Ernst et al., 2011). Data were publicly available by UCSC Genome Browser and were generated by the ENCODE (Encyclopedia of DNA Elements) consortium.

Results

Genome-wide Exploration of Sex-specific Effects

All SNPs that were genotyped on the Affymetrix 500K Array and passed the quality criteria (287,224 SNPs) were analysed in a logistic regression analysis. 2,041 SNPs had a $p < 0.05$ in the main effect of sex, SNP or the gene-sex interaction term. The lowest p-values were observed for an intergenic region on chromosome 7 (chr.7:24143041-24149638; NCBI build 36/hg18), spanning 6.597 kilobasepairs (kb; **Supplementary Table 1**). This region is located 140 kb upstream the gene neuropeptide Y (*NPY*) (chr7:24,290,334-24,298,002) and was covered by eleven successfully genotyped SNPs with an average distance of 0.44kb (± 0.43 kb). Of these SNPs, ten showed a p value in the gene-sex interaction term less than 5×10^{-5} (**Table 1**) and were in high linkage disequilibrium ([LD] $r^2 > 0.85$; **Figure 1a**). The lead SNP was rs198712 with a $p = 5.4 \times 10^{-6}$ for the interaction and $p = 9.8 \times 10^{-6}$ and $p = 0.0240$

Accepted Article

for SNP and sex, respectively (**Table 1**). The sex-specific ORs for the additive effect were 1.629 for males and 0.689 for females. Corroborating this finding, this SNP showed the strongest interaction effect with sex in the genome-wide MB-MDR analysis. The MAF of the male cases and controls was 48% and 36%, and of female cases and controls 30% and 39%, respectively. The MAF in the controls (males and females) as well as in the cases (males and females) was MAF=37% (genotypes are shown in **Table 2**). Accordingly, in the ordinary association tests that considered the cases and controls without taking sex into account, these SNPs showed no significant effect upon AgP.

Replication of the Association on Chromosome 7 Upstream *NPY*.

SNP rs198712 was selected for replication in an additional sample of 382 German and Austrian AgP cases and of 489 German controls. In the replication, the p value for interaction was p=0.0222, and the p values for the SNP and sex were p=0.024 and p=0.203, respectively (**Table 1**). The sex-specific ORs were OR=1.304 for males and OR=0.832 for females, and the MAF was comparable to the values in the initial sample (43% in male cases, 37% in male controls, 38% in female cases, 48% in female controls). After pooling of the explorative and the replication sample, the interaction p value was p=4.03x10⁻⁶.

***In silico* Analysis of the Chromatin State of the *NPY* Region**

To gain insight into the nature of the associated chromosomal region, we analyzed the annotation of chromatin elements of different human cell types from ENCODE data (Hoffman et al., 2013, Yang et al., 2010). The

The associated chromosomal region that was tagged by rs198712 showed tissue specific transcription and possessed a poised promoter (**Figure 1a**). Further 54 kb downstream within this intergenic region, a genome-wide association with childhood asthma was reported for SNP rs886448 (Ding et al., 2013). This SNP was not in LD with the AgP associated region ($r^2=0$, $D'=0.58$), but in accordance to this region, tissue specific transcription was shown for the chromosomal position tagged by the GWAS lead SNP for childhood asthma (**Figure 1a**). Likewise, at the intergenic region downstream of *NPY* that was reported to be strongly associated with severe CP in a GWAS (rs2521634) and not in LD with the AgP associated region ($r^2=0$, $D'=0.13$) (Divaris et al., 2013), tissue specific transcription and poised promoters were shown, as was observed for the AgP associated region (**Figure 1b**). The CP associated region was also described to confer risk of early-onset atherosclerosis in the Framingham SHARe data, i.e. SNP rs10487606, 40bp upstream of rs2521634, was associated with increased coronary artery calcium (Shah et al., 2009).

Discussion

In the current study, we investigated interactions of SNP genotypes with male and female sex with respect to the risk for aggressive periodontitis on a genome-wide scale. Our main finding was an associated intergenic region 140 kb upstream of the gene *NPY* that conferred an increased risk for AgP in men, but a decreased risk in women.

Following the initial observation of the association of this chromosomal region with AgP, we hypothesized that a genetic region of strong linkage disequilibrium ($r^2>0.85$) upstream *NPY* shows gene-sex interaction. This hypothesis was subsequently verified in a second independent sample. Because all ten SNPs that showed the strongest associations in the explorative GWAS analysis were linked on the same haplotype block, rs198712 was

Accepted Article

replicated as a single tagging SNP. Thus, no correction for multiple testing was required in the replication. In the replication, the association was weaker compared with that observed in the explorative GWAS sample. This could be due to the less severe phenotype of the cases in the replication, with >30% alveolar bone loss at ≥ 2 teeth compared to > 50% bone loss in the cases of the GWAS sample at the same age of disease onset, which can result in a decreased statistical power. Likewise, and corresponding to the higher severity of the cases in the GWAS sample compared with the cases of the replication, the male cases of the GWAS sample had a MAF= 48% (MAF= 36% male controls) compared to a MAF= 43% in the male case group of the replication (MAF= 37% male controls). Alternatively or in addition, the true genetic effect size could be overestimated in the explorative sample as a consequence of the winner's curse phenomenon (Lohmueller et al., 2003).

The associated SNPs are located in a chromosomal region, which is tissue specifically transcribed. Interestingly, a recent large GWAS on severe CP described strong association downstream from *NPY* as a main finding (Divaris et al., 2013). Another study that relates *NPY* to CP established the presence of NPY Y1 receptors in the gingival tissue and of NPY protein in human gingival crevicular fluid (GCF), an immunologically relevant exudate of the gingival crevice. This study showed that in the GCF from healthy sites, significantly higher NPY levels were observed compared with periodontitis-affected sites (Lundy et al., 2009). Interestingly, NPY is also the most abundant neuropeptide in bone (Ahmed et al., 1994) and has recently been shown to have a role in maintaining the balance between hard tissue formation and resorption, processes that are relevant to the definition of periodontitis (Haug & Heyeraas, 2006). The immunomodulatory effects of NPY are thought to alter the pro-inflammatory T-helper type 1 (Th1):anti-inflammatory T-helper type 2 (Th2) balance and binding of NPY to Y1 receptors on a variety of immune cells is thought to be responsible for

Accepted Article

promoting the anti-inflammatory Th2 response. NPY is therefore potentially important in the coordination of inflammation and bone metabolism, both of which are central to the pathogenesis of PD (Lundy et al., 2009).

To our knowledge, the current study is the first to observe a sexually dimorphic role of *NPY* in humans that is associated with a complex disease. Interestingly, sex-dependent effects of NPY were previously described in mice. NPY loss-of-function mice showed different anxiogenic responses in behavioral tests in males and females, indicating a sexually dimorphic role of NPY in stress response (Painsipp et al., 2011). Also, gastrointestinal inflammation, known to enhance anxiety in a sex-dependent manner, produced different behavioral responses to stress challenges in female and male NPY knockout mice (Painsipp et al., 2011). Another study on NPY knockout mice showed sex-dependent responses in food intake, upper gastrointestinal transit and faecal pellet output induced by restrained and novel environment stresses (Forbes et al., 2012).

NPY activates the hypothalamic–pituitary–adrenal (HPA) axis and modulates the visceral stress responses. Additionally, NPY is potently anxiolytic (Karl et al., 2008). In accordance with the function in mice, NPY influences many physiological processes in humans, including stress response (Zhou et al., 2008) and stress-induced obesity (Kuo et al., 2007). Often, stress-associated and eating-disorders, which showed sex-dependent effects in NPY knockout mice, also have a different prevalence among women and men in humans (Kessler et al., 1995, Laughlin et al., 2000). A nonsynonymous SNP (Leu/Pro transition) within *NPY* was associated with serum triglyceride concentrations and birth weight, high serum cholesterol and LDL cholesterol levels (Karvonen et al., 1998), as well as alcohol dependence population samples from the United States (Lappalainen et al., 2002) and Finland (Kauhanen

et al., 2000). Stress, obesity and alcohol consumption are considered as risk factors of periodontitis.

In summary, the described findings on the function of NPY add confidence in the validity of our results.

There is a well-known limitation of the logistic regression model in general (Witte & Greenland, 1997) as it can describe only multiplicative effects of odds ratios. In the present study, it was observed that a specific factor, i.e. sex, masked the genetic effect in a way that the genetic influence could be observed only in sex stratified analyses. Without taking gene-sex interactions into account, this association would have been missed. This showed that adding an interaction term is a first step towards a more comprehensive model attempting to describe more precisely the interplay of environmental and genetic factors more precisely.

In conclusion, we identified and replicated the NPY region as conferring an increased risk of AgP in men and a decreased risk in women. Additional studies are warranted to explore the molecular mechanism behind the observed sex-specific effect of genetic variation at *NPY* on the risk of PD.

References

- Ahmed, M., Srinivasan, G. R., Theodorsson, E., Bjurholm, A. & Kreicbergs, A. (1994) Extraction and quantitation of neuropeptides in bone by radioimmunoassay. *Regul Pept* **51**, 179-188.
- Buchwald, S., Kocher, T., Biffar, R., Harb, A., Holtfreter, B. & Meisel, P. (2013) Tooth loss and periodontitis by socio-economic status and inflammation in a longitudinal population-based study. *J Clin Periodontol* **40**, 203-211. doi:10.1111/jcpe.12056.
- Calle, M. L., Urrea, V., Malats, N. & Van Steen, K. (2010) mbmdr: an R package for exploring gene-gene interactions associated with binary or quantitative traits. *Bioinformatics* **26**, 2198-2199. doi:btq352 [pii]

- 10.1093/bioinformatics/btq352.
- Corey, L. A., Nance, W. E., Hofstede, P. & Schenkein, H. A. (1993) Self-reported periodontal disease in a Virginia twin population. *J Periodontol* **64**, 1205-1208.
- Ding, L., Abebe, T., Beyene, J., Wilke, R. A., Goldberg, A., Woo, J. G., Martin, L. J., Rothenberg, M. E., Rao, M., Hershey, G. K., Chakraborty, R. & Mersha, T. B. (2013) Rank-based genome-wide analysis reveals the association of ryanodine receptor-2 gene variants with childhood asthma among human populations. *Hum Genomics* **7**, 16. doi:1479-7364-7-16 [pii]
- 10.1186/1479-7364-7-16.
- Divaris, K., Monda, K. L., North, K. E., Olshan, A. F., Reynolds, L. M., Hsueh, W. C., Lange, E. M., Moss, K., Barros, S. P., Weyant, R. J., Liu, Y., Newman, A. B., Beck, J. D. & Offenbacher, S. (2013) Exploring the genetic basis of chronic periodontitis: a genome-wide association study. *Hum Mol Genet* **22**, 2312-2324. doi:ddt065 [pii]
- 10.1093/hmg/ddt065.
- Eke, P. I., Dye, B. A., Wei, L., Thornton-Evans, G. O. & Genco, R. J. (2012) Prevalence of periodontitis in adults in the United States: 2009 and 2010. *J Dent Res* **91**, 914-920. doi:0022034512457373 [pii]
- 10.1177/0022034512457373.
- Ernst, J., Kheradpour, P., Mikkelsen, T. S., Shores, N., Ward, L. D., Epstein, C. B., Zhang, X., Wang, L., Issner, R., Coyne, M., Ku, M., Durham, T., Kellis, M. & Bernstein, B. E. (2011) Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature* **473**, 43-49. doi:nature09906 [pii]
- 10.1038/nature09906.
- Forbes, S., Herzog, H. & Cox, H. M. (2012) A role for neuropeptide Y in the gender-specific gastrointestinal, corticosterone and feeding responses to stress. *Br J Pharmacol* **166**, 2307-2316. doi:10.1111/j.1476-5381.2012.01939.x.
- Hart, T. C., Marazita, M. L., Schenkein, H. A., Brooks, C. N., Gunsolley, J. G. & Diehl, S. R. (1991) No female preponderance in juvenile periodontitis after correction for ascertainment bias. *J Periodontol* **62**, 745-749. doi:10.1902/jop.1991.62.12.745.
- Haug, S. R. & Heyeraas, K. J. (2006) Modulation of dental inflammation by the sympathetic nervous system. *J Dent Res* **85**, 488-495. doi:85/6/488 [pii].
- Hoffman, M. M., Ernst, J., Wilder, S. P., Kundaje, A., Harris, R. S., Libbrecht, M., Giardine, B., Ellenbogen, P. M., Bilmes, J. A., Birney, E., Hardison, R. C., Dunham, I., Kellis, M. & Noble, W. S. (2013) Integrative annotation of chromatin elements from ENCODE data. *Nucleic Acids Res* **41**, 827-841. doi:gks1284 [pii]
- 10.1093/nar/gks1284.
- Karl, T., Duffy, L. & Herzog, H. (2008) Behavioural profile of a new mouse model for NPY deficiency. *Eur J Neurosci* **28**, 173-180. doi:EJN6306 [pii]
- 10.1111/j.1460-9568.2008.06306.x.
- Karvonen, M. K., Pesonen, U., Koulu, M., Niskanen, L., Laakso, M., Rissanen, A., Dekker, J. M., Hart, L. M., Valve, R. & Uusitupa, M. I. (1998) Association of a leucine(7)-to-proline(7) polymorphism in the signal peptide of neuropeptide Y with high serum cholesterol and LDL cholesterol levels. *Nat Med* **4**, 1434-1437. doi:10.1038/4027.
- Kauhanen, J., Karvonen, M. K., Pesonen, U., Koulu, M., Tuomainen, T. P., Uusitupa, M. I. & Salonen, J. T. (2000) Neuropeptide Y polymorphism and alcohol consumption in middle-aged men. *Am J Med Genet* **93**, 117-121. doi:10.1002/1096-8628(20000717)93:2<117::AID-AJMG7>3.0.CO;2-# [pii].
- Kessler, R. C., Sonnega, A., Bromet, E., Hughes, M. & Nelson, C. B. (1995) Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* **52**, 1048-1060.
- Krawczak, M., Nikolaus, S., von Eberstein, H., Croucher, P. J., El Mokhtari, N. E. & Schreiber, S. (2006) PopGen: population-based recruitment of patients and controls for the analysis of complex genotype-phenotype relationships. *Community Genet* **9**, 55-61.

- Kuo, L. E., Kitlinska, J. B., Tilan, J. U., Li, L., Baker, S. B., Johnson, M. D., Lee, E. W., Burnett, M. S., Fricke, S. T., Kvetnansky, R., Herzog, H. & Zukowska, Z. (2007) Neuropeptide Y acts directly in the periphery on fat tissue and mediates stress-induced obesity and metabolic syndrome. *Nat Med* **13**, 803-811. doi:nm1611 [pii]
10.1038/nm1611.
- Lappalainen, J., Kranzler, H. R., Malison, R., Price, L. H., Van Dyck, C., Rosenheck, R. A., Cramer, J., Southwick, S., Charney, D., Krystal, J. & Gelernter, J. (2002) A functional neuropeptide Y Leu7Pro polymorphism associated with alcohol dependence in a large population sample from the United States. *Arch Gen Psychiatry* **59**, 825-831. doi:yoa10023 [pii].
- Laughlin, G. A., Barrett-Connor, E., Kritz-Silverstein, D. & von Muhlen, D. (2000) Hysterectomy, oophorectomy, and endogenous sex hormone levels in older women: the Rancho Bernardo Study. *J Clin Endocrinol Metab* **85**, 645-651. doi:10.1210/jcem.85.2.6405.
- Lohmueller, K. E., Pearce, C. L., Pike, M., Lander, E. S. & Hirschhorn, J. N. (2003) Meta-analysis of genetic association studies supports a contribution of common variants to susceptibility to common disease. *Nat Genet* **33**, 177-182. doi:10.1038/ng1071
ng1071 [pii].
- Lundy, F. T., El Karim, I. A. & Linden, G. J. (2009) Neuropeptide Y (NPY) and NPY Y1 receptor in periodontal health and disease. *Arch Oral Biol* **54**, 258-262. doi:S0003-9969(08)00279-3 [pii]
10.1016/j.archoralbio.2008.10.002.
- Marazita, M. L., Burmeister, J. A., Gunsolley, J. C., Koertge, T. E., Lake, K. & Schenkein, H. A. (1994) Evidence for autosomal dominant inheritance and race-specific heterogeneity in early-onset periodontitis. *J Periodontol* **65**, 623-630.
- Marcenes, W., Kassebaum, N. J., Bernabe, E., Flaxman, A., Naghavi, M., Lopez, A. & Murray, C. J. (2013) Global burden of oral conditions in 1990-2010: a systematic analysis. *J Dent Res* **92**, 592-597. doi:0022034513490168 [pii]
10.1177/0022034513490168.
- McGrath, C. & Bedi, R. (2003) Measuring the impact of oral health on quality of life in Britain using OHQoL-UK(W). *J Public Health Dent* **63**, 73-77.
- Mealey, B. L. & Oates, T. W. (2006) Diabetes mellitus and periodontal diseases. *J Periodontol* **77**, 1289-1303. doi:10.1902/jop.2006.050459.
- Michalowicz, B. S., Aeppli, D., Virag, J. G., Klump, D. G., Hinrichs, J. E., Segal, N. L., Bouchard, T. J., Jr. & Pihlstrom, B. L. (1991) Periodontal findings in adult twins. *J Periodontol* **62**, 293-299.
- Michalowicz, B. S., Diehl, S. R., Gunsolley, J. C., Sparks, B. S., Brooks, C. N., Koertge, T. E., Califano, J. V., Burmeister, J. A. & Schenkein, H. A. (2000) Evidence of a substantial genetic basis for risk of adult periodontitis. *J Periodontol* **71**, 1699-1707.
- Painsipp, E., Herzog, H., Sperk, G. & Holzer, P. (2011) Sex-dependent control of murine emotional-affective behaviour in health and colitis by peptide YY and neuropeptide Y. *Br J Pharmacol* **163**, 1302-1314. doi:10.1111/j.1476-5381.2011.01326.x.
- Sangwan, A., Tewari, S., Singh, H., Sharma, R. K. & Narula, S. C. (2013) Periodontal status and hyperlipidemia: statin users versus non-users. *J Periodontol* **84**, 3-12. doi:10.1902/jop.2012.110756.
- Schaefer, A. S., Bochenek, G., Manke, T., Nothnagel, M., Graetz, C., Thien, A., Jockel-Schneider, Y., Harks, I., Staufienbiel, I., Wijmenga, C., Eberhardt, J., Guzeldemir, E., Cine, N., Folwaczny, M., Noack, B., Meyle, J., Eickholz, P., Trombelli, L., Scapoli, C., Nohutcu, R., Bruckmann, C., Doerfer, C., Jepsen, S., Loos, B. G. & Schreiber, S. (2013) Validation of reported genetic risk factors for periodontitis in a large-scale replication study. *J Clin Periodontol*. doi:10.1111/jcpe.12092.
- Schaefer, A. S., Richter, G. M., Nothnagel, M., Manke, T., Dommisch, H., Jacobs, G., Arlt, A., Rosenstiel, P., Noack, B., Groessner-Schreiber, B., Jepsen, S., Loos, B. G. & Schreiber, S. (2010) A genome-wide association study identifies GLT6D1 as a susceptibility locus for periodontitis. *Hum Mol Genet* **19**, 553-562. doi:ddp508 [pii]

10.1093/hmg/ddp508.

Schenkein, H. A. & Loos, B. G. (2013) Inflammatory mechanisms linking periodontal diseases to cardiovascular diseases. *J Clin Periodontol* **40 Suppl 14**, S51-69. doi:10.1111/jcpe.12060.

Shah, S. H., Freedman, N. J., Zhang, L., Crosslin, D. R., Stone, D. H., Haynes, C., Johnson, J., Nelson, S., Wang, L., Connelly, J. J., Muehlbauer, M., Ginsburg, G. S., Crossman, D. C., Jones, C. J., Vance, J., Sketch, M. H., Granger, C. B., Newgard, C. B., Gregory, S. G., Goldschmidt-Clermont, P. J., Kraus, W. E. & Hauser, E. R. (2009) Neuropeptide Y gene polymorphisms confer risk of early-onset atherosclerosis. *PLoS Genet* **5**, e1000318. doi:10.1371/journal.pgen.1000318.

Shiau, H. J. & Reynolds, M. A. (2010) Sex differences in destructive periodontal disease: exploring the biologic basis. *J Periodontol* **81**, 1505-1517. doi:10.1902/jop.2010.100045.

Wichmann, H. E., Gieger, C. & Illig, T. (2005) KORA-gen--resource for population genetics, controls and a broad spectrum of disease phenotypes. *Gesundheitswesen* **67 Suppl 1**, S26-30. doi:10.1055/s-2005-858226.

Witte, J. S. & Greenland, S. (1997) A nested approach to evaluating dose-response and trend. *Ann Epidemiol* **7**, 188-193. doi:10.1016/S1047279796001597 [pii].

Yang, C., Wan, X., Yang, Q., Xue, H. & Yu, W. (2010) Identifying main effects and epistatic interactions from large-scale SNP data via adaptive group Lasso. *BMC Bioinformatics* **11 Suppl 1**, S18. doi:10.1186/1471-2105-11-S1-S18 [pii]

10.1186/1471-2105-11-S1-S18.

Zhou, Z., Zhu, G., Hariri, A. R., Enoch, M. A., Scott, D., Sinha, R., Virkkunen, M., Mash, D. C., Lipsky, R. H., Hu, X. Z., Hodgkinson, C. A., Xu, K., Buzas, B., Yuan, Q., Shen, P. H., Ferrell, R. E., Manuck, S. B., Brown, S. M., Hauger, R. L., Stohler, C. S., Zubieta, J. K. & Goldman, D. (2008) Genetic variation in human NPY expression affects stress response and emotion. *Nature* **452**, 997-1001. doi:nature06858 [pii]

10.1038/nature06858.

Acknowledgements

We thank the technicians Tanja Wesse and Sanaz Sedghpour Sabet and the information specialists Lukas Tittmann and Michael Wittig.

Table Legends

Table 1. Top 10 results of gene-sex interaction from logistic regression models of GWAS-data and of rs198712 in the replication

Table 2. Genotypes of the ten most significant SNPs of gene-sex interaction from logistic regression models of GWAS-data and in the replication of tag SNP rs108712

Figure legends

Figure 1. Chromosomal region at *NPY*

The top diagram shows the nominal $-\log^{10}$ P for the plotted SNPs at this chromosomal region. The $-\log^{10}$ P-values are plotted as a function of the genomic SNP position (NCBI build 36). The second panel shows the position of SNP rs886448, which showed genome-wide association with childhood asthma. The following panel (a) shows annotations of chromatin elements from ENCODE data for two cell types (human skeletal muscle myotubes [HSMM] and H1 human embryonic stem cells [H1-hESC], TXN=transcription). The lowest panel (b) shows the annotations of chromatin elements across the chromosomal region of *NPY*, shown for the two cell types as in panel a. The included area covers the chromosomal region of panel a and extends to the GWAS lead SNP of chronic periodontitis (rs2521634). Tissue specific transcription (TXN) of the three regions, which are associated with AgP, childhood asthma (rs886448), and chronic periodontitis are shown in light green. The plotted SNPs are from left to right: 1=rs1494194, 2=rs17287854, 3=rs2813838, 4=rs272709, 5=rs17288003, 6=rs17148810, 7=rs272665, 8=rs272662, 9=rs1859289, 10=rs198733, 11=rs198731, 12=rs198728, 13=rs115063, 14=rs198727, 15=rs115062, 16=rs198720, 17=rs198712, 18=rs198711, 19=rs6461782, 20=rs198701, 21=rs17290092, 22=rs156270, 23=rs156318, 24=rs156308, 25=rs156294, 26=rs156293, 27=rs13245518, 28=rs963830, 29=rs2158342.

Table 1. Top 10 results of gene-sex interaction from logistic regression models of GWAS-data and in the replication of tag SNP rs198712

SNPs	p SNP	P sex	OR	95 % CI	p value	cases		controls	
						MAF (males)	MAF (females)	MAF (males)	MAF (females)
rs198733	1.74E-05	0.112	2.17	1.55-3.16	2.02E-05	50.38	34.01	37.11	40.95
rs198731	7.04E-05	0.149	2.07	1.44-2.97	9.02E-05	49.61	34.38	37.25	40.50
rs198728	9.83E-06	0.075	2.21	1.55-3.16	1.26E-05	50.76	34.01	37.11	40.96
rs115063	1.43E-05	0.096	2.17	1.52-3.09	2.06E-05	50.76	34.52	37.23	41.06
rs198727	9.98E-06	0.076	2.21	1.55-3.16	1.27E-05	50.76	34.01	37.13	40.96
rs115062	5.94E-06	0.054	2.27	1.59-3.25	7.09E-06	50.76	33.67	36.91	40.96
rs198720	2.77E-05	0.088	2.18	1.53-3.12	1.80E-05	51.14	35.03	38.90	42.98
rs198712	9.76E-06	0.024	2.36	1.63-3.42	5.41E-06	48.09	29.89	35.73	38.58
rs198711	1.64E-05	0.071	2.20	1.54-3.15	1.64E-05	49.24	32.23	36.35	39.47
rs198701	8.32E-06	0.042	2.27	1.58-3.25	9.45E-06	49.24	32.12	35.83	39.42
Replication:									
rs198712	0.024	0.2033	1.5	1.07-2.3	0.0222	43.39	38.22	36.99	42.80

Table 2. Genotypes of the ten most significant SNPs of gene-sex interaction of GWAS-data and in the replication of tag SNP rs198712

SNPs	Cases						controls					
	males			females			males			females		
	11	12	22	11	12	22	11	12	22	11	12	22
rs198733	35	59	36	90	80	27	212	220	80	170	208	86
rs198731	33	63	32	87	78	27	205	220	77	170	211	82
rs198728	34	62	36	90	80	27	212	220	80	172	211	87
rs115063	34	62	36	89	80	28	211	222	80	172	210	88
rs198727	34	62	36	90	80	27	212	221	80	172	211	87
rs115062	34	62	36	90	80	26	212	222	78	172	211	87
rs198720	35	59	38	85	86	26	199	224	86	158	220	92
rs198712	36	64	31	92	81	16	216	221	71	182	206	76
rs198711	36	62	34	92	83	22	216	221	76	182	205	83
rs198701	37	60	35	91	80	22	215	222	71	181	205	82
Replication rs198712	55	87	32	81	95	32	99	112	35	82	114	47

GWAS

Replication

