

Short Communications

N₂-Fixation by Chemoautotrophic Hydrogen Bacteria

J. N. Gogotov and H. G. Schlegel

Institut für Mikrobiologie der Universität Göttingen and Institut für Mikrobiologie der Gesellschaft für Strahlen- und Umweltforschung mbH, München, in Göttingen

Received March 20, 1974

Abstract. The Gram-positive coryneform bacteria strains *14g* and *7C* were found to be able to grow with N₂ as sole nitrogen source when incubated under microaerobic conditions. Nitrogenase activity in whole cells was assayed by acetylene reduction. High rates of ethylene production (50–120 nmole/h × mg cell protein) were observed in N₂ or glutamate grown cell suspensions shaken in an atmosphere of 2.5% O₂, 10% acetylene and 87.5% argon.

Key words: Hydrogen-Bacteria — Nitrogen Fixation — Acetylene Reduction — Coryneform Bacteria Strains *14g* and *7C*.

None of the characterized chemolithoautotrophic bacteria has been previously reported to be able to fix nitrogen. Some evidence has been obtained for the utilization of molecular nitrogen as the sole nitrogen source by a bacterium growing under an atmosphere of nitrogen, hydrogen, oxygen and carbon dioxide (Ooyama, 1971). But the bacterium, which was isolated from an oily soil in northern Japan, was not identified.

This report describes a study of acetylene reduction in the following strains of hydrogen oxidizing bacteria: *Hydrogenomonas eutropha* H 16 (Wilde, 1962), *Nocardia opaca* 1b (Aggag and Schlegel, 1973) and the coryneform bacteria strain *7C* (Siebert, 1969; Tunail and Schlegel, 1972; Tunail, 1973) and strain *14g* (Schneider *et al.*, 1973). Only the coryneform bacteria were capable of nitrogen fixation.

Cells were grown in the mineral medium of Schlegel *et al.* (1961) modified by the omission of NH₄Cl and containing trace elements (1 ml/l; Ormerod *et al.*, 1961) in which the ferrous-ammonium citrate was replaced by ferrous citrate. The pH was adjusted to 7.0–7.2. The mineral medium was supplemented with the following sources of nitrogen, oxygen, hydrogen and carbon (the associated figures are % v/v, gas phase, for gases and % w/v, liquid phase, for solids): A) NH₄Cl, 0.1, + O₂, 5–10, + H₂, 85–80, + CO₂, 10; B) N₂, 60, + O₂, 10, + H₂, 20, + CO₂, 10; C) NH₄Cl, 0.1, + N₂, 85, + O₂, 10, + CO₂, 5, + succinate 0.3; D) Gluta-

mate, 0.05, +N₂,85, +O₂,10, +CO₂,5, +succinate, 0.3; E) N₂,85, +O₂,10, +CO₂,5, +succinate, 0.3; F) N₂,85, +O₂,10, +CO₂,5, +fructose, 0.3.

Cultures (1 or 3 l) were grown in round, flat-bottomed flasks (2 or 6 l respectively) at 30° C and gassed by a magnetically driven, plastic coated stirring bar (7 cm, 400 rpm; Schlegel *et al.*, 1961). The appropriate gas mixture was supplied from containers (10 l) connected to the growth flask, and the gas mixture over the culture was flushed and renewed every 6–10 h. Growth was measured turbidimetrically at 436 nm (*Hydrogenomonas eutropha* H 16 and *Nocardia opaca* 1b) or 546 nm (strains 14g and 7C).

The rate of acetylene reduction by resting cells was used as a measure of nitrogenase activity (Dilworth, 1966). Assays (2 ml final volume) were done in Warburg vessels (18 ml) at 30° C using conventional Warburg apparatus (Type P 166, Braun, Melsungen, BRD) with reciprocal shaking (120 strokes/min, amplitude 4 cm). Cell suspension (0.5 ml, 9 mg cell protein) was added to the main vessel, which also contained 0.067 M phosphate buffer pH 7.2; substrate (20 μ mole) was added to the side arm. The Warburg vessels were then evacuated and filled with a gas mixture containing O₂ (2.5–10%), acetylene (10%) and argon (87.5–80%); when hydrogen was present, the argon concentration was correspondingly reduced. At intervals after tipping the Warburg vessels, gas samples (0.2 ml) were removed by syringe for analysis by gas chromatography and replaced by argon (0.2 ml). The gas samples were injected into a Perkin Elmer F 11 gas chromatograph filled with a 45 cm Porapak R column and coupled to a hydrogen flame ionization detector.

Linear rates of ethylene production were obtained when rates did not exceed 700 nmole ethylene/h $\times$ mg cell protein. The protein content of intact cells was measured using the modified Biuret method of Schmidt *et al.* (1963).

When *Hydrogenomonas eutropha* H 16, which had been grown in media containing combined nitrogen, was inoculated into the supplemented mineral media without a source of combined nitrogen (supplements B or D), a limited increase in turbidity was observed (two- to threefold). No further turbidity increase was observed and further subcultures in media without combined nitrogen did not grow. Under no condition was acetylene reduction detected. This absence of nitrogenase was also observed in *Nocardia opaca* 1b.

In contrast, both strains of coryneform hydrogen bacteria grew rather fast under the microaerobic conditions used. With succinate as a substrate, the turbidity increased 30-fold with media supplements D and E.

Both strains *14g* and *7C* have been subcultured several times in media free from combined nitrogen. The cells exhibited high nitrogenase activities.

Using strains *14g* and *7C* the gas phase for optimal acetylene reduction was studied. The rate of ethylene production increased with increasing acetylene concentrations; 10% acetylene was saturating. Oxygen, the hydrogen acceptor for aerobic respiration and energy generation, inhibited acetylene reduction. With the relatively high cell density used, the maximum rates of acetylene reduction were measured at oxygen concentrations of 10% (strain *7C*) or 2.5% (strain *14g*); 40% and 20% O₂, respectively, gave approximately 95% inhibition.

The highest rates of acetylene reduction (120 nmole ethylene/h × mg cell protein) were measured with cells grown in the presence of glutamate or with N₂ as the sole nitrogen source (media supplements D, E, F). Nitrogenase activity was not detected in cells grown in the media containing ammonia (supplements A and C). Pyruvate, succinate or malate served as substrates for acetylene reduction in the presence of oxygen. Only pyruvate gave rise to a modest rate of acetylene reduction in the absence of oxygen (about 5% of the rate observed under aerobic conditions). Hydrogen supported acetylene reduction under micro-aerobic conditions, but only at low partial pressures (2.5% for *7C* and 5% H₂ for *14g*).

References

- Aggag, M., Schlegel, H. G.: Studies on a Gram-positive hydrogen bacterium *Nocardia opaca* 1b. I. Description and physiological characterization. Arch. Mikrobiol. **88**, 299–318 (1973)
- Dilworth, M. J.: Acetylene reduction by nitrogen-fixing preparations from *Clostridium pasteurianum*. Biochim. biophys. Acta (Amst.) **127**, 285–294 (1966)
- Ooyama, J.: Simultaneous fixation of CO₂ and N₂ in the presence of H₂ and O₂ by a bacterium. Rep. Ferment. Res. Inst. **39**, 41–44 (1971)
- Ormerod, J. G., Ormerod, K. S., Gest, H.: Light dependent utilization of organic compounds and photoproduction of molecular hydrogen by photosynthetic bacteria; relationships with nitrogen metabolism. Arch. Biochem. Biophys. **94**, 449–463 (1961)
- Schlegel, H. G., Kaltwasser, H., Gottschalk, G.: Ein Submersverfahren zur Kultur wasserstoffoxydierender Bakterien: Wachstumsphysiologische Untersuchungen. Arch. Mikrobiol. **38**, 209–222 (1961)
- Schmidt, K., Liaaen Jensen, S., Schlegel, H. G.: Die Carotinoide der Thiorhodaceae. I. Okenon als Hauptcarotinoid von *Chromatium okenii* Perty. Arch. Mikrobiol. **46**, 117–126 (1963)
- Schneider, K., Rudolph, V., Schlegel, H. G.: Description and physiological characterization of a coryneform hydrogen bacterium, strain *14g*. Arch. Mikrobiol. **93**, 179–193 (1973)
- Siebert, D.: Über propanverwertende, wasserstoffoxidierende Bakterien und die Charakterisierung eines Förderungsfaktors. Diss., Univ. Göttingen 1969

- Tunail, N.: Charakterisierung des Wasserstoffbacteriums *Arthrobacter 7C* und der Regulation des Kohlenhydratstoffwechsels. Diss., Univ. Göttingen 1973
- Tunail, N., Schlegel, H. G.: Phosphoenolpyruvate, a new inhibitor of glucose-6-phosphate dehydrogenase. *Biochem. biophys. Res. Commun.* **49**, 1554—1560 (1972)
- Wilde, E.: Untersuchungen über Wachstum und Speicherstoffsynthese von *Hydrogenomonas*. *Arch. Mikrobiol.* **43**, 109—137 (1962)

Prof. Dr. H. G. Schlegel
Institut für Mikrobiologie der Universität
D-3400 Göttingen, Grisebachstr. 8
Federal Republic of Germany