Polymorphism rs11085226 in the Gene Encoding Polypyrimidine Tract-Binding Protein 1 Negatively Affects Glucose-Stimulated Insulin Secretion

Martin Heni¹, Caroline Ketterer¹, Robert Wagner¹, Katarzyna Linder¹, Anja Böhm¹, Silke A. Herzberg-Schäfer¹, Fausto Machicao^{1,2}, Klaus-Peter Knoch⁴, Andreas Fritsche^{1,2,3}, Harald Staiger^{1,2}, Hans-Ulrich Häring^{1,2*}, Michele Solimena⁴

1 Division of Endocrinology, Diabetology, Angiology, Nephrology and Clinical Chemistry, Department of Internal Medicine, Eberhard Karls University Tübingen, Tübingen, Germany, 2 Institute for Diabetes Research and Metabolic Diseases of the Helmholtz Center Munich at the University of Tübingen, Partner in the German Center for Diabetes Research (DZD), Tübingen, Germany, 3 Department of Internal Medicine, Nutritional and Preventive Medicine, Eberhard Karls University Tübingen, Tübingen, Tübingen, Germany, 4 Molecular Diabetology, Paul Langerhans Institute Dresden, University Hospital and Medical School Carl Gustav Carus, Dresden University of Technology, Partner in the German Center for Diabetes Research (DZD), Dresden, Germany

Abstract

Objective: Polypyrimidine tract-binding protein 1 (PTBP1) promotes stability and translation of mRNAs coding for insulin secretion granule proteins and thereby plays a role in β -cells function. We studied whether common genetic variations within the *PTBP1* locus influence insulin secretion, and/or proinsulin conversion.

Methods: We genotyped 1,502 healthy German subjects for four tagging single nucleotide polymorphisms (SNPs) within the *PTBP1* locus (rs351974, rs11085226, rs736926, and rs123698) covering 100% of genetic variation with an $r^2 \ge 0.8$. The subjects were metabolically characterized by an oral glucose tolerance test with insulin, proinsulin, and C-peptide measurements. A subgroup of 320 subjects also underwent an IVGTT.

Results: PTBP1 SNP rs11085226 was nominally associated with lower insulinogenic index and lower cleared insulin response in the OGTT ($p \le 0.04$). The other tested SNPs did not show any association with the analyzed OGTT-derived secretion parameters. In the IVGTT subgroup, SNP rs11085226 was accordingly associated with lower insulin levels within the first ten minutes following glucose injection (p = 0.0103). Furthermore, SNP rs351974 was associated with insulin levels in the IVGTT (p = 0.0108). Upon interrogation of MAGIC HOMA-B data, our rs11085226 result was replicated (MAGIC p = 0.018), but the rs351974 was not.

Conclusions: We conclude that common genetic variation in *PTBP1* influences glucose-stimulated insulin secretion. This underlines the importance of PTBP1 for beta cell function *in vivo*.

Citation: Heni M, Ketterer C, Wagner R, Linder K, Böhm A, et al. (2012) Polymorphism rs11085226 in the Gene Encoding Polypyrimidine Tract-Binding Protein 1 Negatively Affects Glucose-Stimulated Insulin Secretion. PLoS ONE 7(10): e46154. doi:10.1371/journal.pone.0046154

Editor: Kathrin Maedler, University of Bremen, Germany

Received June 27, 2012; Accepted August 28, 2012; Published October 15, 2012

Copyright: © 2012 Heni et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The study was supported by a grant from the German Research Foundation to A.F. (DFG FR1561/5-1), by the Kompetenznetz Diabetes mellitus (Competence Network for Diabetes mellitus) funded by the Federal Ministry of Education and Research (FKZ 01G10803-04) (H.-U.H.), and by the BMBF (German Federal Ministry of Education of Beta Cells and Diabetes, 01G10810) (M.S.). Furthermore, the study was supported by a grant from the German Federal Ministry of Education and Research (BMBF) to the German Center for Diabetes Research (DZD e.V.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: hans-ulrich.haering@med.uni-tuebingen.de

Introduction

Glucose stimulation of pancreatic β -cells induces the extracellular release of insulin by inducing the fusion of insulin secretory granules with the plasma membrane [1]. This process is additionally enhanced by incretins (e.g. glucagon-like peptide 1 [GLP-1]) via the second messenger cyclic adenosine monophosphate (cAMP) [2]. In addition to insulin release, glucose rapidly up-regulates the production of new insulin secretory granules. This is due to the enhanced synthesis of most insulin secretory granule components [3–5], like pre-proinsulin, the prohormoneconvertasesPCSK1 and PCSK2 [5–7], chromogranins [8] and the islet cell antigen (ICA) 512/IA-2 [9]. This rapid induction of insulin secretory granule biogenesis is mainly caused by post-transcriptional mechanisms [7,8,10–12].

Polypyrimidine tract-binding protein 1 (PTBP1), formerly known as heterogeneous nuclear ribonucleoprotein I (hnRNP I), enhances mRNA stability and translation, in addition to regulating other processes [13]. For instance, PTBP1 mediates splicing repression in a long list of pre-mRNAs and is thereby important for the regulation of alternative splicing in a tissue-specific manner [14,15]. In pancreatic β -cells and chromaffin cells, protein kinase A-induced phosphorylation of PTBP1 promotes its rapid shift from the nucleus into the cytoplasm [13,16,17]. Glucose stimulation of β -cells induces a similar nucleocytoplasmic translocation of PTBP1 [4,13,16]. By interacting with binding sites within the 3'- and 5'-UTRs (untranslated regions), cytosolic PTBP1, in turn, regulates mRNA localization, stability and the initiation of translation [14,18–20]. Specifically, PTBP1 stabilizes the mRNA coding for insulin and other major insulin secretory granule components such as PCSK1, PCSK2, chromogranin A and ICA512 [4,13,16]. Accordingly, knockdown of PTBP1 in insulinoma INS-1 cells leads to the depletion of insulin secretory granule stores [4].

Impaired insulin secretion together with insulin resistance is a key factor in the pathogenesis of type 2 diabetes mellitus. Various mechanisms could be responsible for impairment of insulin secretion, including a decrease in the number of pancreatic β cells due to apoptosis as well as impaired exocytosis or biogenesis of insulin secretory granules [21,22]. Besides insulin transcription itself, the conversion of proinsulin into insulin and the cleavage of its C-terminal residues by carboxypeptidase E (CPE) are critical for insulin maturation [5,7,23,24]. An increased proinsulin/insulin ratio is a characteristic finding at the onset of type 2 diabetes [25,26]. Notably, PTBP1 promotes the translation of PCSK1 and PCSK2, but not those of CPE, whose translation is not regulated by glucose [4,8,13].

In view of these considerations, genetic variation in *PTBP1* might affect insulin secretion or maturation in humans. To test this hypothesis, we examined the association of common genetic variants (minor allele frequency [MAF]>5%) in *PTBP1* with both insulin secretion and proinsulin-to-insulin conversion.

Methods

Participants

The participants were selected from the ongoing Tübingen Family Study, which currently includes ~2,000 individuals at increased risk of diabetes [27]. Individuals on medication affecting glucose metabolism were excluded. Inclusion of the participants in the present study was based on availability of: (i) DNA samples for genotyping (n = 1,750); and (ii) complete OGTT data (glucose, insulin, C-peptide and proinsulin levels available for all time points during the OGTT, n = 1,502).

The anthropometric characteristics of the study population are shown in Table 1. Most (69%) of these subjects had a family history of diabetes, i.e. at least one second-degree relative with type 2 diabetes mellitus. All participants were genotyped for the following SNPs: *PTBP1* rs351974, rs11085226, rs736926, and rs123698. Informed written consent was obtained from all participants and the local Ethics Committee (University of Tübingen) approved the protocol.

Selection of tagging SNPs and genotyping

Using the publically available phase II data of the International HapMap Project derived from a population of Utah residents with ancestry from Northern and Western Europe (release 24, November 2008, http://www.hapmap.org, [28]), we screened *in silico* the complete *PTBP1* gene (14.92 kb, 15 exons, 14 introns, located on human chromosome 19p13.3) and 5 kb of both the 3' and 5' flanking regions (Figure 1).

Twelve informative SNPs with minor allele frequency (MAF) > 5% are located in this region. Their HapMap linkage disequilibrium data (D' and r-squared values) are shown in Figure 1. Among these, we selected four tagging SNPs that cover 100% of the common genetic variation (MAF>5%) within this locus for genotyping: *PTBP1* rs351974 (A/C) is located in the 5' flanking region of the gene, rs11085226 (A/G) and rs736926 (C/

Table 1. Clinical characteristics of the study population.

| Gender (female/male) | 997/505 |
|----------------------------|------------------|
| NGT/IFG/IGT/(IFG+IGT) | 1095/151/146/110 |
| Age (y) | 39±13 |
| BMI (kg/m ²) | 28.5±7.8 |
| Waist circumference (cm) | 93±17 |
| Fasting glucose (mM) | 5.09±0.54 |
| Glucose, 120 min OGTT (mM) | 6.26±1.65 |
| Fasting insulin (pM) | 62.4±52.4 |
| Insulin, 30 min OGTT (pM) | 430±460 |

Data are given as means \pm SD. BMI – body mass index; IFG – impaired fasting glucose; IGT – impaired glucose tolerance; NGT – normal glucose tolerance; OGTT – oral glucose tolerance test.

doi:10.1371/journal.pone.0046154.t001

T) are located within intron 2, while rs123698 (C/G) is located in intron 10.

DNA from whole blood was isolated using a commercial DNA isolation kit (NucleoSpin, Macherey& Nagel, Düren, Germany). Genotyping was done using the TaqMan assay (Applied Biosystems, Forster City, CA, USA). The TaqMan genotyping reaction was amplified on a GeneAmp PCR system 7000 and fluorescence was detected on an ABI PRISM 7000 sequence detector (Applied Biosystems). The genotyping success rate for *PTBP1* rs351974 was 99.9%, for rs11085226 was 99.9%, for rs736926 was 99.8% and for rs123698 was 99.4%. Genotypes were verified in 50 randomly selected subjects by bidirectional sequencing, and both methods gave identical results.

Oral glucose tolerance test (OGTT)

After an overnight fast of 10 h, subjects ingested a solution containing 75 g glucose at 08:00 hours. Venous blood samples were obtained at 0, 30, 60, 90, and 120 min and plasma glucose, insulin, C-peptide, and proinsulin concentrations were determined. Insulin sensitivity during the OGTT was estimated according to Matsuda and DeFronzo [29].

Intravenous glucose tolerance test (IVGTT) and hyperinsulinemic-euglycemic clamp

In a subgroup of informed subjects (n = 320), an IVGTT was performed after a 10-h overnight fast, as described earlier [30]. After baseline samples had been collected (-10, -5, and 0 min), a glucose dose of 0.3 g/kg body weight was given at time point 0. Blood samples for the measurement of plasma glucose and insulin were obtained at 2, 4, 6, 8, and 10 minutes. In these subjects, a hyperinsulinemic-euglycemic clamp for the determination of insulin sensitivity was performed as well [31].

Analytical procedures

Blood glucose was determined using a bedside glucose analyzer (glucose oxidase method; Yellow Springs Instruments, Yellow Springs, OH, USA). Plasma insulin and proinsulin were determined by microparticle enzyme immunoassay (Abbott Laboratories, Tokyo, Japan, and IBL, Hamburg, Germany, respectively). The proinsulin assay has 0% cross-reactivity with human insulin and C-peptide. The insulin assay has 0% cross-reactivity with proinsulin.



Figure 1. Genomic region of human chromosome 19p13.3 harboring the PTBP1 gene and HapMap linkage disequilibrium data of common (MAF>5%) informative SNPs within this locus. The PTBP1 gene consists of 15 exons and 14 introns. It spans 14.92 kb from nucleotide 748,411 to 763,327. The analyzed locus included 5 kb of both the 5'- and the 3'-flanking regions. The locations of the four tagging SNPs are marked by frames. The upper panel shows D' values and the lower panel shows r² values (black diamonds without values: $r^2 = 1.0$). doi:10.1371/journal.pone.0046154.g001

Calculations and statistical analyses

The area under the curve (AUC) $AUC_{Ins0-30}/AUC_{Glc0-30}$ was calculated as (Ins0+Ins 30)/(Glc0+Glc30). Insulinogenic index (IGI) was calculated as (Ins30 – Ins0)/(Glc30 – Glc 0). Cleared insulin response (CIR) was (100×Ins30)/Glc30×(Glc30 – 3.89). AUC was calculated according to the trapezoid method as: $0.5\times(C_0/2+C_{30}+C_{60}+C_{90}+C_{120}/2)$ with c = concentration.

Unless otherwise stated, data are given as mean \pm SD. Data that were not normally distributed were logarithmically transformed prior to statistical analysis. A Bonferroni-corrected p-value<0.0127 was considered statistically significant according to the number of SNPs tested.

Hardy-Weinberg equilibrium was tested using χ^2 test.

Our study was sufficiently powered $(1-\beta{>}0.8, \,\alpha{=}0.05)$ to detect effect sizes ${\geq}0.18$ for the SNP with the lowest minor allele frequency (rs11085226). Power calculations were performed using G*power software (www.psycho.uni-duesseldorf.de/aap/projects/gpower/). Pairwise linkage disequilibrium (D', r²) was determined using the Java linkage disequilibrium plotter (http://www.genepi. com.au/projects/jlin). For all other statistical analyses, the software package JMP 7.0.1 (SAS Institute, Cary, NC, USA) was used.

MAGIC genome-wide association (GWA) data

GWA-data on glycaemic traits have been made publicly available by the MAGIC consortium [32] and have been downloaded from www.magicinvestigators.org. We interrogated data from the meta-analysis results for the homeostatic model assessment of β -cell function (HOMA-B).

Analysis of linkage between polymorphisms

A search for SNPs in linkage disequilibrium with SNP rs123698was performed on the Broad Institute's SNAP(SNP Annotation and Proxy Search) website (http://www.broadinstitute.org/mpg/snap/ldsearch.php) using the CEU panel from both the *1000 genomes project pilot 1* dataset as well as well as *HapMap release 22*.

Analysis of putative transcription factor binding sites

For these analyses, the JASPAR (http://jaspar.genereg.net) database [33] "homo sapiens" was used with a relative profile score threshold of 80%. We analyzed the 50 bp around each of the two SNPs that showed associations with insulin levels during IVGTT and all linked SNPs with $r^2>0.8$ (1000 genomes project pilot 1: SNPs rs10420407 and rs10420953 for SNP rs11085226; SNPs rs351976, rs8103323, rs8100561, and rs351995 for SNP rs351974).

Results

Subject characteristics

The participants' clinical characteristics are presented in Table 1. In this group of relatively young subjects 66% were females and 34% males. They were slightly overweight and non-diabetic. 10% of the subjects had impaired fasting glucose, 10% had impaired glucose tolerance and 7% had both.

Genotyping

We analyzed the following tagging SNPs: rs351974, rs11085226, rs736926, and rs123698 in *PTBP1*. All SNPs were in Hardy-Weinberg equilibrium (all $p \ge 0.3$) and showed MAFs very close to those reported by the HapMap project (http://www.hapmap.org, [28], Table 2).

Table 2. Linkage disequilibrium statistics (D', r^2) among the four tagging SNPs rs351974, rs11085226, rs736926, and rs123698 covering the 14.92 kb genomic locus harbouring the *PTBP1* gene.

| SNP | rs351974 | rs11085226 | rs736926 | rs123698 | | |
|------------------------------|---------------|---------------|---------------|---------------|--|--|
| rs351974 | - | 0.995 | 0.842 | 0.996 | | |
| rs11085226 | 0.238 | - | 1.000 | 0.985 | | |
| rs736926 | 0.279 | 0.019 | - | 0.976 | | |
| rs123698 | 0.254 | 0.071 | 0.096 | - | | |
| MAF (MAF _{HapMap}) | 0.293 (0.292) | 0.106 (0.092) | 0.140 (0.123) | 0.382 (0.405) | | |

D' values above empty cells; r² values below empty cells. PTBP1– polypyrimidine-tract-binding protein, SNP-single nucleotide polymorphism, MAF-minor allele frequency, MAF_{HapMap}-minor allele frequency from theInternationalHapMap Project. doi:10.1371/journal.pone.0046154.t002

Insulin secretion during the OGTT

The minor allele of SNP rs11085226 in *PTBP1* was nominally associated with lower insulin secretion measured as IGI and CIR during the OGTT in the dominant inheritance model ($p\leq0.04$; Table 3). Whereas the third insulin-based parameter, i.e., AUC_{Ins0-30}/AUC_{Glc0-30} revealed a trend for association, there was no association with C-peptide-based measures (Table 3). The three other tested SNPs in the *PTBP1* gene did not show any association with any of the tested parameters of insulin secretion (all p>0.07; Table 3 and 4). None of the analyzed SNPs was associated with proinsulin levels or proinsulin-to-insulin ratio. In our data, we detected no significant association with HOMA-B.

Insulin secretion during the IVGTT

The minor allele of SNP rs11085226 showed significant association with lower insulin levels within the first ten minutes following glucose injection (AUC_{0-10 min} p = 0.0103, dominant inheritance model; Figure 2 A), while glucose levels did not diverge (AUC_{0-10 min} p = 0.3; Figure 2 B).

Neither SNP rs736926, nor SNP rs123698 were associated with insulin levels during the IVGTT (AUC₀₋₁₀ min p = 0.4 and p = 0.7, respectively), in a model adjusted for gender, age, BMI, and clamp-derived insulin sensitivity index. Performing the same adjustment, SNP rs351974 was associated with insulin levels during the first 10 minutes of the IVGTT (AUC_{Insulin 0-10} min p = 0.0108, dominant inheritance model). SNP rs351974 was furthermore associated with glucose levels during the same time period after adjustment for gender, age, and BMI (AUC_{Glucose 0-10} min p = 0.0172, dominant inheritance model). Both, glucose and insulin levels were lower in carriers of the minor allele of SNP rs351974. The association of this SNP with AUC_{Insulin 0-10} min remained present after adjustment for AUC_{Glucose 0-10} min in addition to gender, age, BMI, and clamp-derived insulin sensitivity index (p = 0.0377).

Replication of the association of SNP rs11085226 with insulin secretion in MAGIC GWA data

In the publicly available data on glycaemic traits from the MAGIC consortium on 46,186 non-diabetic participants [32], SNP rs11085226 was associated with the provided measure of insulin secretion, HOMA-B (p = 0.01815) with higher insulin secretion in carriers of the A-allele. Of the other three investigated *PTBP1* SNPs, rs736926 showed association with HOMA-B in MAGIC, too (p = 0.04164), and SNP rs351974 did not show such

Table 3. Associations of *PTBP1* SNPs rs351974 and rs11085226 with insulin secretion and proinsulin conversion in non-diabetic subjects (N = 1502).

| SNP | rs351974 | | | | | rs11085226 | | | | |
|---|-------------|-------------------|-------------|------------|----------------|-------------------|---------------------|---------------------|----------------|-----------------------|
| Genotype | AA | AC | сс | p 1 | p ₂ | AA | AG | GG | p ₁ | p ₂ |
| N | 760 | 604 | 137 | - | - | 1205 | 273 | 23 | - | - |
| НОМА-В | 131.2±116.1 | 136.7±132.1 | 130.7±126.4 | 0.8 | 0.9 | 132.7±126.2 | 137.5±116.4 | 118.6±60.0 | 0.8 | 0.8 |
| AUC _{Ins0-30} /AUC _{Glc0-30} (×10 ⁻⁹) | 40.8±29.6 | 40.4±31.7 | 43.0±31.6 | 0.7 | 0.5 | 41.2±30.7 | 39.5±29.8 | 42.0±33.8 | 0.07 | 0.08 |
| IGI (×10 ⁻⁹) | 150±299 | 149±137 | 147±116 | 0.9 | 0.7 | 153±253 | 136±113 | 147±135 | 0.08 | 0.0378 |
| CIR (I×mmol ⁻¹ ×10 ⁻⁹) | 1530±1368 | 1516±1378 | 1491±1205 | 0.9 | 0.6 | 1551±1393 | 1395±1188 | 1511±1325 | 0.07 | 0.0429 |
| C-pep30 (pmol/l) | 2062±880 | 2019±889 | 2067±939 | 1.0 | 1.0 | 2055±890 | 2001±894 | 2085±783 | 0.2 | 0.2 |
| AUC _{C-pep} /AUC _{Glc} (×10 ⁻⁹) | 320±103 | 318±108 | 317±108 | 0.8 | 0.9 | 320±103 | 316±115 | 332±100 | 0.2 | 0.2 |
| Proins0 (pmol/l) | 5.80±6.67 | 5.92±6.86 | 4.91±4.14 | 0.5 | 0.7 | 5.78±6.83 | 5.81±5.44 | 5.00±3.71 | 0.3 | 0.1 |
| Proins30 (pmol/l) | 11.6±9.6 | 12.0±15.5 | 11.8±8.4 | 0.7 | 0.8 | 11.7±12.8 | 12.0±9.9 | 12.1±8.6 | 0.8 | 0.6 |
| AUC _{Proins} (pmol/l) | 35.4±27.7 | 36.9±38.5 | 37.6±28.7 | 0.8 | 0.9 | 35.7±31.3 | 38.5±37.8 | 37.3±26.5 | 0.9 | 0.7 |
| Proins0/Ins0 | 0.124±0.173 | 0.136±0.234 | 0.112±0.156 | 0.6 | 0.8 | 0.128±0.207 | 0.128±0.162 | 0.120±0.137 | 0.4 | 0.2 |
| Proins30/Ins30 | 0.032±0.033 | 0.033 ± 0.054 | 0.049±0.035 | 0.8 | 0.5 | 0.032 ± 0.043 | 0.034±0.041 | 0.033±0.031 | 0.4 | 0.2 |
| AUC _{Proins} /AUC _{Ins} | 0.049±0.041 | 0.052 ± 0.059 | 0.049±0.035 | 0.9 | 1.0 | 0.050 ± 0.050 | $0.052 {\pm} 0.043$ | $0.056 {\pm} 0.050$ | 0.8 | 0.5 |

Data represent means \pm SD. For statistical analysis, data were ln-transformed and adjusted for gender, age, BMI, and insulin sensitivity. $p_1 - p$ -values for the additive inheritance model; $p_2 - p$ -values for the dominant inheritance model. AUC – area under the curve; BMI – body mass index; CIR – cleared insulin response; IGI – insulinogenic index; SNP – single nucleotide polymorphism.

doi:10.1371/journal.pone.0046154.t003

an association (p = 0.7). Since SNP rs123698 is not included in the provided MAGIC data set, we searched for linked polymorphisms ($r^2>0.8$)in 1000 genomes pilot 1, as well as HapMap release 22 but found no such linked SNPs. Since we were unable to identify another polymorphism as a proxy for *PTBP1*rs123698, we could not check for associations between this SNP and HOMA-B in MAGIC data.

Discussion

In the present study, we found that the common variant rs11085226 in the *PTBP1* gene is associated with significantly lower glucose-stimulated insulin secretion, but not with altered proinsulin-to-insulin ratio. This finding could be replicated in the huge data set from the MAGIC consortium. We furthermore found SNP rs351974 to be associated with insulin levels during the

Table 4. Associations of *PTBP1* SNPs rs736926 and rs123698 with insulin secretion and proinsulin conversion in non-diabetic subjects (N = 1502).

| SNP | rs736926 | | | | rs123698 | | | | | |
|---|-------------|-------------------|-------------------|----------------|----------------|-------------------|---------------------|---------------------|------------|-----------------------|
| Genotype | сс | ст | π | p ₁ | p ₂ | сс | CG | GG | p 1 | p ₂ |
| N | 1112 | 357 | 30 | - | - | 577 | 692 | 224 | - | - |
| НОМА-В | 132.1±126.1 | 135.2±115.9 | 157.8±127.9 | 0.2 | 0.5 | 133.8±123.3 | 135.3±130.5 | 128.0±103.6 | 0.8 | 0.4 |
| AUC _{Ins0-30} /AUC _{Glc0-30} (×10 ⁻⁹) | 40.6±30.3 | 40.9±31.5 | 47.7±30.8 | 0.3 | 0.5 | 39.8±29.4 | 42.4±32.6 | 38.7±27.5 | 0.2 | 0.2 |
| lGI (×10 ⁻⁹) | 147±257 | 157±141 | 164±115 | 0.2 | 0.1 | 138±147 | 168±252 | 121±330 | 0.07 | 0.1 |
| CIR (I×mmol ⁻¹ ×10 ⁻⁹) | 1521±1406 | 1501 ± 1198 | 1678±1319 | 0.3 | 0.3 | 1443±1231 | 1610±1476 | 1452 ± 1288 | 0.1 | 0.2 |
| C-pep30 (pmol/l) | 2050±880 | 2008±891 | 2280±1178 | 0.7 | 0.6 | 2011±864 | 2080±908 | 2024±894 | 0.4 | 0.2 |
| AUC _{C-pep} /AUC _{Glc} (×10 ⁻⁹) | 320±106 | $313\!\pm\!103$ | 344±122 | 0.4 | 0.6 | 316±105 | 322±107 | 318±102 | 0.6 | 0.3 |
| Proins0 (pmol/l) | 5.82±6.55 | 5.51±6.68 | 6.70±5.66 | 0.09 | 0.3 | 5.79±7.30 | 5.76±6.08 | 5.81±6.02 | 0.4 | 0.2 |
| Proins30 (pmol/l) | 11.8±9.9 | 11.6±17.6 | 14.2±11.7 | 0.3 | 0.3 | 11.8±15.8 | 11.9±9.8 | 11.3±7.9 | 0.5 | 0.2 |
| AUC _{Proins} (pmol/l) | 36.0±30.5 | 36.1±38.2 | 42.1±28.2 | 0.7 | 1.0 | 37.5±37.1 | 35.4±30.7 | 35.1±23.9 | 0.5 | 0.8 |
| Proins0/Ins0 | 0.125±0.167 | 0.137±0.277 | 0.141 ± 0.150 | 0.1 | 0.2 | 0.131±0.244 | 0.124±0.169 | $0.132 {\pm} 0.149$ | 0.3 | 0.4 |
| Proins30/Ins30 | 0.032 0.035 | 0.033±0.061 | 0.031 ± 0.024 | 0.4 | 0.2 | 0.033 ± 0.055 | $0.032 {\pm} 0.031$ | 0.033 ± 0.034 | 0.6 | 0.5 |
| AUC _{Proins} /AUC _{Ins} | 0.049±0.041 | 0.053 ± 0.068 | 0.051 ± 0.038 | 0.9 | 1.0 | 0.052 ± 0.060 | 0.049 ± 0.040 | 0.051 ± 0.038 | 0.9 | 1.0 |

Data represent means \pm SD. For statistical analysis, data were ln-transformed and adjusted for gender, age, BMI, and insulin sensitivity. $p_1 - p$ -values for the additive inheritance model; $p_2 - p$ -values for the dominant inheritance model. AUC – area under the curve; BMI – body mass index; CIR – cleared insulin response; IGI – insulinogenic index; SNP – single nucleotide polymorphism.

doi:10.1371/journal.pone.0046154.t004



Figure 2. Metabolic data in carriers of the PTBP1 SNP rs11085226 during the IVGTT. Shown are plasma insulin (A) and glucose (B) concentrations. Data are given as means +/- SEM. P-values for AUC of the ten minutes following glucose injection adjusted for gender, age, BMI, and clamp-derived insulin sensitivity are given. Arrows mark the time-point of glucose infusion. Black circles: homozygous carriers of the major allele; white circles: heterozygous and homozygous carrier of the minor allele. doi:10.1371/journal.pone.0046154.g002

IVGTT, but not with any of the OGTT-derived parameters in our study, or in the analyzed MAGIC data.

The influence of *PTBP1* rs11085226 in the IVGTT especially on the insulin levels within few minutes after glucose injection with diminished influence at later time points might explain why we could only detect effects on OGTT-derived parameters of insulin secretion that cover the first 30 minutes. The large sample size of MAGIC [32] was necessary to detect also effects of this SNP on fasting insulin secretion (HOMA-B), most likely because glucosestimulated insulin secretion plays only a minor role in the fasting state. Notably, the effect was not detected in C-peptide-based parameters (C-pep30, AUC_{C-pep}/AUC_{Glc}). This discrepancy could be due to effects of the SNP on insulin clearance. So, we examined the association of *PTBP1* rs11085226 with two estimates of insulin clearance (Cpep0/Ins0 and AUC_{Cpep}/AUC_{Ins}). Both parameters were not associated with this SNP (p \geq 0.3).

The association of *PTBP1* SNP rs351974 with insulin levels (independently of its concomitant association with glucose) was only detectable in the IVGTT data but neither in OGTT data nor in MAGIC. Similar discrepancy between insulin levels after oral and i.v. glucose administration was noted for a polymorphism in *SLC30A8* [34]. The underlying mechanism for this observation still

remains open but one possible explanation might be that the incretin effect that is present only after oral glucose uptake compensates for the SNP effects and, thus, making them undetectable during an OGTT.

SNP *PTBP1* rs11085226 is in linkage with two other SNPs, one in intron 3 (rs10420407 [A/G]) and one in exon 5 (rs10420953) (figure 1). The exonic variation is a silent mutation (N108N), thus does not change the amino acid sequence. Therefore, these variations are suggested to influence the expression of *PTBP1* at the transcriptional level rather than altering the protein functions. For instance, they could influence the binding of transcription factors or repressors, however this is unlikely since the JASPAR database [33] contains no putative transcription factor binding sites around the SNPs or linked SNPs. Furthermore, the silent mutation in exon 5 could also affect RNA processing or protein translation rate. SNP rs11085226 could possibly affect alternative splicing, since it is located in an intron next to an alternative splicing site [14,35].

Since PTBP1 influences the expression of proinsulin convertases, we expected that genetic variations could affect proinsulin/ insulin ratios. However, such effects were not found. Interestingly, PTBP1 SNPs rs11085226 and rs351974 reduced insulin secretion very early after glucose administration but not at later time points. One might speculate therefore that the variation influences the number of insulin secretory granules within the readily releasable pool, which are mainly responsible for the first phase of insulin secretion [36]. Of interest, PTBP1 seems to be involved in impairment of this mechanism in type 2 diabetes [37]. The hypothesis of alterations in the pool of readily releasable granules is supported by the fact that PTBP1 supports the biogenesis of insulin secretory granules, while data in rodent β -cells indicate that newly-synthesized insulin secretory granules undergo preferential exocytosis ([38,39]; Ivanova and Solimena, unpublished data). Moreover, PTBP1 may regulate the expression of proteins that are directly involved in the machinery for exocytosis of insulin secretory granules.

Our study could only detect an effect size $\geq 18\%$ and was therefore underpowered to detect smaller effect sizes, e.g. on proinsulin conversion, and thus larger cohorts should be studied to assess such possible smaller effects. Further on, we focused on common variants with a MAF>5%. Therefore we might have overlooked effects of rare variants or mutations that could more dramatically influence PTBP1 levels or functions. Studying such rare variants will need much larger cohorts and direct sequencing strategies. Further clarification about the influence of *PTBP1* SNPs on insulin secretory granule exocytosis will also require additional studies at the molecular level.

Taken together, our data suggest that the common genetic variation in *PTBP1* affects very early glucose-stimulated insulin secretion. Thus, the effects of PTBP1 on insulin secretion that were discovered in model systems are detectable and relevant in humans *in vivo*, too.

Acknowledgments

We thank all study participants for their cooperation. We gratefully acknowledge the excellent technical assistance of Anna Bury, Alke Guirguis, Melanie Weisser, and Roman Werner.

Author Contributions

Conceived and designed the experiments: MH KPK AF HS HUH MS. Performed the experiments: MH CK RW KL AB SHS FM AF. Analyzed the data: MH KK AF HS MS. Contributed reagents/materials/analysis tools: FM. Wrote the paper: MH KK HS AF MS.

References

- Straub SG, Sharp GW (2002) Glucose-stimulated signaling pathways in biphasic insulin secretion. Diabetes Metab Res Rev 18: 451–463.
- Doyle ME, Egan JM (2007) Mechanisms of action of glucagon-like peptide 1 in the pancreas. Pharmacol Ther 113: 546–593.
- Hinke SA, Hellemans K, Schuit FC (2004) Plasticity of the beta cell insulin secretory competence: preparing the pancreatic beta cell for the next meal. J Physiol 558: 369–380.
- Knoch KP, Bergert H, Borgonovo B, Saeger HD, Altkruger A, et al. (2004) Polypyrimidine tract-binding protein promotes insulin secretory granule biogenesis. Nat Cell Biol 6: 207–214.
- 5. Steiner DF (1998) The proprotein convertases. Curr Opin Chem Biol 2: 31-39.
- Martin SK, Carroll R, Benig M, Steiner DF (1994) Regulation by glucose of the biosynthesis of PC2, PC3 and proinsulin in (ob/ob) mouse islets of Langerhans. FEBS Lett 356: 279–282.
- Goodge KA, Hutton JC (2000) Translational regulation of proinsulin biosynthesis and proinsulin conversion in the pancreatic beta-cell. Semin Cell Dev Biol 11: 235–242.
- Guest PC, Rhodes CJ, Hutton JC (1989) Regulation of the biosynthesis of insulin-secretory-granule proteins. Co-ordinate translational control is exerted on some, but not all, granule matrix constituents. Biochem J 257: 431–437.
- Ort T, Voronov S, Guo J, Zawalich K, Froehner SC, et al. (2001) Dephosphorylation of beta2-syntrophin and Ca2+/mu-calpain-mediated cleavage of ICA512 upon stimulation of insulin secretion. EMBO J 20: 4013–4023.
- Itoh N, Okamoto H (1980) Translational control of proinsulin synthesis by glucose. Nature 283: 100–102.
- Steiner DF (1977) The Banting Memorial Lecture 1976. Insulin today. Diabetes 26: 322–340.
- Alarcon C, Wicksteed B, Rhodes CJ (2006) Exendin 4 controls insulin production in rat islet beta cells predominantly by potentiation of glucosestimulated proinsulin biosynthesis at the translational level. Diabetologia 49: 2920–2929.
- Knoch KP, Meisterfeld R, Kersting S, Bergert H, Altkruger A, et al. (2006) cAMP-dependent phosphorylation of PTB1 promotes the expression of insulin secretory granule proteins in beta cells. Cell Metab 3: 123–134.
- Sawicka K, Bushell M, Spriggs KA, Willis AE (2008) Polypyrimidine-tractbinding protein: a multifunctional RNA-binding protein. Biochem Soc Trans 36: 641–647.
- Auweter SD, Allain FH (2008) Structure-function relationships of the polypyrimidine tract binding protein. Cell Mol Life Sci 65: 516–527.
- Suss C, Czupalla C, Winter C, Pursche T, Knoch KP, et al. (2009) Rapid changes of mRNA-binding protein levels following glucose and 3-isobutyl-1methylxanthine stimulation of insulinoma INS-1 cells. Mol Cell Proteomics 8: 393–408.
- Xie J, Lee JA, Kress TL, Mowry KL, Black DL (2003) Protein kinase A phosphorylation modulates transport of the polypyrimidine tract-binding protein. Proc Natl Acad Sci U S A 100: 8776–8781.
- Jang SK, Wimmer E (1990) Cap-independent translation of encephalomyocarditis virus RNA: structural elements of the internal ribosomal entry site and involvement of a cellular 57-kD RNA-binding protein. Genes Dev 4: 1560– 1572.
- Song Y, Tzima E, Ochs K, Bassili G, Trusheim H, et al. (2005) Evidence for an RNA chaperone function of polypyrimidine tract-binding protein in picornavirus translation. RNA 11: 1809–1824.

- Cote CA, Gautreau D, Denegre JM, Kress TL, Terry NA, et al. (1999) A Xenopus protein related to hnRNP I has a role in cytoplasmic RNA localization. Mol Cell 4: 431–437.
- Ahren B (2005) Type 2 diabetes, insulin secretion and beta-cell mass. Curr Mol Med 5: 275–286.
- Kahn SE (2003) The relative contributions of insulin resistance and beta-cell dysfunction to the pathophysiology of Type 2 diabetes. Diabetologia 46: 3–19.
- Steiner DF, James DE (1992) Cellular and molecular biology of the beta cell. Diabetologia 35 Suppl 2: S41–S48.
- Heni M, Haupt A, Schafer SA, Ketterer C, Thamer C, et al. (2010) Association of obesity risk SNPs in PCSK1 with insulin sensitivity and proinsulin conversion. BMC Med Genet 11: 86.
- Stumvoll M, Fritsche A, Stefan N, Hardt E, Haring H (2001) Evidence against a rate-limiting role of proinsulin processing for maximal insulin secretion in subjects with impaired glucose tolerance and beta-cell dysfunction. J Clin Endocrinol Metab 86: 1235–1239.
- Larsson H, Ahren B (1999) Relative hyperproinsulinemia as a sign of islet dysfunction in women with impaired glucose tolerance. J Clin Endocrinol Metab 84: 2068–2074.
- Stefan N, Kantartzis K, Machann J, Schick F, Thamer C, et al. (2008) Identification and characterization of metabolically benign obesity in humans. Arch Intern Med 168: 1609–1616.
- 28. (2003) The International HapMap Project. Nature 426: 789-796.
- Matsuda M, DeFronzo RA (1999) Insulin sensitivity indices obtained from oral glucose tolerance testing: comparison with the euglycemic insulin clamp. Diabetes Care 22: 1462–1470.
- Tripathy D, Wessman Y, Gullstrom M, Tuomi T, Groop L (2003) Importance of obtaining independent measures of insulin secretion and insulin sensitivity during the same test: results with the Botnia clamp. Diabetes Care 26: 1395– 1401.
- Stefan N, Machicao F, Staiger H, Machann J, Schick F, et al. (2005) Polymorphisms in the gene encoding adiponectin receptor 1 are associated with insulin resistance and high liver fat. Diabetologia 48: 2282–2291.
- Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, et al. (2010) New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. Nat Genet 42: 105–116.
- Sandelin A, Alkema W, Engstrom P, Wasserman WW, Lenhard B (2004) JASPAR: an open-access database for eukaryotic transcription factor binding profiles. Nucleic Acids Res 32: D91–D94.
- Staiger H, Machicao F, Stefan N, Tschritter O, Thamer C, et al. (2007) Polymorphisms within novel risk loci for type 2 diabetes determine beta-cell function. PLoS ONE 2: e832.
- Fontana L. Atlas of Genetics and Cytogenetics in Oncology and Haematology: PTBP1 (polypyrimidine tract binding protein 1). Available: http:// AtlasGeneticsOncology.org/Genes/PTBP1ID46504ch19p13.html. Accessed 2012 February 1.
- Rorsman P, Řenstrom E (2003) Insulin granule dynamics in pancreatic beta cells. Diabetologia 46: 1029–1045.
- Ehehalt F, Knoch K, Erdmann K, Krautz C, Jager M, et al. (2010) Impaired insulin turnover in islets from type 2 diabetic patients. Islets 2: 30–36.
- Halban PA (1982) Differential rates of release of newly synthesized and of stored insulin from pancreatic islets. Endocrinology 110: 1183–1188.
- Grodsky GM (1999) The importance of rapid insulin secretion: revisited. Diabetes Technol Ther 1: 259–260.